



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Slow and steady wins the race: Better walking than running. The turtle's lesson in the times of COVID-19

S. Sciomer, MD^a, S. Gallina, MD, PhD^b, A.V. Mattioli, MD, PhD^c, P.G. Agostoni, MD, PhD^{d,e},
F. Moscucci, MD^{a,*}

^a Department of Clinical, Internal, Anesthesiological and Cardiovascular Sciences, University of Rome "Sapienza", Policlinico Umberto I, Viale del Policlinico n.155, Rome 00186, Italy

^b Department of Neuroscience, Imaging and Clinical Sciences, "G. d'Annunzio" University, Chieti 66100, Italy

^c Surgical, Medical and Dental Department of Morphological Sciences Related to Transplant, Oncology and Regenerative Medicine, University of Modena and Reggio Emilia, Via Del Pozzo, 71, Modena 41124, Italy

^d Centro Cardiologico Monzino, IRCCS, Milan, Italy

^e Department of Clinical Sciences and Community Health, University of Milan, Milan, Italy

ARTICLE INFO

Article History:

Received 1 March 2021

Revised 7 April 2021

Accepted 12 April 2021

Available online 2 June 2021

Dear Editor,—as recently hypothesized,^{1, 2} pollution and particulate matter could have played a crucial role in the spread and clinical severity of patients infected with SARS-CoV2.

Therefore, some specific groups of exposed population could be more affected by the inhalation of particulates. Among these, those who practice outdoor aerobic physical activity could be the most severely affected by the SARS-CoV2. Infection, constantly and chronically exposed to inhaled pollutants, during outdoor exercises.^{3,4}

Oxygen is indispensable to the metabolic processes and it allows the survival of our cells. The air we breathe, however, contains only about 20% of Oxygen, which is mixed with Nitrogen (78%), Carbon dioxide (0.04%) and Argon (0.9%) and other substances that vary according to of the place and season, such as pollen, dust of all kinds from domestic to siliceous, micro-particulate, microbial particles, chlorine and many others.^{5,6} Every time we inhale, it is not only oxygen that enters our respiratory tract, but the mixture of gases and volatile substances that are found in the air around us. These acquisitions have been connected to different pathological conditions.^{7–9}

Each quiet inspiration brings between 300 and 500 ml of air into our respiratory tract; a part of it stops in the upper airways (nose, trachea, bronchus and terminal bronchioles), while about 150–300 ml reach the alveolar membrane where the gas exchange takes place.¹⁰

During exercise, the need to obtain oxygen and eliminate carbon dioxide increases, sympathetic bronchodilation causes a reduction in resistance and an increase in flow. The respiratory rate of 15 acts/minute (12 L of air) at rest can reach 40–60 acts/minute (100 L of air) during the exercise phase.¹¹

Furthermore, breathing is generally oral and non-nasal during prolonged exercise. The air that reaches the lower respiratory tract will therefore be colder and unfiltered.¹² Exercise always causes greater airflow to the lung, but the exercises are not all the same. Endurance sports such as marathon, cross-country skiing, cycling involve greater exposure to ambient air and therefore a greater risk of lung contact with toxic substances and harmful agents.¹²

In addition, exercise is universally recognized as a protective factor for cardiovascular and metabolic diseases, although it is important to know the environmental conditions open-air sports are practiced.^{12,13}

Physical activity enables us to increase our interface with the external environment, in particular with the invisible substances that surround us and that can be harmful.^{5,6} A walk lasting twenty minutes, for example, involves a flow of about 240 L of air in our respiratory system, while with a run of the same duration you can reach 2000 L of flow.¹¹ Considering that one thousand liters of air equals one meter³, we can compare these values with the measurements of PM2.5 and PM10 of the ARPA Agency in Lombardy.¹⁴

In Milan (ARPA Verziere station) the daily average of PM10 corresponds to 53 $\mu\text{g} / \text{m}^3$, a run would make inhale about 106 μg of this substance to the runner. In Bergamo (ARPA Via Garibaldi station) the average daily quantity of PM2.5 is 34 $\mu\text{g} / \text{m}^3$, and 41 $\mu\text{g} / \text{m}^3$ of PM10. Thus, a 20 min run would expose to the inhalation of approximately 68 μg of PM 2.5 and 81 μg of PM10. To reach this amount of inhaled pollutants, it is necessary a 3 h walk.¹⁵

Multiple are the effects on amateur athletes' body. The particulate matter has a substantially pro-inflammatory effect, and correlates with acute effects such as asthma crisis, cough and hospitalizations for respiratory disease. Chronic exposure instead is associated with chronic obstructive pulmonary disease, respiratory failure, cardiovascular diseases.³ The polluting micromolecules are capable of

* Corresponding author.

E-mail address: federica.moscucci@uniroma1.it (F. Moscucci).

weakening and destroying the alveolar membrane, making it more susceptible to any kind of insult, including the infections.^{1,16}

The damage mediated by these substances has been demonstrated, as well as there is no real threshold value below whom there is no danger according to the WHO, and the concentration must be kept as low as possible.^{17,18}

All considered, running in this period in Milan, as in Bergamo and in all the places where the levels of PM10 and PM2.5 are close to or above the threshold values can basically mean making a feast of pro-inflammatory polluting agents that could make the lungs more susceptible to infections and complications, including those from SARS-CoV2.

Although the certain advantages on the cardiovascular system of constant physical activity remain firm and certain,¹⁹ outdoor activities must be carefully chosen, in order to expose the individual as little as possible to the effects of intense ventilation in highly polluted areas. We hope that health authorities and individuals will take due account of air pollution levels when choosing the type of physical activity to undertake in the near future.

Declaration of Competing Interest

None.

CRediT authorship contribution statement

S. Sciomer: Conceptualization, Writing - review & editing. **S. Gallina:** Conceptualization, Writing - review & editing. **A.V. Mattioli:** Conceptualization, Writing - review & editing. **P.G. Agostoni:** Conceptualization, Writing - review & editing. **F. Moscucci:** Conceptualization, Writing - review & editing, Writing - original draft.

Acknowledgment

All authors deny any personal or industrial interest about this paper.

References

1. Sciomer S, Moscucci F, Magri D, Badagliacca R, Piccirillo G, Agostoni P. SARS-CoV-2 spread in Northern Italy: what about the pollution role? *Environ Monit Assess.* 2020;192:325.
2. Gatti RC, Velichevskaya A, Tateo A, Amoroso N, Monaco A. Machine learning reveals that prolonged exposure to air pollution is associated with SARS-CoV-2 mortality and infectivity in Italy. *Environ Pollut.* 2020;267: 115471. Aug 21.
3. Slezakova K, Pereira MC, Morais S. Ultrafine particles: levels in ambient air during outdoor sport activities. *Environ Pollut.* 2020;258: 113648. Mar.
4. Cavalcante de Sá M, Nakagawa NK, Saldiva de André CD, et al. Aerobic exercise in polluted urban environments: effects on airway defense mechanisms in young healthy amateur runners. *J Breath Res.* 2016;10:(4) 046018. Dec 21.
5. Kardel F, Wuyts K, De Wael K, Samson R. Assessing atmospheric dry deposition via water-soluble ionic composition of roadside leaves. *J Environ Sci Health A Toxic Hazard Subst Environ Eng.* 2020;21:1–9. Apr.
6. Gaston CJ. Re-examining dust chemical aging and its impacts on Earth's climate. *Acc Chem Res.* 2020. <https://doi.org/10.1021/acs.accounts.0c00102>. Apr 29[Epub ahead of print].
7. Liu H, Zhang X, Zhang H, et al. Effect of air pollution on the total bacteria and pathogenic bacteria in different sizes of particulate matter. *Environ Pollut.* 2018;233:483–493. Feb.
8. Yue S, Wang Y, Wang J, Chen J. Relationships between lung cancer incidences and air pollutants. *Technol Health Care.* 2017;25(S1):411–422. Jul 20.
9. Cai Y, Zhang B, Ke W, et al. Associations of short-term and long-term exposure to ambient air pollutants with hypertension: a systematic review and meta-analysis. *Hypertension.* 2016;68(1):62–70. Jul.
10. Ochs M, Hegermann J, Lopez-Rodriguez E, et al. On top of the alveolar epithelium: surfactant and the glycocalyx. *Int J Mol Sci.* 2020;21(9). Apr 27.
11. European Lung Foundation, T. Troosters, L. Dupont, J. Bott, K. Hansen. Polmoni e attività fisica. Available at: www.european-lung-foundation.org.
12. Sue-Chu M. Winter sports athletes: long-term effects of cold air exposure. *Br J Sports Med.* 2012;46(6):397–401. May.
13. Nasi M, Patrizi G, Pizzi C, et al. The role of physical activity in individuals with cardiovascular risk factors: an opinion paper from Italian Society of Cardiology-Emilia Romagna-Marche and SIC-Sport. *J Cardiovasc Med.* 2019 Oct;20(10):631–639. (Hagerstown).
14. Rundell KW, Anderson SD, Sue-Chu M, Bougault V, Boulet LP. Air quality and temperature effects on exercise-induced bronchoconstriction. *Compr Physiol.* 2015 Apr;5(2):579–610.
15. Arpa Regione Lombardia, Inquinanti. Available at: <https://www.arpalombardia.it/Pages/Aria/Inquinanti/PM10-PM2,5.aspx?firstlevel=Inquinanti>.
16. Carugno M, Dentali F, Mathieu G, et al. PM10 exposure is associated with increased hospitalizations for respiratory syncytial virus bronchiolitis among infants in Lombardy, Italy. *Environ Res.* 2018;166:452–457. Oct.
17. Air Quality Guidelines for Europe, World Health Organization, Available at: http://www.euro.who.int/_data/assets/pdf_file/0005/74732/E71922.pdf.
18. Ministero della salute, Particolato. Available at: http://www.salute.gov.it/imgs/C_17_opuscoliPoster_283_ulterioriallegati_ulterioreallegato_7_alleg.pdf.
19. Ricci F, Izzicupo P, Moscucci F, et al. Recommendations for physical inactivity and sedentary behavior during the coronavirus disease (COVID-19) pandemic. *Front Public Health.* 2020;8:199. May 12.