



ORAL PRESENTATION

Open Access

Long term efficacy and safety of canakinumab in children with systemic juvenile idiopathic arthritis with and without fever

G Horneff^{1*}, N Ruperto¹, H Brunner², P Quartier³, T Constantin¹, E Alexeeva¹, I Kone-Paut¹, K Marzan², N Wulffraat¹, R Schneider², S Padeh¹, V Chasnyk¹, C Wouters¹, J Kummerle Deschner¹, T Kallinich¹, B Lauwerys⁴, E Haddad², E Nasonov¹, M Trachana¹, O Vougiouka¹, K Abrams⁵, K Leon⁵, K Lheritier⁶, A Martini¹, D Lovell²

From 8th International Congress of Familial Mediterranean Fever and Systemic Autoinflammatory Diseases Dresden, Germany. 30 September - 3 October 2015

Introduction

Rapid and sustained efficacy of canakinumab (CAN) were previously demonstrated in patients with systemic juvenile idiopathic arthritis (SJIA) [1]. However, little is known about potential differences in response to CAN treatment between patients with vs. without SJIA-associated fever at the time of the first CAN administration.

Objectives

To evaluate the long-term efficacy and safety profile of CAN-naïve SJIA patients with and without SJIA-associated fever.

Patients and methods

patients aged 2-20 years with SJIA with and without SJIA associated fever at enrollment received open-label CAN 4mg/kg s.c. every 4 wks. Every 3 months, response to CAN was measured by adapted JIA ACR response criteria (aACR/JIA); juvenile arthritis disease activity score (JADAS); clinical inactive disease; clinical remission on medication (CRM, 6 months continuous clinical inactive disease). Safety was assessed monthly.

Results

Data on 122/267 patients, 53 (43%) with and 69 (57%) without SJIA associated fever, were available for analysis with a median 94 wk study duration. At Wk4, ~75% of both subgroups had responded (\geq aACR/JIA30), increasing to 90% at Wk12. At Wk2, ~21% of both subgroups had inactive disease; 44% at Wk8; 60% at Wk20 and

then 60-70% for the remainder of the trial. CRM was achieved in about 29% of patients in both subgroups with ~22% maintaining it for \geq 12 consecutive months. At baseline, the median JADAS score was 21.5 with 8 (7.5%) and 99 (92.5%) patients meeting the criteria for moderate (JADAS $>$ 3.8 and \leq 10.5) and high disease activity (JADAS $>$ 10.5), respectively. At Day 15, the median JADAS was 6.8 and 1.5 at the last assessment. At the last assessment, 53 (48%) patients had inactive disease (JADAS \leq 1); 10 (9%) with low active disease activity (JADAS $>$ 1 and \leq 3.8); while 14 (13%) had moderate and 31 (28%) with high disease activity. Infection (0.56 infections/100 patient-days), typically involving upper respiratory tract was the most common type of adverse event. Fifteen patients discontinued due to an AE and 40 had $>$ 1 SAE (mostly infections, macrophage activation syndrome (MAS), or flare-associated) and no deaths. Eight cases of MAS (0.013 events/100 patient-days) were reported.

Conclusion

Canakinumab provides similar efficacy in SJIA patients with and without SJIA-associated fever at treatment onset. The long-term safety profile was acceptable and similar to the pivotal program in SJIA children with fever at enrollment.

Authors' details

¹PRINTo-Istituto Gaslini, Genova, Italy. ²PRCSG, Cincinnati, OH, United States. ³Necker-Enfant Malades Hospital, Paris, France. ⁴Cliniques Universitaires Saint-Luc and Université Catholique de Louvain, Brussels, Belgium. ⁵Novartis Pharmaceuticals Corporation, East Hanover, NJ, USA. ⁶Novartis Pharma AG, Basel, Switzerland.

¹PRINTo-Istituto Gaslini, Genova, Italy

Full list of author information is available at the end of the article

Published: 28 September 2015

Reference

1. Groom JR, Luster AD: CXCR3 ligands: redundant, collaborative and antagonistic functions. *Immunol Cell Biol* 2011, **89**(2):207-15.

doi:10.1186/1546-0096-13-S1-O83

Cite this article as: Horneff *et al.*: Long term efficacy and safety of canakinumab in children with systemic juvenile idiopathic arthritis with and without fever. *Pediatric Rheumatology* 2015 **13**(Suppl 1):O83.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

