

SCIENTIFIC REPORTS



OPEN

Honokiol suppresses TNF- α -induced neutrophil adhesion on cerebral endothelial cells by disrupting polyubiquitination and degradation of I κ B α

Received: 28 September 2015

Accepted: 05 May 2016

Published: 23 May 2016

Po-Jen Chen^{1,2,*}, Yu-Ling Wang^{1,2,*}, Liang-Mou Kuo^{3,4,*}, Chwan-Fwu Lin⁵, Chun-Yu Chen^{3,6}, Yung-Fong Tsai^{3,6}, Jiann-Jong Shen^{1,2} & Tsong-Long Hwang^{1,2,5,6}

Adhesion molecules expressed on cerebral endothelial cells (ECs) mediate leukocyte recruitment and play a significant role in cerebral inflammation. Increased levels of adhesion molecules on the EC surface induce leukocyte infiltration into inflammatory areas and are thus hallmarks of inflammation. Honokiol, isolated from the Chinese medicinal herb *Magnolia officinalis*, has various pharmacological activities, including anti-inflammatory effects, yet the nature of honokiol targeting molecules remains to be revealed. Here, we investigated the inhibitory effect of honokiol on neutrophil adhesion and vascular cell adhesion molecule-1 (VCAM-1) expression, which underlie its molecular target, and mechanisms for inactivating nuclear factor κ enhancer binding protein (NF- κ B) in mouse cerebral ECs. Honokiol inhibited tumour necrosis factor- α (TNF- α)-induced neutrophil adhesion and VCAM-1 gene expression in cerebral ECs. The inflammatory transcription factor NF- κ B was downregulated by honokiol. Honokiol significantly blocked TNF- α -induced NF- κ B p65 nuclear translocation and degradation of the proteasome-dependent inhibitor of NF- κ B α (I κ B α). From docking model prediction, honokiol directly targeted the ubiquitin-ubiquitin interface of Lys48-linked polychains. Moreover, honokiol prevented the TNF- α -induced Lys48-linked polyubiquitination, including I κ B α -polyubiquitin interaction. Honokiol has protective anti-inflammatory effects on TNF- α -induced neutrophil adhesion and VCAM-1 gene expression in cerebral ECs, at least in part by directly inhibiting ubiquitination-mediated I κ B α degradation and then preventing NF- κ B nuclear translocation.

Cerebral endothelial cells (ECs) are the major components of the blood-brain barrier (BBB). Cerebral ECs are distinct from peripheral ECs because they have fewer vesicles, higher mitochondrial volume fraction, higher electrical resistance, and unique transport systems¹⁻³. They act as a barrier to the central nervous system (CNS) and limit the entry of monocytes, lymphocytes or other leukocytes under physiological conditions.

CNS inflammation is associated with disrupted BBB integrity followed by activated leukocyte transmigration^{4,5}. Adhesion molecules such as vascular cell adhesion molecule-1 (VCAM-1), intercellular adhesion molecule-1 (ICAM-1) and E-selectin are hallmarks of inflammation because their increased expression on the EC surface plays a critical role in monocyte and leukocyte infiltration into inflammation areas^{6,7}. Increased expression of particular adhesion molecules in various inflammatory diseases may suggest opportunities for the development of new therapeutic strategies⁸. VCAM-1 expression is low in normal cerebral ECs but high

¹Graduate Institute of Natural Products, School of Traditional Medicine, College of Medicine, Chang Gung University, Taoyuan 333, Taiwan. ²Chinese Herbal Medicine Research Team, Healthy Aging Research Center, Chang Gung University, Taoyuan 333, Taiwan. ³Graduate Institute of Clinical Medical Sciences, College of Medicine, Chang Gung University, Taoyuan 333, Taiwan. ⁴Department of General Surgery, Chang Gung Memorial Hospital, Chiayi 613, Taiwan. ⁵Research Center for Industry of Human Ecology and Graduate Institute of Health Industry Technology, Chang Gung University of Science and Technology, Taoyuan 333, Taiwan. ⁶Department of Anaesthesiology, Chang Gung Memorial Hospital, Taoyuan 333, Taiwan. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to T.-L.H. (email: htl@mail.cgu.edu.tw)

in ECs during CNS inflammatory disorders such as stroke, multiple sclerosis, and brain tumor metastasis^{3,9–11}. A therapeutic strategy may be to repress VCAM-1 expression and diminish infiltration of inflammatory cells into the brain.

Tumour necrosis factor- α (TNF- α) is a critical cytokine during the progression of CNS diseases and induces the expression of the adhesion molecules VCAM-1, ICAM-1 and E-selectin contributing to these complex processes^{12–14}. The inflammatory transcription factor, nuclear factor κ enhancer binding protein (NF- κ B), is responsible for upregulating these pro-inflammatory genes. In the canonical pathway, NF- κ B is rapidly activated by various inflammatory cytokines in a wide variety of biological phenomena. With TNF- α stimulation, the inhibitor of NF- κ B α (I κ B α) is phosphorylated and sequentially conjugated with K48-linked polyubiquitin, which leads to degradation of I κ B α by the 26S proteasome. Such degradation allows NF- κ B to translocate into nucleus and sequentially stimulate the expression of pro-inflammatory target genes^{15–17}. TNF- α -mediated NF- κ B activation is tightly controlled by ubiquitination, especially in the I κ B pathway^{18,19}. Hence, interfering with ubiquitination in NF- κ B signalling is an interesting strategy for preventing excessive inflammation.

Honokiol (C₁₈H₁₈O₂) (Fig. 1A), a natural product isolated from the Chinese medicinal herb *Magnolia officinalis*, has several pharmacological effects because of its anti-angiogenesis, anti-tumor, neuroprotection, and anti-inflammatory properties^{20–26}. Honokiol can easily penetrate the BBB and blood–cerebrospinal fluid barrier²⁷, so it may be an effective drug for treating CNS disorders. Honokiol has been extensively studied for cancer prevention and therapy because of its multiple targeted pathologically relevant pathways, including epidermal growth factor receptor, signal transducer and activator of transcription 3, and NF- κ B^{28–30}. Honokiol can block NF- κ B-controlled gene expression such as matrix metalloproteinase-9, TNF- α , interleukin-8, ICAM-1, and monocyte chemoattractant protein-1 in various cell types^{30–32}. Recently, honokiol was found a natural inhibitor of NF- κ B³³. Honokiol could repress the TNF- α -induced NF- κ B activation through an extracellular signal-regulated kinase pathway in rat aortic smooth muscle cells³⁴. Although honokiol is a potential natural product for treating CNS disorders, how it terminates NF- κ B signalling downstream of the TNF receptor in inflammatory cerebral ECs is unknown.

To elucidate the possible mechanism(s) of the anti-inflammatory effect of honokiol in cerebral ECs, we aimed to investigate how honokiol attenuates a TNF- α -induced inflammatory effect via repressing NF- κ B-activated VCAM-1 expression. Honokiol suppressed TNF- α -induced neutrophil adhesion by inhibiting VCAM-1 gene expression in ECs. Furthermore, it inhibited NF- κ B p65 nuclear translocation and prohibited it from targeting inflammatory genes by blocking I κ B α degradation. Honokiol targeted the ubiquitin–ubiquitin interface of K48-linked polychains, then repressed the TNF- α -induced polyubiquitination of I κ B α . Our study reveals another mechanism of honokiol in protecting ECs against inflammation by modulating the polyubiquitination of I κ B α to fine-tune NF- κ B-activated transcriptional adaptation, with potential as an agent for preventing and treating CNS diseases.

Results

Honokiol suppressed TNF- α -induced neutrophil adhesion and VCAM-1 expression in ECs. To investigate the anti-inflammatory effects of honokiol on ECs, we first explored whether honokiol inhibited neutrophil adhesion to ECs. Stimulation of ECs with TNF- α (5, 10, 25, and 50 ng/ml) dose-dependently promoted neutrophil adhesion (Fig. 1B). On pre-treating ECs with honokiol (1, 3, and 10 μ M) for 30 min, then TNF- α (10 ng/ml), honokiol significantly decreased in the amount of neutrophils on ECs (Fig. 1C,D). Honokiol did not alter EC cell viability with TNF- α (10 ng/ml) treatment for 6 h as demonstrated by viability assay (Fig. 1E), suggesting that honokiol exerted anti-inflammatory effects in ECs without affecting cell viability.

The cell adhesion molecule VCAM-1 has a critical role in leucocyte adhesion to ECs at the site of CNS inflammation. We treated ECs with TNF- α (10 ng/ml) for various times to investigate VCAM-1 mRNA expression. TNF- α significantly increased VCAM-1 mRNA expression at 1, 3, and 6 h, with peak response at 3 h (Fig. 2A). Honokiol pre-treatment dose-dependently reversed TNF- α -induced VCAM-1 mRNA levels (Fig. 2B). Flow cytometry of EC-surface VCAM-1 protein expression confirmed the decreased VCAM-1 expression in ECs (Fig. 2C). Therefore, honokiol suppressed the TNF- α -induced neutrophil adhesion to ECs, which should depend on downregulation of VCAM-1 expression in ECs.

Honokiol prevented NF- κ B nuclear translocation by inhibiting I κ B α degradation. Because the VCAM-1 gene promoter region contains NF- κ B binding sites³⁵ and previous studies indicated that honokiol inhibited NF- κ B activity^{30,31,33,34}, we examined whether honokiol regulated NF- κ B signalling in cerebral ECs. NF- κ B is a heterodimeric transcription factor including p65 (RelA), RelB, c-Rel, p105/p50, and p100/p52 and is inactivated in the cytoplasm when binding to the inhibitor protein I κ B¹⁷. On treatment with stimuli such as TNF- α , I κ B is degraded and causes NF- κ B to translocate into the nucleus¹⁷. Correspondingly, we found that treating ECs with honokiol (3 μ M) suppressed TNF- α -induced p65 nuclear translocation (Fig. 3A). p65 and I κ B α were co-localized in the cytoplasm of control ECs (Fig. 3B). TNF- α (10 ng/ml) treatment for 15 min produced I κ B α degradation and p65 nuclear translocation; however, honokiol (3 μ M) could prevent the TNF- α -induced I κ B α degradation and p65 nuclear translocation (Fig. 3B). Similar results were obtained by western blot analysis: TNF- α -induced I κ B α degradation at 10 and 15 min was significantly abolished by honokiol treatment (3 μ M) (Fig. 4A,B). Additionally, the TNF- α -induced I κ B α phosphorylation was accumulated by honokiol treatment (Fig. 4A,B), suggesting that honokiol may mainly alter on the I κ B α stability instead of phosphorylation. To further determine whether the TNF- α -induced I κ B α degradation or phosphorylation was affected by honokiol, ECs were treated with MG132, a proteasome inhibitor preventing proteasomal degradation of I κ B α ³⁶. MG132 (5 μ M) inhibited TNF- α -induced I κ B α degradation, thus the TNF- α -activated I κ B α phosphorylation was increased. Honokiol had no effect on I κ B α expression and slightly suppressed the TNF- α -induced I κ B α phosphorylation although there was no statistical significance (Fig. 4C,D). We suggest that honokiol may exert dual




Figure 1. Honokiol repressed tumor necrosis factor α (TNF- α)-induced adhesion of neutrophils to endothelial cells (ECs). (A) Chemical structure of honokiol. (B) bEnd.3 ECs were incubated with TNF- α at various concentrations (5, 10, 25, and 50 ng/ml) for 3 h. (C) Human neutrophils were then incubated with ECs for 30 min. Adherent neutrophils on ECs were detected by microscopy. ECs were incubated with DMSO or honokiol (1, 3, and 10 μ M) for 30 min, then stimulated with TNF- α (10 ng/ml) for 3 h. DiI-labelled neutrophils were incubated with ECs for 30 min. Adherent neutrophils on ECs were detected by microscopy. Bars, 200 μ m. (D) Adherent neutrophils were counted and quantified. (E) Cell viability measured by WST-1 reagent in ECs incubated with or without honokiol (1, 3, and 10 μ M) for 30 min, then treated with TNF- α (10 ng/ml) for 6 h. Data are mean \pm SEM from 5 experiments performed in duplicate. * P < 0.05 vs untreated control cells.

effects on TNF- α -controlled I κ B α -NF κ B pathway through the main polyubiquitin targeting and minor phosphorylation signalling in our model. Additionally, it has been reported that honokiol acts on the GABA_A receptor^{20,37}. To explore whether the effect of honokiol in ECs is dependent on GABA_A receptor, the GABA_A receptor antagonist, SR95531, or GABA_A agonist, muscimol, were tested. As the results shown in Fig. 4E, muscimol did not alter the TNF- α -induced I κ B α degradation in ECs. Furthermore, SR95531 did not affect the inhibitory effect of honokiol on TNF- α -induced I κ B α degradation in ECs. These results suggested that honokiol-blocked I κ B α degradation is not changed by the GABA_A receptor antagonist as well as GABA_A agonist.

Honokiol targeted the ubiquitin–ubiquitin interface of K48-linked polychains. Because I κ B α degradation is tightly controlled by ubiquitination¹⁸, we wondered whether honokiol blocked the ubiquitination of I κ B α . Hence, we investigated honokiol docking with K48-linked polyubiquitin. After molecule docking investigations with SwissDock³⁸, 31 clusters of binding models were generated. Honokiol moieties were located near the interface of di-ubiquitin in most of the binding models (Fig. 5A). The best energy binding model is in Fig. 5B; the




Figure 2. Honokiol suppressed TNF- α -induced adhesion molecule expression. (A) RT-qPCR analysis of vascular cell adhesion molecule-1 (VCAM-1) mRNA level in bEnd.3 ECs incubated with DMSO or honokiol (1 μ M) for 30 min, then stimulated with TNF- α (10 ng/ml) for 1, 3, and 6 h. Normalization was to β -actin mRNA level. (B) RT-qPCR analysis of VCAM-1 mRNA level in ECs incubated with honokiol at 1, 3, and 10 μ M for 30 min, then stimulated with TNF- α (10 ng/ml) for 1 h. (C) ECs were incubated with DMSO or honokiol (3 or 10 μ M) for 30 min, then stimulated with TNF- α (10 ng/ml) for 1 h. ECs were stained with rat IgG or anti-VCAM-1 antibodies for 30 min. The unbound antibodies were removed, and then Alexa Fluor 488 anti-rat IgG was added for 30 min. Fluorescein-labelled ECs were analysed for surface VCAM-1 expression by flow cytometry. ECs incubated with rat IgG and Alexa Fluor 488 dye-conjugated antibodies were rat IgG controls. Data are mean \pm SEM from 3 independent experiments. * P < 0.05 vs untreated control. ** P < 0.05 vs corresponding control (TNF- α alone).

aromatic ring of honokiol can interact with hydrophobic residues F45 and A46 in ubiquitin chain A. R74, located at C-terminal ubiquitin chain B was close to the -OH group of honokiol (Fig. 5B). Therefore, honokiol may bind to the ubiquitin-ubiquitin interface of K48-linked polychains and interfere with the polyubiquitin linkage.

Honokiol blocked K48-linked polyubiquitination of I κ B α in ECs. TNF- α -mediated NF- κ B signal transduction is regulated by ubiquitination of various proteins via K48-, K63- or M1 (linear)-linked polychains^{39,40}. To further confirm our docking models in Fig. 5, we determined whether honokiol inhibited I κ B α degradation by directly blocking K48-linked polyubiquitination of I κ B α in ECs. By western blot analysis, pre-treating ECs with honokiol (3 μ M) reduced the TNF- α -induced K48- but not K63- and linear-linked ubiquitin conjugation (Fig. 6A and Supplemental Fig. S1). TNF- α induced the K48-linked polyubiquitin formation with a 2-fold increase that was repressed by honokiol (Fig. 6A). As well, honokiol markedly blocked the 2-fold increase in conjugation of K48-linked polyubiquitin to I κ B α induced by TNF- α on immunoprecipitation assay (Fig. 6B). Additionally, the TNF- α signalling triggers the polyubiquitination of I κ B α through the ubiquitin E3 ligase complex SCF- β -TrCP⁴¹. Honokiol also repressed the 2-fold increase in conjugation of β -TrCP to I κ B α induced by TNF- α (Fig. 6C). Together, we suggest that honokiol would block TNF- α -induced I κ B α degradation by directly disrupting K48-linked polyubiquitination in ECs.




Figure 3. Honokiol blocked the TNF- α -induced NF- κ B p65 nuclear translocation and inhibitor of NF- κ B α (I κ B α) degradation. (A) Inverted microscopy of intracellular localization of NF- κ B p65 (green) in ECs incubated with DMSO or honokiol (3 μ M) for 30 min, then with TNF- α (10 ng/ml) for 15 min. Nuclei were stained with Hoechst 33342 (blue). (B) Confocal microscopy of localization and expression of I κ B α (Red) and p65 (green) in ECs preincubated with DMSO or honokiol (3 μ M) for 30 min, then with TNF- α (10 ng/ml) for 15 min. Bars, 10 μ m. Representative images from 3 or 4 experiments are shown.

Discussion

CNS inflammation is associated with disrupted cerebral EC homeostasis and BBB integrity, followed by activated leukocyte transmigration. The critical pro-inflammatory factor, TNF- α , upregulates adhesion molecules in ECs and facilitates leukocyte extravasation into the CNS^{12,13}. Thus, an attractive therapeutic approach for treating CNS disorders is to limit leukocyte extravasation into the CNS by targeting endothelial adhesion molecules. NF- κ B is considered a critical inflammatory transcription factor upregulating adhesion molecules^{13,14}. In the present study, honokiol diminished TNF- α -induced adhesion between neutrophils and ECs by blocking VCAM-1 mRNA expression and NF- κ B activation in cerebral ECs. This inhibition resulted from directly targeting K48-linked polyubiquitination to prevent I κ B α degradation. Such a unique mechanism explains why honokiol may be a potential natural compound for treating CNS inflammation.

The pro-inflammatory cytokine, TNF- α , can induce neutrophil infiltration into inflammatory regions^{12,13}. Here, we show that TNF- α triggered neutrophil adhesion on ECs via NF- κ B activation and VCAM-1 upregulation in ECs (Figs 1–3). However, TNF- α is also an important cytokine activating neutrophils during inflammation and triggering intracellular signal transduction, including the NF- κ B pathway, which plays a leading role in the function of neutrophils^{42–44}. In Fig. 1D, although the ECs were pre-treated with TNF- α and honokiol before neutrophils adhesion, the effects of TNF- α and honokiol on neutrophils still should be noted. VCAM-1 in ECs of BBB mediates leukocyte transmigration during inflammation by interacting with $\alpha_4\beta_1$ integrin on leukocytes^{45,46}. Also, co-stimulating integrin with cytokines induces NF- κ B activation and I κ B degradation in neutrophils⁴⁷, so TNF- α -induced cell adhesion may be regulated in ECs and neutrophils. Honokiol has an anti-inflammatory effect in neutrophils by inhibiting reactive oxygen species production and neutrophil infiltration *in vivo*^{48–50}. Whether




Figure 4. Honokiol prevented the TNF- α -induced I κ B α proteosomal degradation. (A) Western blot analysis of I κ B α , phospho-I κ B α and tubulin (loading control) protein expression in ECs pre-treated with DMSO or honokiol (3 μ M) for 30 min, then stimulated with TNF- α (10 ng/ml) at different times (5, 10, 15, and 30 min). (B) I κ B α and phospho-I κ B α signals were quantified, expressed as a relative ratio (normalized with tubulin) and plotted against time. (C) Western blot analysis of phospho-I κ B α , I κ B α and tubulin protein expression in ECs treated with MG132 (5 μ M) for 1 h, then incubated with DMSO or honokiol (3 μ M) for 30 min and stimulated with TNF- α (10 ng/ml). (D) Phospho-I κ B α and I κ B α signals were quantified, expressed as a relative ratio (normalized with tubulin) and plotted against time. (E) EC cells were exposure to DMSO or the GABA_A receptor antagonist, SR95531 (20 μ M), for 30 min and sequentially treated with DMSO, honokiol (3 μ M), or GABA_A agonist, muscimol (100 μ M) for 30 min. The ECs were stimulated with TNF- α (10 ng/ml) for 15 min and analyzed using western blot. The I κ B α signals were quantified expressed as a relative ratio (normalized with tubulin). Representative images from 3 experiments are shown. Data are mean \pm SEM from 3 independent experiments. * P < 0.05; ** P < 0.01 versus untreated control cells. All the Western blotting experiments were performed under the following condition. After transferring the blots onto nitrocellulose membranes, we immediately cropped the targeted blots according to referenced indicating markers, and then targeted proteins were immunoblotted with its specific monoclonal antibody.




Figure 5. Docking models of honokiol-targeted ubiquitin. (A) Surface presentation demonstrates the structure of di-ubiquitin (gray). Honokiol moieties are coloured yellow and rendered in stick representation. (B) Close-up of honokiol docking site (best energy mode) in the ubiquitin–ubiquitin interface of Lys48-linked chains. The figures were prepared by using Chimera software. The crystal structure of ubiquitin was obtained from the PDB (accession code 1F9J)⁶². The honokiol 3D structure was obtained from the Zinc database (code 1536, ZINC 12)⁶³.

honokiol blocks neutrophil-associated diseases by repressing inflammation-induced NF- κ B activation could be addressed.

How TNF- α activates NF- κ B pathways in cells has been established^{14,18,19}. In general, the binding of TNF to its receptor (TNFR1) leads to recruitment of several ubiquitin E3 ligases to catalyze K63-linked polyubiquitination of receptor-interacting protein (RIP1), which clusters the transforming growth factor- β -activated kinase 1 (TAK1) and I κ B kinase (IKK) complex. TAK1 phosphorylation of IKK α / β triggers I κ B α phosphorylation and sequential K48-linked polyubiquitination and degradation of I κ B α to activate NF- κ B. Here, we showed that honokiol may inhibit I κ B α degradation by directly binding to and disrupting K48-linked polyubiquitin, thereby suppressing substrate digestion by the proteasome (Figs 4–6). However, honokiol has been found to suppress NF- κ B activation by regulating IKK–I κ B α phosphorylation signalling^{30,31}. RIP1 is conjugated with K63-linked polyubiquitin following TNF stimulation, then becomes deubiquitinated by A20 through the N-terminal OTU domain of A20. RIP1 is then convertibly conjugated with K48-linked polyubiquitin by the C-terminal zinc finger domain of A20, thereby leading to the proteasomal degradation of RIP1 and inactivation of IKK^{14,51}. Thus, IKK–I κ B α phosphorylation signalling should be affected by upstream ubiquitination. If honokiol also interferes with K48-linked polyubiquitin of RIP1, it should prevent RIP1 degradation and cause IKK activation and I κ B α degradation, which would conflict with honokiol-repressed I κ B α degradation (Fig. 4). Thus, we focused on the ubiquitination rather than phosphorylation of I κ B α in this study, and the detailed association needs further elucidation. In addition, the IKK complex is also activated by K63/M1-linked hybrid ubiquitin chains^{39,40}. However, cells that cannot generate K63-linked polyubiquitin can still activate IKK and NF- κ B with TNF stimulation⁵², which suggests that K63 polyubiquitination-mediated phosphorylation and activation of IKK–I κ B α pathway may be not essential for NF- κ B activity. As we show in Figs 5 and 6, TNF- α mainly induced K48- but not K63- or linear (M1)-linked polyubiquitin, and honokiol also interfered with K48- but not K63- or linear (M1)-linked polyubiquitin, so honokiol-targeted K48 polyubiquitination may be more dominant in TNF-induced CNS inflammation.

Furthermore, honokiol has been reported to act on the GABA_A receptor^{20,37}. In Fig. 4E, we found that muscimol did not alter the TNF- α -induced I κ B α degradation and the GABA_A receptor antagonist, SR95531, also did not affect the inhibitory effect of honokiol on TNF- α -induced I κ B α degradation in ECs, suggesting that honokiol-blocked I κ B α degradation is not changed by the GABA_A receptor antagonist as well as GABA_A agonist, although muscimol and SR95531 does not address the benzodiazepine binding site of the GABA_A receptor. In addition, we found that TNF- α -induced NF- κ B p65 nuclear translocation was not completely blocked by honokiol in Fig. 3, however, honokiol significantly inhibited the TNF- α -triggered I κ B α degradation at this time point (Figs 3B and 4A). It has been reported that the p65 nuclear translocation was controlled by phosphorylation at Ser536 of itself which was independent on I κ B α ⁵³, suggesting that honokiol may not totally control the TNF- α -regulated NF- κ B activity because of I κ B α -independent regulation. On the other hand, the TNF- α still caused the I κ B α degradation in the presence of honokiol after longer 30 min stimulation in Fig. 4A, suggesting that the effect of honokiol on TNF- α -mediated NF- κ B activation may be a delayed response rather than a merely suppressed response, although honokiol would repress the VCAM-1 expression or neutrophil adhesion after 1–3 h TNF- α stimulation (Figs 1 and 2).

Other small compounds, ubistatins, could prevent the polyubiquitination and degradation of Sic1; NMR structures demonstrated that ubistatins blocked proteasome-dependent degradation by binding the ubiquitin–ubiquitin interface of K48-linked polychains⁵⁴. Similarly, our docking models revealed that honokiol may target the ubiquitin–ubiquitin interface of K48-linked chains (Fig. 5A). The hydrophobic interaction between the aromatic ring of honokiol and F45 or A46 may promote a conformational change of the ubiquitin loop.




Figure 6. Honokiol blocked TNF- α -induced K48-linked polyubiquitination in ECs. (A) Western blot analysis of K48-linked polyubiquitination in the presence and absence of honokiol or TNF- α . The whole ubiquitin signals were quantified expressed as a relative ratio (normalized with tubulin). (B,C) Immunoprecipitation and western blot assay of the interaction between K48-linked polyubiquitin (B) or β -TrCP (C) and I κ B α in ECs pre-treated with honokiol (3 μ M) for 30 min, then stimulated with TNF- α (10 ng/ml) for 15 min in the presence of MG132 (5 μ M). The I κ B α -precipitated K48-linked polyubiquitin (100–170 kDa) (B) or β -TrCP (C) signals were quantified expressed as a relative ratio (normalized with input I κ B α). Representative images from 3 experiments are shown.

Simultaneously, R74 located at ubiquitin chain B may connect to the hydroxyl group of honokiol by a hydrogen bond, for an orientation change at the C-terminal ubiquitin chain A (Fig. 5B). Honokiol may induce a conformational change causing K48 not to bind to G76 and disrupt K48-linked polyubiquitin linkage, thereby directly interfering with K48-linked ubiquitination in ECs. Furthermore, co-crystallizing the complex of honokiol with various linked ubiquitins including K48- and K63-linked and linear chain ubiquitins may help to understand the mechanistic basis of the biological action of honokiol on ubiquitination. Moreover, honokiol may block the TNF- α signalling-triggered polyubiquitination of I κ B α through the ubiquitin E3 ligase complex SCF- β -TrCP⁴¹ (Fig. 6B,C). The detail mechanism of honokiol-blocked polyubiquitin via affecting the interaction between I κ B α and ubiquitin ligase as well as deubiquitin enzyme may need further investigation.

Currently, there is great interest in developing new drugs from medicinal plants; however, which protein or protein group can be targeted by drugs is difficult to determine. In this study, we demonstrate that the small natural product honokiol is a unique example of disrupting the ubiquitin-ubiquitin interface of K48-linked polyubiquitins. Thus, honokiol could prevent I κ B α degradation after TNF- α treatment, then the constitutively expressed I κ B α prohibit NF- κ B from translocating to the nucleus and activating inflammatory genes such as VCAM-1 in ECs. Our study supports an additional novel pharmacological mechanism of honokiol in anti-inflammation and understanding the molecular mechanism of honokiol in cells to enhance its efficiency. Toxicological studies of honokiol treatment have demonstrated no pathologic changes in organs, including liver, lung, kidney, spleen, brain, heart, pancreas, intestines, or bone marrow, after systemic or oral administration⁵⁵. Honokiol also shows high ability to penetrate the BBB²⁷. Honokiol is a potential natural product for preventing and treating CNS inflammatory diseases.

In summary, we reveal how honokiol prevents TNF- α -induced neutrophil adhesion on cerebral ECs by disrupting the polyubiquitination and degradation of I κ B α to block NF- κ B-controlled VCAM-1 expression.

Methods

Extraction and isolation of honokiol. The dried bark of *Magnolia officinalis* cortex (3 kg) was repeatedly extracted with MeOH (20 L × 3) at 70 °C for 4 h. The methanol extracts were evaporated and partitioned between CH₂Cl₂ and H₂O. The CH₂Cl₂ layer was subjected to column chromatography over silica gel and gradient elution with hexane-EtOAc and EtOAc-MeOH mixtures used as solvent systems. The fraction eluted with hexane-EtOAc (10:1–4:1) was repeatedly chromatographed with silica gel and Sephadex LH-20 to obtain honokiol (6.8 g)⁵⁶. The structure of honokiol was determined by ¹H NMR (CDCl₃, 500 MHz) spectrum analysis (Supplemental Fig. S1) and compared with reference data⁵⁷. When honokiol was dissolved in dimethyl sulfoxide (DMSO), the final concentration of DMSO in the cell experiments is 0.1% and did not affect the parameters measured.

Reagents. Rat monoclonal antibodies (mAbs) against VCAM-1 (NBP1-26587) were from Novus biologicals (Littleton, Colorado). Rabbit mAbs against NF-κB p65 (#8242), K48-linkage specific polyubiquitin (#8081), β-TrCP (#4394) and α/β-Tubulin (#2148) as well as mouse mAbs against IκBα (#4814) and phospho-IκBα (Ser32/36) (#9246) were from Cell Signaling Technology (Beverly, MA). Rabbit mAbs against K63-linkage specific polyubiquitin (05–1308) and linear ubiquitin (MABS199) were from Millipore (Temecula, CA). Recombinant mouse TNF-α, rat IgG and mouse IgG₁ was from R&D systems (Minneapolis, MN). Muscimol was from Tocris Bioscience (Ellisville, MO). All other reagent-grade chemicals were from Sigma (St Louis, MO), including MG132 and SR95531.

Cell culture. Mouse brain microvascular ECs (bEnd.3) were purchased from the Bioresource Collection and Research Centre (Hsinchu, Taiwan). Cultures were maintained in a humidified atmosphere (37 °C, 5% CO₂) with Dulbecco's modified Eagle's medium (DMEM, Gibco, Grand Island, NY) supplemented with 10% fetal bovine serum (FBS; Biological Industries, Israel) and 100 U/ml penicillin/streptomycin (Gibco). ECs were grown to confluence and passaged every 3 days at 1 × 10⁵ cells/ml.

Neutrophil isolation. Human neutrophils were isolated from heparinised peripheral venous blood by using a standard method of dextran sedimentation and Ficoll–Hypaque centrifugation^{58–61}. After hypotonic lysis of contaminating erythrocytes, isolated neutrophils were suspended in RPMI 1640 containing 0.1% FBS. Blood was collected from healthy volunteers (ages 20–30 years). This study protocol was investigated and approved by the Institutional Review Board at Chang Gung Memorial Hospital, and written informed consent was obtained from every volunteer. The methods were carried out in accordance with the approved guidelines.

Neutrophil adhesion experiment. Neutrophils were labelled with 1,1'-dioctadecyl-3,3',3',3'-tetramethylindocarbocyanine (DiI; Molecular Probes, Eugene, OR) for 20 min in RPMI 1640 containing 0.1% FBS. ECs were treated with DMSO or honokiol (1, 3, and 10 μM) for 30 min before TNF-α (5, 10, 25, and 50 ng/ml) for 3 h. The labelled neutrophils (1 × 10⁵ cells/mL) were then incubated with ECs for 30 min. Non-adherent neutrophils were removed by gentle washing with RPMI. Adherent neutrophils on ECs were counted in 6 randomly selected areas (0.572 mm²) under a motorised inverted microscope (IX81, Olympus, Japan) with 10X objective.

Cell viability assay. Cell viability was measured by WST-1 assay (Roche, Mannheim, Germany). ECs were seeded into 96 wells for 24 h at 5 × 10⁴ cells/well in 100 μl culture medium and treated with or without honokiol (1, 3, and 10 μM) for 30 min before TNF-α (10 ng/ml) for 6 h, then WST-1 (10 μl) was added and absorbance was measured at 450 nm. The reference wavelength was 600 nm.

RNA isolation and quantitative real-time PCR. Total RNA was extracted from the ECs by using TRIzol reagent (Invitrogen, Carlsbad, CA) and used as a template for cDNA synthesis. cDNA, obtained with High Capacity cDNA Reverse Transcription Kits (Applied Biosystems, Foster City, CA), was amplified by PCR on a CFX Connect (Bio-Rad) with Power SYBR Green PCR Master Mix (Applied Biosystems) and primers for VCAM-1 (100 nM, sense: 5'-ACGTCAGAACACCGAATCC-3'; antisense: 5'-GTGGTGCTGTGACAATGACC-3') and β-actin (sense: 5'-CTGGGTCATCTTTTCACGGT-3'; antisense: 5'-TGTTACCAACTGGGACGACA-3') used as an internal control. The real-time PCR was conducted at 95 °C for 10 min followed by 40 cycles of denaturation at 95 °C for 10 sec, annealing/extension at 60 °C for 1 min. PCR conditions were optimised to achieve a single peak by melting curve analysis on CFX Connect (Bio-Rad). Raw data collected by use of CFX Manager were analysed and quantified by the relative standard curve method.

Immunofluorescence with flow cytometry. ECs (5 × 10⁶/mL) were detached with Versene buffer containing EDTA and suspended in phosphate buffered saline (PBS) containing rat IgG control or rat monoclonal antibodies against VCAM-1 (10 μg/mL) at 4 °C for 30 min. The unbound antibodies were removed, then Alexa Fluor 488 anti-rat IgG (Invitrogen) was added for 30 min at 4 °C in PBS containing 10% FBS. Fluorescein-labelled ECs were analysed for surface VCAM-1 expression by flow cytometry (BD Accuri C6, BD Biosciences, San Jose, CA). ECs incubated with rat IgG and Alexa Fluor 488 dye-conjugated antibodies were rat IgG controls.

Immunofluorescence and microscopy. ECs were grown on coverslips and fixed in 4% paraformaldehyde at room temperature for 15 min. Cells were blocked and permeabilised for 30 min with a solution of PBS containing 0.2% Triton X-100 and 1% bovine serum albumin. After incubation with primary antibodies (p65; 1:200 and IκBα; 1:100, Cell Signaling) for 1 h, ECs were washed and incubated with secondary antibodies (Alexa Fluor 488 or 594; 1:500, Molecular Probes) for 30 min. After the addition of Hoechst 33342 (1 μg/ml, Invitrogen) for another 5 min, cells were washed and mounted with prolong antifade reagent (Invitrogen) on coverslips. Fluorescence

was captured under a motorised inverted microscope (IX81, Olympus) and confocal microscope (LSM510, Carl Zeiss). We used a 63×/NA 1.4 or 100×/NA 1.4 oil objectives (Carl Zeiss). Images were obtained by using LSM 510 META (Carl Zeiss).

Immunoprecipitation and immunoblot analysis. Cell lysates were prepared in lysis buffer (50 mM Tris-HCl, 150 mM NaCl, 0.5% Nonidet P-40, 5 mM EDTA, 50 mM NaF containing protease and phosphatase inhibitors (Roche) and rotated at 4 °C for 30 min. Protein concentration were determined by using a BCA protein assay kit (Pierce, Rockford, IL). Immunoprecipitation involved 200 µg cell lysates and 0.5 µg mouse IgG1 or anti-IκBα antibodies with overnight incubation at 4 °C, then 100 µl protein G plus/protein A agarose (Calbiochem) added for 12 h with rotation at 4 °C. Beads were washed 4 times with lysis buffer. Bound proteins were eluted by boiling for 10 min in sample buffer, and eluted samples were loaded onto an 8% SDS-PAGE gel. The corresponding lysate (30 µg protein) was loaded onto an SDS-PAGE gel (10% running, 4% stacking) and transferred to PVDF membranes (Millipore), which were blocked with 5% milk in TBS, 0.1% Tween 20 (TBST), then incubated with the designated antibodies and horseradish peroxidase-conjugated secondary antibodies (Cell Signaling). Peroxidase activity was evaluated by using an enhanced chemiluminescence kit (Millipore). The reactive band intensities were analysed by using UVP Biospectrum (UVP LLC, Upland, CA).

Molecular docking. Small molecules were docked and analyzed on target proteins by using SwissDock, a small molecule-protein docking web service established on EADock DSS³⁸. The binding of honokiol and ubiquitin with the most favourable energy was estimated with FACTS and clustered. The crystal structure of ubiquitin was obtained from the PDB (accession code 1F9J)⁶². The 3D structure of honokiol was obtained from the Zinc database (code 1536, ZINC 12)⁶³. The predicted binding modes were presented by using UCSF Chimera (a visualisation system for exploratory research and analysis)⁶⁴.

Statistical analysis. Data are expressed as mean ± SEM. Statistical analysis involved Student's *t* test for two groups and one-way ANOVA followed by Scheffé's test for multiple comparisons. *P* < 0.05 was considered statistically significant.

References

- Combes, V., Guillemin, G. J., Chan-Ling, T., Hunt, N. H. & Grau, G. E. The crossroads of neuroinflammation in infectious diseases: endothelial cells and astrocytes. *Trends Parasitol* **28**, 311–9 (2012).
- Abbott, N. J., Patabendige, A. A., Dolman, D. E., Yusof, S. R. & Begley, D. J. Structure and function of the blood-brain barrier. *Neurobiol Dis* **37**, 13–25 (2010).
- Petty, M. A. & Lo, E. H. Junctional complexes of the blood-brain barrier: permeability changes in neuroinflammation. *Prog Neurobiol* **68**, 311–23 (2002).
- Chavarria, A. & Alcocer-Varela, J. Is damage in central nervous system due to inflammation? *Autoimmun Rev* **3**, 251–60 (2004).
- Rossi, B., Angiari, S., Zenaro, E., Budui, S. L. & Constantin, G. Vascular inflammation in central nervous system diseases: adhesion receptors controlling leukocyte-endothelial interactions. *J Leukoc Biol* **89**, 539–56 (2011).
- Muller, W. A. Mechanisms of transendothelial migration of leukocytes. *Circ Res* **105**, 223–30 (2009).
- Langer, H. F. & Chavakis, T. Leukocyte-endothelial interactions in inflammation. *J Cell Mol Med* **13**, 1211–20 (2009).
- Hua, S. Targeting sites of inflammation: intercellular adhesion molecule-1 as a target for novel inflammatory therapies. *Front Pharmacol* **4**, 127 (2013).
- Supanc, V., Biloglav, Z., Kes, V. B. & Demarin, V. Role of cell adhesion molecules in acute ischemic stroke. *Ann Saudi Med* **31**, 365–70 (2011).
- Serres, S. *et al.* VCAM-1-targeted magnetic resonance imaging reveals subclinical disease in a mouse model of multiple sclerosis. *FASEB J* **25**, 4415–22 (2011).
- Soto, M. S., Serres, S., Anthony, D. C. & Sibson, N. R. Functional role of endothelial adhesion molecules in the early stages of brain metastasis. *Neuro Oncol* **16**, 540–51 (2014).
- Sharief, M. K. & Hentges, R. Association between tumor necrosis factor-α and disease progression in patients with multiple sclerosis. *N Engl J Med* **325**, 467–72 (1991).
- Kallmann, B. A. *et al.* Cytokine-induced modulation of cellular adhesion to human cerebral endothelial cells is mediated by soluble vascular cell adhesion molecule-1. *Brain* **123** (Pt 4), 687–97 (2000).
- Walczak, H. TNF and ubiquitin at the crossroads of gene activation, cell death, inflammation, and cancer. *Immunol Rev* **244**, 9–28 (2011).
- Iwai, K. Diverse ubiquitin signaling in NF-κB activation. *Trends Cell Biol* **22**, 355–64 (2012).
- Napetschnig, J. & Wu, H. Molecular basis of NF-κB signaling. *Annu Rev Biophys* **42**, 443–68 (2013).
- Hayden, M. S. & Ghosh, S. Shared principles in NF-κB signaling. *Cell* **132**, 344–62 (2008).
- Wullaert, A., Heynink, K., Janssens, S. & Beyaert, R. Ubiquitin: tool and target for intracellular NF-κB inhibitors. *Trends Immunol* **27**, 533–40 (2006).
- Bianchi, K. & Meier, P. A tangled web of ubiquitin chains: breaking news in TNF-R1 signaling. *Mol Cell* **36**, 736–42 (2009).
- Fried, L. E. & Arbiser, J. L. Honokiol, a multifunctional antiangiogenic and antitumor agent. *Antioxid Redox Signal* **11**, 1139–48 (2009).
- Chen, C. M., Liu, S. H. & Lin-Shiau, S. Y. Honokiol, a neuroprotectant against mouse cerebral ischaemia, mediated by preserving Na⁺, K⁺-ATPase activity and mitochondrial functions. *Basic Clin Pharmacol Toxicol* **101**, 108–16 (2007).
- Harada, S. *et al.* Honokiol suppresses the development of post-ischemic glucose intolerance and neuronal damage in mice. *J Nat Med* **66**, 591–9 (2012).
- Chen, Y. J. *et al.* Honokiol, a low molecular weight natural product, prevents inflammatory response and cartilage matrix degradation in human osteoarthritis chondrocytes. *J Orthop Res* **32**, 573–80 (2014).
- Singh, T. & Katiyar, S. K. Honokiol inhibits non-small cell lung cancer cell migration by targeting PGE(2)-mediated activation of beta-catenin signaling. *PLoS One* **8**, e60749 (2013).
- Chuang, D. Y. *et al.* Magnolia polyphenols attenuate oxidative and inflammatory responses in neurons and microglial cells. *J Neuroinflammation* **10**, 15 (2013).
- Bai, X. *et al.* Honokiol, a small molecular weight natural product, inhibits angiogenesis *in vitro* and tumor growth *in vivo*. *J Biol Chem* **278**, 35501–7 (2003).
- Wang, X. *et al.* Honokiol crosses BBB and BCSFB, and inhibits brain tumor growth in rat 9L intracerebral gliosarcoma model and human U251 xenograft glioma model. *PLoS One* **6**, e18490 (2011).

28. Leeman-Neill, R. J. *et al.* Honokiol inhibits epidermal growth factor receptor signaling and enhances the antitumor effects of epidermal growth factor receptor inhibitors. *Clin Cancer Res* **16**, 2571–9 (2010).
29. Rajendran, P. *et al.* Honokiol inhibits signal transducer and activator of transcription-3 signaling, proliferation, and survival of hepatocellular carcinoma cells via the protein tyrosine phosphatase SHP-1. *J Cell Physiol* **227**, 2184–95 (2012).
30. Tse, A. K., Wan, C. K., Shen, X. L., Yang, M. & Fong, W. F. Honokiol inhibits TNF- α -stimulated NF- κ B activation and NF- κ B-regulated gene expression through suppression of IKK activation. *Biochem Pharmacol* **70**, 1443–57 (2005).
31. Ahn, K. S. *et al.* Honokiol potentiates apoptosis, suppresses osteoclastogenesis, and inhibits invasion through modulation of nuclear factor- κ B activation pathway. *Mol Cancer Res* **4**, 621–33 (2006).
32. Zhang, P. *et al.* Honokiol inhibits the inflammatory reaction during cerebral ischemia reperfusion by suppressing NF- κ B activation and cytokine production of glial cells. *Neurosci Lett* **534**, 123–7 (2013).
33. Nabekura, T., Hiroi, T., Kawasaki, T. & Uwai, Y. Effects of natural nuclear factor- κ B inhibitors on anticancer drug efflux transporter human P-glycoprotein. *Biomed Pharmacother* **70**, 140–5 (2015).
34. Zhu, X., Wang, Z., Hu, C., Li, Z. & Hu, J. Honokiol suppresses TNF- α -induced migration and matrix metalloproteinase expression by blocking NF- κ B activation via the ERK signaling pathway in rat aortic smooth muscle cells. *Acta Histochem* **116**, 588–95 (2014).
35. Iademaro, M. F., McQuillan, J. J., Rosen, G. D. & Dean, D. C. Characterization of the promoter for vascular cell adhesion molecule-1 (VCAM-1). *J Biol Chem* **267**, 16323–9 (1992).
36. Lee, D. H. & Goldberg, A. L. Proteasome inhibitors: valuable new tools for cell biologists. *Trends Cell Biol* **8**, 397–403 (1998).
37. Ai, J., Wang, X. & Nielsen, M. Honokiol and magnolol selectively interact with GABAA receptor subtypes *in vitro*. *Pharmacology* **63**, 34–41 (2001).
38. Grosdidier, A., Zoete, V. & Michielin, O. SwissDock, a protein-small molecule docking web service based on EADock DSS. *Nucleic Acids Res* **39**, W270–7 (2011).
39. Emmerich, C. H. *et al.* Activation of the canonical IKK complex by K63/M1-linked hybrid ubiquitin chains. *Proc Natl Acad Sci USA* **110**, 15247–52 (2013).
40. Iwai, K. Diverse roles of the ubiquitin system in NF- κ B activation. *Biochim Biophys Acta* **1843**, 129–36 (2014).
41. Kanarek, N. & Ben-Neriah, Y. Regulation of NF- κ B by ubiquitination and degradation of the I κ Bs. *Immunol Rev* **246**, 77–94 (2012).
42. Futosi, K., Fodor, S. & Mocsai, A. Reprint of Neutrophil cell surface receptors and their intracellular signal transduction pathways. *Int Immunopharmacol* **17**, 1185–97 (2013).
43. Bastien, D. & Lacroix, S. Cytokine pathways regulating glial and leukocyte function after spinal cord and peripheral nerve injury. *Exp Neurol* **258**, 62–77 (2014).
44. Liu, S. F. & Malik, A. B. NF- κ B activation as a pathological mechanism of septic shock and inflammation. *Am J Physiol Lung Cell Mol Physiol* **290**, L622–L645 (2006).
45. Luster, A. D., Alon, R. & von Andrian, U. H. Immune cell migration in inflammation: present and future therapeutic targets. *Nat Immunol* **6**, 1182–90 (2005).
46. Shulman, Z. *et al.* Lymphocyte crawling and transendothelial migration require chemokine triggering of high-affinity LFA-1 integrin. *Immunity* **30**, 384–96 (2009).
47. Kettritz, R., Choi, M., Rolle, S., Wellner, M. & Luft, F. C. Integrins and cytokines activate nuclear transcription factor- κ B in human neutrophils. *J Biol Chem* **279**, 2657–65 (2004).
48. Liou, K. T., Shen, Y. C., Chen, C. F., Tsao, C. M. & Tsai, S. K. Honokiol protects rat brain from focal cerebral ischemia-reperfusion injury by inhibiting neutrophil infiltration and reactive oxygen species production. *Brain Res* **992**, 159–66 (2003).
49. Liu, J. *et al.* Honokiol downregulates Kruppel-like factor 4 expression, attenuates inflammation, and reduces histopathology after spinal cord injury in rats. *Spine (Phila Pa 1976)* **40**, 363–8 (2015).
50. Liou, K. T., Shen, Y. C., Chen, C. F., Tsao, C. M. & Tsai, S. K. The anti-inflammatory effect of honokiol on neutrophils: mechanisms in the inhibition of reactive oxygen species production. *Eur J Pharmacol* **475**, 19–27 (2003).
51. Heyninck, K. & Beyaert, R. A20 inhibits NF- κ B activation by dual ubiquitin-editing functions. *Trends Biochem Sci* **30**, 1–4 (2005).
52. Xu, M., Skaug, B., Zeng, W. & Chen, Z. J. A ubiquitin replacement strategy in human cells reveals distinct mechanisms of IKK activation by TNF α and IL-1 β . *Mol Cell* **36**, 302–14 (2009).
53. Sasaki, C. Y., Barberi, T. J., Ghosh, P. & Longo, D. L. Phosphorylation of RelA/p65 on serine 536 defines an I κ B α -independent NF- κ B pathway. *J Biol Chem* **280**, 34538–47 (2005).
54. Verma, R. *et al.* Ubistatins inhibit proteasome-dependent degradation by binding the ubiquitin chain. *Science* **306**, 117–20 (2004).
55. Arora, S. *et al.* Honokiol: a novel natural agent for cancer prevention and therapy. *Curr Mol Med* **12**, 1244–52 (2012).
56. Lin, C. F. *et al.* Maximizing dermal targeting and minimizing transdermal penetration by magnolol/honokiol methoxylation. *Int J Pharm* **445**, 153–62 (2013).
57. Li, G. H. *et al.* Nematicidal Activity of Honokiol and Magnolol Isolated from Magnolia tripetala. *J Phytopathology* **157**, 390–392 (2009).
58. Hwang, T. L. *et al.* Potent inhibition of superoxide anion production in activated human neutrophils by isopedicin, a bioactive component of the Chinese medicinal herb *Fissistigma oldhamii*. *Free Radic Biol Med* **46**, 520–8 (2009).
59. Hwang, T. L. *et al.* Soluble guanylyl cyclase activator YC-1 inhibits human neutrophil functions through a cGMP-independent but cAMP-dependent pathway. *Mol Pharmacol* **64**, 1419–27 (2003).
60. Yu, H. P. *et al.* 2-(2-Fluorobenzamido)benzoate ethyl ester (EFB-1) inhibits superoxide production by human neutrophils and attenuates hemorrhagic shock-induced organ dysfunction in rats. *Free Radic Biol Med* **50**, 1737–48 (2011).
61. Yang, S. C. *et al.* Propofol inhibits superoxide production, elastase release, and chemotaxis in formyl peptide-activated human neutrophils by blocking formyl peptide receptor 1. *J Immunol* **190**, 6511–9 (2013).
62. Phillips, C. L., Thrower, J., Pickart, C. M. & Hill, C. P. Structure of a new crystal form of tetraubiquitin. *Acta Crystallogr D Biol Crystallogr* **57**, 341–4 (2001).
63. Irwin, J. J. & Shoichet, B. K. ZINC—a free database of commercially available compounds for virtual screening. *J Chem Inf Model* **45**, 177–82 (2005).
64. Pettersen, E. F. *et al.* UCSF Chimera—a visualization system for exploratory research and analysis. *J Comput Chem* **25**, 1605–12 (2004).

Acknowledgements

This research was financially supported by the grants from the Ministry of Science Technology (NSC 102-2628-B-182-002-MY3, MOST 102-2628-B-255-003-MY3 and MOST 104-2320-B-255-004-MY3), Ministry of Education (EMRPD1F0311), and Chang Gung Memorial Hospital (CMRPD1B0281~3, CMRPF1D0442~3, CMRPF1F0011~3, CMRPF1F0061~3, CMRPG6D0083 and BMRP450), Taiwan. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

P.-J.C. and Y.-L.W. designed and performed most experiments. L.-M.K., C.-Y.C., Y.-F.T., C.-F.L. and J.-J.S. helped to perform experiments and analyzed the data. P.-J.C., Y.-L.W. and T.-L.H. wrote and completed the manuscript. T.-L.H. supervised the entire study.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Chen, P.-J. *et al.* Honokiol suppresses TNF- α -induced neutrophil adhesion on cerebral endothelial cells by disrupting polyubiquitination and degradation of I κ B α . *Sci. Rep.* **6**, 26554; doi: 10.1038/srep26554 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>