




REVIEW

Research advances in plant–insect molecular interaction

[version 1; peer review: 2 approved]

Chun-Yu Chen^{1,2}, Ying-Bo Mao ¹

¹Chinese Academy of Sciences (CAS) Key Laboratory of Insect Developmental and Evolutionary Biology, CAS Center for Excellence in Molecular Plant Sciences, Shanghai Institute of Plant Physiology and Ecology, University of CAS, Chinese Academy of Sciences, Shanghai, China

²National Key Laboratory of Plant Molecular Genetics, CAS Center for Excellence in Molecular Plant Sciences, Shanghai Institute of Plant Physiology and Ecology, University of CAS, Chinese Academy of Sciences, Shanghai, China

V1 First published: 19 Mar 2020, 9(F1000 Faculty Rev):198 (<https://doi.org/10.12688/f1000research.21502.1>)

Latest published: 19 Mar 2020, 9(F1000 Faculty Rev):198 (<https://doi.org/10.12688/f1000research.21502.1>)

Abstract



Acute and precise signal perception and transduction are essential for plant defense against insects. Insect elicitors—that is, the biologically active molecules from insects' oral secretion (which contains regurgitant and saliva), frass, ovipositional fluids, and the endosymbionts—are recognized by plants and subsequently induce a local or systematic defense response. On the other hand, insects secrete various types of effectors to interfere with plant defense at multiple levels for better adaptation. Jasmonate is a main regulator involved in plant defense against insects and integrates with multiple pathways to make up the intricate defense network. Jasmonate signaling is strictly regulated in plants to avoid the hypersensitive defense response and seems to be vulnerable to assault by insect effectors at the same time. Here, we summarize recently identified elicitors, effectors, and their target proteins in plants and discuss their underlying molecular mechanisms.

Keywords

Plant defense, Insect herbivory, Jasmonate (JA), Elicitor, Effector

Open Peer Review

Reviewer Status  

	Invited Reviewers	
	1	2
version 1 19 Mar 2020		

F1000 Faculty Reviews are written by members of the prestigious F1000 Faculty. They are commissioned and are peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

1 **Gary Felton**, Pennsylvania State University, University Park, USA

2 **Gregg Howe**, Michigan State University, East Lansing, USA

Any comments on the article can be found at the end of the article.

Corresponding author: Ying-Bo Mao (ybmao@sibs.ac.cn)

Author roles: Chen CY: Writing – Original Draft Preparation; Mao YB: Writing – Original Draft Preparation

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Ministry of Science and Technology of China grant 2016YFA0500803, the Ministry of Agriculture of China grant 2016ZX08009001-009, National Natural Sciences of China grants 31772177 and 31788103, and Chinese Academy of Sciences grant QYZDY-SSW-SMC026.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2020 Chen CY and Mao YB. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Chen CY and Mao YB. **Research advances in plant–insect molecular interaction [version 1; peer review: 2 approved]** F1000Research 2020, 9(F1000 Faculty Rev):198 (<https://doi.org/10.12688/f1000research.21502.1>)

First published: 19 Mar 2020, 9(F1000 Faculty Rev):198 (<https://doi.org/10.12688/f1000research.21502.1>)

Introduction

There are about 1 million insects and over 300,000 plants on our planet, and plant–insect interactions are the driving force of biodiversity. With long-term co-evolution, plants and insects have developed sophisticated mechanisms for adaptation¹. In general, plants can recognize herbivore-/damage-/microbe-associated molecular patterns (HAMPs/DAMPs/MAMPs) and make the right defense. The early defense responses contain depolarization of the plasma transmembrane potential, changes of cytosolic Ca²⁺, reactive oxygen species (ROS) burst, and mitogen-activated protein kinase (MAPK)^{2,3}. Most of these reactions are able to activate jasmonate (JA)-mediated plant defense^{4,5}. JA is a main regulator of plant defense and its synthesis and regulation have been extensively studied^{6–9}. Recent studies reveal new insights in JA oxidative metabolism and their negative regulation in the JA pathway^{10,11}. In most plants, JA-Ile is the active signal recognized by the COI1 and promotes JAZ–COI1 interaction leading to JAZ degradation. This relieves the JAZ-interacting transcription factors to activate downstream defense gene expressions^{12–16}. However, in *Marchantia polymorpha*, MpCOI1 recognized OPDA-Ile instead of JA-Ile. That work revealed the ligand-receptor co-evolution of the JA signaling pathway in land plants¹⁷. MYC2 is a well-studied transcription factor in JA signaling and can interact with both JAZ and MED25, the subunit of the mediator complex. The JAZ proteins recruit TOPLESS scaffold protein to inhibit gene transcription, whereas MED25 brings COI1 to MYC2 targeting promoters¹⁸. In this model, COI1 is thought to be the nuclear receptor. JAT1, which localizes at the nuclear envelope and plasma membrane, is the transporter responsible for the influx of JA-Ile into nucleus¹⁹. To balance the tradeoff between growth and defense, plants strictly regulate JA signaling to avoid a hypersensitive defense response^{20,21}. Some development regulators, including SPLs and DELLAs, target JAZ or MYC transcription factors to modulate JA signaling output^{22–26}. Interestingly, some insects use similar strategies to attenuate plant defense for fitness.

Herbivorous insects have different mouthparts and feeding habits. Active molecules from insects' oral secretion (OS) (which contains regurgitant and saliva), frass, ovipositional fluids, and the endosymbionts of insects have a large impact on plant defense. Some of these molecules used by plants to trigger specialized defense are called elicitors, and those to weaken the plant defense response are defined as effectors. Plant–insect recognition is the first and also the key step of an effective defense in plants^{27,28}. In this review, we discuss recent research advances in insect elicitors and effectors and their roles in plant–insect interactions.

Plant perceptions of insect herbivory

Plant perception of an insect attack is the first step of defense. Insect herbivory raised diverse active molecules such as damage-associated molecules, insect-derived elicitors, and the plant endogenous molecules activated by insect digestive enzymes (Figure 1). The specific and efficient recognition of these active molecules guarantees the timely priming of plant defense^{29,30}.

Plant-derived signal molecules activated by herbivory

Wounding damage caused by insect herbivory will quickly trigger plant defense signaling. The first reported damage-related peptide signal was systemin, an 18-amino acid polypeptide cleaved from prosystemin (inactive form) in tomato upon wounding stimulus³¹. Systemin promotes JA accumulation and activates the expression of genes encoding proteinase inhibitors which have insecticide activity³². Whereas systemin had been reported long before, its receptor SYR1, a leucine-rich repeat receptor kinase (LRR-RK), was identified recently. The introgression line, which lacks SYR1 expression, is highly sensitive to *Spodoptera littoralis*³³. Besides systemin, other wound-induced peptides had been identified in plants, including *Arabidopsis*, maize, and rice. The application of synthetic 23-amino acid maize Peps could mimic the *Spodoptera exigua* attack, and similar Peps were found in rice recently^{34,35}. In *Arabidopsis*, AtPeps, which is generated from PROPEPs under the catalyzation of the cysteine protease METACASPASE4 (MC4), acts as signals to trigger both JA and SA signaling pathways^{36,37}. Like the systemin-SYR1 module, the reported receptors of AtPeps—AtPEPR1 and AtPEPR2—are also classified in the LRR-RK family^{38,39}.

From *Spodoptera frugiperda* larval OSs, researchers isolated a disulfide-bridged peptide (+ICDINGVCVDA–), termed inceptin, that can induce the accumulation of defense hormones such as ethylene, JA, and SA in cowpea plants. Inceptin is the proteolytic fragment of chloroplastic ATP synthase γ -subunit of cowpea plants digested by *S. frugiperda* larvae^{40,41}. Recently, on BioRxiv, it was reported that the receptor of inceptin in plants was a leucine-rich repeat receptor-like protein, INR, which is distinguished from LRR-RKs by lacking an intracellular kinase domain⁴². These findings expand the paradigm of plant surface recognition of insect herbivory.

Elicitors secreted by insects

Besides plant signal molecules activated by insect feeding, a number of reported elicitors are derived from insects themselves and most of them belong to HAMPs⁴³. It had been reported that the OS, the oviposition and the honeydew of insects could induce a plant defense response, including the accumulation of JA and secondary metabolism⁴⁴. These insect-derived elicitors can be classified as fatty acid derivatives, enzymes, and some other proteins⁴³.

The first identified fatty acid–amino acid conjugate (FAC) elicitor was volicitin, which was isolated from *S. exigua* larval OSs. Volicitin can induce the emission of volatiles in maize to attract predators⁴⁵. After volicitin, other forms of FACs from various insect OSs had been found in succession^{46,47}. In *Nicotiana attenuate*, FACs from *Manduca sexta* activate the MAPK pathway⁴⁸. Besides FACs, califerins, the sulfoxy fatty acids that exist in OSs of grasshopper (*Schistocerca americana*) larvae, also have elicitor activity⁴⁹. Glucose oxidases (GOXs) and β -glucosidase are enzyme-like elicitors. GOX is identified from *Helicoverpa zea* and specifically activates defense response in tomato^{50,51}. The β -glucosidase in *Pieris brassicae* larval OSs triggers the emission of volatiles from wounded cabbage




Figure 1. Schematic diagram of herbivory-associated elicitors and effectors manipulating plant defense. Receptors (SYR, PEPR, INR, and LecRKs) located on the plant cell surface recognize small peptides (systemin, inceptin, and Peps) and, together with the co-receptors (SERKs/BAK1 and SOBIR1), trigger downstream defense signaling. Also, elicitors derived from insects, including FACS, β -Glu, and GOX, are able to activate plant defense with the unknown mechanisms. On the other hand, insects secrete effectors to weaken plant defenses. Some effectors interfere with jasmonate (JA) signaling directly (HARP1, 2b, C2, β C1, and SSGP-71) or indirectly (Armet and Bt56) by enhancing salicylic acid (SA) accumulation to compromise JA signaling. Some effectors (Mp1 and Me10) target plant proteins (VPS52 and TFT7) that are directly involved in defense. The DNase II eliminates the extracellular DNA which is released by damaged cells to trigger plant defense. MIF and C002 from aphids are of benefit to insects living on the host plants, but the underlying mechanisms remain elusive. Notably, some elicitors/effectors are plant-specific. Here, the GOX from *Helicoverpa zea* acts as an effector, inhibiting nicotine accumulation in tobacco, and, on the other hand, acts as an elicitor specifically inducing plant response in tomato.

leaves and this attracts predators such as parasitic wasp^{52,53}. Lipase and phospholipase C are other types of salivary enzyme-like elicitors. Lipase of *Schistocerca gregaria* OS elevates oxylipin accumulation and defense response in *Arabidopsis*⁵⁴. Phospholipase C of *Spodoptera frugiperda* induces the accumulation of proteinase inhibitors in corn⁵⁵.

The above-mentioned elicitors are from chewing insects. The elicitors from the piercing-sucking insects are isolated largely from salivary glands. The mucin-like salivary protein (NIMLP) of planthopper (*Nilaparvata lugens*) is a double-edged sword. On one hand, it contributes to the formation of salivary sheaths for successful feeding; on the other hand, it was used by plants to trigger a defense response, like Ca²⁺ mobilization, the MEK2 MAPK cascades, and JA signaling transduction, thereby reducing the performance of *N. lugens*⁵⁶. Tetranins is another characterized elicitor identified from *Tetranychus urticae*. Tetranins increases the expression of defense genes and activates JA, salicylic acid (SA), and abscisic acid biosynthesis in plant. It also promotes volatile emission to attract predatory mites⁵⁷.

Some elicitors are from endosymbionts. MAMPs could be released through herbivory OSs and recognized by plants to induce pattern-triggered immunity (PTI)^{58,59}. The chaperon GroEL from the endosymbionts *Buchnera* of potato aphids (*Macrosiphum euphorbiae*) induces oxidative bursts and PTI in *Arabidopsis*⁶⁰. From the *S. littoralis* larval OSs, the porin-like proteins most likely of bacterial origin can induce the early response of plant defense⁶¹. A recent report reveals that some elicitors are from honeydew-associated microbes in sucking arthropods⁶².

Insect effectors twist plant defense

To adapt to their host plants, insects have developed multilayered means for fitness. Besides releasing elicitors, the insect releases effectors that disturb host plant defense response for successful feeding⁶³. The reported insect effectors are identified from both the herbivory itself and insect-related microbiomes (Figure 1).

The first reported insect effector was GOX from the chewing insect, *H. zea*, which inhibits nicotine accumulation and elevates the SA-mediated PR-1a protein level in tobacco^{64,65}. Notably, the same GOX protein induces plant response in tomato^{50,51}, which we discussed in the 'Elicitors secreted by insects' section. This suggests that the same protein acts as the effector or as the elicitor depending on their interacted host plant. Another piece of evidence in support of insect effectors is that the *S. littoralis* larvae that fed on OS pretreated plants had a greater weight increase⁶⁶.

The direct interaction with JA signaling-related components is an efficient way for herbivory effectors to inhibit plant defense. In our recent work, we isolated a venom-like protein termed HARP1, which is identified from the OS of *Helicoverpa armigera*. HARP1 can interact with multiple JAZ proteins of *Arabidopsis* and cotton plants to prevent COI1-mediated JAZ degradation, thereby blocking the JA signaling output⁶⁷. SSGP-71 is an E3 ubiquitin ligase-mimicking protein in

Hessian fly (*Mayetiola destructor*). It allows the insect to hijack the plant proteasome and block the basal immunity⁶⁸. These studies fill in the gap of the working mechanism about how insects manipulate effectors to block plant defense for better adaptation.

Some insect effectors inhibit plant defense by interfering with the crosstalk between SA and JA. For example, Bt56 from the whitefly (*Bemisia tabaci*) enhanced the performance of the whitefly on tobacco by decreasing JA signaling through the antagonism between JA and SA. Bt56 could directly interact with KNOTTED 1-like homeobox transcription factor NTH202 and eliminate the negative modulation of NTH202 on SA accumulation⁶⁹. Armet, the effector of pea aphid (*Acyrtosiphon pisum*) protein, induced SA accumulation by blocking SA methylation and enhanced the pathogen resistance in plants, reflecting a novel tripartite interaction of insect-plant-pathogen^{70,71}.

The extracellular DNA and hydrogen peroxide that are released by damaged cells can trigger plant defense³⁰. Therefore, some insects secrete effectors to eliminate the production of these DAMPs. The planthopper (*Laodelphax striatellus*) secretes salivary DNase II, which acts as an effector by erasing extracellular DNA, and the *Trichoplusia ni* salivary catalase functions as an ROS scavenger to reduce hydrogen peroxide, thus inhibiting ROS burst and other plant defense responses^{72,73}.

Moreover, some effectors were reported to target other defense-related proteins in plants. A set of saliva proteins in aphids were proven to have effector activity through proteomic combined RNA sequencing (RNA-seq) analysis^{63,74-79}. A macrophage migration inhibitory factor (MIF) from pea aphid saliva inhibits immune response in *N. benthamiana* and improves aphid performance. Interestingly, the MIFs in vertebrates are also involved in the immune pathway, suggesting the highly conserved function of MIF^{80,81}. Vacuolar protein sorting-associated protein 52 (VPS52) in potato (*Solanum tuberosum*) has negative impacts on green peach aphid (*Myzus persicae*) infection. *M. persicae* saliva-secreted protein Mp1 targets the VPS52 as an effective virulence strategy^{77,82}. Me10 from *M. euphorbiae* interacts with tomato TFT7, a 14-3-3 isoform involved in aphid resistance, and enhances aphid longevity and fecundity⁸³. Some effectors can target the host cell wall. Expansin-like protein (HaEXPB2) from the nematode (*Heterodera avenae*) binds to cellulose of tobacco, thereby increasing nematode infectivity⁸⁴.

The effectors mentioned above are generated from the insect itself. Other effectors are also derived from insect-borne microbe. Although the exact effector components need to be explored, it was found that Colorado potato beetle (*Leptinotarsa decemlineata*) larvae suppress tomato defense response by exploiting bacteria in their OSs and gut^{85,86}. Besides bacteria, some active molecules from vector-borne pathogens are reported to interfere with plant defense and are of benefit for their insect vectors living on host plants⁸⁷. The phytoplasm protein SAP11 and SAP54 of aster yellows phytoplasma strain witches' broom was proposed to promote aphid colonization

and also interfere with plant development^{88–90}. The β C1 of tomato yellow leaf curl China virus directly interacts with MYC2 protein to decrease the MYC2-regulated terpene synthase, thereby reducing plant resistance to the whitefly⁹¹. The 2b protein of the aphid-borne cucumber mosaic virus (CMV) stabilizes JAZ proteins by direct interaction, thus blocking JA signaling output, and this benefits aphid (*M. persicae*) performance on the host plant⁹². The C2 protein of tomato yellow leaf curl virus can also compromise JA signaling in tobacco by interacting with plant ubiquitin to block JAZ1 protein degradation, thereby reducing plant resistance to the insect vector whitefly⁹³. These studies reveal the intricate interaction of plant–virus–insect vector. In Table 1, we summarize the reported insect-associated elicitors and effectors from different species and their probable roles.

Prospects

JA is a conserved defense regulator in the plant kingdom. On one hand, various elicitors can be recognized by plants to trigger JA signaling. On the other hand, the JA pathway tends to be targeted by a diverse range of attackers for fitness (Figure 1). Some insect effectors have a mechanism similar to that of the virus proteins in blocking JA signaling^{67,91–93}. It would be interesting to study whether there are relationships between the phylogeny of insect effectors and viral proteins. Although numerous elicitors and effectors were identified, their target proteins, the underlying mechanisms, and the transportation mechanisms of the effectors entering plant cells are largely unknown and deserve further investigation. In plants, JA is integrated with multiple signaling to

Table 1. Herbivory-associated elicitors and effectors.

	Name	Origin	Biofunction	References	
Elicitors	Plant-derived	Systemin	Wounded tomato plants	Perceived by SYR1, induce accumulation of proteinase inhibitor and ethylene, and induce oxidative bursts	31,33
		PEPs	Wounded plants (<i>Arabidopsis</i> , maize, rice)	Induce defensin and burst of hydrogen peroxide (H ₂ O ₂) after perceiving by PEPRs	36,38,39
		Inceptin	Degradation of cowpea ATP synthase by <i>Spodoptera frugiperda</i> during herbivory	Increase the concentration of JA and SA by interacting with INR	40,42
	Derived from insect	Volicitin	<i>Spodoptera exigua</i>	Induce volatiles emission in corn	45
		Caeliferins	<i>Schistocerca americana</i>		49
		GOX	<i>Helicoverpa zea</i> , <i>Ostrinia nubilalis</i>	Specifically promote defense response in tomato	50,51
		β -glucosidase	<i>Pieris brassicae</i>	Increase volatile emission in cabbage	52
		Lipase	<i>Schistocerca gregaria</i>	Elevate the oxylipins accumulation in <i>Arabidopsis</i>	54
		Phospholipase C	<i>S. frugiperda</i>	Trigger proteinase inhibitors accumulation in corn	55
		Bruchins	<i>Bruchus pisorum</i>	Induce neoplasms formation beneath the insect egg in pea	94,95
		NIMLP	<i>Nilaparvata lugens</i>	Induce plant defense response in rice	56
		Tetranins	<i>Tetranychus urticae</i>	Cytosolic calcium influx and membrane depolarization induce biosynthesis of JA. SA and ABA in kidney bean	57
		GroEL	<i>Buchnera</i> in <i>Macrosiphum euphorbiae</i>	Induce PTI and ROS accumulation in <i>Arabidopsis</i>	60
		Porin-like proteins	<i>Bacteria</i> in <i>Spodoptera littoralis</i>	Trigger membrane potential changes and cytosolic Ca ²⁺ elevations in <i>Arabidopsis</i> and <i>Vicia faba</i>	61
		Unidentified	Gut-associated bacteria in <i>H. zea</i>	Increase salivary GOX to induce defense in tomato	96,97
Unidentified	Honeydew-associated microbes <i>N. lugens</i>	Induce accumulation of phytoalexins and volatile emission in rice	62,98		

	Name	Origin	Biofunction	References	
Effectors	GOX	<i>H. zea</i>	Decrease nicotine accumulation in tobacco	64	
	HARP1	<i>Helicoverpa armigera</i>	Interact with and stabilize JAZs, depress JA signaling in <i>Arabidopsis</i>	67	
	SSGP-71	<i>Mayetiola destructor</i>	Interact with Skp, decrease plant proteasome activity, thus block hormone signaling in wheat	68	
	Bt56	<i>Bemisia tabaci</i>	Interact with NTH202 to increase SA biosynthesis, thus decrease JA response in tobacco	69	
	Armet	<i>Acyrtosiphon pisum</i>	Help feeding of insect, induce SA accumulation and pathogen response in <i>N. benthamiana</i> and <i>Medicago truncatula</i>	70,71	
	DNase II	<i>Laodelphax striatellus</i>	Erase extracellular DNA released by damaged cell in rice	72	
	Catalase	<i>Trichoplusia ni</i>	Reduce H ₂ O ₂ in tomato	73	
	C002	<i>A. pisum, Myzus persicae</i>	ApC002 and MpC002 help insect foraging and feeding on fava bean and <i>N. benthamiana</i> , respectively	63,74	
	MIF	<i>A. pisum, M. persicae</i>	Improve aphid performance, inhibit immune response in <i>N. benthamiana</i>	80	
	Mp1	<i>M. persicae</i>	Interact with VPS52 to relocate to vesicle-like structures and enhance insect virulence in <i>Arabidopsis</i> and potato	82	
	Me10	<i>M. euphorbiae</i>	Interact with TFT7, enhance the longevity and fecundity on tomato	83	
	Mp42, Mp55 Me23	<i>M. persicae, M. euphorbiae</i>	Increase aphid reproduction, suppress <i>N. benthamiana</i> defenses	99,100	
	HaEXPB2	<i>Heterodera avenae</i>	Bind to cellulose and target cell wall when parasitizing <i>N. benthamiana</i>	84	
	Phosphatase 2C	<i>M. destructor</i>	Interfere with the wheat signal transduction pathway possibly by phosphatase ability	101	
	Endo-beta-1,4-Glucanase (NIEG1)	<i>N. lugens</i>	Degrade celluloses in plant cell wall, enable insect stylet to reach the rice phloem	102	
	NcSP75	<i>Nephotettix cincticeps</i>	Help successful ingestion from sieve elements of rice	103	
	NcSP84	<i>N. cincticeps</i>	Suppress accumulation of Ca ²⁺ and H ₂ O ₂ and sieve element clogging in rice	104	
	NISEF1	<i>N. lugens</i>	Help successful ingestion from sieve elements of rice	105	
	Derived from insect-borne microbe	Unidentified	Gut and oral secretion-associated bacteria in Colorado potato beetle	Suppress tomato defense response Bind and destabilize TCPs, reduce plant defense in <i>Arabidopsis</i>	85,86
		SAP11	Aster yellows witches' broom in <i>Macrosteles quadrilineatus</i>		106
SAP54		Aster yellows witches' broom in <i>M. quadrilineatus</i>	Degrade MTFs through interacting with RAD23, influence floral development in <i>Arabidopsis</i>	90	
βC1		Tomato yellow leaf curl China virus in <i>B. tabaci</i>	Interact with MYC2 and suppress MYC2-regulated terpene synthesis in <i>Arabidopsis</i>	91	
2b		Cucumber mosaic virus (CMV) in <i>M. persicae</i>	Interact with and stable JAZ protein, blocking JA signaling in <i>Arabidopsis</i>	92	
C2		Tomato yellow leaf curl virus in <i>B. tabaci</i>	Interact with plant ubiquitin, blocking JA signaling in tobacco	93	

form a complex and flexible defense network. Recent research has revealed the intricate defense network shaped by insect herbivory^{69,107–109}. Studies have also shown that insects can use plant defense metabolites to find their host plants and to fend off

predators^{110,111}; this gives new insight into plant–insect interactions. Further investigations will greatly enrich our knowledge of the complex and flexible interactions between plants and insects and will also be helpful for breeding insect-proof crops^{112,113}.

References



- Wu J, Baldwin IT: **New insights into plant responses to the attack from insect herbivores.** *Annu Rev Genet.* 2010; **44**: 1–24.
[PubMed Abstract](#) | [Publisher Full Text](#)
- F** Li J, Liu X, Wang Q, *et al.*: **A Group D MAPK Protects Plants from Autotoxicity by Suppressing Herbivore-Induced Defense Signaling.** *Plant Physiol.* 2019; **179**(4): 1386–401.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Ye M, Glauser G, Lou Y, *et al.*: **Molecular Dissection of Early Defense Signaling Underlying Volatile-Mediated Defense Regulation and Herbivore Resistance in Rice.** *Plant Cell.* 2019; **31**(3): 687–98.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Zebelo SA, Maffei ME: **Role of early signalling events in plant-insect interactions.** *J Exp Bot.* 2015; **66**(2): 435–48.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Erb M, Meldau S, Howe GA: **Role of phytohormones in insect-specific plant reactions.** *Trends Plant Sci.* 2012; **17**(5): 250–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Howe GA, Major IT, Koo AJ: **Modularity in Jasmonate Signaling for Multistress Resilience.** *Annu Rev Plant Biol.* 2018; **69**: 387–415.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Howe GA: **Plant hormones: Metabolic end run to jasmonate.** *Nat Chem Biol.* 2018; **14**(2): 109–10.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Huang H, Liu B, Liu L, *et al.*: **Jasmonate action in plant growth and development.** *J Exp Bot.* 2017; **68**(6): 1349–59.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Browse J: **Jasmonate passes muster: a receptor and targets for the defense hormone.** *Annu Rev Plant Biol.* 2009; **60**: 183–205.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Smirnova E, Marquis V, Poirier L, *et al.*: **Jasmonic Acid Oxidase 2 Hydroxylates Jasmonic Acid and Represses Basal Defense and Resistance Responses against *Botrytis cinerea* Infection.** *Mol Plant.* 2017; **10**(9): 1159–73.
[PubMed Abstract](#) | [Publisher Full Text](#)
- F** Caarls L, Elberse J, Awwanah M, *et al.*: ***Arabidopsis* JASMONATE-INDUCED OXYGENASES down-regulate plant immunity by hydroxylation and inactivation of the hormone jasmonic acid.** *Proc Natl Acad Sci U S A.* 2017; **114**(24): 6388–93.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Chini A, Gimenez-Ibanez S, Goossens A, *et al.*: **Redundancy and specificity in jasmonate signalling.** *Curr Opin Plant Biol.* 2016; **33**: 147–56.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Xie DX, Feys BF, James S, *et al.*: ***COI1*: an *Arabidopsis* gene required for jasmonate-regulated defense and fertility.** *Science.* 1998; **280**(5366): 1091–4.
[PubMed Abstract](#) | [Publisher Full Text](#)
- F** Sheard LB, Tan X, Mao H, *et al.*: **Jasmonate perception by inositol-phosphate-potentiated *COI1*-JAZ co-receptor.** *Nature.* 2010; **468**(7322): 400–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- F** Chini A, Fonseca S, Fernández G, *et al.*: **The JAZ family of repressors is the missing link in jasmonate signalling.** *Nature.* 2007; **448**(7154): 666–71.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- F** Thines B, Katsir L, Melotto M, *et al.*: **JAZ repressor proteins are targets of the SCF^{COI1} complex during jasmonate signalling.** *Nature.* 2007; **448**(7154): 661–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- F** Monte I, Ishida S, Zamarréno AM, *et al.*: **Ligand-receptor co-evolution shaped the jasmonate pathway in land plants.** *Nat Chem Biol.* 2018; **14**(5): 480–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- An C, Li L, Zhai Q, *et al.*: **Mediator subunit MED25 links the jasmonate receptor to transcriptionally active chromatin.** *Proc Natl Acad Sci U S A.* 2017; **114**(42): E8930–E8939.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- F** Li Q, Zheng J, Li S, *et al.*: **Transporter-Mediated Nuclear Entry of Jasmonoyl-Isoleucine Is Essential for Jasmonate Signaling.** *Mol Plant.* 2017; **10**(5): 695–708.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- F** Guo Q, Yoshida Y, Major IT, *et al.*: **JAZ repressors of metabolic defense promote growth and reproductive fitness in *Arabidopsis*.** *Proc Natl Acad Sci U S A.* 2018; **115**(45): E10768–E10777.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- F** Guo Q, Major IT, Howe GA: **Resolution of growth-defense conflict: mechanistic insights from jasmonate signaling.** *Curr Opin Plant Biol.* 2018; **44**: 72–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- Hou X, Lee LY, Xia K, *et al.*: **DELLAs modulate jasmonate signaling via competitive binding to JAZs.** *Dev Cell.* 2010; **19**(6): 884–94.
[PubMed Abstract](#) | [Publisher Full Text](#)
- F** Yang DL, Yao J, Mei CS, *et al.*: **Plant hormone jasmonate prioritizes defense over growth by interfering with gibberellin signaling cascade.** *Proc Natl Acad Sci U S A.* 2012; **109**(19): E1192–200.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | **F1000 Recommendation**
- Hong GJ, Xue XY, Mao YB, *et al.*: ***Arabidopsis* MYC2 interacts with DELLA proteins in regulating sesquiterpene synthase gene expression.** *Plant Cell.* 2012; **24**(6): 2635–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wild M, Davière JM, Cheminant S, *et al.*: **The *Arabidopsis* DELLA RGA-LIKE3 is a direct target of MYC2 and modulates jasmonate signaling responses.** *Plant Cell.* 2012; **24**(8): 3307–19.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mao YB, Liu YQ, Chen DY, *et al.*: **Jasmonate response decay and defense metabolite accumulation contributes to age-regulated dynamics of plant insect resistance.** *Nat Commun.* 2017; **8**: 13925.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Howe GA, Jander G: **Plant immunity to insect herbivores.** *Annu Rev Plant Biol.* 2008; **59**: 41–66.
[PubMed Abstract](#) | [Publisher Full Text](#)
- F** Aljborg Z, Chen MS: **Indirect plant defense against insect herbivores: a review.** *Insect Sci.* 2018; **25**(1): 2–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- F** Erb M, Reymond P: **Molecular Interactions Between Plants and Insect Herbivores.** *Annu Rev Plant Biol.* 2019; **70**: 527–57.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- Gust AA, Pruitt R, Nürnberger T: **Sensing Danger: Key to Activating Plant Immunity.** *Trends Plant Sci.* 2017; **22**(9): 779–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pearce G, Strydom D, Johnson S, *et al.*: **A polypeptide from tomato leaves induces wound-inducible proteinase inhibitor proteins.** *Science.* 1991; **253**(5022): 895–7.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Orozco-Cardenas M, McGurl B, Ryan CA, *et al.*: **Expression of an antisense prosystemin gene in tomato plants reduces resistance toward *Manduca sexta* larvae.** *Proc Natl Acad Sci U S A.* 1993; **90**(17): 8273–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- F** Wang L, Einig E, Almeida-Trapp M, *et al.*: **The systemin receptor SYR1 enhances resistance of tomato against herbivorous insects.** *Nat Plants.* 2018; **4**(3): 152–6.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- F** Shinya T, Yasuda S, Hyodo K, *et al.*: **Integration of danger peptide signals with herbivore-associated molecular pattern signaling amplifies anti-herbivore defense responses in rice.** *Plant J.* 2018; **94**(4): 626–37.
[PubMed Abstract](#) | [Publisher Full Text](#) | **F1000 Recommendation**
- Huffaker A, Dafoe NJ, Schmelz EA: **ZmPep1, an ortholog of *Arabidopsis* elicitor peptide 1, regulates maize innate immunity and enhances disease resistance.**

- Plant Physiol.* 2011; **155**(3): 1325–38.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
36. Huffaker A, Pearce G, Ryan CA: **An endogenous peptide signal in *Arabidopsis* activates components of the innate immune response.** *Proc Natl Acad Sci U S A.* 2006; **103**(26): 10098–103.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
37. **F** Hander T, Fernández-Fernández ÁD, Kumpf RP, *et al.*: **Damage on plants activates Ca²⁺-dependent metacaspases for release of immunomodulatory peptides.** *Science.* 2019; **363**(6433): pii: eaar7486.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
38. Yamaguchi Y, Pearce G, Ryan CA: **The cell surface leucine-rich repeat receptor for AtPep1, an endogenous peptide elicitor in *Arabidopsis*, is functional in transgenic tobacco cells.** *Proc Natl Acad Sci U S A.* 2006; **103**(26): 10104–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
39. **F** Yamaguchi Y, Huffaker A, Bryan AC, *et al.*: **PEPR2 is a second receptor for the Pep1 and Pep2 peptides and contributes to defense responses in *Arabidopsis*.** *Plant Cell.* 2010; **22**(2): 508–22.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
40. **F** Schmelz EA, Carroll MJ, LeClere S, *et al.*: **Fragments of ATP synthase mediate plant perception of insect attack.** *Proc Natl Acad Sci U S A.* 2006; **103**(23): 8894–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
41. Schmelz EA, LeClere S, Carroll MJ, *et al.*: **Cowpea chloroplastic ATP synthase is the source of multiple plant defense elicitors during insect herbivory.** *Plant Physiol.* 2007; **144**(2): 793–805.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
42. Steinbrenner AD, Muñoz-Amatrián M, Aguilar Venegas JM, *et al.*: **A receptor for herbivore-associated molecular patterns mediates plant immunity.** *bioRxiv.* 2019; 679803.
[Publisher Full Text](#)
43. Bonaventure G, VanDoorn A, Baldwin IT: **Herbivore-associated elicitors: FAC signaling and metabolism.** *Trends Plant Sci.* 2011; **16**(6): 294–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
44. Stam JM, Kroes A, Li Y, *et al.*: **Plant interactions with multiple insect herbivores: from community to genes.** *Annu Rev Plant Biol.* 2014; **65**: 689–713.
[PubMed Abstract](#) | [Publisher Full Text](#)
45. Alborn HT, Turlings TCJ, Jones TH, *et al.*: **An elicitor of plant volatiles from beet armyworm oral secretion.** *Science.* 1997; **276**(5314): 945–9.
[Publisher Full Text](#)
46. Halitschke R, Schittko U, Pohnert G, *et al.*: **Molecular interactions between the specialist herbivore *Manduca sexta* (Lepidoptera, Sphingidae) and its natural host *Nicotiana attenuata*. III. Fatty acid-amino acid conjugates in herbivore oral secretions are necessary and sufficient for herbivore-specific plant responses.** *Plant Physiol.* 2001; **125**(2): 711–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
47. Yoshinaga N, Aboshi T, Ishikawa C, *et al.*: **Fatty acid amides, previously identified in caterpillars, found in the cricket *Teleogryllus taiwanemma* and fruit fly *Drosophila melanogaster* larvae.** *J Chem Ecol.* 2007; **33**(7): 1376–81.
[PubMed Abstract](#) | [Publisher Full Text](#)
48. **F** Wu J, Hettenhausen C, Meldau S, *et al.*: **Herbivory rapidly activates MAPK signaling in attacked and unattacked leaf regions but not between leaves of *Nicotiana attenuata*.** *Plant Cell.* 2007; **19**(3): 1096–122.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
49. Alborn HT, Hansen TV, Jones TH, *et al.*: **Disulfoxy fatty acids from the American bird grasshopper *Schistocerca americana*, elicitors of plant volatiles.** *Proc Natl Acad Sci U S A.* 2007; **104**(32): 12976–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
50. Tian D, Peiffer M, Shoemaker E, *et al.*: **Salivary glucose oxidase from caterpillars mediates the induction of rapid and delayed-induced defenses in the tomato plant.** *PLoS One.* 2012; **7**(4): e36168.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
51. Louis J, Peiffer M, Ray S, *et al.*: **Host-specific salivary elicitor(s) of European corn borer induce defenses in tomato and maize.** *New Phytol.* 2013; **199**(1): 66–73.
[PubMed Abstract](#) | [Publisher Full Text](#)
52. Mattiacci L, Dicke M, Posthumus MA: **beta-Glucosidase: an elicitor of herbivore-induced plant odor that attracts host-searching parasitic wasps.** *Proc Natl Acad Sci U S A.* 1995; **92**(6): 2036–40.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
53. Hopke J, Donath J, Bleichert S, *et al.*: **Herbivore-induced volatiles: the emission of acyclic homoterpenes from leaves of *Phaseolus lunatus* and *Zea mays* can be triggered by a beta-glucosidase and jasmonic acid.** *FEBS Lett.* 1994; **352**(2): 146–50.
[PubMed Abstract](#) | [Publisher Full Text](#)
54. Schäfer M, Fischer C, Meldau S, *et al.*: **Lipase activity in insect oral secretions mediates defense responses in *Arabidopsis*.** *Plant Physiol.* 2011; **156**(3): 1520–34.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
55. **F** Acevedo FE, Peiffer M, Ray S, *et al.*: **Intraspecific differences in plant defense induction by fall armyworm strains.** *New Phytol.* 2018; **218**(1): 310–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
56. **F** Shangguan X, Zhang J, Liu B, *et al.*: **A Mucin-Like Protein of Planthopper Is Required for Feeding and Induces Immunity Response in Plants.** *Plant Physiol.* 2018; **176**(1): 552–65.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
57. **F** Iida J, Desaki Y, Hata K, *et al.*: **Tetranins: new putative spider mite elicitors of host plant defense.** *New Phytol.* 2019; **224**(2): 875–85.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
58. Kaloshian I: **Gene-for-gene disease resistance: bridging insect pest and pathogen defense.** *J Chem Ecol.* 2004; **30**(12): 2419–38.
[PubMed Abstract](#) | [Publisher Full Text](#)
59. **F** Noman A, Aqeel M, Qasim M, *et al.*: **Plant-insect-microbe interaction: A love triangle between enemies in ecosystem.** *Sci Total Environ.* 2020; **699**: 134181.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
60. Chaudhary R, Atamian HS, Shen Z, *et al.*: **GroEL from the endosymbiont *Buchnera aphidicola* betrays the aphid by triggering plant defense.** *Proc Natl Acad Sci U S A.* 2014; **111**(24): 8919–24.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
61. Guo H, Wielsch N, Hafke JB, *et al.*: **A porin-like protein from oral secretions of *Spodoptera littoralis* larvae induces defense-related early events in plant leaves.** *Insect Biochem Mol Biol.* 2013; **43**(9): 849–58.
[PubMed Abstract](#) | [Publisher Full Text](#)
62. **F** Wari D, Kabir MA, Mujiono K, *et al.*: **Honeydew-associated microbes elicit defense responses against brown planthopper in rice.** *J Exp Bot.* 2019; **70**(5): 1683–96.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
63. Bos JIB, Prince D, Pitino M, *et al.*: **A functional genomics approach identifies candidate effectors from the aphid species *Myzus persicae* (green peach aphid).** *PLoS Genet.* 2010; **6**(11): e1001216.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
64. Musser RO, Hum-Musser SM, Eichenseer H, *et al.*: **Herbivory: caterpillar saliva beats plant defences.** *Nature.* 2002; **416**(6881): 599–600.
[PubMed Abstract](#) | [Publisher Full Text](#)
65. Musser RO, Cipollini DF, Hum-Musser SM, *et al.*: **Evidence that the caterpillar salivary enzyme glucose oxidase provides herbivore offense in solanaceous plants.** *Arch Insect Biochem Physiol.* 2005; **58**(2): 128–37.
[PubMed Abstract](#) | [Publisher Full Text](#)
66. Consales F, Schweizer F, Erb M, *et al.*: **Insect oral secretions suppress wound-induced responses in *Arabidopsis*.** *J Exp Bot.* 2012; **63**(2): 727–37.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
67. Chen CY, Liu YQ, Song WM, *et al.*: **An effector from cotton bollworm oral secretion impairs host plant defense signaling.** *Proc Natl Acad Sci U S A.* 2019; **116**(28): 14331–8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
68. Zhao C, Escalante LN, Chen H, *et al.*: **A massive expansion of effector genes underlies gall-formation in the wheat pest *Mayetiola destructor*.** *Curr Biol.* 2015; **25**(5): 613–20.
[PubMed Abstract](#) | [Publisher Full Text](#)
69. **F** Xu HX, Qian LX, Wang XW, *et al.*: **A salivary effector enables whitefly to feed on host plants by eliciting salicylic acid-signaling pathway.** *Proc Natl Acad Sci U S A.* 2019; **116**(2): 490–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
70. **F** Cui N, Lu H, Wang T, *et al.*: **Armet, an aphid effector protein, induces pathogen resistance in plants by promoting the accumulation of salicylic acid.** *Philos Trans R Soc Lond B Biol Sci.* 2019; **374**(1767): 20180314.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
71. Wang W, Dai H, Zhang Y, *et al.*: **Armet is an effector protein mediating aphid-plant interactions.** *FASEB J.* 2015; **29**(5): 2032–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
72. **F** Huang HJ, Cui JR, Xia X, *et al.*: **Salivary DNase II from *Laodelphax striatellus* acts as an effector that suppresses plant defence.** *New Phytol.* 2019; **224**(2): 860–74.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
73. **F** Rivera-Vega LJ, Stanley BA, Stanley A, *et al.*: **Proteomic analysis of labial saliva of the generalist cabbage looper (*Trichoplusia ni*) and its role in interactions with host plants.** *J Insect Physiol.* 2018; **107**: 97–103.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
74. Mutti NS, Louis J, Pappan LK, *et al.*: **A protein from the salivary glands of the pea aphid, *Acyrtosiphon pisum*, is essential in feeding on a host plant.** *Proc Natl Acad Sci U S A.* 2008; **105**(29): 9965–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
75. Mutti NS, Park Y, Reese JC, *et al.*: **RNAi knockdown of a salivary transcript leading to lethality in the pea aphid, *Acyrtosiphon pisum*.** *J Insect Sci.* 2006; **6**: 1–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
76. **F** Pitino M, Coleman AD, Maffei ME, *et al.*: **Silencing of aphid genes by dsRNA feeding from plants.** *PLoS One.* 2011; **6**(10): e25709.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
77. Pitino M, Hogenhout SA: **Aphid protein effectors promote aphid colonization in a plant species-specific manner.** *Mol Plant Microbe Interact.* 2013; **26**(1): 130–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
78. Ollivier M, Legeai F, Rispe C: **Comparative analysis of the *Acyrtosiphon pisum***

- genome and expressed sequence tag-based gene sets from other aphid species. *Insect Mol Biol.* 2010; 19(Suppl 2): 33–45.
[PubMed Abstract](#) | [Publisher Full Text](#)
79. Ji R, Wang Y, Cheng Y, *et al.*: Transcriptome Analysis of Green Peach Aphid (*Myzus persicae*): Insight into Developmental Regulation and Inter-Species Divergence. *Front Plant Sci.* 2016; 7: 1562.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
80. Naessens E, Dubreuil G, Giordanengo P, *et al.*: A Secreted MIF Cytokine Enables Aphid Feeding and Represses Plant Immune Responses. *Curr Biol.* 2015; 25(14): 1898–903.
[PubMed Abstract](#) | [Publisher Full Text](#)
81. Calandra T, Roger T: Macrophage migration inhibitory factor: a regulator of innate immunity. *Nat Rev Immunol.* 2003; 3(10): 791–800.
[PubMed Abstract](#) | [Publisher Full Text](#)
82. **F** Rodriguez PA, Escudero-Martinez C, Bos JI: An Aphid Effector Targets Trafficking Protein VPS52 in a Host-Specific Manner to Promote Virulence. *Plant Physiol.* 2017; 173(3): 1892–903.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
83. **F** Chaudhary R, Peng HC, He J, *et al.*: Aphid effector Me10 interacts with tomato TFF7, a 14-3-3 isoform involved in aphid resistance. *New Phytol.* 2019; 221(3): 1518–28.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
84. Liu J, Peng H, Cui J, *et al.*: Molecular Characterization of A Novel Effector Expansin-like Protein from *Heterodera avenae* that Induces Cell Death in *Nicotiana benthamiana*. *Sci Rep.* 2016; 6: 35677.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
85. **F** Chung SH, Rosa C, Scully ED, *et al.*: Herbivore exploits orally secreted bacteria to suppress plant defenses. *Proc Natl Acad Sci U S A.* 2013; 110(39): 15728–33.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
86. **F** Chung SH, Scully ED, Peiffer M, *et al.*: Host plant species determines symbiotic bacterial community mediating suppression of plant defenses. *Sci Rep.* 2017; 7: 39690.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
87. **F** Jiang Y, Zhang CX, Chen R, *et al.*: Challenging battles of plants with phloem-feeding insects and prokaryotic pathogens. *Proc Natl Acad Sci U S A.* 2019; 116(47): 23390–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
88. **F** Pecher P, Moro G, Canale MC, *et al.*: Phytoplasma SAP11 effector destabilization of TCP transcription factors differentially impact development and defence of *Arabidopsis* versus maize. *PLoS Pathog.* 2019; 15(9): e1008035.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
89. **F** Sugio A, Kingdom HN, MacLean AM, *et al.*: Phytoplasma protein effector SAP11 enhances insect vector reproduction by manipulating plant development and defense hormone biosynthesis. *Proc Natl Acad Sci U S A.* 2011; 108(48): E1254–63.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
90. **F** MacLean AM, Orlovskis Z, Kowitwanich K, *et al.*: Phytoplasma effector SAP54 hijacks plant reproduction by degrading MADS-box proteins and promotes insect colonization in a RAD23-dependent manner. *PLoS Biol.* 2014; 12(4): e1001835.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
91. Li R, Weldegergis BT, Li J, *et al.*: Virulence factors of geminivirus interact with MYC2 to subvert plant resistance and promote vector performance. *Plant Cell.* 2014; 26(12): 4991–5008.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
92. **F** Wu D, Qi T, Li WX, *et al.*: Viral effector protein manipulates host hormone signaling to attract insect vectors. *Cell Res.* 2017; 27(3): 402–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
93. **F** Li P, Liu C, Deng WH, *et al.*: Plant begomoviruses subvert ubiquitination to suppress plant defenses against insect vectors. *PLoS Pathog.* 2019; 15(2): e1007607.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
94. Hilker M, Meiners T: Early herbivore alert: insect eggs induce plant defense. *J Chem Ecol.* 2006; 32(7): 1379–97.
[PubMed Abstract](#) | [Publisher Full Text](#)
95. Hilker M, Fatouros NE: Resisting the onset of herbivore attack: plants perceive and respond to insect eggs. *Curr Opin Plant Biol.* 2016; 32: 9–16.
[PubMed Abstract](#) | [Publisher Full Text](#)
96. Wang J, Peiffer M, Hoover K, *et al.*: *Helicoverpa zea* gut-associated bacteria indirectly induce defenses in tomato by triggering a salivary elicitor(s). *New Phytol.* 2017; 214(3): 1294–306.
[PubMed Abstract](#) | [Publisher Full Text](#)
97. **F** Wang J, Yang M, Song Y, *et al.*: Gut-Associated Bacteria of *Helicoverpa zea* Indirectly Trigger Plant Defenses in Maize. *J Chem Ecol.* 2018; 44(7–8): 690–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
98. **F** Wari D, Alamgir KM, Mujiono K, *et al.*: Brown planthopper honeydew-associated symbiotic microbes elicit momilactones in rice. *Plant Signal Behav.* 2019; 14(11): 1655335.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
99. Eizinga DA, de Vos M, Jander G: Suppression of plant defenses by a *Myzus persicae* (green peach aphid) salivary effector protein. *Mol Plant Microbe Interact.* 2014; 27(7): 747–56.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
100. Atamian HS, Chaudhary R, Cin VD, *et al.*: In planta expression or delivery of potato aphid *Macrosiphum euphorbiae* effectors Me10 and Me23 enhances aphid fecundity. *Mol Plant Microbe Interact.* 2013; 26(1): 67–74.
[PubMed Abstract](#) | [Publisher Full Text](#)
101. Zhao C, Shukle R, Navarro-Escalante L, *et al.*: *Avirulence* gene mapping in the Hessian fly (*Mayetiola destructor*) reveals a protein phosphatase 2C effector gene family. *J Insect Physiol.* 2016; 84: 22–31.
[PubMed Abstract](#) | [Publisher Full Text](#)
102. Ji R, Ye W, Chen H, *et al.*: A Salivary Endo- β -1,4-Glucanase Acts as an Effector That Enables the Brown Planthopper to Feed on Rice. *Plant Physiol.* 2017; 173(3): 1920–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
103. **F** Matsumoto Y, Hattori M: The green rice leafhopper, *Nephotettix cincticeps* (Hemiptera: Cicadellidae), salivary protein NcSP75 is a key effector for successful phloem ingestion. *PLoS One.* 2018; 13(9): e0202492.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
104. Hattori M, Nakamura M, Komatsu S, *et al.*: Molecular cloning of a novel calcium-binding protein in the secreted saliva of the green rice leafhopper *Nephotettix cincticeps*. *Insect Biochem Mol Biol.* 2012; 42(1): 1–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
105. Ye W, Yu H, Jian Y, *et al.*: A salivary EF-hand calcium-binding protein of the brown planthopper *Nilaparvata lugens* functions as an effector for defense responses in rice. *Sci Rep.* 2017; 7: 40498.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
106. Sugio A, MacLean AM, Hogenhout SA: The small phytoplasma virulence effector SAP11 contains distinct domains required for nuclear targeting and CIN-TCP binding and destabilization. *New Phytol.* 2014; 202(3): 838–48.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
107. **F** Ma F, Yang X, Shi Z, *et al.*: Novel crosstalk between ethylene- and jasmonic acid-pathway responses to a piercing-sucking insect in rice. *New Phytol.* 2019; 225(1): 474–87.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
108. Liu L, Sonbol FM, Huot B, *et al.*: Salicylic acid receptors activate jasmonic acid signalling through a non-canonical pathway to promote effector-triggered immunity. *Nat Commun.* 2016; 7: 13099.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
109. Li R, Llorca LC, Schuman MC, *et al.*: ZEITLUPE in the Roots of Wild Tobacco Regulates Jasmonate-Mediated Nicotine Biosynthesis and Resistance to a Generalist Herbivore. *Plant Physiol.* 2018; 177(2): 833–46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
110. **F** Ye M, Veyrat N, Xu H, *et al.*: An herbivore-induced plant volatile reduces parasitoid attraction by changing the smell of caterpillars. *Sci Adv.* 2018; 4(5): eaar4767.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#) | [F1000 Recommendation](#)
111. **F** Hu L, Mateo P, Ye M, *et al.*: Plant iron acquisition strategy exploited by an insect herbivore. *Science.* 2018; 361(6403): 694–7.
[PubMed Abstract](#) | [Publisher Full Text](#) | [F1000 Recommendation](#)
112. Zhao Y, Huang J, Wang Z, *et al.*: Allelic diversity in an NLR gene *BPH9* enables rice to combat planthopper variation. *Proc Natl Acad Sci U S A.* 2016; 113(45): 12850–5.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
113. Liu Y, Wu H, Chen H, *et al.*: A gene cluster encoding lectin receptor kinases confers broad-spectrum and durable insect resistance in rice. *Nat Biotechnol.* 2015; 33(3): 301–5.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Peer Review Status:  

Editorial Note on the Review Process

F1000 Faculty Reviews are written by members of the prestigious F1000 Faculty. They are commissioned and are peer reviewed before publication to ensure that the final, published version is comprehensive and accessible. The reviewers who approved the final version are listed with their names and affiliations.

The reviewers who approved this article are:

Version 1

- Gregg Howe**
MSU-DOE Plant Research Laboratory, Michigan State University, East Lansing, MI, 48824, USA
Competing Interests: No competing interests were disclosed.
- Gary Felton**
Department of Entomology, Pennsylvania State University, University Park, PA, USA
Competing Interests: No competing interests were disclosed.

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research