

A Systematic Review on methods of evaluate sentence production deficits in agrammatic aphasia patients: Validity and Reliability issues

Azar Mehri¹, Shohreh Jalaie²

¹Speech Therapy Department, Faculty of Rehabilitation, Tehran University of Medical Sciences, ²PhD of Biostatistics, Physiotherapy Department, Faculty of Rehabilitation, Tehran University of Medical Sciences, Tehran, Iran

Background: The grammar assessment in aphasia has been done by few standard tests, but today these tests cannot precise evaluate the sentence production in agrammatic patients. In this study, we review structures and contents of tests or tasks designed to find more frequent methods for sentence production ability in aphasia patients. **Materials and Methods:** We searched the Cochrane library, Medline by PubMed, Science Direct, Scopus, and Google Scholar from 1980 to October 1, 2013 and evaluated all of exist tests or tasks included in the articles and systematic reviews. The sentence production has been studied in three methods. It contains the use of sentence production in spontaneous speech, tasks designed and both methods. The quality of studies was assessed using Critical Appraisal Skills Program. **Results:** The 160 articles were reviewed and 38 articles were studied according to inclusion and exclusion criteria. They were classified into three categories based on assessment methods of sentence production. In 39.5% studies, researchers have used tasks designed, 7.9% articles have applied spontaneous speech and 52.6% articles have used both methods for evaluation production. Inter-rater reliability was between 90% and 100% and intra-rater reliability was between 96% and 98% in studied. **Conclusion:** Agrammatic aphasia has syntax disorders, especially in sentence production. Most researchers and clinicians used both methods for evaluation production.

Key words: Agrammatism, sentence production, syntax, validity, reliability

How to cite this article: Mehri A, Jalaie Sh. A Systematic Review on methods of evaluate sentence production deficits in agrammatic aphasia patients: Validity and Reliability issues. *J Res Med Sci* 2014;19:885-98.

INTRODUCTION

Aphasia is an acquired communication disorder caused by brain damage and it result in impairments in speaking, listening, reading, and writing.^[1] According to the National Institute for Deafness and Other Communication Disorders (1999), about 1 million people in the United States currently have aphasia.^[1]

In aphasiology, “grammar” means the sentence structure, such as verb inflection as well as the relationship between sentence constituents. Agrammatism refers to pattern of sentence production that reflects an absence of grammatical structure.^[2] Individuals with agrammatism show difficulty producing closed-class words that is, free-standing grammatical morphemes such as preposition, articles, pronouns, and auxiliary verbs.^[2] Traditionally, it was argued that Broca’s aphasia has a grammatical deficit and Wernicke’s aphasia has a lexical semantic deficit. Furthermore, speakers with Broca’s aphasia have more severe deficit in verb retrieval.^[3] Because various kinds of sentence production may be involved in these patients, an in-depth assessment of the impairment is imperative.^[2]

For the assessment of agrammatism in aphasia is to administer a standardized aphasia test battery such as the Western Aphasia Battery (Kertesz, 1982) or the Boston Diagnostic Aphasia Examination (Goodglass and Kaplan, 1983).^[2] Standard tests to grammar assessment have three areas include testing sentence comprehension, testing verbs and verb arguments, testing sentence production and examining spontaneous discourse. Sentence comprehension tests are revised token test (McNeil and Prescott, 1978), auditory comprehension test for sentences (Shewan, 1981). However, they do not address all types of sentence-level problems. The most aspect of testing for agrammatic aphasia is the examination of verbs and verb arguments structures. Few published test is presently available for doing this, although several research laboratories around the world have developed protocols for this purpose.^[2] One measure that has been used clinically for testing sentence production is the story completion test (Gleason *et al.*, 1975; Goodglass *et al.*, 1972). The examiner presents the model of the target sentence using the semantically reversed picture and then asking the patient to produce a similar sentence the target picture.^[2]

Address for correspondence: Dr. Shohreh Jalaie, PhD of Biostatistics, Physiotherapy Department, Faculty of Rehabilitation, Tehran University of Medical Sciences, Pich-e-Shemiran, Enghelab Avenue, Tehran 1148965141, Iran. E-mail: jalaieish@sina.tums.ac.ir

Received: 09-10-2013; **Revised:** 13-03-2014; **Accepted:** 08-06-2014

Agrammatic output is commonly sampled in the clinic by asking an individual to produce a series of researcher made sentences that describe a nonstandard picture or sequence of pictures. Other regularly employed methods of data collection include a speaker re-telling a well-known story or engaging in dialogue on a topic suggested by the clinician. In contrast, the sampling of spontaneous speech produced for real-life communication with key conversational partners is still relatively rare.^[4] In general, the tests can assess various aspects of grammar that this is not possible in assessing spontaneous speech and pictures description. Also, speakers with aphasia produce shorter syntactic units than the speakers without aphasia.^[5] Therefore, clinicians cannot assessment all of grammar or syntax aspects perfectly. Mentioned standard tests above cannot evaluate sentence production detect therefore researchers begin to make individually tests or tasks to evaluate sentence production for agrammatic aphasia patients.

It seems a combination of all of exist tests for assessment data combined with quantitative and qualitative evaluating is necessary to provide a relatively complete process to evaluation^[5] and assessment of treatment progress of sentence production in aphasia. A variety of different languages tests or tasks have studied of syntax or morphology in the world that are designed in sentence production. Authors did not find systematic review in diagnostic or assessment tests to sentence production assessment. In this paper in order to help speech-language pathologists clinicians and researchers, all of exist syntax tests or tasks in this patients based on their structure and methods are studied, classified and discussed their assessing areas The main goal of this study is to determine more frequent methods in sentence production assessment in agrammatic Broca aphasia.

MATERIALS AND METHODS

Literature search

We searched the Cochrane library, Medline by PubMed, Science Direct, Scopus and Google Scholar from 1980 to October 1, 2013. All studies were searched with this keyword: “(Sentence or syntax or syntactic) and production test (or assessment or evaluation or examination) and agrammatic aphasia” in these sites.

Study selection and eligibility criteria

In this study, all articles were studied included sentence assessment tests or tasks in adult agrammatic aphasia.

All of the syntax or sentence production articles or ways to determine sentence production assessment, all systematic review about them and all articles that were targeted therapy; but, it was used for the evaluation of sentence production were collected.

Eligibility criteria were:

1. The studies included test or task to diagnosis and assessment of sentence production abilities,
2. The test or task performed on agrammatic aphasia patients,
3. The test or task to determine of the methods to evaluate sentence production,
4. The articles were designed to evaluate sentence production ability before and after treatment (speech therapy).

Exclusion criteria were if:

1. The paper was reported a case study,
2. The language of test or task was not English,
3. The test or task did not assess sentence production specifically (such as they were evaluated verb production by verb naming or complete sentences, verb agreement or tense and they did not assess making sentences),
4. The paper was review article of previous study, and
5. The article was assessed bilingual aphasia patients.

Data extraction and abstraction

In this paper, the content of each test or task was evaluated and classified in separate tables. Then they were reported based on:

1. General characteristics of study (first author’s name, publication year or study year, study design, and sampling method) and then were discussed on:
2. Characteristics of the study population (age and sex of studied participants and sample size, follow-up), and were completed
3. Type and duration of the interventional study, and finally were reported
4. Main finding of studies (validity and reliability of tasks or tests).

RESULTS

The data collected from the databanks showed that many studies have investigated the production of sentence.

A total of 160 articles were found in the sites that have studied sentence production. In this review, 122 articles were removed, and 38 articles were evaluated in detail.

In this section, articles were designed on three methods of sentence production assessment regardless of the purpose of studies. The results are presented in separate tables. Table 1 summarizes the features of tasks and tests designed for sentence production in agrammatic aphasia.^[6-20] Table 2 includes the characteristics of the tasks designed for evaluate of spontaneous speech.^[21-23] Table 3 contains the tasks to evaluate of both sentence production and spontaneous speech.^[24-43] All studies were of two types of

Table 1: The characteristics of tasks and tests designed for sentence production in agrammatic aphasia

First author's name (reference number)	Title	Publication year	Study design	Items of assessment or intervention	Contents of assessment or intervention	Age	Sex	Sample size	Duration of intervention	Follow-up	Validity (%)	Reliability (%)
Saffran, et al. ^[6]	The word order problem in agrammatism. II. Production	1980	(assessment)	Task=24 item (picture-description)	Active sentence	Not reported	Both sexes	P=5 N=22 (experiment 1, 2) N=17 (experiment 3)	-	-	Not reported (normal and patient is assessed)	Not reported
Schwartz, et al. ^[7]	Mapping therapy: A treatment program for agrammatism	1994	Single-subject multiple baseline design	Task=not tell (picture-description)	Active transitives with action verbs; active transitives with experiential verbs and noncanonical sentence types (e.g., passives) with action verbs (type C)	Not available	Both sexes	P=8	Not available	Not available	Not available	Not available
Caplan and Hanna ^[8]	Sentence production by aphasic patients in a constrained task	1998	(assessment)	Task=25 item (picture description)	Active, passive, dative, dative-passive, subject-object relative	P=20-81 years N=41-81 years	Both sexes	P=60 N=55	-	-	Not reported (normal and patient is assessed)	Not reported
Ihara and Fujita ^[9]	On the production of irrelevant arguments in agrammatism	2003	(assessment)	Task=50 verbs (picture-description)	Action verbs in active sentence	P=37-72 years N=39, 58 years	Both sexes Males	P=5 N=2	-	-	Not reported	Not reported
Bastiaanse and Thompson ^[10]	Verb and auxiliary movement in agrammatic Broca's aphasia	2003	(assessment)	Task=40 items (sentence production priming using picture)	(a) Active declarative sentences with a finite lexical verb, (b) active declarative sentences with a finite auxiliary, (c) yes/no question, in which the finite auxiliary moves to C	P=27-65 years	Both sexes	P=15	-	-	Not reported	Not reported
Lee and Thompson ^[11]	Agrammatic aphasic production and comprehension of unaccusative verbs in sentence contexts	2004	(assessment)	Task=22 items (picture description)	Unaccusative=12 and unergative verb=10	P=58.8 years (mean age) N=19.7 years (mean age)	Both sexes	P=8 N=5	-	-	Not reported (normal and patient is assessed)	Independent scorer=99%
Bastiaanse and van Zonneveld ^[12]	Sentence production with verbs of alternating transitivity in agrammatic Broca's aphasia	2005	(assessment)	Task=30 items (sentence elicitation task using picture)	Transitive; intransitive or unaccusative	P=20-70 years N=Matched to patients	Both sexes	P=16 N=6	-	-	Not reported (normal and patient is assessed)	Not reported

(Continued)

Table 1: (Continued)

First author's name (reference number)	Title	Publication year	Study design	Items of assessment or intervention	Contents of assessment or intervention	Age	Sex	Sample size	Duration of intervention up	Follow-up	Validity (%)	Reliability (%)
Schroder et al. ^[13]	Does training-induced improvement of noncanonical sentence production in agrammatic aphasia generalize to comprehension? A multiple single case study	2005	A-B-A-C-A design	Task=80 items (sentence elicitation task using picture)	ORC, object derived who-questions (whoq)	P=33-67 years	Both sexes	P=7	12 sessions in each treatment phases	-	Not reported	Not reported
Duman et al. ^[14]	Object scrambling and finiteness in Turkish agrammatic production	2007	(assessment)	Task=60 items (sentence completion task using picture)	Word order and verb inflection	P=24-74 years N=Matched to patients	Both sexes	P=8 N=8	-	-	Not reported (normal and patient is assessed)	Not reported
Burchert et al. ^[15]	Production of noncanonical sentences in agrammatic aphasia: Limits in representation or rule application?	2008	(assessment)	Task=60 items (sentence completion task using picture)	Scrambled and nonscrambled sentences	P=25-69 years	Both sexes	P=9	-	-	Not reported (normal and patient is assessed)	Not reported
Neuhaus and Penke ^[16]	Production and comprehension of wh-questions in German Broca's aphasia	2008	(assessment)	Task=54 items (elicitation sentence=54 items using picture and sentence repetition task=30 items)	Wh-subject, wh-object or wh-adjunct questions	P=53-68 years	Both sexes	P=9	-	-	Not reported	Not reported
Thompson and Lee ^[17]	Psych verb production and comprehension in agrammatic Broca's aphasia	2009	(assessment)	Tasks=24 items (sentence production using picture)	SubjExp and ObjExp psych verbs	P=58.8 years (mean age) N=19.7 years (mean age)	Both sexes	P=8 N=5	-	-	Not reported (normal and patient is assessed)	Inter-coder reliability=for comprehension task=100% and for production task=97.8%
Cho and Thompson ^[18]	What goes wrong during passive sentence production in agrammatic aphasia: An eye tracking study	2010	(assessment)	Task=40 items (sentence production priming using picture)	Active and passive	P=38-66 years	Both sexes	P=9	-	-	Not reported	Not reported

(Continued)

Table 1: (Continued)

First author's name (reference number)	Title	Publication year	Study design	Items of assessment or intervention	Contents of assessment or intervention	Age	Sex	Sample size	Duration of Follow-up intervention	Validity (%)	Reliability (%)
Dragoy and Bastiaanse ^[9]	Verb production and word order in Russian agrammatic speakers	2010	(assessment)	Task=60 items (sentence production priming using picture)	Six conditions=sentences with different numbers of verb arguments (one or two), different types of thematic role mapping (direct or indirect) and different word order (basic or scrambled)	P=24-68 years N=Matched to patients	Both sexes	P=16 N=16	-	Not reported (normal and patient is assessed)	Not reported
Thompson <i>et al.</i> ^[20]	Training verb argument structure production in agrammatic aphasia: Behavioral and neural recovery patterns	2013	Single-subject multiple baseline design	Task=not tell (sentence production using picture)	Active	Not available	Both sexes	P=8 (4=experimental and 4=control participants)	Not available	Not reported (normal and patient is assessed)	Not available

P = Number of patients; N = Number of normal adults; ORC = Object relative clauses

Table 2: The characteristics of tasks designed for evaluate of spontaneous speech in agrammatic aphasia

First author's name (reference number)	Title	Publication year	Study design	Items of assessment or intervention	Contents of assessment or intervention	Age	Sex	Sample size	Duration of Follow-up intervention	Validity (%)	Reliability (%)
Rochon <i>et al.</i> ^[21]	Quantitative analysis of aphasic sentence production: Further development and new data	2000	(assessment)	Spontaneous speech (in story telling)	The story of "Cinderella" or "Little Red Riding Hood", "Jack and the Beanstalk"	P=22-74 years N=20-73 years	Both sexes	P=29 N=12	-	Not reported (normal and patient is assessed)	Interrater reliability=for patients=97% and for normal=98% and test-retest Reliability=between 53 and 92% in elements
Webster <i>et al.</i> ^[22]	An investigation of the interaction between thematic and phrasal structure in nonfluent agrammatic subjects	2001	(assessment)	Spontaneous speech (in story telling)	The story of Cinderella	P=40-72 years N=18-90 Years	Both sexes	P=14 N=20	-	Not reported (normal and patient is assessed)	Not reported
Milman <i>et al.</i> ^[23]	A psychometric analysis of functional category production in English agrammatic narratives	2008	(assessment)	Spontaneous speech=in story telling	Production of clausal functional categories=in tell the story of "Cinderella"	P=29-78 years N=28-86 years	Both sexes	P=18 N=18	-	Not reported (normal and patient is assessed)	Inter-coder reliability=96%

P = Number of patients; N = Number of normal adults

Table 3: The characteristics of tasks and tests designed for sentence production and for evaluate of spontaneous speech in agrammatic aphasia

First author's name (reference number)	Publication year	Title	Study design	Items of assessment or intervention	Contents of assessment or intervention	Age	Sex	Sample size	Duration of intervention	Follow-up	Validity (%)	Reliability (%)
Thompson and McReynolds ^[24]	1986	Wh interrogative in agrammatic aphasia: An experimental analysis of auditory-visual stimulation and direct-production treatment	Single-subject multiple baseline design	Spontaneous speech (in picture description) Tasks=30 items (sentence modeling tasks using picture)	Cookie theft-coming home-cart 1 from the WAB What, where, who, why questions sequences	P=35-58 years	Both sexes	P=4	80% correct responses on two consecutive sessions or 15 sessions	5 weeks after treatment	Not reported (normal and patient is assessed)	Interobserver reliability for dependent variable = 98% interjudge reliability for independent variable = 98% production = 98% and comprehension = 99%
Goodglass et al. ^[25]	1993	Comparison of morphology and syntax in free narrative and structured tests: Fluent versus nonfluent aphasics	(assessment) Single-subject multiple baseline design	Spontaneous speech (in picture description) Tasks=not tell (the cross-modal MSB=part word order)	Cookie theft and four cartoon strip sequences Active SVO sentences and simple passives sequences	P=55-81 years N=Matched to patients	Both sexes	P=14 N=18	-	-	Not reported	Not reported
Thompson and Shapiro ^[26]	1995	Training sentence production in agrammatism: Implications for normal and disordered language	Single-subject multiple baseline design	Spontaneous speech (in picture description) telling and conversational conditions) Tasks=60 items (sentence production priming using picture)	The story of Cinderella and conversational conditions Wh-question-object cleft-passive sequences	Not reported	Not reported	P=17	Not reported	Not reported	Not reported (normal and patient is assessed)	Not reported
Thompson et al. ^[27]	1996	Training wh-question production in agrammatic aphasia: Analysis of argument and adjunct movement	Single-subject multiple baseline design	Spontaneous speech (in picture description) telling and conversational conditions) Tasks=80 items (sentence production priming using picture)	The story of Cinderella and conversational conditions Who, what, when and where questions sentences	P=39-79 years N=Matched to patients	Both sexes	P=7 N=5	Between 10 and 17.5 weeks	4 and 6 weeks after treatment	Not reported	Three trained coders=87% Primary examiner and an independent observer=90%

(Continued)

Table 3: (Continued)

First author's name (reference number)	Title	Publication year	Study design	Items of assessment or intervention	Contents of assessment or intervention	Age	Sex	Sample size	Duration of intervention	Follow-up	Validity (%)	Reliability (%)
Thompson <i>et al.</i> ^[26]	Training and generalized production of wh -and NP-movement structures in agrammatic aphasia	1997	Single-subject multiple baseline design	Spontaneous speech (in the story telling) Tasks=60 items (sentence production priming using picture)	The stories of Cinderella and Little Red Riding Hood Wh-question - object cleft- Passive - subject raising	Not reported	Males	P=2	Between 18 and 21 weeks	4 weeks after treatment	Not reported	Two trained coders=82% Primary examiner and an independent observer=95%
Van De Sandt-Koenderman <i>et al.</i> , ^[29]	Stimulating sentence production in agrammatic patients: The effect of the visual cue program on spontaneous speech	1997	A-B-A-B design	Spontaneous speech (not available) Tasks=not available (a sentence-processing test battery)	Not available Not available	Not available	Not available	P=2	Not available	Not available	Not available	Not available
Hartsuiker and Kolk ^[30]	Syntactic facilitation in agrammatic sentence production	1998	(assessment)	Spontaneous speech (in conversational conditions Tasks=54 items (picture description without priming and picture description with priming)	Describe their illness history, former occupation, family and daily activities= 10 min Active and its passive sentences	P=28-65 years N=53 years (mean age)	Both sexes	P=12 N=12	-	-	Not reported (normal and patient is assessed)	Not reported
Thompson <i>et al.</i> ^[31]	The role of syntactic complexity in training wh-movement structures in agrammatic aphasia: Optimal order for promoting generalization	1998	Single-subject multiple baseline design	Spontaneous speech (in the story telling) Tasks=45 items (sentence production priming using picture)	The stories of Cinderella Object clefts and/or object extracted who-questions and passive sentences	P=29-68 years N=Matched to patients	Both sexes	P=3 N=5	100% correct in each treatment phase	2 weeks after treatment	Not reported (normal and patient is assessed)	Not reported Primary examiner and an independent observer=97%

(Continued)

Table 3: (Continued)

First author's name (reference number)	Title	Publication year	Study design	Items of assessment or intervention	Contents of assessment or intervention	Age	Sex	Sample size	Duration of intervention	Follow-up	Validity (%)	Reliability (%)
Ballard and Thompson ^[32]	Treatment and generalization of complex sentence production in agrammatism	1999	Single-subject multiple baseline design	Spontaneous speech (in the story telling) Tasks=50 items (sentence production priming using picture)	The stories of Cinderella Object clefts extracted matrix and embedded questions (which are noncanonical with wh-movement) and embedded actives (which are canonical with no overt movement)	P=38-69 years N=25.4 years (mean age)	Both sexes	P=5 N=10	Phase 1 = □60% correct over three consecutive sessions or 9 sessions Phase 2 = 100% correct over three consecutive sessions or 36 sessions	4 weeks after treatment	Not reported (normal and patient is assessed)	1. Inter reliability = 90.8% and intrater reliability = 96% 2. Primary examiner and an independent observer=interrater = 94.8% and intrater reliability = 97%
Jacobs and Thompson ^[33]	Cross-modal generalization effects of training noncanonical sentence comprehension and production in agrammatic aphasia	2000	Single-subject multiple baseline design	Spontaneous speech (in the story telling and conversational conditions) Task=20 items (sentence production priming and 2 written production using picture)	(a) The story of Cinderella, (b) Selection of part of discuss between participant and a family member (5-min), (c) Explain of prerecorded ABC "American Agenda" news segments Passive and object cleft	P=39-79 years	Both sexes	P=4	80% correct in two of three consecutive sessions or 15 sessions	2 weeks after treatment	Not reported	Interjudge reliability of responses=93%
Weinrich et al. ^[34]	Training agrammatic subjects on passive sentences: Implications for syntactic deficit theories	2001	Single-subject multiple baseline design	Spontaneous speech (in the story telling) Task=28 items (select the word and arrange them in a separate C-VIC template)	The story of Cinderella Passive	P=68, 71 years	Males	P=2	In each phase=80% correct in three consecutive sessions	-	Not reported	Not reported

(Continued)

Table 3: (Continued)

First author's name (reference number)	Title	Publication year	Study design	Items of assessment or intervention	Contents of assessment or intervention	Age	Sex	Sample size	Duration of intervention	Follow-up	Validity (%)	Reliability (%)
Friedmann ^[35]	Question production in agrammatism: The tree pruning hypothesis	2002	(assessment)	Spontaneous speech (in free conversation) Task=48 items (constrained question and question elicitation task repetition task using picture)	Free conversation between patient and examiner Wh question and yes/no question	P=16-70 years	Both sexes	P=15	-	-	Not reported	Two examiner reliability=95%
Schneider and Thompson ^[36]	Verb production in agrammatic aphasia: The influence of semantic class and argument structure properties on generalization	2003	Single-subject multiple baseline design	Spontaneous speech (in story telling) Task=102 items (making sentence using picture)	Tell the Cinderella story Active sentence with 2-place and 3-place verbs	P=Not reported	Both sexes	P=7	90% correct over three consecutive sessions or 12 sessions per treatment condition	3 weeks after treatment	Not reported (normal and patient is assessed)	1. Inter-observer reliability for the dependent variable=94% 2. Inter-judge reliability on the independent variable=100% 3. Inter-rater reliability for narrative=87%
Thompson et al. ^[37]	The role of syntactic complexity in treatment of sentence deficits in agrammatic aphasia: The CATE	2003	Single-subject multiple baseline design	Spontaneous speech (in the story telling) Tasks=60 items (sentence production priming using picture)	The story of Cinderella Object relative clause, object-cleft sentences and object-extracted who-questions sentences	P=Not reported	Not reported	P=4	10 sessions in each treatment phase	4 weeks after treatment	Not reported	Both the examiner and an independent reliability=97%
Thompson et al. ^[38]	Neural plasticity and treatment-induced recovery of sentence processing in agrammatism	2010	Single-subject multiple baseline design	Spontaneous speech (in the story telling) Test SPPT using picture	The story of Cinderella Active, subject wh-question, subject relative, object wh-question, object relative sentences	P=38-66 years N=32-79 years	Both sexes	P=6 N=12	80% correct or production on two consecutive probes or 20 sessions	-	Not reported (normal and patient is assessed)	Inter-rater reliability (by two independent rater)=for production=99.1% and for comprehension = 100%

(Continued)

Table 3: (Continued)

First author's name (reference number)	Title	Publication year	Study design	Items of assessment or intervention	Contents of assessment or intervention	Age	Sex	Sample size	Duration of intervention	Follow-up	Validity (%)	Reliability (%)
Thompson <i>et al.</i> ^[39]	Sentactics®: Computer-automated treatment of underlying forms	2010	Single-subject multiple baseline design	Spontaneous speech (in the story telling) Task=30 items (sentence-production priming)	Tell the story of Cinderella, Object relative, object cleft and object-extracted wh-question sentences	P=35-68 years	Both sexes	P=12	80% correct performance on 4 consecutive days or 20 sessions	-	Not reported	Intra-rater reliability=98%
Links <i>et al.</i> ^[40]	Training verb and sentence production in agrammatic Broca's aphasia	2010	Single-subject multiple baseline design	Spontaneous speech (in verbal communication between patient and examiner) Tasks=60 item (making sentence)	Explain the illness and speech problem in daily living and verbal communication Verbs	P=31-68 years N=Matched to patients	Both sexes	P=11 N=40	12 weeks	1 and 3 months posttreatment	Not reported (normal and patient is assessed)	3 rater reliability=high score
Lee and Thompson ^[41]	Real-time production of unergative and unaccusative sentences in normal and agrammatic speakers: An eye tracking study	2011	(assessment)	Spontaneous speech (in the story telling) Tasks=20 items and 40 filler items (making sentence)	The story of Cinderella Unergative and unaccusative verbs	P=35-56 years N=19-21 years	Both sexes	P=9 N=12	-	-	Not reported (normal and patient is assessed)	Not reported
Cho-Reyes and Thompson ^[42]	Verb and sentence production and comprehension in aphasia: NAVS	2012	(assessment)	Spontaneous speech (in the story telling) The NAVS=30 items (sentence and passives, production priming)	The story of Cinderella Active, SWQ	P=33-71 years	Both sexes	P=59	-	-	Not reported (normal and patient is assessed)	The examiner and an independent rater reliability=99.8%
Thompson <i>et al.</i> ^[43]	Syntactic and morphosyntactic processing in stroke-induced and primary progressive aphasia	2013	(assessment)	Spontaneous speech (in the story telling) NAVS test=30 items (sentence and passives, production priming)	Tell the story of Cinderella Active, SWQ and SR clause and passives, OWQ and OR clause	P=Experiment 1: 35-79 years Experiment 3: 50-74 years N=Experiment 3: 32-81 years	Both sexes	P=Experiment 1: 66 P=Experiment 2: 46 P=Experiment 3: 32 N=15	-	-	Not reported	Not reported

P = Number of patients; N = Number of normal adults; CATE = Complexity account of treatment efficacy; NAVS = Northwestern Assessment of Verbs and Sentences; MSB=Morphosyntax battery; SPPT = Sentence Production Priming Test; WAB = Western Aphasia Battery, SVO = Subject-Verb-Object; SWQ = Subject-extracted wh-question; SR = Subject relative; OWQ = Object-extracted wh-question; OR = Object relative; C-VIC = Computerized visual communication; WHQ=who-questions; Subj Exp=Subject Experience; Obj Exp=Object Experience.

assessment and intervention, and it was the kind of speech therapy intervention. This therapy is a type of noninvasive interventions.

In 15 studies, researchers have used tasks designed to evaluate the sentence production. All studies have used the picture description for this purpose, but one study also has repetition task. Some articles have studied canonical (active) sentence only, but 9 articles (60%) has evaluated canonical and noncanonical sentences, 2 articles has evaluated unaccusative and unergative verb and 1 article has assessed the psych verb production. Duration of the test was not reported in articles.

Finding shows the 3 studies were used the spontaneous speech for the assessment of sentence production. In these articles was evaluated narrative speech by the story telling of Cinderella. Duration of the story telling was not reported in articles, and 1 article has studied test-retest reliability.

20 articles have used both sentence production tasks and spontaneous speech for sentence production evaluation in agrammatic aphasia patients. Three studies have assessed free conversation between patient and examiner, but 12 studies (60%) have evaluated story re-telling. Furthermore, 2 articles have applied the story telling, and free conversation and two of them have applied picture description in aphasia tests. The content of 4 tasks in these articles was about verbs and its assessment and 16 tasks have evaluated noncanonical sentences in agrammatic patients. Duration of storytelling and tasks was not reported. The validity of tasks or tests was calculated in 10 articles by assessing normal adults and the others were not reported. Ten articles were calculated the inter-rater reliability that the most of them were evaluated sentence production before and after treatment.

Figure 1 reports studies that have been used tasks or spontaneous speech or both methods for sentence




Figure 1: Number of articles in three methods for sentence production assessment in 1980–2013 (1 = 1980, 2 = 1984–1988, 3 = 1989–1993, 4 = 1994–1998, 5 = 1999–2003, 6 = 2004–2008, 7 = 2009–2013)

production evaluation. It shows that most studies have used tasks in 2004-2008, but in the years 1999-2003 and 2009-2013 using both methods were common.

The result of evaluate the quality of articles using the Critical Appraisal Skills Program (CASP) for diagnostic test study showed based on data presented in articles could not response to the second question in CASP. So in all of articles response of the second question was “No.” It was impossible to continue to evaluate based on CASP guidelines.

DISCUSSION

The tests or tasks sentence production are useful tools for evaluating of speech output in aphasia patients. These tools can help clinicians in early diagnosis problems so it will not lose golden time of treatment. The correct selection of tasks can be very important based on the content and its quality, testing time and the scoring methods. In this study, we review all tasks and tests based on scientific methods of sentence production in the various language constructs.

Findings of this review suggest that there are different ways to evaluate the sentence production in agrammatic patients. It seems that before 1980, no study to use the scientific and systematic method for this purpose. From 1980, the sentence production is assessed by the different tasks or collecting spontaneous speech in story telling or free conversation. The researchers based their study’s aim are used each of these two methods or both them. In this study, we found articles that have used specific tasks to evaluate better and more accurate sentence deficits in agrammatic aphasia. And almost there are 2 articles every year in this field. Tests or tasks design process has been going on continuously, and researchers in recent years have suggested the more accurate methods for the analysis of samples from patients. As mentioned above, the most of the research were designed and used of tests or tasks, but they have not done anything to standardize the tests or tasks. Some authors in later years, her tasks or tests have performed on the greater number of patients; they have solved their problems and have used other tasks (other lexical and syntactic skills) then have developed the standard test. It should be noted that very few standardized tests have designed and have published to evaluate the types of sentence production in this patients. In most studies was applied researcher made test. In this review, there were a standardized test designed to assess the comprehension and perception of various types of sentences and during the years 1986-2011 were used in different studies. Finally, it was published in 2011, and its psychometric properties have investigated.

The researcher has used 2 methods for sentence production evaluation in articles: “Tests or tasks” and “spontaneous

speech.” The finding show that 39.5% of articles have used the method of design tests or tasks, the 7.9% have used spontaneous speech, and half of articles (52.6%) has applied both methods. Most studies have used both methods in the years 2009-2013. It can be inferred that the use of it can better demonstrate production deficits.

The 23 articles have examined the spontaneous speech totally, and speech sample have collected by the conversation between the patient and the examiner or the patient’s family (34. 8%) or by re-telling stories (65.2% of studies). It seems the stories telling can be extracting more information of lexical, structural (syntactic) and morphological characteristics compared with free conversation in these patients. Because the patient uses stereotype and short sentences to communicate and cannot be determined what kind of grammatical elements has been removed in the speech? But in telling the story, the patients to describe the events happened and have to use grammatical elements such as a subject, objects, adjectives, adverbs, pronouns, and verbs in their sentences. Thus, examiner can be deleted or identified nongrammatical elements in the patient’s speech and to judge it properly.

Review of articles show the researcher has used a variety of tasks or tests to evaluate of sentence production. These tasks include sentence completion, sentence repetition, making the sentence with the words given and sentence production priming. The few studies have used the combination of tasks in order to accurately speech analyze. By reviewing all studies (35 articles has used tasks or tests) it was found that the sentence production priming tasks was applied in 45.7% of studies (16 articles), making sentence was used 34.3% (12 studies) and sentence completion in the 11.3% of them (4 articles). In recent years, researchers tend to use of sentence production priming task. In this task, the examiner show one picture to the patient and produce a grammatical sentence after determining the role of elements in a sentence. The examiner asks the patient to perform the same process and produce a correct sentence. In this task is an emphasis on the self-production in patient. This task can be a good way to show one’s ability or inability on sentence production.

The Broca aphasia patients have the brain injury in the frontal lobe (in 44 and 45 Broadman area), and they have the most frequent syntactic errors that are processed in these areas. Therefore, processing and production of noncanonical sentences than canonical sentences is difficult for them because the noncanonical sentences are complex. These sentences contain two types of syntactic movement: NP-movement and wh-movement. The wh-movement sentences is more complex the NP-movement sentences because in this type of sentence is necessary the grammatical

elements to move to a higher nodes in the syntactic tree. The articles are considered complex sentences evaluating in these patients and in their tasks have assessed these sentences. The review of studies shows that two types of syntax movements have been studied in articles from 1986. The 25% of the article have studied the NP-movement and NP-movement sentences production. According to the main study, 28% of them have evaluated the type of NP-movement sentences, 48% have assessed the production of wh-movement sentences and 12% of articles have evaluated both types of syntactic movement. Furthermore, it is noted that 12% of them have assessed the scrambling. It is should be noted that in all studies was used the white and black line pictures for assessing of the sentence production.

Since the main objective of articles was to differentiate between normal subjects and patients; therefore, the finding showed the differential validity was calculated by the different statistical analysis. Half of the articles (55.3% of studies) have used the differential validity, but the other studies were not examined the kinds of validity. The reason could be that the pictures and tasks are clear for researcher, or the validity was not reported in the paper. Since any change in tasks or items (depend on easy or difficult items or type of the requested task) has a direct impact on patient response and the result of the test will affect directly, so the examiner through the assessment of validity of items and pictures in tasks can be sure the selected items is correct for evaluating of sentence production precisely. Hence, researchers should be evaluating the kind of validity of tests or tasks to select the most appropriate items.

There are three types of reliability in studies. The first type is the test-retest reliability. It was performed to ensure repeatability and consistency of responses in each test and calculated standard error of the mean in the test. This reliability has two types of intra-and inter-reliability. Inter-reliability reveals the examiner effects on the test. This reliability has been done in treatment studies in order to avoid the researcher bias. The test-retest reliability was calculated in 1 article. The second type of reliability measures the internal consistency of test, and it show that the test has integrity. This type of reliability is not reported in any of the articles. The third type is the editor’s reliability. It is performed in tests that report qualitative scores and tests are done based on examiner judgment. Since the correct answers are clear in all of the tasks and tests and identify the correct answer is not dependence on the examiner view, so this reliability is not necessary in any of the tests and tasks. Nevertheless, it seems to evaluate the reliability of the tasks or tests increases the value of them and researchers can use it more confidently in other studies.

The sentences production is investigated in English structure in most studies. Only few articles were studied on other

languages such as German, Dutch, Turkish, Arabic, Hebrew, and Russian. Since every language have the different structure therefore it is necessary are designed tests or tasks based on the language structure. The cross-linguistic studies can be useful to researchers. Therefore, it is necessary to study other languages such as Asian and African languages. It is noted that it is impossible examine the quality of the test or tasks based on CASP criteria in none of the articles. Because the second condition for this diagnostic or assessment study (which is the basic condition) was not reported in any of the articles.

CONCLUSION

The speech production evaluation is necessary the early stages of the disorder. By this way, we can assess the patient's syntax and morphology ability and resolve their problems using appropriate methods of treatment in early intervention. Without careful assessment of sentence production problems and treatment program will be based on early symptoms and it possible delayed the recovery in patients. In general, it can be stated that the sentence production have been studied in different methods. It can be inferred the use of both method of sentence production- narrative speech (free speech conversation or the story telling) and tasks designed- can get more information about the used or omitted various grammatical elements in agrammatic patients. Expert researchers and clinicians can use both of these methods and collected the complete language samples and tasks designed to extract more accurate data. Furthermore, the tests or tasks shall be designed the way that clinicians can do and analyze it, and Instructions of tests or tasks explained well and unambiguous. Therefore, it is essential that tasks and tests be standardized.

AUTHOR'S CONTRIBUTION

AM contributed in the conception of the work, conducting the study, revising the draft, approval of the final version of the manuscript, and agreed for all aspects of the work. ShJ contributed in the conception of the work, drafting and revising the draft, approval of the final version of the manuscript, and agreed for all aspects of the work.

REFERENCES

1. Chapey R, Hallowell B. Introduction to language intervention strategies in adult aphasia. In: Chapey R, editor. *Language Intervention Strategies in Aphasia and Related Neurogenic Communication Disorders*. 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2001. p. 3-10.
2. Thompson CK. Treatment of underlying forms: A linguistic specific approach for sentence production deficits in agrammatic aphasia. In: Chapey R, editor. *Language Intervention Strategies in Aphasia and Related Neurogenic Communication Disorders*. 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2001. p. 605, 612-4.
3. Bastiaanse R, Edwards S. Word order and finiteness in Dutch and English Broca's and Wernicke's aphasia. *Brain Lang* 2004;89:91-107.
4. Beeke S, Maxim J, Wilkinson R. Rethinking agrammatism: Factors affecting the form of language elicited via clinical test procedures. *Clin Linguist Phon* 2008;22:317-23.
5. Lind M, Kristoffersen KE, Moen I, Simonsen HG. Semi-spontaneous oral text production: Measurements in clinical practice. *Clin Linguist Phon* 2009;23:872-86.
6. Saffran EM, Schwartz MF, Marin OS. The word order problem in agrammatism. II. Production. *Brain Lang* 1980;10:263-80.
7. Schwartz MF, Saffran EM, Fink RB, Myers JL, Martin N. Mapping therapy: A treatment programme for agrammatism. *Aphasiology* 1994;8:19-54.
8. Caplan D, Hanna JE. Sentence production by aphasic patients in a constrained task. *Brain Lang* 1998;63:184-218.
9. Ihara H, Fujita I. On the production of irrelevant arguments in agrammatism. *Brain Lang* 2003;84:273-85.
10. Bastiaanse R, Thompson CK. Verb and auxiliary movement in agrammatic Broca's aphasia. *Brain Lang* 2003;84:286-305.
11. Lee M, Thompson CK. Agrammatic aphasic production and comprehension of unaccusative verbs in sentence contexts. *J Neurolinguistics* 2004;17:315-30.
12. Bastiaanse R, van Zonneveld R. Sentence production with verbs of alternating transitivity in agrammatic Broca's aphasia. *J Neurolinguistics* 2005;18:57-66.
13. Schroder A, Stadie N, Postler J, Lorenz A, Swoboda-Moll M, Burchert F, *et al*. Does training-induced improvement of noncanonical sentence production in agrammatic aphasia generalize to comprehension? A multiple single case study. *Brain Lang* 2005;95:202-3.
14. Duman TY, Aygen G, O'zgin N, Bastiaanse R. Object scrambling and finiteness in Turkish agrammatic production. *J Neurolinguistics* 2007;20:306-31.
15. Burchert F, Meissner N, De Bleser R. Production of non-canonical sentences in agrammatic aphasia: Limits in representation or rule application? *Brain Lang* 2008;104:170-9.
16. Neuhaus E, Penke M. Production and comprehension of wh-questions in German Broca's aphasia. *J Neurolinguistics* 2008;21:150-76.
17. Thompson CK, Lee M. Psych verb production and comprehension in agrammatic Broca's aphasia. *J Neurolinguistics* 2009;22:354-69.
18. Cho S, Thompson CK. What goes wrong during passive sentence production in agrammatic aphasia: An eyetracking study. *Aphasiology* 2010;24:1576-92.
19. Dragoy O, Bastiaanse R. Verb production and word order in Russian agrammatic speakers. *Aphasiology* 2010;24:28-55.
20. Thompson CK, Riley EA, Den Ouden DB, Meltzer-Asscher A, Lukic S. Training verb argument structure production in agrammatic aphasia: Behavioral and Neural recovery patterns. *Cortex* 2013; in press.
21. Rochon E, Saffran EM, Berndt RS, Schwartz MF. Quantitative analysis of aphasic sentence production: Further development and new data. *Brain Lang* 2000;72:193-218.
22. Webster J, Franklin S, Howard D. An investigation of the interaction between thematic and phrasal structure in nonfluent agrammatic subjects. *Brain Lang* 2001;78:197-211.
23. Milman LH, Dickey MW, Thompson CK. A psychometric analysis of functional category production in English agrammatic narratives. *Brain Lang* 2008;105:18-31.
24. Thompson CK, McReynolds LV. Wh interrogative production in agrammatic aphasia: An experimental analysis of auditory-visual stimulation and direct-production treatment. *J Speech Hear Res* 1986;29:193-206.
25. Goodglass H, Christiansen JA, Gallagher R. Comparison of morphology and syntax in free narrative and structured tests: Fluent vs. nonfluent aphasics. *Cortex* 1993;29:377-407.
26. Thompson CK, Shapiro LP. Training sentence production in agrammatism: Implications for normal and disordered language. *Brain Lang* 1995;50:201-24.

27. Thompson CK, Shapiro LP, Tait ME, Jacobs BJ, Schneider SL. Training wh-question production in agrammatic aphasia: Analysis of argument and adjunct movement. *Brain Lang* 1996;52:175-228.
28. Thompson CK, Shapiro LP, Ballard KJ, Jacobs BJ, Schneider SS, Tait ME. Training and generalized production of wh-and NP-movement structures in agrammatic aphasia. *J Speech Lang Hear Res* 1997;40:228-44.
29. Van de Sandt-Koenderman WM, Bonta E, Wielaert SM, Visch-Brink EG. Stimulating sentence production in agrammatic patients: The effect of the visual cue programme on spontaneous speech. *Aphasiology* 1997;11:735-59.
30. Hartsuiker RJ, Kolk HH. Syntactic facilitation in agrammatic sentence production. *Brain Lang* 1998;62:221-54.
31. Thompson CK, Ballard KJ, Shapiro LP. The role of syntactic complexity in training wh-movement structures in agrammatic aphasia: Optimal order for promoting generalization. *J Int Neuropsychol Soc* 1998;4:661-74.
32. Ballard KJ, Thompson CK. Treatment and generalization of complex sentence production in agrammatism. *J Speech Lang Hear Res* 1999;42:690-707.
33. Jacobs BJ, Thompson CK. Cross-modal generalization effects of training noncanonical sentence comprehension and production in agrammatic aphasia. *J Speech Lang Hear Res* 2000;43:5-20.
34. Weinrich M, Boser KI, McCall D, Bishop V. Training agrammatic subjects on passive sentences: Implications for syntactic deficit theories. *Brain Lang* 2001;76:45-61.
35. Friedmann N. Question production in agrammatism: The tree pruning hypothesis. *Brain Lang* 2002;80:160-87.
36. Schneider SL, Thompson CK. Verb production in agrammatic aphasia: The influence of semantic class and argument structure properties on generalisation. *Aphasiology* 2003;17:213-41.
37. Thompson CK, Shapiro LP, Kiran S, Sobecks J. The role of syntactic complexity in treatment of sentence deficits in agrammatic aphasia: The complexity account of treatment efficacy (CATE). *J Speech Lang Hear Res* 2003;46:591-607.
38. Thompson CK, den Ouden DB, Bonakdarpour B, Garibaldi K, Parrish TB. Neural plasticity and treatment-induced recovery of sentence processing in agrammatism. *Neuropsychologia* 2010;48:3211-27.
39. Thompson CK, Choy JJ, Holland A, Cole R. Sentactics®: Computer-Automated Treatment of Underlying Forms. *Aphasiology* 2010;24:1242-266.
40. Links PA, Hurkmans JB, Bastiaanse R. Training verb and sentence production in agrammatic Broca's aphasia. *Aphasiology* 2010;24:1303-25.
41. Lee J, Thompson CK. Real-time production of unergative and unaccusative sentences in normal and agrammatic speakers: An eyetracking study. *Aphasiology* 2011;25:813-25.
42. Cho-Reyes S, Thompson CK. Verb and sentence production and comprehension in aphasia: North western assessment of verbs and sentences (NAVS). *Aphasiology* 2012;26:1250-77.
43. Thompson CK, Meltzer-Asscher A, Cho S, Lee J, Wieneke C, Weintraub S, *et al.* Syntactic and morphosyntactic processing in stroke-induced and primary progressive aphasia. *Behav Neurol* 2013;26:35-54.

Source of Support: Nil, **Conflict of Interest:** None declared.