DOI: 10.1111/ijcp.13524

E D I T O R I A L INFECTIOUS DISEASE

CLINICAL PRACTICE WILEY

When public health crises collide: Social disparities and COVID-19

In *To Have or to Be*?, psychoanalyst Erich Fromm writes about pursuit after domination of nature, material abundance and unlimited happiness, which made modern society become more interested in *having* than in *being*. Income, in his view, should not be as accentuated as to create different experiences of life for different groups.¹ Of the concepts that Fromm presents, the domination of nature, which facilitates zoonotic spillover events by increasing the overlap between the habitat of various species with that of humans,²⁻⁵ and the gap between the rich and the poor, which recently has become the widest in years,⁶ become particularly relevant in context of the COVID-19 pandemic.

Even though susceptibility to COVID-19 does not know socioeconomic boundaries, a critical and worrisome finding is emerging from preliminary data and may re-shape infectious disease outbreak management strategies for the future. An early analysis of COVID-19 data from several jurisdictions in the United States found that counties with a majority of African American residents had threetimes higher infection rates and six-times higher mortality rates than counties with a majority of Caucasian residents.⁷ Another analysis, of March 2020 COVID-19 hospitalisation data from 14 states in the United States, found more African American individuals among hospitalised patients whose race or ethnicity was recorded.⁸ These and other findings reveal a disproportionately higher risk of serious or fatal COVID-19 in minorities. What makes these observations remarkable is that hypertension, diabetes and obesity, which are risk factors for more severe or fatal COVID-19.9-13 are exactly the chronic conditions that have long been recognised as disproportionately affecting racial/ethnic minorities and socioeconomically disfavoured individuals and groups.¹⁴

Obesity affects minorities and low-socioeconomic-status groups disproportionately at all ages,¹⁵ a finding that was reported in several countries.¹⁶⁻¹⁹ Some of the risk factors that account for disparities in obesity include low socioeconomic status,²⁰ food insecurity, restricted access to healthy diet and recreational facilities,²¹⁻²⁴ residence in areas with fast food restaurants,²⁵ a high neighbourhood density of small grocery stores,²⁶ distance to a store,²⁷ exposure to obesogenic environments,^{28,29} shift work ³⁰ and irregular sleep patterns.³¹⁻³³

Obesity increases the risk for other chronic diseases,¹² including diabetes and hypertension.³⁴ African American adults in the United States have among the highest rates of hypertension worldwide.³⁵ Several factors were implicated in disparities in hypertension, including socioeconomic status,³⁶ differences in awareness,³⁷ residence in a food desert,³⁸ chronic stress,^{39,40} fewer healthcare resources⁴¹ and income.⁴² Disparities for diabetes were described in minority populations in terms of increased prevalence,^{43,44} worse management and control^{45,46} and higher rates of complications.^{45,47} Over the past three decades the socioeconomic disparities for type 2 diabetes have widened.⁴⁸

Racial, ethnic and socioeconomic disparities also shape inequities in the access to mental health care.⁴⁹⁻⁵² This is very relevant for COVID-19, in context of the guarantine that was implemented in many countries in various forms, including school closures, allowing non-essential personnel to work from home, closure of mass transit systems, cancellation of public events and restrictions on the assembly of groups of people.⁵³⁻⁵⁵ Social isolation negatively impacts mental health and, with >70% of the young people and adults not receiving adequate mental health treatment from health care personnel worldwide,⁵⁶ the implications in the wake of COVID-19 are extensive and far-reaching. The 2002-2003 SARS pandemic revealed that a substantial proportion of the guarantined individuals may display PTSD and depression symptoms, with longer duration of the quarantine being associated with more severe PTSD.⁵⁷ During the same pandemic, hospital employees from Beijing who were guarantined had higher PTSD levels than those who were not, even 3 years later.⁵⁸ Among individuals from South Korea isolated for 2 weeks during the 2015 MERS outbreak, anxiety and anger were still present 4-6 months after the quarantine.⁵⁹

The disproportionately higher suffering of socio-economically disadvantaged individuals at a moment of crisis is, unfortunately, nothing new. In the 14th century, in the Black Death pandemic, the poorest populations were also the most extensively impacted ones in terms of mortality,^{60,61} and low-income individuals had a considerably worse outcome after the 1918 flu pandemic.⁶² The disproportionate effect on socio-economically disadvantaged individuals was also apparent in the wake of natural disasters, such as Hurricane Katrina⁶³ or the Deepwater Horizon oil spill.⁶⁴ One aspect that makes COVID-19 different is that several segments of the population become more vulnerable not simply because of socioeconomic disparities, but as a result of chronic medical conditions that these disparities have at least partly fueled over decades. The partial overlap between the risk factors for these two groups of diseases is reminiscent of debates on whether the broad classification of diseases into non-communicable and communicable ones is a meaningful one, considering that the two groups often overlap and interact markedly with one another.⁶⁵⁻⁶⁷ LEY-CLINICAL PRACTICE

Another aspect that sets COVID-19 aside from other pandemics in recent history is the extent and the duration of the quarantine and the resulting increase in unemployment rates,^{68,69} which only promise to prolong and exacerbate the extent of social inequities and the burden of chronic diseases.

COVID-19 provides a steep and perplexing learning curve that underscores the imperative need to envision infectious diseases not simply from a biomedical perspective, but as part of a complex framework that incorporates ethnic, socioeconomic and political dimensions. Racial/ethnic and socioeconomic disparities are conducive to the development of chronic medical conditions that could increase the risk of severe COVID-19, widening the disparities and accentuating the chronic disease burden and, as a result, further marginalising already vulnerable individuals and groups. The implications of this positive feedback loop for individuals, groups and society, extend beyond COVID-19 and beyond infectious diseases in general. The current pandemic eloquently demonstrates, albeit at a high cost, that societies function on the basis of a social contract, as described by Jean-Jacques Rousseau and, undoubtedly, offers an important moment to reflect on the profound, far-reaching and multi-layered consequences of disparities in society.

> Richard A. Stein^{1,2} D Oana Ometa³

¹Department of Chemical and Biomolecular Engineering, NYU Tandon School of Engineering, Brooklyn, NY, USA ²Department of Natural Sciences, LaGuardia Community College, City University of New York, New York, NY, USA ³Journalism and Digital Media Department, Faculty of Political, Administrative and Communication Sciences Babes-Bolyai, Babes-Bolyai University, Cluj-Napoca, Romania

Correspondence

Richard A. Stein, Department of Chemical and Biomolecular Engineering, NYU Tandon School of Engineering, Brooklyn, NY, USA.

Email: steinr01@nyu.edu; richardastein@gmail.com

ORCID

Richard A. Stein D https://orcid.org/0000-0002-5944-8008

REFERENCES

- Fromm E. Have or to be. Continuum: New York 1977; Originally published in The World Perspective Series. Harper & Row; 1976.
- Olival KJ, Hosseini PR, Zambrana-Torrelio C, Ross N, Bogich TL, Daszak P. Host and viral traits predict zoonotic spillover from mammals. *Nature*. 2017;546:646-650.
- Hayman DT, Bowen RA, Cryan PM, et al. Ecology of zoonotic infectious diseases in bats: current knowledge and future directions. *Zoonoses Public Health*. 2013;60:2-21.
- Cheng VC, Lau SK, Woo PC, Yuen KY. Severe acute respiratory syndrome coronavirus as an agent of emerging and reemerging infection. *Clin Microbiol Rev.* 2007;20:660-694.

- Malta M, Rimoin AW, Strathdee SA. The coronavirus 2019-nCoV epidemic: Is hindsight 20/20? EClinicalMedicine. 2020;20:100289.
- Dabla-Norris E, Kochhar K, Ricka F, Suphaphiphat N, Tsounta E. Causes and consequences of income inequality: a global perspective. *Staff Discuss Notes*. 2015;15:1.
- Thebault R, Ba Tran A, Williams V. The coronavirus is infecting and killing black Americans at an alarmingly high rate. The Washington Post; 2020. https://www.ashingtonpostcom/nation/2020/04/07/ coronavirus-is-infecting-killing-black-americans-an-alarmingly -high-rate-post-analysis-shows/?arc404=true. Accessed April 20, 2020.
- Garg S, Kim L, Whitaker M, et al. Hospitalization rates and characteristics of patients hospitalized with laboratory-confirmed coronavirus disease 2019 – COVID-NET, 14 States, March 1–30, 2020. MMWR Morb Mortal Wkly Rep. 2020;69:458–464.
- 9. Lippi G, Wong J, Henry BM. Hypertension and its severity or mortality in Coronavirus Disease 2019 (COVID-19): a pooled analysis. *Pol Arch Intern Med.* 2020;130:304–309.
- Muniyappa R, Gubbi S. COVID-19 pandemic, corona viruses, and diabetes mellitus. Am J Physiol Endocrinol Metab. 2020;318:E736-E741.
- Guo W, Li M, Dong Y, et al. Diabetes is a risk factor for the progression and prognosis of COVID-19. *Diabetes Metab Res Rev.* 2020:e3319.
- 12. Ryan DH, Ravussin E, Heymsfield S. COVID 19 and the patient with obesity - the editors speak out. *Obesity (Silver Spring)*. 2020;28:847-847.
- Lighter J, Phillips M, Hochman S, et al. Obesity in patients younger than 60 years is a risk factor for Covid-19 hospital admission. *Clin Infect Dis.* 2020:ciaa415.
- Pellowski JA, Kalichman SC, Matthews KA, Adler N. A pandemic of the poor: social disadvantage and the U.S. HIV epidemic. *Am Psychol.* 2013;68:197-209.
- Wang Y, Beydoun MA. The obesity epidemic in the United Statesgender, age, socioeconomic, racial/ethnic, and geographic characteristics: a systematic review and meta-regression analysis. *Epidemiol Rev.* 2007;29:6-28.
- Kumanyika SK. Unraveling common threads in obesity risk among racial/ethnic minority and migrant populations. *Public Health*. 2019;172:125-134.
- Willows N, Dyck Fehderau D, Raine KD. Analysis Grid for Environments Linked to Obesity (ANGELO) framework to develop community-driven health programmes in an Indigenous community in Canada. *Health Soc Care Community*. 2016;24:567-575.
- Rosas LG, Stafford RS. Practical research strategies for reducing social and racial/ethnic disparities in obesity. Int J Obes (Lond). 2012;2012:S16-S22.
- 19. Jamil HJ, Rajan AK, Grzybowski M, Fakhouri M, Arnetz B. Obesity and overweight in ethnic minorities of the Detroit metropolitan area of Michigan. *J Community Health*. 2014;39:301-309.
- 20. Osborn CY, de Groot M, Wagner JA. Racial and ethnic disparities in diabetes complications in the northeastern United States: the role of socioeconomic status. *J Natl Med Assoc.* 2013;105:51-58.
- 21. Gordon-Larsen P, Nelson MC, Page P, Popkin BM. Inequality in the built environment underlies key health disparities in physical activity and obesity. *Pediatrics*. 2006;117:417-424.
- 22. Fernandes MM, Sturm R. The role of school physical activity programs in child body mass trajectory. *J Phys Act Health*. 2011;8:174-181.
- Bower KM, Thorpe RJ Jr, Rohde C, Gaskin DJ. The intersection of neighborhood racial segregation, poverty, and urbanicity and its impact on food store availability in the United States. *Prev Med*. 2014;58:33-39.
- Casagrande SS, Whitt-Glover MC, Lancaster KJ, Odoms-Young AM, Gary TL. Built environment and health behaviors among African Americans: a systematic review. Am J Prev Med. 2009;36:174-181.

- 25. Morland KB, Evenson KR. Obesity prevalence and the local food environment. *Health Place*. 2009;15:491-495.
- Gibson DM. The neighborhood food environment and adult weight status: estimates from longitudinal data. *Am J Public Health*. 2011;101:71-78.
- Ghosh-Dastidar B, Cohen D, Hunter G, et al. Distance to store, food prices, and obesity in urban food deserts. *Am J Prev Med*. 2014;47:587-595.
- Bell CN, Kerr J, Young JL. Associations between obesity, obesogenic environments, and structural racism vary by county-level racial composition. *Int J Environ Res Public Health.* 2019;16:861.
- Unal ER, Lynn T, Neidich J, et al. Racial disparity in maternal and fetal-cord bisphenol A concentrations. J Perinatol. 2012;32:844-850.
- Givens ML, Malecki KC, Peppard PE, et al. Shiftwork, sleep habits, and metabolic disparities: results from the survey of the health of Wisconsin. Sleep Health. 2015;1:115-120.
- Piccolo RS, Yang M, Bliwise DL, Yaggi HK, Araujo AB. Racial and socioeconomic disparities in sleep and chronic disease: results of a longitudinal investigation. *Ethn Dis.* 2013;23:499-507.
- Curtis DS, Fuller-Rowell TE, El-Sheikh M, Carnethon MR, Ryff CD. Habitual sleep as a contributor to racial differences in cardiometabolic risk. *Proc Natl Acad Sci USA*. 2017;114:8889-8894.
- Donat M, Brown C, Williams N, et al. Linking sleep duration and obesity among black and white US adults. Clin Pract (Lond). 2013;10:661-667.
- Pappachan JM, Chacko EC, Arunagirinathan G, Sriraman R. Management of hypertension and diabetes in obesity: non-pharmacological measures. *Int J Hypertens*. 2011;2011:398065.
- Roger VL, Go AS, Lloyd-Jones DM, et al. Heart disease and stroke statistics-2011 update: a report from the American Heart Association. *Circulation*. 2011;123:e18-e209.
- Ferdinand KC, Yadav K, Nasser SA, et al. Disparities in hypertension and cardiovascular disease in blacks: the critical role of medication adherence. J Clin Hypertens (Greenwich). 2017;19:1015-1024.
- Al Kibria GM. Racial/ethnic disparities in prevalence, treatment, and control of hypertension among US adults following application of the 2017 American College of Cardiology/American Heart Association guideline. *Prev Med Rep.* 2019;14:100850.
- Suarez JJ, Isakova T, Anderson CA, Boulware LE, Wolf M, Scialla JJ. Food access, chronic kidney disease, and hypertension in the U.S. *Am J Prev Med.* 2015;49:912-920.
- Hicken MT, Lee H, Morenoff J, House JS, Williams DR. Racial/ethnic disparities in hypertension prevalence: reconsidering the role of chronic stress. *Am J Public Health*. 2014;104:117-123.
- Dolezsar CM, McGrath JJ, Herzig AJM, Miller SB. Perceived racial discrimination and hypertension: a comprehensive systematic review. *Health Psychol.* 2014;33:20-34.
- Thorpe RJ Jr, Bowie JV, Smolen JR, et al. Racial disparities in hypertension awareness and management: are there differences among African Americans and Whites living under similar social conditions? *Ethn Dis.* 2014;24:269-275.
- Kelli HM, Hammadah M, Ahmed H, et al. Association between living in food deserts and cardiovascular risk. *Circ Cardiovasc Qual Outcomes*. 2017;10:e003532.
- Link CL, McKinlay JB. Disparities in the prevalence of diabetes: is it race/ethnicity or socioeconomic status? Results from the Boston Area Community Health (BACH) survey. Ethn Dis. 2009;19:288-292.
- Rodriguez JE, Campbell KM. Racial and ethnic disparities in prevalence and care of patients with type 2 diabetes. *Clin Diabetes*. 2017;35:66-70.
- Peek ME, Cargill A, Huang ES. Diabetes health disparities: a systematic review of health care interventions. *Med Care Res Rev.* 2007;64:101S-156S.
- Thackeray R, Merrill RM, Neiger BL. Disparities in diabetes management practice between racial and ethnic groups in the United States. *Diabetes Educ.* 2004;30:665-675.
- 47. Cummings DM, Doherty L, Howard G, et al. Blood pressure control in diabetes: temporal progress yet persistent racial

disparities: national results from the REasons for Geographic And Racial Differences in Stroke (REGARDS) study. *Diabetes Care*. 2010;33:798-803.

- Mezuk B, Li X, Cederin K, et al. Beyond access: characteristics of the food environment and risk of diabetes. *Am J Epidemiol.* 2016;183:1129-1137.
- Alegria M, Canino G, Rios R, et al. Inequalities in use of specialty mental health services among Latinos, African Americans, and non-Latino whites. *Psychiatr Serv.* 2002;53:1547-1555.
- Cook BL, Trinh NH, Li Z, Hou SS-Y, Progovac AM. Trends in racial-ethnic disparities in access to mental health care, 2004–2012. *Psychiatr Serv.* 2017;68:9-16.
- Alegria M, Vallas M, Pumariega AJ. Racial and ethnic disparities in pediatric mental health. *Child Adolesc Psychiatr Clin N Am.* 2010;19:759-774.
- Fryers T, Melzer D, Jenkins R. Social inequalities and the common mental disorders: a systematic review of the evidence. Soc Psychiatry Psychiatr Epidemiol. 2003;38:229-237.
- 53. Usher K, Bhullar N, Jackson D. Life in the pandemic: social isolation and mental health. *J Clin Nurs*. 2020.
- 54. Viner RM, Russell SJ, Croker H, et al. School closure and management practices during coronavirus outbreaks including COVID-19: a rapid systematic review. *Lancet Child Adolesc Health*. 2020;4:397-404.
- 55. Gostin LO, Friedman EA, Wetter SA. Responding to covid-19: how to navigate a public health emergency legally and ethically. *Hastings Cent Rep.* 2020;50:8-12.
- Henderson C, Evans-Lacko S, Thornicroft G. Mental illness stigma, help seeking, and public health programs. *Am J Public Health*. 2013;103:777-780.
- Hawryluck L, Gold WL, Robinson S, Pogorski S, Galea S, Styra R. SARS control and psychological effects of quarantine, Toronto, Canada. *Emerg Infect Dis.* 2004;10:1206-1212.
- Wu P, Fang Y, Guan Z, et al. The psychological impact of the SARS epidemic on hospital employees in China: exposure, risk perception, and altruistic acceptance of risk. *Can J Psychiatry*. 2009;54:302-311.
- Jeong H, Yim HW, Song YJ, et al. Mental health status of people isolated due to Middle East Respiratory Syndrome. *Epidemiol Health*. 2016;38:e2016048.
- Age DSN. Patterns of mortality during the black death in London, A.D. 1349-1350. J Archaeol Sci. 2010;37:3394-3400.
- DeWitte SN, Wood JW. Selectivity of black death mortality with respect to preexisting health. Proc Natl Acad Sci USA. 2008;105:1436-1441.
- Sydenstricker E. The incidence of influenza among persons of different economic status during the epidemic of 1918. 1931. Public Health Rep. 2006;121:191-204; discussion 190.
- Sastry N, VanLandingham M. One year later: mental illness prevalence and disparities among New Orleans residents displaced by Hurricane Katrina. Am J Public Health. 2009;99:S725-S731.
- 64. Lichtveld M. Disasters through the lens of disparities: elevate community resilience as an essential public health service. *Am J Public Health*. 2018;108:28-30.
- 65. Blundell HJ, Hine P. Non-communicable diseases: ditch the label and recapture public awareness. *Int Health*. 2019;11:5-6.
- Adjaye-Gbewonyo K, Vaughan M. Reframing NCDs? An analysis of current debates. *Glob Health Action*. 2019;12:1641043.
- 67. Oni T, Unwin N. Why the communicable/non-communicable disease dichotomy is problematic for public health control strategies: implications of multimorbidity for health systems in an era of health transition. *Int Health*. 2015;7:390-399.
- Woolhandler S, Himmelstein DU. Intersecting U.S. epidemics: COVID-19 and lack of health insurance. Ann Intern Med. 2020.M20-1491.
- 69. Karnon J. The case for a temporary COVID-19 income tax levy now, during the crisis. *Appl Health Econ Health Policy*. 2020.