

Research article

Open Access

In silico method for systematic analysis of feature importance in microRNA-mRNA interactions

Jiamin Xiao¹, Yizhou Li¹, Kelong Wang^{1,2}, Zhining Wen¹, Menglong Li^{*1}, Lifang Zhang¹ and Xuanmin Guang¹

Address: ¹College of Chemistry, Sichuan University, Chengdu 610064, PR China and ²Department of Chemistry and Biochemistry, Texas Tech University, Lubbock, Texas 79409-1061, USA

Email: Jiamin Xiao - xiaojiamin07@126.com; Yizhou Li - liyizhou_415@163.com; Kelong Wang - clong_w@hotmail.com; Zhining Wen - w_zhining@163.com; Menglong Li* - liml@scu.edu.cn; Lifang Zhang - zlffang2008@163.com; Xuanmin Guang - fly005304@163.com

* Corresponding author

Published: 16 December 2009

Received: 12 August 2009

BMC Bioinformatics 2009, 10:427 doi:10.1186/1471-2105-10-427

Accepted: 16 December 2009

This article is available from: <http://www.biomedcentral.com/1471-2105/10/427>

© 2009 Xiao et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: MicroRNA (miRNA), which is short non-coding RNA, plays a pivotal role in the regulation of many biological processes and affects the stability and/or translation of mRNA. Recently, machine learning algorithms were developed to predict potential miRNA targets. Most of these methods are robust but are not sensitive to redundant or irrelevant features. Despite their good performance, the relative importance of each feature is still unclear. With increasing experimental data becoming available, research interest has shifted from higher prediction performance to uncovering the mechanism of microRNA-mRNA interactions.

Results: Systematic analysis of sequence, structural and positional features was carried out for two different data sets. The dominant functional features were distinguished from uninformative features in single and hybrid feature sets. Models were developed using only statistically significant sequence, structural and positional features, resulting in area under the receiver operating curves (AUC) values of 0.919, 0.927 and 0.969 for one data set and of 0.926, 0.874 and 0.954 for another data set, respectively. Hybrid models were developed by combining various features and achieved AUC of 0.978 and 0.970 for two different data sets. Functional miRNA information is well reflected in these features, which are expected to be valuable in understanding the mechanism of microRNA-mRNA interactions and in designing experiments.

Conclusions: Differing from previous approaches, this study focused on systematic analysis of all types of features. Statistically significant features were identified and used to construct models that yield similar accuracy to previous studies in a shorter computation time.

Background

MicroRNAs (miRNAs) are short non-coding RNAs of approximately 22 nucleotides with some differences in one or two nucleotides in the 3' terminus. In eukaryotes, miRNA affects the stability and/or translation of mRNAs

and is involved in the regulation of various biological processes, such as development, differentiation, and apoptosis [1-5]. It has been reported that more than one-third of human genes can be targeted by miRNA and miRNAs have been linked to conditions such as lymphoma,

Results and Discussion

Feature extraction

All features were classified as sequence, structural or positional features. Studies have demonstrated that thermodynamic stability plays an important role in miRNA target prediction and machine learning algorithms have confirmed this [30,31]. In our feature extraction procedure, several features relate to thermodynamic feature indirectly, because the RNA secondary structure prediction was based on thermodynamic properties. Target sites for seed and non-seed segments were treated artificially in previous studies, which destroy the integrity of the target site and leads to underinvestigation of relevant biological properties. Here, a miRNA-target duplex was treated as a unit, which overcomes the disadvantage of previous methods, and feature extraction was parsed using *Perl*.

Sequence features (SEQ, Table 1) mainly include base frequencies and compositions. Background frequencies ($p_A = 0.34$, $p_C = 0.19$, $p_G = 0.18$, $p_U = 0.29$) have typically been used to produce artificial negative samples [22,32]. In microarray data analysis, researchers discovered that the four base frequencies were significantly different for candidate downregulated and normal genes. Furthermore, many dinucleotide sequences are statistically significant in miRNA target prediction [27,29].

Structural features (STRU, Table 2), including folding information for miRNA-target duplexes, should have the necessary information and should be crucial for recognition of miRNA-target interactions. Three main types of structural features, stems, loops and bulges, were used to represent RNA secondary structure. A well-known perfect or near-perfect pairing seed region is fairly important for these interactions (G:U wobble base-pairing is allowed). In addition, various types of loops and bulges in both seed and non-seed regions also affect the interactions [33].

Positional features (POSI, Table 3) reflect the mechanism of miRNA-target interactions. Saunders *et al.* investigated polymorphism of miRNA-target duplexes using single nucleotide polymorphism data, which revealed that a base mutation in the target sequence affects the regulation

function of miRNA [34]. These studies suggest that position-specific states (whether matching or not) may be important for miRNA target identification. Here only 21 nucleotides from the 5' terminus were considered. Four binary numbers were used to represent a position. If the sequence length is < 21 nt, the RF algorithm can automatically set the missing positions as missing values and replace them with the most frequent non-missing value (see the Methods section).

Prediction performance with RF

We used the RF algorithm and optimized the parameters. Two parameters, number of trees to grow *ntree* and number of variables randomly sampled as candidates at each split *mtry* were optimized using a grid search approach. During the grid search, the values of *ntree* = {500, 2000, 500} and *mtry* = {0, mdim, 1} were optimized based on 10-fold cross-validation (where the first number indicates the initial value, the second indicates is the final value, and the third is the increment used to generate values; mdim is number of features), which is partitions the original sample into 10 subsamples, 9 subsamples are to train model and the remainder one to test model and this process repeats 10 times. We then selected the value for the best-performing parameters to estimate the performance of the training model. It has been reported that an out-of-bag (OOB) error is very similar to the classification error for cross-validation (see Methods), which is a built-in measure of performance [35]. Table S1 lists the predictor performance comparison between based on cross-validation and OOB estimate (see Additional file 1).

RF models were constructed using SEQ, STRU, POSI, and the total feature set for two difference negative data sets; 200 samples from each class were randomly selected and used for training. This procedure was repeated 100 times and the average fraction of true positive (sensitivity) and true negative (specificity) predictions were used to determine the accuracy (The 200 negative samples always contain 35 experimental data in every randomly selection). The average prediction results for the models are listed in Table 4. Models based on the negative data set Neg_1 yielded higher accuracy than those based on Neg_2,

Table 1: Sequence features

Feature	Description
N_frac	Four features, percentage of A U C G nucleotides in the target sequence
GC_frac	Percentage of GC content in the target sequence
nt_match	Percentage of matching bases in the target sequence
nt_GUmatch	Percentage of GU matching bases in the target sequence
nt_mismatch	Percentage of mismatch bases in the target sequence
Dinucleotides	16 features, number of dinucleotide counts in the target sequence
Trinucleotides	64 features, number of trinucleotide counts in the target sequence

Table 2: Structural features

Feature	Description
Stem	Number of stems
Overhang	Number of overhangs
Stem_max	Length of the maximal stem in the target sequence
Loop_mismatch	Number of loops that only contain two mismatch bases
Loop_symm	Number of symmetric loops that contain more than two bases
Loop_asym	Number of asymmetric loops that contain more than two bases
Stems ^a	8 features, number of stems of length 6-10 and those with length < 6 and > 10
Loops ^b	8 features, the number of loops with length of 1-7 and those with length > 7
Bulges ^c	8 features, number of bulges with length of 1-7 and those with length > 7

^a A stem is defined as a set of consecutive pairs separated by unpaired bases.

^b A loop is defined as a set of unpaired bases between two strands.

^c A bulge is defined as a set of unpaired bases only in one strand

except for specificity for SEQ and POSI, probably because of the different data sources. The performance using sequence or structural features was worse than that for positional features. Our models achieved sensitivity of 0.947 and specificity of 0.916 for Neg_1 and sensitivity of 0.917 and specificity of 0.949 for Neg_2 using positional features. These results indicate that positional features yield a low number of false positive predictions and good model performance. Several studies have clearly shown an increase in the accuracy of prediction on combining numerous features. However, the hybrid of different feature types did not yield the highest accuracy (sensitivity 0.870 and specificity 0.922) for Neg_2. This indicates that negative correlation occurs in hybrids of different features. Hence, we examined the interactions among different features in each set using correlation analysis (see Additional file 2).

Feature importance measures

Current classification tasks need a measure of feature importance rather than only predicting the response using "black-box" models. Here, two different strategies were applied to measure feature importance in the prediction of miRNA-target interactions.

Permutation importance analysis of RF

RF is a classification method that also provides feature importance measures. It can distinguish significant predictor features from uninformative features and reduces interactions among features as much as possible. Three

measures of feature importance, the selection frequency, Gini importance and permutation importance, are available. In the present study, permutation importance was used as to measure feature importance in miRNA-target interactions separately for three feature sets to distinguish significant functional predictor features. The process was repeated 100 times with random resampling of constructed models and the feature measure scores were calculated. The distributions of these scores are shown in Figures 2, 3 and 4.

For sequence features (Figure 2), the percentage of match bases greatly contributes to recognition of interactions for both Neg_1 and Neg_2. The results are consistent with those of previous studies [27,29,36]. In addition, several dinucleotide and trinucleotide sequences were also statistically significant, such as UC, GG, CUC, and AGG. However, probably because of the negative data sources, base frequencies for A, G, and U in Neg_2 were ranked in the top 10. The results indicate that sequence-based feature importance depends on the data source.

The most informative structural feature was the stem, which indicated matching conditions for both seed and non-seed regions. Bulges provide more insight into miRNA-target interactions than loops do because more informative features are related to bulges than to loops (Figure 3). The overhang also has a significant effect on classification, which depends on the nature of the miRNA-target duplex. Features related to stems revealed

Table 3: Positional features

First two binary numbers	Meaning	Last two binary numbers	Meaning
00	A	00	Mismatch
11	U	11	GU match
10	G	10	Match
01	C		

Table 4: RF prediction results

Feature set	Neg_1			Neg_2		
	Optimal parameters	Se	Sp	Optimal parameters	Se	Sp
SEQ	ntree ^a = 2000, mtry ^b = 16	0.873	0.828	ntree = 1000, mtry = 66	0.821	0.885
STRU	ntree = 1500, mtry = 16	0.852	0.826	ntree = 1500, mtry = 7	0.807	0.808
POSI	ntree = 1000, mtry = 5	0.947	0.916	ntree = 1000, mtry = 4	0.917	0.949
Total	ntree = 2000, mtry = 6	0.971	0.918	ntree = 500, mtry = 37	0.870	0.922

Cross-validation was used to estimate the predictor performance of SEQ, STRU, POSI sets and the total feature set for two different negative data sets. Neg_1 comprises all experimental samples and inferred negative samples and Neg_2 comprises all experimental samples and artificial negative samples from miRanda. Sensitivity (Se) was calculated as TP/(TP+FN) and specificity (Sp) as TN/(TN+FP), where TP is the number of true positives, TN is the number of true negatives, FP is the number of false positives and FN is the number of false negatives.

^a number of trees to grow.

^b number of variables randomly sampled as candidates at each split.

that a length of 8 nt may be a more suitable definition of the seed region, although false miRNA targets usually had a match of < 6 nt in the seed region. Moreover, 1-nt bulges and 2-nt loops were ranked top and thus might greatly affect miRNA-target interactions.

Positional features suggested that non-seed regions also play an important role in miRNA-target interactions as shown by Figure 4, in which matching/mismatching serve as the positive/negative class, respectively. It has been shown in many studies that perfect or near-perfect base pairing in the seed region greatly contributes to the performance of models. Our results confirm that most of the dominant functional positional features are in the seed region.

A new feature importance measure strategy using hybrid models

A further investigation was implemented for a combination of the three feature sets. In this section, a novel conditional feature importance strategy was used to evaluate the total features. This strategy was implemented using the *cforest* function of the *party* package in R language, which can generate additional information on feature interactions [37-39].

A combination of three feature sets was considered to evaluate the feature importance using conditional variable importance strategy. Training data were randomly sampled and the procedure was repeated 100 times as above. Some of the significant features are shown in Figure 5. The measure strongly indicates that positional features greatly contribute to miRNA-target interactions because most statistically significant features are positional features in Neg_1. However, sequence features are dominant in the statistically significant features in Neg_2. These results are in agreement with the RF permutation importance. In addition, this strategy based on conditional inference trees seems to produce less noise than a permutation importance strategy.

Prediction performance of using only statistically significant features

To demonstrate the statistically significant features for predicting miRNA-target interactions, we compared the performance of different feature sets using only the significant features for two different negative data sets. The procedure is summarized below.

- (1) Calculate the feature importance score for each feature and rank the features according to these scores.
- (2) Eliminate the last feature and use the remaining features to construct a model.
- (3) Repeat step 2 until a remarkable decrease in accuracy occurs.
- (4) The remaining features are considered statistically significant and are used to construct a model. Receiver operating characteristic (ROC) are used to evaluate the prediction sensitivity and specificity.

We used this procedure to analyze the contribution of statistically significant features to the prediction accuracy (see Additional file 3). Finally, the top 10, 11, 5, and 10 statistically significant features were used to construct models for SEQ, STRU, POSI and total feature sets, which yielded AUC values of 0.919, 0.922, 0.969, and 0.978 for Neg_1 and 0.926, 0.874, 0.954, and 0.970 for Neg_2, respectively. Figure 6 shows ROC curves for each feature set for the two data sets. The highest accuracy was obtained by combining three types of feature sets. It is clear that the positional feature set exhibits higher accuracy than the other two feature types. The positional features were first investigated using miTarget, in which there are five position-based features ranked in the top 10, all belonging to the seed region. However, our positional features include two pieces of information: the type of nucleotide and whether it matches or not.




Figure 2
Feature importance measure for sequence features. The discriminatory power of each feature was determined by calculating the importance value, with larger values indicating more relevant properties. The importance distribution is shown for each sequence feature as a box plot in which the middle bar is the median, the outer edges are the 10 and 90 percentiles and the edges of the box are the 25 and 75 percentiles. Outliers are shown as circles. Neg_1 (all experimental samples and inferred negative samples) and Neg_2 (all experimental samples and artificial negative samples from miRanda) were analyzed separately.




Figure 3
Feature importance measure for structural features. The three rectangles denote features related to stems, bulges and loops.




Figure 4
Feature importance measure for positional features.




Figure 5

Comprehensive evaluation results. The plot displays the distribution for conditional feature importance using the *cforest* function for median importance values > 0.005. Sample without replacement and return unscaled measurement.




Figure 6
ROC curves for miRNA-target interactions identified using Neg_1 and Neg_2. Models were developed using only 10, 11, 5 and 10 statistically significant sequences, structural and positional features and total features, respectively. ROC curves are used to evaluate and compare the performance of miRNA-target interactions identified for four different feature sets.

In NBmiRTar, most statistically significant features were associated with loops and bulges, similar to the structural features of our method. Its motif features correspond to dinucleotide and trinucleotide sequences in our method. Our results prove that stems greatly contribute to recognition of miRNA-target interactions. More systematic analysis of dinucleotide and trinucleotide sequences was carried out in this study. MiRTif uses various k -gram frequencies as features for a triplet SVM classifier to predict pre-miRNA [40,41]. It is thought that these features represent the real environment for miRNA-target interactions. However, they might not be suitable for guiding experimental procedures.

Conclusions

MiRNA investigation not only sheds new light on RNA function, but can also reveal the mechanism involved in cell function and regulation. The actual correlation and mechanism for miRNA-target interactions are still unclear. However, the best solution might involve a combination of experimental and computational approaches. Our results demonstrate that this method yields good prediction and is robust. Moreover, the results will be useful in designing experimental procedures. As more experimental and unbiased data become available, our

approach could be improved and used to identify more reliable predictor features reflecting real miRNA-target interactions.

Methods

Machine learning, a broad subfield of artificial intelligence, can be used to automatically extract general rules from data sets through experience. Random forest is one of the most accurate prediction tools currently available for classification and regression. It is briefly described in this section.

Random forest

Random forest (RF) contains a number of unpruned decision trees. Each tree is trained and gives a classification using a different bootstrap sample from the original data. RF does not need a separate test set to obtain an unbiased estimate of the test set error because when using bootstrap sample from the original data, approximately one-third of the samples are left for internal estimation, which is called OOB data. However, if measures are based on the predictors' performance in the training set, there is no way of knowing whether the predictors are over-fitted to the training set. Instead, cross-validation should be used to test the performance of predictors. The RF algorithm is

widely used for classification and regression. It has been applied in complicated interactions and for data sets with many features, or so-called "small n , large p " problems [35,42]. Based on a tree structure, it has some advantages, such as interpretable classification rules and additional information that measures the importance of features. The important feature extraction strategy is a difficult issue owing to the complexity of feature interaction with other features. However, prediction of the model response cannot be achieved for many applications. Furthermore, RF can replace missing values by computing the median of all values of a variable in class j when the m th variable is not categorical, then using this value to replace all missing values for the m th variable in class j . If the m th variable is categorical, replacement is for the most frequent non-missing value in class j . These replacement values are called fills [35].

The rationale for permutation importance is random permutation of the predictor variable X_j , so its original association with the response Y disappeared. When the permuted variable X_j , together with the remaining non-permuted predictor variables, is used to predict the response, the prediction accuracy (i.e. the number of observations classified correctly) decreases greatly if the original variable X_j is associated with the response. Thus, a reasonable measure for variable importance is the difference in prediction accuracy before and after permutation of X_j . As an improvement, conditional importance can be considered [37,39]. Feature importance was our main focus, which is suitable for feature selection in many applications. In this study we used the *randomForest* and *party* packages in R language.

Data sets and performance evaluation

All experimental data were downloaded from TarBase 4.0 [11], which records experimentally validated target data via manual collection. The criteria for selection of training data were as follows:

- (1) Cleaved target data were eliminated because they might be different from translation repressed targets.
- (2) A miRNA-target duplex binding picture must be available (Figure 1).
- (3) Each target site sequence should not contain any unknown nucleotide (i.e. N can represent any nucleotide).

Consequently, a total of 294 miRNA-target pairs (259 positive and 35 negative) were collected for six species: *Drosophila*, *Caenorhabditis elegans*, human, mouse, rat and zebrafish. These data contained folding information

for duplexes, and truly biologically relevant simulation adapts to feature importance measures. However, the current set of 35 validated negative samples might not be enough to represent the negative class (Table S2 in Additional file 4). Therefore, two artificial negative sets were generated, as described below.

An inferred negative sample set and all negative experimental samples comprised Neg_1. It has been reported that *let-7* miRNA cannot repress expression after deletion of target sites on *lin-41* [23]. In other words, the remaining regions on the *lin-41* 3' UTR are not targeted by *let-7* miRNA [33]. Thus, if all the actual target sites on *lin-41* are masked, then the other remaining regions with favorable seed pairings are apposite as negative samples. The same is true for *miR-126** [43]. In practice, we used RNAhybrid to predict the binding duplex, using only duplexes with a match > 4 nt and discarding the other pairs to improve the quality of the data set. Thus, 167 inferred negative samples were obtained, 65 from *let-7* and 102 from *miR-126**.

We also generated 1000 artificial mature miRNAs (20-24 nt long) with A, C, G and U frequencies of 0.34, 0.19, 0.18 and 0.29, respectively. These base frequencies are not consistent with those in true miRNA. MiRanda was then used to predict target sites for the 1000 artificial miRNA from the 12,102 human 3'UTR sequences [44]. All these target sites are presumed to be false positive predictions since the query sequence did not include true miRNA. In practice, only 50 random artificial miRNA sequences were used to generate artificial negative samples from the 1000 sequences, because use of all 1000 artificial sequences yielded a set of predictions that was too large to be manageable. This artificial negative set was produced using the default MFE and SC value. In this case, MiRanda produced 78,169 false target sites, which together with the experimental negative samples comprised NEG_2. TargetBoost and NBmiRTar also generated a large negative class with this method [22,24].

Finally, the sensitivity and specificity were evaluated. The sensitivity for positive prediction, specificity for negative prediction and ROC plots of the true positive rate versus the false positive rate for varying decision cutoffs were used as measures of the model performance.

Availability and requirements

Source code and binaries freely available for download at http://cic.scu.edu.cn/bioinformatics/miRNA_code.zip

Programming language: Perl, R language

License: none

Abbreviations

miRNA: microRNA, mRNA, messenger RNA; RF: Random forest; SEQ: sequence features; STRU: structural features; POSI: positional features.

Authors' contributions

JX wrote the programs, designed the experiments, and drafted the manuscript. YL, LZ, XG helped in analysis and discussion, YL, KW, ZW refined the manuscript and gave useful comments. ML initialized and supervised the whole project. All authors read and approved the final manuscript.

Additional material

Additional file 1

Table S1. Comparison of OOB and CV error estimate.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2105-10-427-S1.DOC>]

Additional file 2

The interactions among different features in each set using correlation analysis.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2105-10-427-S2.ZIP>]

Additional file 3

The analysis on contribution of statistically significant features.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2105-10-427-S3.XLS>]

Additional file 4

Table S2. Prediction results of RF using experimental data. A very imbalanced experimental dataset (259 positive and 35 negative samples) was used for model training.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2105-10-427-S4.DOC>]

Acknowledgements

We gratefully acknowledge Dr. Peng Jiang and the anonymous reviewers for their constructive comments on this article. This work was supported by National Nature Science Foundation of China (No. 20775052).

References

- Bartel DP: **MicroRNAs: Genomics, biogenesis, mechanism, and function.** *Cell* 2004, **116**(2):281-297.
- Claverie JM: **Fewer genes, more noncoding RNA.** *Science* 2005, **309**(5740):1529-1530.
- Nelson P, Kiriakidou M, Sharma A, Maniatakis E, Mourelatos Z: **The microRNA world: small is mighty.** *Trends Biochem Sci* 2003, **28**(10):534-540.
- Mattick JS, Makunin IV: **Non-coding RNA.** *Hum Mol Genet* 2006, **15**:R17-R29.
- Wang GF, Mao WM, Zheng S: **MicroRNA-183 regulates Ezrin expression in lung cancer cells.** *FEBS Lett* 2008, **582**(25-26):3663-3668.
- He L, Thomson JM, Hemann MT, Hernando-Monge E, Mu D, Goodson S, Powers S, Cordon-Cardo C, Lowe SW, Hannon GJ, Hammond SM: **A microRNA polycistron as a potential human oncogene.** *Nature* 2005, **435**(7043):828-833.
- Iorio MV, Visone R, Di Leva G, Donati V, Petrocca F, Casalini P, Tac-cioli C, Volinia S, Liu CG, Alder H, Calin GA, Menard S, Croce CM: **MicroRNA signatures in human ovarian cancer.** *Cancer Res* 2007, **67**(18):8699-8707.
- Lau NC, Lim LP, Weinstein EG, Bartel DP: **An abundant class of tiny RNAs with probable regulatory roles in Caenorhabditis elegans.** *Science* 2001, **294**(5543):858-862.
- Lagos-Quintana M, Rauhut R, Lendeckel W, Tuschl T: **Identification of novel genes coding for small expressed RNAs.** *Science* 2001, **294**(5543):853-858.
- Lee RC, Ambros V: **An extensive class of small RNAs in Caenorhabditis elegans.** *Science* 2001, **294**(5543):862-864.
- Rhoades MW, Reinhart BJ, Lim LP, Burge CB, Bartel DP: **Prediction of plant microRNA targets.** *Cell* 2002, **110**(4):513-520.
- Griffiths-Jones S, Saini HK, van Dongen S, Enright AJ: **miRBase: tools for microRNA genomics.** *Nucleic Acids Res* 2008, **36**:D154-D158.
- Sethupathy P, Corda B, Hatzigeorgiou AG: **TarBase: A comprehensive database of experimentally supported animal microRNA targets.** *RNA* 2006, **12**(2):192-197.
- Lewis BP, Shih IH, Jones-Rhoades MW, Bartel DP, Burge CB: **Prediction of mammalian microRNA targets.** *Cell* 2003, **115**(7):787-798.
- Lall S, Grun D, Krek A, Chen K, Wang YL, Dewey CN, Sood P, Colombo T, Bray N, MacMenamin P, Kao HL, Gunsalus KC, Pachter L, Piano F, Rajewsky N: **A genome-wide map of conserved microRNA targets in C. elegans.** *Curr Biol* 2006, **16**(5):460-471.
- Miranda KC, Huynh T, Tay Y, Ang YS, Tam WL, Thomson AM, Lim B, Rigoutsos I: **A pattern-based method for the identification of microRNA binding sites and their corresponding heteroduplexes.** *Cell* 2006, **126**(6):1203-1217.
- Kiriakidou M, Nelson PT, Kouranov A, Fitziev P, Bouyioukos C, Mourelatos Z, Hatzigeorgiou A: **A combined computational-experimental approach predicts human microRNA targets.** *Genes Dev* 2004, **18**(10):1165-1178.
- Enright AJ, John B, Gaul U, Tuschl T, Sander C, Marks DS: **MicroRNA targets in Drosophila.** *Genome Biol* 2004, **5**(1):R1.
- Lai EC: **Predicting and validating microRNA targets.** *Genome Biol* 2004, **5**(9):115.
- Sethupathy P, Megraw M, Hatzigeorgiou AG: **A guide through present computational approaches for the identification of mammalian microRNA targets.** *Nat Methods* 2006, **3**(11):881-886.
- Berezikov E, Thuemmler F, van Laake LW, Kondova I, Bontrop R, Cuppen E, Plasterk RHA: **Diversity of microRNAs in human and chimpanzee brain.** *Nat Genet* 2006, **38**(12):1375-1377.
- Saetrom O, Snove O, Saetrom P: **Weighted sequence motifs as an improved seeding step in microRNA target prediction algorithms.** *RNA* 2005, **11**(7):995-1003.
- Kim SK, Nam JW, Rhee JK, Lee WJ, Zhang BT: **miTarget: microRNA target gene prediction using a support vector machine.** *BMC Bioinformatics* 2006, **7**:411.
- Yousef M, Jung S, Kossenkov AV, Showe LC, Showe MK: **Nave Bayes for microRNA target predictions machine learning for microRNA targets.** *Bioinformatics* 2007, **23**(22):2987-2992.
- Yan XQ, Chao TF, Tu K, Zhang Y, Xie L, Gong YH, Yuan JG, Qiang BQ, Peng XZ: **Improving the prediction of human microRNA target genes by using ensemble algorithm.** *FEBS Lett* 2007, **581**(8):1587-1593.
- Yang YC, Wang YP, Li KB: **MiRTif: a support vector machine-based microRNA target interaction filter.** *BMC Bioinformatics* 2008, **9**:S4.
- Grimson A, Farh KKH, Johnston WK, Garrett-Engle P, Lim LP, Bartel DP: **MicroRNA targeting specificity in mammals: Determinants beyond seed pairing.** *Mol Cell* 2007, **27**(1):91-105.
- Nielsen CB, Shomron N, Sandberg R, Hornstein E, Kitzman J, Burge CB: **Determinants of targeting by endogenous and exogenous microRNAs and siRNAs.** *RNA* 2007, **13**:1894-1910.
- Wang XW, El Naqa IM: **Prediction of both conserved and non-conserved microRNA targets in animals.** *Bioinformatics* 2008, **24**(3):325-332.

30. Long D, Lee R, Williams P, Chan CY, Ambros V, Ding Y: **Potent effect of target structure on microRNA function.** *Nat Struct Mol Biol* 2007, **14(4)**:287-294.
31. Kertesz M, Iovino N, Unnerstall U, Gaul U, Segal E: **The role of site accessibility in microRNA target recognition.** *Nat Genet* 2007, **39(10)**:1278-1284.
32. Rajewsky N, Sockanathan ND: **Computational identification of microRNA targets.** *Dev Biol* 2004, **267(2)**:529-535.
33. Vella MC, Reinert K, Slack FJ: **Architecture of a validated MicroRNA :: target interaction.** *Chem Biol* 2004, **11(12)**:1619-1623.
34. Saunders MA, Liang H, Li WH: **Human polymorphism at microRNAs and microRNA target sites.** *Proc Natl Acad Sci USA* 2007, **104**:3300-3305.
35. Breiman L: **Random forest.** *Mach Learning* 2001, **45**:5-32.
36. Doench JG, Sharp PA: **Specificity of microRNA target selection in translational repression.** *Genes Dev* 2004, **18(5)**:504-511.
37. Strobl C, Boulesteix AL, Kneib T, Augustin T, Zeileis A: **Conditional variable importance for random forests.** *BMC Bioinformatics* 2008, **9**:11.
38. Hothorn T, Hornik K, Zeileis A: **A laboratory for recursive partitioning.** *R package version 09-98 2008* [<http://cran.stat.auckland.ac.nz/web/packages/party/>].
39. Strobl C, Boulesteix AL, Zeileis A, Hothorn T: **Bias in random forest variable importance measures: Illustrations, sources and a solution.** *BMC Bioinformatics* 2007, **8**:21.
40. Xue CH, Li F, He T, Liu GP, Li YD, Zhang XG: **Classification of real and pseudo microRNA precursors using local structure-sequence features and support vector machine.** *BMC Bioinformatics* 2005, **6**:301.
41. Jiang P, Wu H, Wang W, Ma W, Sun X, Lu Z: **MiPred: classification of real and pseudo microRNA precursors using random forest prediction model with combined features.** *Nucleic Acids Res* 2007, **35**:W339-W344.
42. Liaw A, Wiener M: **Classification and regression by random Forest.** *R News* 2002, **2**:18-22 [<http://www.r-project.org/doc/Rnews/bib/Rnewsbib.html>].
43. Musiyenko A, Bitko V, Barik S: **Ectopic expression of miR-126*, an intronic product of the vascular endothelial EGF-like 7 gene, regulates protein translation and invasiveness of prostate cancer LNCaP cells.** *J Mol Med* 2008, **86(3)**:313-322.
44. Brockman JM, Singh P, Liu DL, Quinlan S, Salisbury J, Graber JH: **PACdb: PolyA cleavage site and 3'-UTR database.** *Bioinformatics* 2005, **21(18)**:3691-3693.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

