



Situation Analysis of Varroosis and Tropilaelaps Infestation of Honeybees in Thailand, 2017–2018

Tawan Thongsawang¹
Putthipanya Rueangsom¹ ²
Khemmapat Boonyo¹
Vilaiporn Wongphruksasoong¹
Rapeepong Suphanchaimat^{1,2,3} 

¹Bureau of Disease Control and Veterinary Services, Department of Livestock Development, Ministry of Agriculture and Cooperatives, Bangkok, Thailand; ²International Health Policy Program, Ministry of Public Health, Nonthaburi, Thailand; ³Division of Epidemiology, Department of Disease Control, Ministry of Public Health, Nonthaburi, Thailand

Background/Aim: To explore the prevalence of *Varroa destructor* and *Tropilaelaps* infestation in honeybees in Thailand and investigate factors associated with those diseases.

Methods: A quantitative cross-sectional design was employed during 2017–2018. We sampled 144 apiaries in 13 provinces from the surveillance database of the Department of Livestock Development. In total, 1,152 bee samples were collected. A microscopic exam was performed to assess if each sample was infested with *Varroa destructor* mites and *tropilaelaps* mites. A chi-square test and multivariable logistic regression were conducted.

Results: The prevalence of *Varroa destructor* and *Tropilaelaps* infestation at the apiary level was 50.69% and 32.64%, respectively. At the beehive level, we found that the prevalence of *Varroa destructor* infestation was 22.74% while that of *Tropilaelaps* infestation was 6.94%. The northern region saw the highest prevalence of *Varroa destructor* and *Tropilaelaps* infestation. Apiaries that received a “Good Agricultural Practice” (GAP) certificate from the Bureau of Livestock Standards and Certification, demonstrated a 42% lower chance of contracting both parasitic infestations; however, no statistically significant difference was reported. Apiaries that had a history of chemical use showed approximately 2.7 times greater odds of *Tropilaelaps* infestation (adjusted odds ratio [AOR] = 2.69; 95% confidence interval [CI] = 1.16–6.21) with statistical significance ($p = 0.02$). The probability of *Varroa destructor* infestation amongst apiaries with apiary movement was approximately 60% lower than amongst those without apiary movement (AOR = 0.40, 95% CI = 0.20–0.80, $p = 0.01$).

Conclusion: *Varroa destructor* and *Tropilaelaps* infestations are a critical concern for beekeeping in Thailand. Apiary movement tended to lower the risk of *Varroa destructor* infestation while chemical use tended to enhance the risk of *Tropilaelaps* infestation. Further studies that allow a more comprehensive collection of determinants of parasitic infestation in honeybees, for instance, apiary cleaning frequency and farm environments (such as temperature and rainfall), are recommended.

Keywords: *Tropilaelaps*, *Varroa destructor*, honeybee, Thailand

Introduction

The apiculture business in Thailand started in 1953 and has grown rapidly since then with support from both government and private sectors.¹ The development of beekeeping in Thailand is based on the intention to meet international standards. At present, the quality of Thai honey and bee products is well-accepted in wider international markets around the world. In 2015, Thailand exported over 19,000 tonnes of honey products and produced revenue of over 1.6 million Thai Baht (US\$ 51,364).²

The rapid expansion of the apiculture market and beekeeping industries comes alongside high-density bee farming practices, which is likely to cause disease

Correspondence: Tawan Thongsawang
Bureau of Disease Control and Veterinary Services, Department of Livestock Development, 69/1 Phayathai Road, Ratchathewi District, Bangkok, 10400, Thailand
Email tawanthongsawang@gmail.com

transmission among beehives/apiaries. *Varroa destructor* and *Tropilaelaps* infestations are amongst many parasitic diseases in beehives/apiaries that potentially undermine honey production and have a detrimental effect on bees' health, and may eventually compromise the honey market. Both diseases were also listed as harmful diseases as specified by the World Organisation for Animal Health (OIE) and Thailand Animal Epidemics Act, B.E. 2558 (2015).^{3,4} *Varroa* spp. is also a carrier of deformed wing virus (DWV) that causes abnormality of the wings in adult bees,⁵ subsequently leading to colony collapse disorder (CCD).⁶

In addition, a decline in the stock of bees negatively affects the whole agricultural cycle as over 70% of food crops require pollination from bees.⁷ Therefore, bees are an essential component of food security for humans and biodiversity of the ecosystem.⁸

For the above reasons, the Department of Livestock Development (DLD) within the Ministry of Agriculture and Cooperatives initiated a survey on diseases, chemicals, and drug residuals in bees during 2002–2003. The survey findings revealed that *Varroa destructor* and *Tropilaelaps* infestations, and European foulbrood were commonly found in many regions of Thailand, particularly the North, the Northeast, and the South.⁹

During 2017–2018, the DLD launched an active surveillance system for bee diseases using bee-apiary data in 13 sentinel provinces in the Thai Good Agricultural Practices (GAP) database. However, data from the survey have not been analysed and presented in a systematic manner until now. Furthermore, apiary-related factors that may contribute to parasitic diseases in bees have also not been scientifically explored.

Thus, the purpose of this study is to describe the prevalence of parasitic mite diseases in Thailand and the correlation between potential risk factors and the presence of these diseases. The results of this study should help provide recommendations for effective surveillance and better management of parasitic diseases in the apiculture areas in Thailand.

Materials and Methods

Study Design

Cross-sectional quantitative study.

Participants and Study Sites

The study target was bee apiaries in 13 provinces registered in the database of the active surveillance system of

the DLD. These provinces consisted of Chaiyaphum, Chiang Mai, Chiang Rai, Chon Buri, Chumphon, Lamphun, Loei, Nakhon Sawan, Nan, Phayao, Phitsanulok, Phrae and Uttaradit.

Sampling and Sample Size Calculation

We used a two-stage cluster sampling technique for sample acquisition. The primary sampling unit was apiaries, and the elementary sampling unit was bees. Sample size estimation was based on the prevalence estimation formula: $n = Z_{\alpha/2}^2 P(1-P)/d^2$, where $Z_{\alpha/2} = 1.96$; P denoted expected prevalence and d was an acceptable error. We assumed that the expected prevalence of parasitic disease in bee samples was 0.2,⁹ and we presumed an acceptable error of 0.05. As we applied the two-stage cluster sampling, the sample size needed to be adjusted for design effect. In this respect, we assumed a design effect of 2.¹⁰ Based on these assumptions, the estimated number of samples amounted to 922 samples. However, to account for the possibility of incomplete information due to errors from sample collection, we increased this sample size by 20% based on the concept of probability proportional to size (PPS) for a final sample size estimate of 1,160 samples.

Initially, we planned to select 145 apiaries from a pool of 318 apiaries in the 13 provinces to obtain the 1,160 bee samples – with eight bee samples randomly collected from each apiary. However, since there was one sampled apiary that we could not approach during fieldwork, we ultimately collected 1,152 bee samples, which still outnumbered the required sample size of 922 as shown in Table 1.

To collect the samples, we firstly collected approximately 100–200 bees in a clear bottle and covered it with a plastic lid. Secondly, we submersed the bees with 70% alcohol and waited for 30–60 minutes. Finally, we filtered the bees with a mesh strainer and collected the liquid to examine the mites under a stereomicroscope. This sample collection procedure was adapted from the OIE and the Thai National Institute of Animal Health.

Data Analysis

Both descriptive statistics and inferential statistics were applied. For descriptive statistics, frequency and percentage were used to assess the prevalence of *Varroa destructor* and *Tropilaelaps* infestations. Prevalence was analysed at two levels: (i) the beehive level and (ii) the apiary level. An apiary was classified as having contracted a disease if at least one of the bee samples tested

Table 1 Number of Bee Samples and Apiaries Selected in Each Province

Province	No. of Existing Apiaries Database	Density of Apiary (Apiary: Orchards and Perennials Area* (km ²))	No. of Apiaries from the Sampling	No. of Actual Acquired Samples
Chon Buri	1	1: 1,121	1	8
Chaiyaphum	1	1: 238	1	8
Loei	2	1: 600	1	8
Chiang Mai	129	1: 9	57	456
Chiang Rai	36	1: 33	16	128
Phrae	38	1: 3	17	136
Nan	58	1: 8	26	208
Phayao	2	1: 289	1	8
Lamphun	3	1: 249	2	16
Uttaradit	9	1: 27	4	32
Phitsanulok	27	1: 21	12	96
Nakhon	4	1: 33	2	16
Sawan				
Chumphon	7	1: 470	4	32

Notes: The average number of colonies per apiary is 214; Source: *Office of Agricultural Economics of Thailand.¹⁷

positive for *Varroa* spp. or *Tropilaelaps* spp. from the microscopic examination. For inferential statistics, we used a Chi-square test for univariable analysis and a multivariable logistic regression for multivariable analysis; the multivariable logistic regression analysis was conducted at the apiary level only. The outcome variable of interest was the presence of *Varroa destructor* and *Tropilaelaps* infestations in the apiary, regardless of the number of infested bees. The independent variables were (i) the acquisition of a GAP certificate, (ii) history of apiary movement within 30 days before sample collection, and (iii) the use of chemicals (repellents and acaricides) within 30 days prior to the fieldwork. The measures of association were presented in terms of crude odds ratio (COR) and adjusted odds ratio (AOR) with a 95% confidence interval (95% CI).

Ethics Consideration

This study was approved by the Research Committee of the Bureau of Disease Control and Veterinary Services, Department of Livestock Development, Thailand (Permit number: 0610.04/272).

Results

Descriptive Analysis

This study found that the prevalence of *Varroa destructor* and *Tropilaelaps* infestations at the apiary level was 50.69% and 32.64%, respectively. The top three provinces with the greatest prevalence of *Tropilaelaps* infestation

were Chon Buri, Phrae, and Uttaradit. For *Varroa destructor* infestation, the top three provinces were Loei, Nan, and Phrae.

At the beehive level, the overall prevalence of *Tropilaelaps* and *Varroa destructor* infestation was 6.94% and 22.74%, respectively. For *Tropilaelaps* infestation the province with the greatest prevalence was Uttaradit, followed by Phrae and Chon Buri. For *Varroa destructor* infestation, Loei had the highest prevalence, followed by Uttaradit and Nan. Chiyaphum, Phayao, Lumphun, and Nakorn Sawan saw no positive bee samples for either *Tropilaelaps* or *Varroa destructor* infestations. More details are presented in Table 2.

Figure 1 provides information about the characteristics of the apiaries analysed in the study. Of the 144 apiaries participating in the survey, 86 received a GAP certificate (59.72%). Just over half of the apiaries had moved within a month prior to the survey (56.20%). Approximately a quarter of the apiaries had used chemicals at some point prior to the survey (24.82%). Note that, for the history of apiary movement and chemical usage, we had complete information from only 137 apiaries.

Inferential Analysis

For *Tropilaelaps* infestations, a univariable analysis by the Chi square test revealed that 61.86% of disease-free apiaries had a GAP certificate; on the other hand, 55.32% of infested apiaries also had a GAP certificate. However, this percentage difference did not show any statistical

Table 2 Prevalence of Honeybee Parasitic Diseases by Provinces

Province	Apiary Prevalence		Beehive Prevalence	
	% of Infested Apiary (Positive Samples/Total Samples)		% of Infested Beehive (Positive Samples/Total Samples)	
	<i>Tropilaelaps</i> spp.	<i>Varroa</i> spp.	<i>Tropilaelaps</i> spp.	<i>Varroa</i> spp.
Chon Buri	100% (1/1)	0% (0/1)	12.50% (1/8)	0% (0/8)
Chiyaphum	0% (0/1)	0% (0/1)	0% (0/8)	0% (0/8)
Loei	0% (0/1)	100% (1/1)	0% (0/8)	62.50% (5/8)
Chiang Mai	22.41% (13/58)	50% (29/58)	4.31% (20/464)	23.28% (108/464)
Chiang Rai	43.75% (7/16)	50% (8/16)	7.81% (10/128)	20.31% (26/128)
Phrae	58.82% (10/17)	52.94% (9/17)	14.71% (20/136)	13.24% (18/136)
Nan	38.46% (10/26)	65.38% (17/26)	8.65% (18/208)	31.25% (65/208)
Phayao	0% (0/1)	0% (0/1)	0% (0/8)	0% (0/8)
Lumphun	0% (0/2)	0% (0/2)	0% (0/16)	0% (0/16)
Uttaradit	50% (2/4)	50% (2/4)	15.63% (5/32)	40.63% (13/32)
Phitsanulok	25% (3/12)	41.67% (5/12)	5.21% (5/96)	25% (24/96)
Nakorn Sawan	0% (0/1)	0% (0/1)	0% (0/8)	0% (0/8)
Chumphon	25% (1/4)	50% (2/4)	3.13% (1/32)	9.38% (3/32)
Total	32.64% (47/144)	50.69% (73/144)	6.94% (80/1,152)	22.74% (262/1,152)

significance. Likewise, there was no statistical significance found in apiary movement. The only variable that showed a statistically significant difference was the use of chemicals, with a COR of 2.27 and 95% CI of 1.02–5.04, [Table 3](#).

In terms of *Varroa destructor* infestations, apiary movement was more concentrated among disease-free apiaries than infested apiaries (67.16% versus 45.71%). This difference exhibited a statistical significance, as evidenced by the P-value of 0.01. Apiaries with a history of movement were about 59% less likely to experience *Varroa*

destructor infestation relative to those with static domiciles. Having a GAP certificate and the use of chemicals were also more common among disease-free apiaries compared with infested apiaries; both factors did not show a statistically significant difference, [Table 4](#).

The multivariable logistic regression demonstrated that chemical usage was positively correlated with *Tropilaelaps* infestations (AOR = 2.69 [95% CI = 1.16–6.21]) with statistical significance (P-value = 0.02). There was no statistically significant difference between apiary movement and *Tropilaelaps* infestations (AOR = 0.59 [95% CI




Figure 1 Characteristics of the participating apiaries.

Table 3 Association of Apiary Characteristics and *Tropilaelaps* infestation from Univariable Analysis

	Case	Non-Case	COR (95% CI)	P-value
GAP certificate (n = 144)	26/47 (55.32%)	60/97 (61.86%)	0.76 (0.38–1.55)	0.45
Apiary movement (n = 137)	22/45 (48.89%)	55/92 (59.78%)	0.64 (0.31–1.32)	0.23
Chemical usage (n = 137)	16/45 (35.56%)	18/92 (19.57%)	2.27 (1.02–5.04)	0.04

Table 4 Association of Apiary Characteristics and *Varroa destructor* Infestation from Univariable Analysis

	Case	Non-Case	COR (95% CI)	P-value
GAP certificate (n = 144)	40/73 (54.79%)	46/71 (64.79%)	0.66 (0.34–1.29)	0.22
Apiary movement (n = 137)	32/70 (45.71%)	45/67 (67.16%)	0.41 (0.21–0.82)	0.01
Chemical usage (n = 137)	16/70 (22.86%)	18/67 (26.87%)	0.81 (0.37–1.75)	0.59

= 0.28–1.25]). A similar result was found for GAP certificate (AOR = 0.58 [95% CI = 0.27–1.25]). For *Varroa destructor*, apiary movement showed a negative association with developing the disease as evidenced by an AOR of 0.40 (95% CI = 0.20–0.80) with a P-value of 0.01. GAP certificate and chemical usage exhibited the same direction of association with apiary movement. Statistical significance was not found in GAP certificate and chemical usage variables (P-value = 0.14 and 0.85, respectively) as shown in Tables 5 and 6.

Discussion

Overall, we found that most of the study samples were collected from the northern region of Thailand. The prevalence of *Tropilaelaps* infestations at the apiary level and at the beehive level was 32.64% and 6.94%, respectively, while the corresponding prevalence of *Varroa destructor* infestation was 50.69% and 22.74%, respectively. Considering the prevalence in each province, the highest apiary prevalence of *Tropilaelaps* infestation was in Chon Buri (100%), followed by Phrae (58.82%), and Uttaradit (50%). For *Varroa destructor* infestation, the provincial prevalence was greatest in Loei (100%), followed by Nan (65.38%), and Phrae (52.94%).

It appears the prevalence of *Tropilaelaps* and *Varroa destructor* infestations was quite high in most provinces in the northern region (such as Nan, Chiang Rai, and Phrae). This might be due to the region being an area where honeybee food sources are abundant (for instance, longan and lychee orchards). Phankaew also points to some specific behaviours of most orchard owners that put beehives under stress, such as the use of pesticides on orchards and prohibiting the movement of apiaries into orchards. Some longan-orchard owners do not allow beekeepers to move honeybee colonies into the orchards due to their belief that honeybees may cause longan flowers to drop and reduce the amount of longans grown.¹¹ This practice causes stress to the honeybees as a result of food shortage in the orchards and ultimately makes honeybees more vulnerable to diseases.

The possession of a GAP certificate appears to be a protective factor for parasitic diseases, although the strength of association did not show a statistical significance. It is possible that the possession of a GAP certificate ensures beekeepers follow good beekeeping practices by introducing prevention and control programmes against honeybee diseases and parasites in every stage of production. The requirements consist of strict beekeeping hygiene, regular cleaning of apiary and beekeeping

Table 5 Association of Apiary Characteristics and *Tropilaelaps* Infestation from Multivariable Analysis

	AOR	95% CI	P-value
GAP certificate (n = 144)	0.58	(0.27–1.25)	0.16
Apiary movement (n = 137)	0.59	(0.28–1.25)	0.17
Chemical usage (n = 137)	2.69	(1.16–6.21)	0.02

Table 6 Association of Apiary Characteristics and *Varroa destructor* Infestation from Multivariable Analysis

	AOR	95% CI	P-value
GAP certificate (n = 144)	0.58	(0.28–1.20)	0.14
Apiary movement (n = 137)	0.40	(0.20–0.80)	0.01
Chemical usage (n = 137)	0.93	(0.41–2.08)	0.85

equipment, and elimination of beehives suspected of being at risk of outbreak.¹² A study in Argentina by Giacobino et al supports this notion; it demonstrated that honeybee apiaries that do not follow standard management guidelines (like GAP) face a 3% greater risk of *Varroa destructor* infestation, compared with those following standard guidelines.¹³

In addition, apiary movement showed a favourable impact against *Varroa destructor* infestations, and this association presented a statistical significance. A possible explanation for this is that during the apiary movement process, the beekeepers usually extract honey from the honeycomb with a centrifuge machine. In this process, honeybee parasites are eliminated from beehives. The movement process also allows beekeepers to observe parasites and clean beehives. Moreover, the majority of beekeepers who move apiaries regularly are a group of commercial beekeepers that pay careful attention to the cleanliness of the beehives prior to movement.

Chemical use also demonstrated a statistically significant association with *Tropilaelaps* infestations, with the odds of infestation greater than one. A previous study by Chantawannakul et al reported that *Tropilaelaps* spp. has a short life cycle and rapid propagation. The use of chemicals over a long duration may increase the resistance of *Tropilaelaps* spp. towards these chemicals.¹⁴ However, chemical use seems to have some protective effect against *Varroa destructor* infestations, although without any statistical significance. Pettis suggested that many chemicals used by beekeepers (such as Amitraz) are likely to offer better protection against *Varroa destructor* infestations than *Tropilaelaps* infestations. However, because of the nature of *Tropilaelaps* spp. and *Varroa* spp. where they tend to remain in the combs and or in brood cells, some chemicals may not be sufficient to get rid of honeybee parasites.¹⁵ On the other hand, there is literature that supports the use of chemicals to protect beehives against parasitic diseases. A study by Haber et al suggested that varroacides, especially Amitraz, help reduce the loss of honey production during winter (winter losses).¹⁶

In terms of recommendations, we recommend that all apiaries in Thailand should follow the standards stipulated by the GAP certificate. This is supported by our studies where meeting the standards of a GAP certificate is likely to provide protective effects against parasitic diseases, although statistical significance was not shown at the 95% confidence level. However, if the confidence level is further relaxed (for instance, at the 80–85% confidence level), the protective effect of GAP certificate will be more obvious. In addition, beekeepers should be crucially aware of the risks and benefits of apiary movement and chemical use. In theory, chemicals may protect against parasites in honeybees but may create the risk of chemical resistance at the same time. Apiary movement enables the beekeepers to clean the beehives but it may cause further spread of disease to a new area in parallel.

This study contains both strengths and weaknesses. One of the strengths is the use of primary data collection on selected apiaries in all regions of Thailand. However, there are also some limitations. Firstly, the nature of a cross sectional study does not allow us to draw out strong evidence of a causal relationship between the factors of interest and parasitic diseases in honeybees. Regular data collection in the future that allows time-series analysis will be of great value. Secondly, identifying the presence of disease is largely subject to the competency of examiners; in this study, we employed more than one examiner but we provided a briefing session to all examiners prior to the fieldwork in order to ensure similar examination protocol in all apiaries. Lastly, as this study was the first national surveillance of honeybee diseases in Thailand, many variables that might be related to *Varroa destructor* and *Tropilaelaps* infestations were not collected from the outset such as apiary location, the cleaning process for each apiary, beekeeping practices, source of queen bees, the reason of chemical use (parasitic protection or elimination), density of beehive per harvest area, and the density of honeybee per beehive. This issue warrants further research to extend the academic value of beehive research in Thailand.

Conclusion

The prevalence of *Varroa destructor* and *Tropilaelaps* infestation at the apiary level was 50.69% and 32.64%, respectively. At the beehive level, the prevalence of *Varroa destructor* infestation was 22.74% while the prevalence of *Tropilaelaps* infestation was much smaller (6.94%). The northern region presented the highest

prevalence of *Varroa destructor* and *Tropilaelaps* infestation. An apiary that acquired a GAP certificate was less likely to face *Varroa destructor* and *Tropilaelaps* infestations, although a statistically significant difference was not exhibited. Apiaries that had a history of chemical use showed approximately three times greater odds of *Tropilaelaps* infestation. The odds of encountering *Varroa destructor* infestation amongst apiaries with a history apiary movement was about 60% lower than those without apiary movement. We recommend that all apiaries in Thailand follow the standards stipulated by the GAP certificate. In addition, beekeepers should be crucially aware of the risk and benefits of apiary movement and chemical use. Further studies to collect comprehensive information on apiaries, including information about cleaning processes, source of queen bees, and density of beehive per harvest area would be of great academic value.

Acknowledgments

This study was supported by the Department of Livestock Development under the Ministry of Agriculture and Cooperatives (Thailand). The authors would like to thank Mr Nopporn Tohmee from the National Institute of Animal Health (NIAH) and Mr Chairaj Pocharoen from the Livestock Administrative Region 5 for extremely helpful advice and immense support on this manuscript.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

The authors received funding for fieldwork from the Department of Livestock Development, Ministry of Agriculture and Cooperatives, Thailand. The publication fee was supported by the International Health Policy Program (IHPP), Ministry of Public Health, Thailand. IHPP received funding support from the Thailand Science Research and Innovation (TSRI) under the Senior Research Scholar on Health Policy and System Research [Contract No. RTA6280007].

Disclosure

The authors declare that there are no conflicts of interest.

References

1. Chantawannakul P, Wongsiri S, Chakpitak N. *Beekeeping Guideline*. Vol. 1. Chantawannakul P, editor. Chiang Mai: Chotana Print Company Limited; 2018.
2. Phanphum P. Report of honey production of Thailand 2014–2016. 2016. Available from: http://www.agriman.doae.go.th/home/news/year.2017/043_breed.bees.pdf. Accessed March 15, 2020.
3. World Organisation for Animal Health. Recommendations applicable to OIE Listed diseases and other diseases of importance to international trade. 2019. Available from: <https://www.oie.int/en/standard-setting/terrestrial-code/access-online/>. Accessed March 15, 2020.
4. Ministry of Agriculture and Cooperatives of Thailand. Notification of the Ministry of agriculture and cooperatives on the prescribing of additional animal epidemics under animal epidemic Act B.E.2558 (2015). *R Thai Gov Gaz*. 2015;132(special part 347 Ngor):10–15.
5. Shimanuki H, Knox DA. *Diagnosis of Honey Bee Diseases*. Beltsville, MD: Agricultural Research Service U.S. Department of Agriculture; 1991.
6. Ryabov EV, Childers AK, Chen Y, et al. Recent spread of *Varroa destructor* virus-1, a honey bee pathogen, in the United States. *Sci Rep*. 2017;7(1). doi:10.1038/s41598-017-17802-3
7. Center of EIA Burapha University. What happens if bees disappear from the earth? 2019. Available from: <https://www.eiaborapha.com/site/2019/04/26/>. Accessed March 15, 2020.
8. Food and Agriculture Organization of United Nations. Declining bee populations pose threat to global food security and nutrition. 2019. Available from: <http://www.fao.org/news/story/en/item/1194910/icode/>. Accessed March 15, 2020.
9. Ratananokorn L, Neramitmansook W, Wongpakorn M. Survey of bee diseases, chemical and drug residues in honey and bee products in Thailand. *Khon Kaen Agric J*. 2008;36:123–130.
10. Shackman G Sample size and design effect. 2001. Available from: <http://faculty.smu.edu/slstokes/stat6380/deff.doc.pdf>. Accessed March 15, 2020.
11. Phankaew C. Apiculture and pollinator industry survey in Thailand. *Int J Agric Ext*. 2016;04(02):95–103.
12. Ministry of Agriculture and Cooperatives of Thailand. Notification of the Ministry of agriculture and cooperatives on the standardization of agricultural products: good agricultural practices for bee farms under Agricultural standards act B.E. 2551 (2008). *R Thai Gov Gaz*. 2016;133(special part 106 Ngor):12.
13. Giacobino A, Cagnolo NB, Merke J, et al. Risk factors associated with the presence of *Varroa destructor* in honey bee colonies from east-central Argentina. *Prev Vet Med*. 2014;115(3–4):280–287. doi:10.1016/j.prevetmed.2014.04.002
14. Chantawannakul P, Natapot W, Khongpinitbunjong K, Buawangpong N, Chaimanee V, Sirisa-ard P. Parasitic mites and nose mites in bees. Chiang Mai: Max Printing; 2011.
15. Pettis JS, Rose R, Chaimanee V, Hong XY. Chemical and cultural control of *Tropilaelaps mercedesae* mites in honeybee (*Apis mellifera*) colonies in Northern Thailand. *PLoS One*. 2017;12(11):e0188063. doi:10.1371/journal.pone.0188063
16. Haber AI, Steinhauer NA, vanEngelsdorp D. Use of chemical and nonchemical methods for the control of *Varroa destructor* (Acari: varroidae) and associated winter colony losses in U.S. beekeeping operations. *J Econ Entomol*. 2019;112(4):1509–1525. doi:10.1093/jee/toz088
17. Office of Agricultural Economics of Thailand. Land utilization for agriculture by province in Thailand, 2017. 2017. Available from: <http://www.oae.go.th/assets/portals/1/files/Land.Utilization2560.pdf>. Accessed March 12, 2021.

Veterinary Medicine: Research and Reports

Dovepress

Publish your work in this journal

Veterinary Medicine: Research and Reports is an international, peer-reviewed, open access journal publishing original research, case reports, editorials, reviews and commentaries on all areas of veterinary medicine. The manuscript management system is completely online

and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/veterinary-medicine-research-and-reports-journal>