

# Effects of Music, Massage, Exercise, or Acupuncture in the Treatment of Depression Among College Students: A Network Meta-Analysis

Fang-bo Li<sup>1,\*</sup>, Ping Lu<sup>2,\*</sup>, Hai-tao Wu<sup>1</sup>, Ming-han Wang<sup>1</sup>, Jing-dong Wang<sup>1</sup>

<sup>1</sup>Graduate School, Guangzhou Sport University, Guangzhou, Guangdong, 510500, People's Republic of China; <sup>2</sup>School of PE and Health, Guangzhou University of Chinese Medicine, Guangzhou, Guangdong, 510006, People's Republic of China

\*These authors contributed equally to this work

Correspondence: Fang-bo Li, Graduate School, Guangzhou Sport University, Linhe West Road, Tianhe District, Guangzhou, Guangdong, 510500, People's Republic of China, Tel +86 18860292906, Email 975394244@qq.com

**Objective:** To assess the therapeutic impacts of exercise, massage, and music interventions on college students experiencing depression by employing a mesh meta-analysis approach. This research intends to offer valuable insights to aid in the development of non-pharmaceutical treatment strategies for depression.

**Methods:** We conducted a thorough search across various databases including Cochrane, PubMed, Embase, Web of Science, CNKI, and Wanfang to explore the effects of music, massage, aerobic exercise, fitness Qigong, yoga, tai chi, ball games, strength training, dance, whole body vibration training, and high-intensity interval training on the treatment of depression in college students. The search period was from January 1, 2023, which marks the establishment of each database. Subsequently, a mesh meta-analysis was performed using the “Stata 15.1” software, incorporating outcome indicators from 24 included literature comprising a total of 1458 patients.

**Results:** Based on the ranking of the optimal intervention effects of various non-pharmaceutical methods, the order, from highest to lowest probability, was as follows: high-intensity interval training (96%), yoga (94.90%), dance (78.30%), music (73.30%), ball games (62.50%), strength training (51.70%), aerobic training (45.30%), tai chi (35.40%), vibration training (27.30%), massage (20.10%), qigong (14.30%), and no intervention (1.00%). This ranking aligns closely with the findings obtained from pairwise comparisons between different interventions.

**Conclusion:** High-intensity interval training is likely to yield the most effective therapeutic results for college students with depression. In the pairwise comparison of different interventions, High-intensity interval training is also better than most interventions. However, to establish its intervention effect more conclusively, further validation through additional high-quality randomized controlled trials is necessary.

**Keywords:** depression, music, exercise, college student, network meta-analysis

## Introduction

Depression is a common and serious mental health concern among college and university students.<sup>1</sup> The 2019 Global Burden of Illness, Injury, and Risk Factors study showed that depression is one of the most disabling mental disorders, and among adolescents, depression ranked fourth on the list of the world's leading causes of burden in 2019.<sup>2</sup> Studies have shown that depression has become a major emotional problem among college students. College students in their early adulthood are prone to emotional dysregulation and despair.<sup>3</sup> In 2021, China's Ministry of Education requested for the first time that depression screening be included among students' physical examinations, demonstrating that the harm of depression in college students has piqued the state and society's attention.<sup>4</sup> In this context, how to provide effective non-drug treatment to improve depressive symptoms for patients with depression has attracted the attention and research of many researchers. Non-drug treatment methods for depression are currently exercise intervention, music intervention,

meditation, acupuncture, massage, and so on. Studies have found that aerobic exercise, traditional Chinese sports (fitness qigong, tai chi, etc.), and meditation can effectively improve the depressive symptoms of college students.<sup>1</sup> Music, massage, and acupuncture have also been proven to be effective in the treatment of depression.<sup>5–9</sup> Several interventions have been evaluated individually or in comparison using randomized controlled trials.<sup>5,10–13</sup> However, there are only a few studies comparing the efficacy of multiple non-pharmacological interventions in network meta-analysis. Therefore, four interventions (exercise, music, massage, and acupuncture) that are regularly utilized in the treatment of depression among college students were selected as research objects. The network meta-analysis method was used to integrate the relevant clinical evidence of direct and indirect comparison. The different interventions of the same evidence were summarized, quantitatively synthesized, and analyzed, the probability was ranked according to the efficacy of the index, and the best therapy was screened.

## Materials and Methods

### Literature Searching and Selection of Articles

Two investigators independently conducted a literature search. The databases were searched until 1 January 2023. The keywords utilized for the searches were “college student depression”, “exercise”, “dance”, “yoga”, “tai chi”, “qi gong”, “music”, “massage” and “acupuncture”. Studies in the Cochrane, PubMed, Embase, Web of Science, CNKI, and Wanfang databases that were written in Chinese or English and were related to the study were retrieved. The terms were grouped into three categories (intervention, outcome measure, and intervention), connected by Boolean logical operators (OR or AND), and applied to three search levels. The retrieved papers were initially screened using the bibliographic management software Endnote, such as deduplication, reading titles, and abstracts and the remaining studies were examined in detail, according to the PICOS set appropriate inclusion criteria.

The following were the research inclusion criteria: (1) The participants of the study were specifically college students who were experiencing depression. It is important to note that the study focused on undergraduate students, excluding high school students; (2) experimental interventions related to exercise, music, massage, and acupuncture; (3) the outcome indicators were the depression scores assessed by the scale, and the before and after indicators were reported in detail, such as mean, standard deviation, sample size, etc. to generate effect sizes; (4) the research type is two-arm experimental research; (5) the baseline data are comparable.

The following were the research exclusion criteria: (1) The literature is in the form of reviews, conference abstracts, case reports, etc.; (2) the data reported in the literature were incomplete, the contact authors were unsuccessful, and the original data could not be obtained; (3) the baseline data is incomplete; and (4) the intervention target receives drug intervention at the same time in addition to several interventions prescribed.

After determining the criteria for inclusion and exclusion, two investigators began screening the literature at the same time, compared the findings of the inclusion of their selected literature, and the third investigator moderated the disagreeable material.

### Assessment of Methodological Quality

The PEDro scale developed by the Center for Evidence-Based Medicine in Physical Therapy was used to assess the quality of the literature.<sup>14</sup> The scale comprises 11 items, including subjects, allocation concealment, blinding, and other main content, and the score  $\leq 3$  is poor quality; the score in 4–5 is medium quality; the score in 6–8 is good quality; and the score of 9 points is high quality. At the same time, the Cochrane risk bias assessment tool was employed to assess the quality of the included studies. The evaluation primarily focused on seven aspects, which included random sequence generation, allocation concealment, blinding of participants, blinding of outcome variable evaluation, loss of outcome data, selective reporting, and other potential biases.

### Data Extraction and Statistical Analysis

For the included indices, the mean, standard deviation, and sample size of the before-and-after test data were extracted from each experimental group and control group. The analytical data utilized the change from baseline as the basis for

comparison. The effect size of different types of intervention on depressive symptoms in college students was explored through a network meta-analysis.

The statistical analysis was performed using the Stata 15.1 software for network meta-analysis. The effect size of the measured data was expressed in weighted mean difference (WMD) or standardized mean difference (SMD) with its 95% confidence interval (CI). In cases where the units, measures, or rating scales of the extracted data were not consistent, SMD was utilized to represent the effect size. If the 95% CIs did not include zero, it indicated a statistically significant difference between the two treatments. Conversely, if the difference fell within the confidence interval, it was considered not statistically significant.<sup>15</sup> The surface under the cumulative ranking (SUCRA) was used to represent the preferred ranking of each treatment measure.<sup>16</sup> A larger area under curve indicates a greater probability that the psychotherapy will be the best treatment. Funnel plots were used to indicate publication bias, and the symmetry of the funnel plots was used to assess publication bias between included studies. The pooled effect size of each included study was a study in which the x-axis intersected on the funnel plot as a perpendicular line, and the points distributed to the left of the perpendicular line represented studies with fewer effect sizes than pooled effect sizes; the dots distributed to the right of the perpendicular line represent studies in which the effect size is greater than the combined effect size: the number of dots on both sides is essentially the same, indicating no publication bias, and conversely, there is publication bias.

## Results

### Literature Search results


We searched Cochrane, PubMed, Embase, Web of Science, CNKI, and Wanfang databases for references 102, 409, 544, 696, 211, and 907. The screening was carried out in accordance with the established inclusion and exclusion criteria. A total of 2845 irrelevant articles were removed, leaving only 24 for evaluation. The selection process is shown in [Figure 1](#).

### Characteristics of the Studies

The 24 included papers came from 5 different countries and were divided into Randomized controlled trial and Controlled trial by experimental category, with 17 randomized controlled trials and 7 non-randomized controlled trials. The intervention targeted all college students with varying degrees of depression symptoms, with a total of 1458 people and about 63% of them being women. After data screening, the experimental group contained 117 cases of music intervention, 21 cases of massage, 98 cases of aerobic exercise, 15 cases of qigong, 61 cases of yoga, 97 cases of Tai Chi, 230 cases of ball games, 15 cases of strength training, 101 cases of dance, 15 cases of whole-body vibration training, and 14 cases of high-intensity interval training (The category “movement” is refined according to its type). Regarding the duration of interventions, one study lasted for two weeks, three studies for four weeks, one study for six weeks, seven studies for eight weeks, one study for ten weeks, seven studies for twelve weeks, and four studies for sixteen weeks. The outcome measures utilized scales to assess the levels of positive response and depression, including SDS, SCL-90, DERS, BDI, PHQ-9, CES-D, among others (refer to [Table 1](#)).

### Quality Assessment Results

[Table 2](#) presents the details of the Quality assessment results, which included a total of 24 articles. These articles were rated on the PEDro scale, with one article scoring 9, fifteen articles scoring 7, and eight articles scoring 6. All the included studies satisfied the eligibility criteria, which encompassed factors such as source credibility, adequate follow-up (85% compliance), intention-to-treat analysis, intergroup statistical comparisons, reporting of point measurements, and variability measurements. Given that the studies employed various self-assessment scales to measure depression, the majority did not employ double-blind trials or blinded outcome evaluators. When considering the results of the Cochrane system evaluation tool, it can be observed that the included literature generally meets the quality requirements, although there is a scarcity of high-quality studies (see [Figure 2](#)).



**Figure 1** Study selection process.

## Results of the Network Meta-Analysis

### Network Evidence Map of the Impact of Different Interventions on Depression

In the network evidence map, dots indicate the sample size of the intervention, with larger dots indicating more sample size included in the intervention; lines represent direct comparisons of interventions, with thicker lines suggesting a higher number of inclusions for direct comparisons of the two interventions. The network evidence map of the efficacy of depression involved 11 treatment measures: music, massage, aerobic exercise, qigong, yoga, tai chi, ball games, strength exercise, dance, whole body vibration training, and high-intensity interval training. The network evidence plot for depression scores showed that direct or indirect comparisons between different psychological interventions were essential for network meta-analysis. A network plot of evidence for the efficacy of depression is shown in [Figure 3](#).

Table I Research Characteristics Table

No	Author, Years	Country	Study Design	Population Characters	Number Analysed at End of Treatment (N)	Classification	Duration	Main Outcomes	Reference
1	Zhang et al, 2014	China	Controlled trial	College students with depression; Gender: 36m/56w	82 (50/32)	Five Elements Music of Traditional Chinese Medicine	6 weeks, 2 times a week	SDS; SCL-90	[17]
2	Ming Zhang et al, 2022	China	RCT	College students with DERS $\geq$ 101, BDI score $\geq$ 10, and depressive symptoms; Gender: 34m/37w	71 (36/35)	Group improvisational music therapy (GIMT)	4 weeks, 1 times a week	DERS;BDI	[3]
3	Hye-Kyung Lee et al, 2014	Korean	Non equivalent control group design	College students with PHQ-9 score $>$ 5; Gender: 12m/29w	41 (21/20)	Self-Aromatherapy Massage	2 weeks, 3 times a week	PHQ-9;	[7]
4	Cheng Xiang et al, 2016	China	RCT	Mildly depressed college students; Gender: 15m/15w	30 (15/15)	Health Qigong Wuqinxi	12 weeks, 3 times a week	BDI	[18]
5	Nasrin Falsafi et al, 2016	USA	RCT	Students who had a diagnosis of depression or anxiety	90 (30/30/30)	Yoga	8 weeks, 75 minutes a week	BDI	[19]
6	Jiayuan Zhang et al, 2018	China	RCT	Freshman and sophomore had subthreshold symptoms of depression; Gender: 21m/41w	62 (32/30)	Mindfulness-based Tai Chi Chuan (MTCC)	8 weeks, 2 times a week	PHQ-9;	[20]
7	Chen et al, 2019	China	RCT	Moderate depression [CES-D score 24–29] female college students	36 (18/18)	Taijiquan exercise	16 weeks, 3 times a week	CES-D	[21]
8	Duan et al, 2014	China	Controlled trial	College students with SDS score $\geq$ 53; Gender: 61m/48w	109 (30/41/38)	Basketball; sports dance; badminton	12 weeks, 3 times a week	SDS	[22]
9	Ma et al, 2017	China	RCT	College students with depression and anxiety scores exceeding the critical value; Gender: 59m/65w	124 (31/31/31/31)	Basketball; dance; badminton (moderate-intensity)	12 weeks, 3 times a week	SDS	[23]

(Continued)

Table I (Continued).

No	Author, Years	Country	Study Design	Population Characters	Number Analysed at End of Treatment (N)	Classification	Duration	Main Outcomes	Reference
10	Wang et al, 2016	China	Controlled trial	College Students with positive anxiety and depression scores;	98 (39/22/37)	Badminton (Singles; doubles); volleyball	16 weeks, 3 times a week	SDS	[24]
11	Li et al, 2017	China	RCT	Gender: 58m/40w College students with depression factor score>3;	64 (32/32)	Volleyball and group psychological guidance	8 weeks, 4times a week	SDS	[25]
12	Mao et al, 2014	China	RCT	Gender: 33m/31w Male college students with mild depression	29 (15/14)	Tennis with different intensity	12 weeks, 3 times a week	SDS	[26]
13	Zhang et al, 2012	China	Controlled trial	College students with mild and moderate depression;	120 (40/40/40)	Sports dance training group, basketball training group and table tennis training group	16 weeks, 3 times a week	SDS	[27]
14	Guan et al, 2008	China	RCT	Gender: 43m/77w Female students with low to moderate levels of depression (CES-D score is 16–29)	30 (15/15)	Progressive Strength Training	8 weeks, 3 times a week	CES-D	[28]
15	Li et al, 2014	China	RCT	Female college students with depression symptoms	45 (15/15/15)	Strength training; aerobic exercise	8 weeks, 3 times a week	CES-D	[29]
16	Tang et al, 2009	China	RCT	College students with moderate and mild depression	104 (39/34/31)	Basketball, badminton, etc; Yoga, aerobics, etc	12 weeks, 3 times a week	SDS	[30]
17	Jiao et al, 2010	China	RCT	College students with depression; Gender: 18m/26w	44 (21/23)	Aerobic Exercise	8 weeks, 3 times a week	SDS	[31]
18	Duk, Oh Sang et al, 2012	Korean	RCT	College students with depressive symptoms; Gender: 0m/16w	16 (8/8)	Moderate aerobic exercise	12 weeks, 3 times a week	BDI	[32]

19	Geetanjali Chawla et al, 2022	India	RCT	College students with Depression, Anxiety Stress Scale (DASS) (depression > 10); Gender: 9m/21w	30 (15/15)	Whole-body vibration (WBV) training	4 weeks, 2 times a week	DASS scores (Depression; Anxiety; Stress)	[33]
20	Zhao et al, 2017	China	RCT	College students with SDS score≥53;Gender:0m/43w	43 (14/14/15)	High intensity intermittent exercise group (HIT); Continuous exercise group	4 weeks, 3 times a week	SDS	[34]
21	Li et al, 2017	China	Controlled trial	College students with SDS score≥50;Gender:15m/23w	38 (20/18)	Taijiquan exercise	12 weeks, 3 times a week	SDS	[35]
22	Song et al, 2017	China	Controlled trial	College students with depression; Gender: 0m/51w	51 (27/24)	Taijiquan exercise, Aerobic Exercise	16 weeks, 3 times a week	BDI	[36]
23	Khirollah Sadeghi et al, 2016	Iran	RCT	Undergraduate students with BDI score>13; Gender: 24m/6w	30 (16/14)	Aerobic Exercise	8 weeks, 2 times a week	BDI	[37]
24	Chen-Jung Chen et al, 2014	China Taiwan	RCT	Nursing students with DMSRIA scores of 8points or higher; Gender: 2m/69w	71 (31/40)	Five Elements Music of Traditional Chinese Medicine	10weeks, 2 times a week	Depression Mood Self Report Inventory for Adolescence (DMSRIA)	[38]

**Notes:** The scores of several scales used in the included study are positively correlated with the degree of depression, that is, the higher the score, the more severe the degree of depression.

**Table 2** The Physiotherapy Evidence Database (PEDro) Scores for the Articles Included

Publication	Item on the PEDro Scale											Score
	a	b	c	d	e	f	g	h	i	j	k	
Zhang et al, 2014 <sup>17</sup>	1	0	0	1	0	0	0	1	1	1	1	6
Ming Zhang et al, 2022 <sup>3</sup>	1	1	0	1	0	0	0	1	1	1	1	7
Hye-Kyung Lee et al, 2014 <sup>7</sup>	1	0	0	1	0	0	0	1	1	1	1	6
CHENG Xiang et al, 2016 <sup>18</sup>	1	1	0	0	0	0	0	1	1	1	1	6
Nasrin Falsafi et al, 2016 <sup>19</sup>	1	1	0	1	0	0	0	1	1	1	1	7
Jiayuan Zhang et al, 2018 <sup>20</sup>	1	1	1	1	0	0	1	1	1	1	1	9
Chen et al, 2019 <sup>21</sup>	1	1	0	1	0	0	0	1	1	1	1	7
Duan et al, 2014 <sup>22</sup>	1	0	0	1	0	0	0	1	1	1	1	6
Ma et al, 2017 <sup>23</sup>	1	1	0	1	0	0	0	1	1	1	1	7
Wang et al, 2016 <sup>24</sup>	1	0	0	1	0	0	0	1	1	1	1	6
Li et al, 2017 <sup>25</sup>	1	1	0	1	0	0	0	1	1	1	1	7
Mao et al, 2014 <sup>26</sup>	1	1	0	1	0	0	0	1	1	1	1	7
Zhang et al, 2012 <sup>27</sup>	1	0	0	1	0	0	0	1	1	1	1	6
Guan et al, 2008 <sup>28</sup>	1	1	0	1	0	0	0	1	1	1	1	7
Li et al, 2014 <sup>29</sup>	1	1	0	1	0	0	0	1	1	1	1	7
Tang et al, 2009 <sup>30</sup>	1	1	0	1	0	0	0	1	1	1	1	7
Jiao et al, 2010 <sup>31</sup>	1	1	0	1	0	0	0	1	1	1	1	7
Duk, Oh Sang et al, 2012 <sup>32</sup>	1	1	0	1	0	0	0	1	1	1	1	7
Geetanjli Chawla et al, 2022 <sup>33</sup>	1	1	0	1	0	0	0	1	1	1	1	7
Zhao et al, 2017 <sup>34</sup>	1	1	0	1	0	0	0	1	1	1	1	7
Li et al, 2017 <sup>35</sup>	1	0	0	1	0	0	0	1	1	1	1	6
Song et al, 2017 <sup>36</sup>	1	0	0	1	0	0	0	1	1	1	1	6
Khirollah Sadeghi et al, 2016 <sup>37</sup>	1	1	0	1	0	0	0	1	1	1	1	7
Chen-Jung Chen et al, 2014 <sup>38</sup>	1	1	0	1	0	0	0	1	1	1	1	7

**Notes:** Item on the PEDro scale (a: Eligibility criteria and source; b: Random allocation; c: Concealed allocation; d: Baseline comparability; e: Blinding of participants; f: Blinding of therapists; g: Blinding of assessors; h: Adequate follow-up (. 85%); i: Intention-to-treat analysis; j: Between-group statistical comparisons; k: Reporting of point measures and measures of variability).

## A Network Meta-Analysis of Efficacy in Depression

Twenty-four studies provided data on depression scores among college students following various treatments. However, the original studies used different depression rating scales, necessitating the use of SMD and its 95% CI to indicate the magnitude of the effect on depression scores. It is important to note that all depression scores on the scales reflected the level of depression positively, meaning that a lower effect size corresponds to a more favorable effect on depressive symptoms. Pairwise comparisons of depression scores by treatment revealed the following results: high-intensity interval training was superior to vibration training [SMD = -13.61, 95% CI (-20.61 to -6.62)]; high-intensity interval training was superior to dance [SMD = -8.13, 95% CI (-13.44 to -2.83)]; high-intensity interval training was superior to strength training [SMD = -10.81, 95% CI (-16.77 to -4.85)]; high-intensity interval training was better than ball games [SMD = -9.39, 95% CI (-14.65 to -4.13)]; high-intensity interval training was better than Tai Chi [SMD = -12.62, 95% CI (-17.81 to -7.43)]; high-intensity interval training was better than qigong [SMD = -14.81, 95% CI (-20.04 to -9.59)]; high-intensity interval training was better than aerobic training [SMD = -11.60, 95% CI (-16.72 to -6.49)]; high-intensity interval training was better than music [SMD = -8.52, 95% CI (-13.80 to -3.25)]; high-intensity interval training was superior to massage [SMD = -14.23, 95% CI (-19.61 to -8.86)]; high-intensity interval training was superior to no intervention [SMD = -16.61, 95% CI (-21.78 to -11.45)].

In the comparison of vibration training intervention with other treatments: vibration training intervention was inferior to dance [SMD = 5.48, 95% CI (0.60 to 10.37)]; vibration training intervention was inferior to yoga [SMD = 13.10, 95% CI (7.68 to 18.51)]; vibration training intervention was inferior to music [SMD = 5.09, 95% CI (0.24 to 9.94)]. For dance compared to other treatments: dance was superior to ball games [SMD = -1.26, 95% CI (-2.13 to -0.38)]; dance was






Figure 2 Results of Cochrane system evaluation.




Figure 3 Network analysis of depression efficacy. The numbers represent the quantity after intervention measures.

superior to Tai Chi [SMD = -4.49, 95% CI (-5.93 to -3.05)]; dance was superior to yoga [SMD = 7.61, 95% CI (4.77 to 10.46)]; dance was superior to qigong [SMD = -6.68, 95% CI (-8.16 to -5.20)]; dance was superior to aerobic training [SMD = -3.47, 95% CI (-5.64 to -1.30)]; dance was superior to massage [SMD = -6.10, 95% CI (-8.04 to -4.16)]; dance was superior to no intervention [SMD = -8.48, 95% CI (-9.72 to -7.25)].

In the comparison of strength training with other treatments: strength training was superior to yoga [SMD = 10.30, 95% CI (6.31 to 14.28)]; strength training was superior to qigong [SMD = -4.00, 95% CI (-7.09 to -0.91)]; strength training was superior to massage [SMD = -3.42, 95% CI (-6.75 to -0.09)]; strength training was superior to no intervention [SMD = -5.80, 95% CI (-8.78 to -2.82)]. For ball games compared to other treatments: ball games were better than Tai Chi [SMD = -3.23, 95% CI (-4.48 to -1.98)]; ball games were inferior to yoga [SMD = 8.87, 95% CI

(6.13 to 11.61)]; ball games were better than qigong [SMD = -5.42, 95% CI (-6.72 to -4.13)]; ball games were better than aerobic training [SMD = -2.21, 95% CI (-4.27 to -0.16)]; ball games were better than massage [SMD = -4.84, 95% CI (-6.64 to -3.05)]; ball games were better than no intervention [SMD = -7.22, 95% CI (-8.23 to -6.22)].

In the comparison of Tai Chi with other treatments: Tai Chi was inferior to yoga [SMD = 12.10, 95% CI (9.36 to 14.84)]; Tai Chi was better than qigong [SMD = -2.19, 95% CI (-3.29 to -1.09)]; Tai Chi was inferior to music [SMD = 4.10, 95% CI (2.79 to 5.40)]; Tai Chi was superior to no intervention [SMD = -3.99, 95% CI (-4.73 to -3.25)]. For yoga compared to other treatments: yoga was superior to qigong [SMD = -14.30, 95% CI (-17.06 to -11.54)]; yoga was superior to aerobic training [SMD = -11.08, 95% CI (-14.27 to -7.90)]; yoga was better than music [SMD = -8.00, 95% CI (-10.85 to -5.16)]; yoga was superior to massage [SMD = -13.72, 95% CI (-16.75 to -10.68)]; yoga was superior to no intervention [SMD = -16.10, 95% CI (-18.73 to -13.46)].

In the comparison of qigong with other treatments: qigong was inferior to aerobic training [SMD = 3.21, 95% CI (1.25 to 5.18)]; qigong was inferior to music [SMD = 6.29, 95% CI (4.94 to 7.64)]; qigong was superior to no intervention [SMD = -1.80, 95% CI (-2.61 to -0.99)]. For aerobic training compared to other treatments: aerobic training was inferior to music [SMD = 3.08, 95% CI (0.99 to 5.17)]; aerobic training was superior to massage [SMD = -2.63, 95% CI (-4.96 to -0.30)]; aerobic training was superior to no intervention [SMD = -5.01, 95% CI (-6.80 to -3.22)].

In the comparison of music with other treatments: music was superior to massage [SMD = -5.71, 95% CI (-7.55 to -3.87)]; music was superior to no intervention [SMD = -8.09, 95% CI (-9.17 to -7.01)]. For massage compared to no intervention: massage was better than no intervention [SMD = -2.38, 95% CI (-3.87 to -0.89)]. In all the comparisons mentioned above, none of the 95% confidence intervals included 0, indicating that the differences were statistically significant. The results of the network meta-analysis of depression scores are shown in Figure 4.

### Ranking of Efficacy

Efficacy ranking: The probability that an intervention will be the best measure is assessed using the area under the cumulative ranking probability plot (SUCRA), with larger areas under the curve indicating a greater likelihood that the intervention will be the best measure. SUCRA results showed that the interventions were ranked in order of efficacy: interval training (96%) > yoga (94.90%) > dance (78.30%) > Music (73.30%) > ball games (62.50%) > strength training (51.70%) > aerobic training (45.30%) > Tai Chi (35.40%) > vibration training (27.30%) > massage (20.10%) > Qigong (14.30%) > no intervention (1.00%), indicating that the probability of interval training is the best treatment for depression in college students. The results of the preferred ranking of various treatment measures for depression are shown in Table 3.





Figure 4 Meta-analysis results of depression score rates.

**Table 3** Probability: Ranking of Different Mind-Body Exercise Therapies to Improve Depression Grade in College Students


Treatment	SUCRA	MeanRank	No
Interval training	96%	1.4	1
Yoga	94.90%	1.6	2
Dance	78.30%	3.4	3
Music	73.30%	3.9	4
Ball	62.50%	5.1	5
Strength exercise	51.70%	6.3	6
Aerobic exercise	45.30%	7	7
Taichi	35.40%	8.1	8
Vibration training	27.30%	9	9
Massage	20.10%	9.8	10
Qigong	14.30%	10.4	11
Placebo	1.00%	11.9	12

### Subgroup Analysis of Different Types of Exercise Interventions

We further categorized and grouped different sports interventions and conducted a conventional meta-analysis based on four subgroups: aerobic exercise, traditional Chinese sports, ball games, and high-intensity interval sports. The results are depicted in Figure 5, where 1, 2, 3, and 4 represent aerobic sports, traditional Chinese sports, ball games, and High-intensity interval training, respectively. In terms of effectiveness, High-intensity interval training (SMD=-2.90, [95% CI: -3.96, -1.84]) exhibited the highest effect, followed by ball games (SMD=-2.05, [95% CI: -2.40, -1.70]), Chinese traditional sports (SMD=-1.80, [95% CI:



**Figure 5** Subgroup analysis. Number: 1, aerobic sports;<sup>31,32,34,37</sup> 2, traditional Chinese sports;<sup>18-21,30,35</sup> 3, ball games;<sup>23,25,30</sup> 4, high-intensity interval sports.<sup>34</sup>



**Figure 6** Calibration comparison funnel diagram. Capital: A, Placebo; B, Music; C, Massage; D, aerobic sports; E, Qigong; F, Yoga; G, Tai chi; H, Ball games; I, strength exercise; J, Dance; K, whole body vibration training; L, high-intensity interval sports.

−2.08, −1.53]), and aerobic sports (SMD=−1.31, [95% CI: −1.73, −0.90]). Heterogeneity testing revealed an I2 value of 75.5%, possibly attributed to factors such as intervention duration and sample size.

### Small Sample Effects or Publication Bias Tests

For the studies included in the network META, small sample effect estimation and publication bias test were performed by correcting the comparison funnel map. The included papers were basically symmetrical, indicating that the possibility of a small sample effect in this analysis was small, and no obvious publication bias was found. (Figure 6) Additionally, an Egger's test was performed on the included studies, yielding a P-value of 0.10, which is greater than 0.05. This suggests a low likelihood of significant publication bias.

## Discussion

Depression in young people is a serious mental health issue all over the world. In the treatment of depression, non-pharmacological interventions do not have problems such as drug resistance and side effects compared to pharmacological interventions.<sup>39</sup> A study found that after 4 months of experimentation, the exercise therapy group could achieve the same antidepressant effect as drug therapy (sertraline).<sup>40</sup> Meanwhile, physical and mental interventions, particularly exercise interventions, are rarely associated with mental illness stigma.<sup>41</sup> Therefore, this review conducted a network meta-analysis of experimental studies of commonly used non-pharmacological interventions for depression in college students, including music interventions, massage, aerobic exercise, qigong, yoga, Tai Chi, ball games, strength training, dance, whole-body vibration training, and high-intensity interval training. The study synthesized information from direct and indirect comparisons of various interventions, compared several interventions, and ranked the effectiveness of various non-pharmacological therapies based on the analysis results. The possibility of becoming the best treatment was explored.

The findings revealed that the 11 non-pharmacological interventions included were beneficial in reducing depression in college students, with high-intensity interval training having the highest probability of having the best effect. The effect of the intervention was statistically significant. According to scholar Zhao Hong-qin, while both gradual high-intensity interval exercise and continuous exercise can reduce depression in college girls, gradual high-intensity interval

exercise has a better positive impact. High-intensity interval exercise, as compared to continuous aerobic exercise, can significantly increase the pleasure of exercise, thereby more effectively stimulating the pituitary-adrenal axis and improving depressive conditions.<sup>34</sup> From the perspective of the mechanism of action, exercise can increase the content of relevant neurotransmitters in the blood of depressed patients and increase the efficiency of transmitters in the synaptic space of the central nervous system. The efficacy of most antidepressants in treatment is also to enhance the level of amine metabolism in the brain, indicating that the mechanisms of action of exercise and antidepressants are similar.<sup>31</sup> Antidepressants are currently widely used, but their clinical efficacy may be affected by poor adherence to medication regimens, which often have some side effects, or it takes up to 6 weeks after initiation to begin to have measurable effects on depressive symptoms; thereby, non-pharmacological treatments such as high-intensity interval training and yoga can be used as primary interventions. The results of SUCRA showed that in addition to the included exercise-based interventions, music interventions also rated highest. Michalak et al concluded that listening to appropriate and favorite music reduced participants' sympathetic nervous system activity while increasing parasympathetic nerve activity, resulting in peaceful breathing, stable blood pressure and heart rate, and the effect of sedation and hypnosis in the release of endorphin substances in the body, thereby significantly reducing anxiety and depression.<sup>42</sup> In addition, massage and Qigong are given low priority, with massage being regarded by researchers to be a synergistic measure to complement aromatic environmental therapy, and the results need to be validated by further studies due to the limited number of participants.<sup>7</sup> Furthermore, qigong and Tai Chi are often classified as traditional Chinese fitness exercises in research.<sup>43,44</sup> In terms of exercise characteristics, it falls within the category of moderate and low-intensity aerobic exercise.<sup>1</sup> According to the findings of this study, the low priority of traditional fitness exercises in China may be due to the relatively low intensity of exercise and the inability to provide suitable stimulation for young people to enhance their psychological condition.

There are still several limitations to this study: (1) From the perspective of study characteristics, the degree of depression of the subjects included in this study was different, and there were certain regional environments and genders. Discrepancies may lead to inconsistency between studies. (2) From the perspective of study design, the intervention period, intensity, frequency, and time of different study designs were also slightly different. (3) From the perspective of literature screening, although this study conducted a comprehensive search of two major domestic databases and four major foreign databases, there was still the possibility of missing detection. At the same time, the number of articles on different interventions varied relatively significantly in the included literature. For example, there is only one paper on massage, which makes objectively explaining its intervention impact on college students with depression difficult, and the acupuncture therapy included in the pre-search was rejected during literature screening, which is regrettable. Additionally, a comparison of the measures is missing from the final result.

(4) There is a potential for "reporting bias" since a research protocol was not registered for this study. (5) Other possible biases should also be considered. Given the aforementioned limitations, it is suggested that future RCT studies: (1) RCT design and reporting should follow international standards in order to increase the quality of research and reporting; (2) RCTs directly compared the efficacy of different interventions on depression to compensate for the shortcomings of indirect comparisons.

## Conclusion

The network meta-analysis revealed that various non-pharmaceutical interventions, including music intervention, massage, aerobic exercise, Qigong, yoga, Tai Chi, ball games, strength training, dance, whole-body vibration training, and high-intensity interval training, had a discernible impact on regulating depression in college students. Among these interventions, high-intensity interval training showed a higher probability of yielding the most favorable effect on depression scores compared to others. However, further high-quality research is necessary to establish its specific effects more conclusively. Considering the limitations of this study, future studies should categorize and evaluate the basic situation of participants (such as depression level), as well as the impact of intervention time and frequency on depression. More high-quality clinical trials are needed to prove this in the future.

## Data Sharing Statement

Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

## Acknowledgments

We recognize effort and great work from all team members and appreciate project coordinators, investigators, and data operators.

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Song J, Liu ZZ, Huang J, et al. Effects of aerobic exercise, traditional Chinese exercises, and meditation on depressive symptoms of college student: a meta-analysis of randomized controlled trials. *Medicine*. 2021;100(1):e23819. doi:10.1097/MD.00000000000023819
- GBD 2019 Diseases and Injuries Collaborators. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*. 2020;396(10258):1204–1222. doi:10.1016/S0140-6736(20)30925-9
- Zhang M, Ding Y, Zhang J, et al. Effect of group impromptu music therapy on emotional regulation and depressive symptoms of college students: a randomized controlled study. *Front Psychol*. 2022;13:851526. doi:10.3389/fpsyg.2022.851526
- China's Ministry of Education. Incorporating depression screening into students' physical examination. Available from: [http://www.moe.gov.cn/jyb\\_xxgk/xxgk\\_jyta/jyta\\_twys/202111/t20211104\\_577685.html](http://www.moe.gov.cn/jyb_xxgk/xxgk_jyta/jyta_twys/202111/t20211104_577685.html). Accessed July 27, 2023.
- Cho EC, Kang H. Effects of positive psychology-based music therapy group counseling program on depression, anxiety and perceived stress in art college student. *Local Global*. 2022;46(2):277–300. doi:10.33071/ssrich.46.2.202205.277
- Chen CJ, Chen YC, Ho CS, et al. Effects of preferred music therapy on peer attachment, depression, and salivary cortisol among early adolescents in Taiwan. *J Adv Nurs*. 2019;75(9):1911–1921. doi:10.1111/jan.13975
- Hye-Kyung L. The effect of self-aromatherapy massage on depression and satisfaction with life in depressed college students. *Korean J Health Educ Promot*. 2014;31(3):63–72. doi:10.14367/kjhpep.2014.31.3.63
- Cai HQ, Su SY, Dai Q, et al. Clinical observation of acupuncture combined with moxibustion on college students with depression. *Zhen Ci Yan Jiu = Acupuncture Research*. 2021;46(4):330–334. doi:10.13702/j.1000-0607.200570
- Wang X, Lin H, Jiang X, et al. Effect of electroacupuncture and counseling on sub-threshold depression: a study protocol for a multicenter randomized controlled trial. *Front Psychiatry*. 2020;11:346. doi:10.3389/fpsyg.2020.00346
- Shen Y. Effect of exercise intervention and rehabilitation on patients with depression. *Revista Brasileira De Medicina Do Esporte*. 2023;29. doi:10.1590/1517-8692202329012022\_0157
- Liang A, Zhao S, Song J, et al. Treatment effect of exercise intervention for female college students with depression: analysis of electroencephalogram microstates and power spectrum. *Sustainability*. 2021;13(12):6822. doi:10.3390/su13126822
- Li L. Experimental study on influence of different physical exercise situation upon anxiety and depression of college students. *J Southwest China Normal Univ*. 2013;38(10):107–111. doi:10.13718/j.cnki.xsxb.2013.10.001
- Tu X, Zhang S, Lai Z, et al. Machine learning-based aerobic exercise recognition and its effect on health status of college students. *Mobile Inform Syst*. 2022;2022:1–10. doi:10.1155/2022/9956753
- Cashin AG, McAuley JH. Clinimetrics: Physiotherapy Evidence Database (PEDro) scale. *J Physiother*. 2020;66(1):59. doi:10.1016/j.jphys.2019.08.005
- Xu L. *A Network Meta-Analysis of the Effects of Different Psychotherapy on Depression and Anxiety of College Students*. Xinxiang Medical University; 2022. doi:10.27434/d.cnki.gxxy.2022.000281
- Salanti G, Ades AE, Ioannidis J. Graphical methods and numerical summaries for presenting results from multiple-treatment meta-analysis: an overview and tutorial. *Journal of Clinical Epidemiology*. 2011;64(2):163–171. doi:10.1016/j.jclinepi.2010.03.016
- Zhang B, Chen Y, Qiu Z. The influence of Chinese medicine five elements music therapy on college students' depression. *J Nanjing Univ Trad Chin Med*. 2014;15(02):115–117.
- Cheng X, Wang D-M, Chen X, et al. 健身气功五禽戏改善轻度抑郁大学生前额叶和海马氢质子磁 [Health Qigong Wuqinxi improves hydrogen proton magnetic resonance spectra in prefrontal cortex and hippocampus in college students with mild depression]. *Nan fang yi ke da xue xue bao*. 2016;36(11):1468–1476. Chinese. doi:10.3969/j.issn.1673-4254.2016.11.04
- Falsafi N. A randomized controlled trial of mindfulness versus yoga: effects on depression and/or anxiety in college students. *J Am Psychiatr Nurses Assoc*. 2016;22(6):483–497. doi:10.1177/1078390316663307
- Zhang J, Qin S, Zhou Y, et al. A randomized controlled trial of mindfulness-based Tai Chi Chuan for subthreshold depression adolescents. *Neuropsychiatr Dis Treat*. 2018;14:2313–2321. doi:10.2147/NDT.S173255
- Chen J, Li Y, Wu Y, et al. Effect of Taijiquan exercise on depression and serum inflammatory factors of female college students. *Chin J School Health*. 2019;40(07):1065–1068. doi:10.16835/j.cnki.1000-9817.2019.07.029
- Duan Y. Study of different ways of exercise intervention on college students' anxiety and depression. *Chin J School Health*. 2014;35(07):1022–1024. doi:10.16835/j.cnki.1000-9817.2014.07.024
- Ma M. Evaluation of intervention effect of different exercise items on depression and anxiety of zhuang college students Chinese. *J School Health*. 2017;38(01):112–114. doi:10.16835/j.cnki.1000-9817.2017.01.035
- Wang J, Xie X. Effects of net separated competition training on anxiety and depression among college students. *Chin J School Health*. 2016;37(01):93–95. doi:10.16835/j.cnki.1000-9817.2016.01.028
- Li Y. Intervention effect of volleyball combined with group psychological counseling on depression college students. *Chin J School Health*. 2017;38(06):935–938. doi:10.16835/j.cnki.1000-9817.2017.06.045

26. Mao X, Shen H. Research on the effect of the intervention with tennis activities on the male college students with mild depression. *J Nanjing Inst Phys Educ.* 2014;13(04):148–152. doi:10.15877/j.cnki.nsin.2014.04.070
27. Zhang R, Li J, Yang M. Intervention effect of different sports on depression in college students. *J Kunming Med Univ.* 2012;33(08):27–29+49. doi:10.3969/j.issn.1003-4706.2012.08.007
28. Guan S, Li Q, Zhang S. Effect of strength training on depression and monoamine transmitters in female college students with depression symptoms. *J Wuhan Inst Phys Educ.* 2008;(10):70–74. doi:10.15930/j.cnki.wtxb.2008.10.013
29. Li Q, Guan S, Yin L. Effect of sport training on depression and monoamine transmitters in female college students with depression symptoms. *China J Modern Med.* 2014;24(08):86–90. doi:10.3969/j.issn.1005-8982.2014.08.018
30. Tang W, Guo Y, Wang X, et al. A study on physical exercises and college students' depression and self-concept of body. *J Beijing Sport Univ.* 2009;32(04):99–101. doi:10.19582/j.cnki.11-3785/g8.2009.04.028
31. Jiao H. Effect of aerobic exercise intervention on depressive mood for normal college students. *China J Health Psychol.* 2010;18(08):939–941. doi:10.13342/j.cnki.cjhp.2010.08.035
32. Duk OS, Cho PS. Effect of treadmill exercise on depressive and neurotransmitters in college women with depressive symptoms. *J Sport Leisure Stud.* 2012;50:923–934. doi:10.51979/KSSLS.2012.12.50.923
33. Chawla G, Azharuddin M, Ahmad I, et al. Effect of whole-body vibration on depression, anxiety, stress, and quality of life in college students: a randomized controlled trial. *Oman Med J.* 2022;37(4):e408. doi:10.5001/omj.2022.72
34. Zhao H. Study on the effect of high intensity intermittent exercise on depression of female college students. *J Liaocheng Univ.* 2017;30(01):88–92. doi:10.3969/j.issn.1672-6634.2017.01.016
35. Li X, Wang F, Du X. Experimental study of taijiquan on treating depression of medical university students. *Sichuan Sports Sci.* 2017;36(04):59–62. doi:10.13932/j.cnki.sctyx.2017.04.13
36. Song X, Du B, Du H. Research on the influence of taijiquan and aerobic jogging on the depression of female normal college students. *Contemp Sports Technol.* 2017;7(15):107–108. doi:10.16655/j.cnki.2095-2813.2017.15.107
37. Sadeghi K, Ahmadi SM, Ahmadi SM, et al. A comparative study of the efficacy of cognitive group therapy and aerobic exercise in the treatment of depression among the students. *Glob J Health Sci.* 2016;8(10):54171. doi:10.5539/gjhs.v8n10p1
38. Chen CJ, Sung HC, Lee MS, et al. The effects of Chinese five-element music therapy on nursing students with depressed mood. *Int J Nurs Pract.* 2015;21(2):192–199. doi:10.1111/ijn.12236
39. Celentano DD, Aramrattana A, Sutcliffe CG, et al. Associations of substance abuse and sexual risks with self-reported depressive symptoms in young adults in northern Thailand. *J Addict Med.* 2008;2(2):66–73. doi:10.1097/ADM.0b013e31816c60c1
40. Hoffman BM, Babyak MA, Craighead WE, et al. Exercise and pharmacotherapy in patients with major depression: one-year follow-up of the SMILE study. *Psychosom Med.* 2011;73(2):127–133. doi:10.1097/PSY.0b013e31820433a5
41. Yeung A, Slipp LE, Niles H, et al. Effectiveness of the relaxation response-based group intervention for treating depressed Chinese American immigrants: a pilot study. *Int J Environ Res Public Health.* 2014;11(9):9186–9201. doi:10.3390/ijerph110909186
42. Michalak J, Troje NF, Fischer J, et al. Embodiment of sadness and depression--gait patterns associated with dysphoric mood. *Psychosomatic Medicine.* 2009;71(5):580–587. doi:10.1097/PSY.0b013e3181a2515c
43. Li R, Liu J. Meta-analysis of the effect of Chinese traditional exercise on motor function and related indexes in patients with Parkinson's disease. *Shandong Sports Sci Technol.* 2022;44(06):57–62. doi:10.14105/j.cnki.1009-9840.2022.06.006
44. Tan X, Yi J, Guo C, et al. Meta-analysis of the effect of traditional Chinese martial arts on exercise tolerance in elderly patients with chronic pulmonary obstruction. *Hubei Sports Sci.* 2022;41(11):986–992. doi:10.3969/j.issn.1003-983X.2022.11.009

## Neuropsychiatric Disease and Treatment

Dovepress

### Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS, and is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>