



Potential Novel Strategies for the Treatment of Dental Pulp-Derived Pain: Pharmacological Approaches and Beyond

Christina M. A. P. Schuh^{1†}, Bruna Benso^{2,3†} and Sebastian Aguayo^{2*}

¹ Centro de Medicina Regenerativa, Facultad de Medicina Clínica Alemana-Universidad del Desarrollo, Santiago, Chile, ² School of Dentistry, Faculty of Medicine, Pontificia Universidad Católica de Chile, Santiago, Chile, ³ Department of Physiology, Faculty of Medicine, Universidad Austral de Chile, Millennium Nucleus of Ion Channels Associated Diseases (MiNICAD), Valdivia, Chile

OPEN ACCESS

Edited by:

Ramón Sotomayor-Zárate,
University of Valparaíso,
Chile

Reviewed by:

Hugo F. Miranda,
University of Chile, Chile
Teresa Pelissier,
University of Chile, Chile

*Correspondence:

Sebastian Aguayo
sebastian.aguayo@uc.cl

[†]These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Translational Pharmacology,
a section of the journal
Frontiers in Pharmacology

Received: 05 July 2019

Accepted: 22 August 2019

Published: 18 September 2019

Citation:

Schuh CMAP, Benso B and
Aguayo S (2019) Potential Novel
Strategies for the Treatment of Dental
Pulp-Derived Pain: Pharmacological
Approaches and Beyond.
Front. Pharmacol. 10:1068.
doi: 10.3389/fphar.2019.01068

The diagnosis and management of pain is an everyday occurrence in dentistry, and its effective control is essential to ensure the wellbeing of patients. Most tooth-associated pain originates from the dental pulp, a highly vascularized and innervated tissue, which is encased within mineralized dentin. It plays a crucial role in the sensing of stimuli from the local environment, such as infections (i.e. dental caries) and traumatic injury, leading to a local inflammatory response and subsequently to an increase in intra-pulp pressure, activating nerve endings. However, thermal, chemical, and mechanical stimuli also have the ability to generate dental pulp pain, which presents mechanisms highly specific to this tissue and which have to be considered in pain management. Traditionally, the management of dental pulp pain has mostly been pharmacological, using non-steroidal anti-inflammatory drugs (NSAIDs) and opioids, or restorative (i.e. removal of dental caries), or a combination of both. Both research areas continuously present novel and creative approaches. This includes the modulation of thermo-sensitive *transient receptor potential cation channels* (TRP) by newly designed drugs in pharmacological research, as well as the use of novel biomaterials, stem cells, exosomes and physical stimulation to obtain pulp regeneration in regenerative medicine. Therefore, the aim of this review is to present an up-to-date account of causes underlying dental pain, novel treatments involving the control of pain and inflammation and the induction of pulp regeneration, as well as insights in pain in dentistry from the physiological, pharmacological, regenerative and clinical perspectives.

Keywords: dentistry, pain, dentin, dental caries, regenerative medicine, pharmacology, collagen, endodontics

INTRODUCTION

Despite many advances in the fields of diagnosis, material sciences and therapeutics, oral diseases continue to burden millions of people worldwide, causing a significant impact on both health costs and patient quality of life (Hall-Stoodley et al., 2004; Logan and Brett, 2013; Römling et al., 2014). According to the Global Burden of Disease Study 2016, it is estimated that more than 3.5 billion people worldwide suffer from oral diseases, with 2.4 billion of those cases being dental caries (Vos et al., 2017). Furthermore, it is also estimated that 743 million people are affected by periodontal

disease, a chronic progressive weakening of the supporting structures of the tooth that leads to tooth loss and dysfunction (Tonetti et al., 2017). Despite the fact that these oral diseases display a wide variety of symptoms, many patients seek dental advice due to the presence of pain in the mouth and/or facial region. Therefore, the diagnosis and management of orofacial pain is an essential need for dentists to ensure the wellbeing of patients, as well as to determine the most appropriate treatment plan for each clinical situation.

In general terms, pain can be defined as an “unpleasant sensory and emotional experience that is associated with actual or potential tissue damage or described in such terms” (Gebhart, 2000). Thus, it is a subjective appreciation that varies between individuals, which can make its clinical diagnosis challenging. In dentistry, pain is a multifactorial experience that also includes strong emotional and previous-experience components (Sessle, 1986). There are many origins of pain in the orofacial region; however, the most frequent pain is the one initiating from within the tooth and dental pulp (Estrela et al., 2011), which can be triggered by a wide and diverse range of stimuli including hot and cold temperatures, air puffs, sugar consumption, and mechanical pressure. Orofacial pain originating from structures surrounding the tooth (such as periodontal tissue, oral mucosa and alveolar bone) will not be covered in this review.

Tooth Anatomy and Its Relevance in Dental Pain Perception


Teeth are highly complex organs located within the oral cavity. They are constituted by three mineralized tissues known as enamel, dentin, and cement, which surround the un-mineralized tissue known as dental pulp (Figure 1A). Mineralized tissues of the tooth are mostly comprised of hydroxyapatite crystals with varying amounts of organic content (e.g. type-I collagen, other proteins), and their main functions are to provide structural integrity to the tooth and to protect the pulp from environmental injury (Shahmoradi et al., 2014). Enamel is the most mineralized

tissue in the body; therefore, it plays a central role in mastication and in protecting dentin and pulp from environmental injury. Cement, on the other hand, is present on the surface of the root, and is involved in the anchoring of the tooth to the alveolar bone by means of the periodontal ligament.

Dentin is a mineralized dental tissue contained within enamel and cementum, constituted by hydroxyapatite (70% weight), organic material (20% weight) and water (10% weight) (Goldberg, 2011). The organic material is mostly comprised of a matrix of type-I collagen, and its presence is crucial for the structural stability as well as the elasticity of the tooth (Butler and Ritchie, 1995; Bertinetti et al., 2015). Dentin is maintained by the cellular prolongations of odontoblasts, which are nested within dentinal tubules- channels extending from the dental pulp until the enamel-dentin junction (Arana-Chavez and Massa, 2004) (Figure 1B). These tubules are flooded with dentinal (interstitial) fluid; thus, dentin receives nutrients by direct perfusion from the dental pulp. Most importantly, as dentin is strongly interconnected with the dental pulp and its innervation, it is crucial in the process of dental pain perception.

Interestingly, the initial portion of the dentinal tubules contains pulp nerve endings and thus dentin exposed to the oral cavity (i.e. by caries or trauma) can actively respond to environmental stimuli and trigger nociception (Figure 1B). Stimulatory sources that can directly or indirectly affect the dental pulp include chemical irritation, dental caries, infiltration of bonding materials, trauma and orthodontic movements, among others (Tikhonova et al., 2018). This phenomenon is also known as *dental hypersensitivity*, as seemingly innocuous stimuli can generate an exaggerated pain response in patients (Migliani et al., 2010). Nerve endings located in the initial portion of dentinal tubules and dental pulp are mainly constituted of A-delta and C fibers—afferent endings of the trigeminal nerve—that can be triggered by both mechanical and thermal stimuli (Jain et al., 2013).

There are three main hypotheses regarding the mechanisms behind dental hypersensitivity. The first hypothesis, also



known as the *neural theory*, states that nerve endings within dentinal tubules are directly able to respond to external stimuli. Nociceptive temperature-sensitive receptors such as TRPV1 and TRPA1 are believed to be involved in pain transduction from hot and cold thermal stimuli (Chung et al., 2013). The second hypothesis or *hydrodynamic theory* proposes that thermal and mechanical stimuli, such as air spraying, can exert pressure changes that cause fluid movement within the dentinal tubules (Andrew and Matthews, 2000; Shibukawa et al., 2015). This movement is believed to trigger A-delta and C fibers, although the presence of specialized mechanoreceptors in these fibers is still debatable. Finally, the *odontoblast transducer theory* states that odontoblast cells themselves are responsible for pain perception. Expression of TRPV channels in odontoblasts as well as their direct excitability support this theory (Won et al., 2018).

Additional to dentinal hypersensitivity, dental pain can also be a consequence of localized inflammation within the dental pulp. This response is at times exacerbated due to the lack of compliance given by the surrounding mineralized tissues, and therefore minor inflammatory changes can result in exaggerated pain perception. Known pro-inflammatory mediators involved in pulpal pain include substance P, calcitonin gene-related peptide, histamine, and cytokines (Caviedes-Bucheli et al., 2008, Caviedes-Bucheli et al., 2011; Sessle, 2011). These molecules are responsible for sensitizing nerve endings as well as increasing intra-pulp pressure by virtue of increasing vascular supply to the dental pulp (Rechenberg et al., 2016). Furthermore, inflammatory mediators released as part of the response system act in the production of vascularized alterations mediated by different receptors such as those from toll-like receptor (TLR) variants 2 and 4 (Nakanishi et al., 2011). It is believed that TLR-2 positive cells are present in the early phases of regulation (including the first 3 h after stimulus), which recognize different pathogen-derived molecules such as triacyl lipopeptides from bacteria and mycobacteria, peptidoglycan, and lipoteichoic acid of gram-positive bacteria and zymosan from fungi (Park et al., 2015; Rechenberg et al., 2016). On the other hand, TLR4 has a central role in the response to co-stimulatory molecules originating from gram-negative bacteria, such as lipopolysaccharides (LPS) (Nakanishi et al., 2011; Park et al., 2015). Considering TLR and the release of pro-inflammatory cytokines like IL-8, IL-6 and chemokines, the succession of stimulatory events can lead to a painful and hyperalgesic process within the dental pulp (ElSalhy et al., 2013). The wide range of factors and stimuli involved leads many to describe the dental pulp as an innervated tissue associated with pain transduction mechanisms that have yet to be fully explained (Mutoh et al., 2007).

Dental Caries and Its Relevance in Dental Pain

There are several known causes responsible for the onset of dental pain including infection, trauma, dental treatment and chemical injury (Alonso-Ezpeleta et al., 2012; Jain et al., 2013; Khan et al., 2019). These situations share the common factor of either exposing dentin or dental pulp to the environment and/or generating an inflammatory process within the dental pulp,

initiating dentinal hypersensitivity, and/or inflammatory pain. Nevertheless, in the clinical setting, the most common source of dental pain is due to dental caries.

Dental caries (also known as tooth decay) affects the vast majority of the population and is one of the main causes of tooth loss in patients of all ages (Petersen et al., 2005; Kassebaum et al., 2014; Dye et al., 2015). Worldwide, recent data shows that most adults have been affected by dental caries, and therefore this pathology represents a large economic burden to individuals and healthcare providers. The Global Burden of Disease Study 2013, reporting data on 301 diseases and injuries, concluded that tooth pain resulting from caries affected over 200 million worldwide, and was the fifth commonest acute condition observed overall (Global Burden of Disease Study, 2013; Rice et al., 2016).

Caries is a biofilm-mediated multifactorial infectious disease characterized by the loss of mineral from the tooth surface due to bacterial colonization, which can affect all mineralized tissues in teeth (Pitts et al., 2017). Although caries pathogenesis is highly complex and multifactorial, it is believed to be initiated by species such as *Streptococcus mutans* and *lactobacilli*, which induce surface demineralization by the production of extracellular acids that destroy the mineralized matrix of the tooth (Beighton et al., 2010; Krzyściak et al., 2014; Wassel and Khattab, 2017). The process of caries formation starts on the surface of the outer layers of the tooth, and if not treated it progresses into the underlying dentin layer (Pitts et al., 2017). Caries can also affect the root portion of teeth once this area becomes exposed to the oral environment. This is common in elderly people suffering from gingival recession, and thus these patients are at higher risk of developing root caries lesions (Wierichs and Meyer-Lueckel, 2015). Therefore, dental caries can trigger dental pain in a two-fold manner: by exposing dentin to the environment with subsequent onset of dentinal hypersensitivity, and by generating a localized pulp inflammation in response to bacterial invasion and bacterial molecules (such as bacterial peptides and LPS).

Pain as a consequence of dental caries is a known occurrence in the clinics, which is also supported by many investigations demonstrating the association between caries and pain in children and adults (Reid et al., 2003; Nomura et al., 2005; Bastos et al., 2006). In children, pain generated by caries has also been associated to reduced child weight, growth, and quality of life (Sheiham, 2006). Several studies have provided insight into the mechanisms behind how caries can trigger dental pain within the dental pulp. Firstly, the progressive demineralization generated by caries dissolves the outer layers of the tooth and exposes dental pulp nerve endings to environmental stimuli, triggering dentinal hypersensitivity. However, there is also evidence suggesting bacterial toxins are able to permeate into the dental pulp *via* dentinal tubules and initiate an inflammatory response within odontoblasts and the surrounding tissue (Farges et al., 2015). Durand et al. (2006) showed that lipoteichoic acid (LTA), an important bacterial wall molecule, increased the expression of chemokines in odontoblasts *via* upregulation of TLR 2. Similarly, Farges et al. (2011) observed increase in pro-inflammatory IL-6 production in odontoblasts upon LTA stimulation. Furthermore, cyclooxygenase (COX) has been shown to be associated to tissue inflammation on several levels. In inflamed tissues there

is an increase in the biosynthesis of prostaglandins, which are generated by COX isoenzymes metabolizing arachidonic acid (Ricciotti and FitzGerald, 2011). Also, while COX-1 is constitutively expressed in many tissues, COX-2 is promoted by inflammatory stimuli and is directly involved with inflammatory pain generation (Dubois et al., 1998). Within this context, Petrini et al. (2012) demonstrated an increase of prostaglandin E2 in teeth with reversible dental pulp inflammation compared to normal controls. In addition, the penetration of bacterial cells into the dental pulp may also be an important stimulus for nociception, as Chiu et al. (2013) demonstrated that bacteria can directly activate nociceptors in a mice model. Therefore, it is necessary to remove all caries-infected dental tissue as a previous step before beginning tooth restoration, in order to completely resolve inflammation in the dental pulp. However, in cases of extreme pulpal inflammation or when the caries lesion enters into the pulp chamber, dental pulp necrosis may occur. In these cases, the dental pulp must be completely removed by means of a root canal procedure (pulpectomy) and replaced by thermoplastic material.

Furthermore, Rodd and Boissonade (2000) demonstrated an increase of substance P in carious teeth compared to healthy samples, and its expression was further increased in painful samples compared to asymptomatic teeth. The levels of another neuropeptide, neuropeptide Y, was also found to be elevated in teeth affected by caries compared to controls (El Karim et al., 2006). TRPV-1 expression was also increased in carious teeth compared to controls, and vascular expression of TRPV-1 within the dental pulp was associated to increased dental pain (Morgan et al., 2005). Recently, Hall et al. (2016) demonstrated an upregulation of TNF- α in patients with caries, which was even more pronounced in patients with dental pain.

CURRENT PHARMACOLOGICAL APPROACHES FOR DENTAL PAIN TREATMENT

The therapeutic management of sensitive and painful regions is one of the most relevant aspects in the dental practice due to its direct influence on patient quality of life (Murray et al., 1996). Current management of pain deriving from inflammation utilizes several approaches including local intervention and adjuvant pharmacological treatments (Ward, 1974; Dal Secco et al., 2006). Local interventions, particularly dental restorations, aim to isolate the dental pulp and dentin from environmental stimuli triggering hypersensitivity as well as remove caries lesions to resolve underlying pulpal inflammatory process. On the other hand, pharmacological treatment contributes to the promotion of analgesia, with frequently used drugs being nonsteroidal anti-inflammatories (NSAIDs), opioids and NMDA receptor blockers (Chou and Huffman, 2007). Considering that the origin of pulpal pain can occur at different points during treatment (i.e. preoperative, perioperative, and postoperative), the drug strategy must be chosen according to the individual conditions of each clinical case (Primosch et al., 1993; Vanegas et al., 2010). For example, analgesic drug treatment can begin before

intervention (known as pre-emptive analgesia) with the aim of inhibiting the inflammatory process prior to dental restoration in asymptomatic patients (Ong et al., 2005). As several investigations have discussed conventional pharmacological management of dental pain in the past (Hargreaves and Abbott, 2005; Becker, 2010; Moore and Hersh, 2014; Berlin et al., 2018), this review will only provide a brief summary of current treatment options utilized in clinics.

Currently, the first line of drugs for the treatment of dental-derived pain are acetaminophen and NSAIDs (Moore et al., 2018). Both have been shown to be essential in the management of minor to moderate post-operative pain in dentistry (Bailey et al., 2013) and proven to be safe and effective, having them as the most favored options among treatments available in the market (Ong et al., 2005). NSAIDs have been demonstrated to inhibit the action of COX-1 and COX 2, thus are important in controlling inflammatory pain in a wide range of tissues (Brune and Patrignani, 2015). On the other hand, the exact mechanism of action for acetaminophen in humans remains unclear, although some authors believe it has an effect on a splice variant of COX-1 (previously believed to be the COX-3 isoenzyme) (Graham and Scott, 2005; Bertolini et al., 2006; Sharma and Mehta, 2013). In conjunction with opioids, acetaminophen and NSAIDs such as ibuprofen have been demonstrated to alleviate moderate to severe postoperative pain (Ashley et al., 2016) as well as acute dental pain (Chang et al., 2001; Bailey et al., 2013). Pain treatment of vulnerable patient groups such as elderly or patients with gastrointestinal issues is challenging, however there is supportive evidence for the use of acetaminophen alone or in combination with codeine (Ong et al., 2007; Ganzberg, 2018). Studies show that associating NSAIDs with opioids may have the additional benefit of controlling post-operative pain (Table 1), however, selective COX-2 inhibitors have been proven to be more effective (Chang et al., 2001; Korn et al., 2004). Additionally, special considerations should be taken to reduce opioid overprescribing and abuse, especially with young patients and patients that can be managed solely with NSAIDs (Noble et al., 2010).

Multimodal drug therapy is achieved by combining drugs with complementary mechanisms of action with the aim of reducing postoperative pain (Brooks et al., 2017; Matute Crespo and Montero Matamala, 2017). When treating moderate to severe pain, combination pharmacotherapy may be an effective alternative to treat pain and improve patient recovery times (Ganda, 2008). Originally, multimodal analgesia was advocated for use in ambulatory anesthesia combining NSAIDs and opioid analgesics, with or without anesthetic infiltration (Devin and McGirt, 2015). Combining oral or IV acetaminophen with selective or non-selective cyclooxygenase-2 inhibitors (COXIBs) has also demonstrated to be a safe option for pain treatment (Ong et al., 2007). These drugs can be administered *via* different routes, however special considerations must be taken into account when treating high-risk patient groups such as the elderly (Becker, 2010), as there are still conflicting reports in the literature regarding their safety and tolerability in these patients (Ganda, 2008). Overall, dental practitioners should make clinical decisions based on optimal described dosages, with non-opioid drugs as the primary treatment option (Becker, 2010).

TABLE 1 | Pharmacological treatment possibilities to control post-operative pain in dentistry with opioids alone and in different combinations.

	Authors*	Study design	Sample size	Pharmacological treatment groups	Outcome instrument	Postoperative measurement	Postoperative assessment of pain	Study conclusions	
Opioid alone	Tramadol	(Mehrvarzfar et al., 2012)	Randomized clinical trial, double-blind and placebo-controlled	100	Placebo Tramadol 100 mg Acetaminophen 325 mg +Ibuprofen 200 mg+ Caffeine 40 mg (Novafen) Naproxen 500 mg	10-point VAS	Placebo 3.2 (2.6–3.9) Tramadol 2.2 (1.2–3.1) Novafen 0.4 (0.1–0.8) Naproxen 0.7 (0.3–1.1)	24 h	A single oral dose of Naproxen, Novafen and Tramadol taken immediately after dental treatment reduced postoperative pain following pulpectomy.
	Codeine	(Sunshine et al., 1983)	Randomized clinical trial, double-blind and placebo-controlled	120	Placebo Propiram Fumarate 50 mg Codeine Sulfate 60 mg	4-point rate scale none = 0; mild = 1; moderate = 2, severe = 3	Placebo 1.27 (0.25) Propiram Fumarate 2.11 (0.22) Codeine 1.72 (0.23)	6 h	Propiram fumarate 50 mg is an effective oral analgesic similar to codeine sulfate 60 mg, with the possibility of a longer effect.
Opioid combination	Tramadol/ Acetaminophen	(Edwards et al., 2002)	Systematic- review with meta-analysis	1376	Acetaminophen 650 mg Ibuprofen 400 mg Tramadol 75 mg Tramadol 75mg/acetaminophen 650 mg	NNT (95% CI) PGE**	4.5 (3.6–6.1) 2.7 (2.3–3.3) 11 (6.9–26) 3.0 (2.5–3.7)	8 h	Overall, this meta-analysis demonstrated analgesic superiority of the combination drug over its components, without additional toxicity.
	Codeine/ Acetaminophen	(Chang et al., 2001)	Randomized clinical trial, double-blind and placebo-active controlled	393	Placebo (1:6:6, respectively) Rofecoxib 50 mg Codeine 60 mg/ Acetaminophen 600 mg	TOPAR6***	Placebo 3.4 (1.0–5.8) Rofecoxib 12.4 (11.3–13.4) Codeine/ Acetaminophen 7.0 (5.9/8.0)	6 h	In this study of moderate to severe postoperative dental pain, the analgesic efficacy of Rofecoxib 50 mg was greater than that of codeine/acetaminophen, with a lower incidence of adverse events and nausea.
	Oxycodone/ Acetaminophen	(Korn et al., 2004)	Randomized clinical trial, double-blind and placebo-active controlled	212	Placebo (1:3:3, respectively) Rofecoxib 50 mg Oxycodone 5mg/ Acetaminophen 325 mg	GLOBAL24#	Placebo 0.3 (0.2) Rofecoxib 2.1 (0.2) Oxycodone/ Acetaminophen 1.3 (0.2)	24 h	The superior efficacy of Rofecoxib 50 mg compared to oxycodone/acetaminophen 5/325 mg support the use of Rofecoxib for the treatment of acute post-surgical pain.

*All studies included in this summarized tables were checked for GRADE criteria; **PGE- Patient Global Evidence; ***TOPAR6- pain relief over 6 h; #GLOBAL24 - Global assessment of treatment at 24 h.

Nevertheless, in the case of persistent acute pain, short-acting opioid drugs may be an interesting alternative for treating this group of patients (Becker and Phero, 2005).

Despite the widespread use of acetaminophen, NSAIDs and opioids, and the benefits they provide for dental pain management, it is important to address several limitations of these drugs. Drug selection criteria is directly related to patient treatment needs and potential risk factors (such as cardiovascular disease), as there is a high number of potential interactions between commonly used NSAIDs and other medications (Ganzberg, 2018). As a general recommendation, patients must be evaluated for safety and possible adverse effects before taking NSAID treatment into consideration, especially in high-risk groups such as medically compromised patients (Wongrakpanich et al., 2018). A number of side effects for NSAIDs and opioids have been reported, such as gastrointestinal problems, nausea, and constipation (Benyamin et al., 2008; Nagi et al., 2015; Ganzberg, 2018). Furthermore, dependence to some of these drugs is currently a critical issue for public health in the US and worldwide, particularly opioid addiction due to misuse and excessive prescription (Nack et al., 2017; Jones et al., 2018). Therefore, several innovative approaches are currently being pursued as alternatives to conventional pharmacological pain management therapy in dentistry, including novel drug targets and regenerative and non-invasive medical approaches.

FUTURE TRENDS AND NOVEL INSIGHTS INTO THE PHARMACOLOGICAL TREATMENT OF PAIN IN DENTISTRY: UNVEILING TRPV1 AS A DRUG TARGET

As previously stated, dental pulp pain has its origin in the stimulation of the nerve fibers acting as a complex defense mechanism against external factors (Magloire et al., 2010; Solé-Magdalena et al., 2018). When dental pulp suffers an injury, the nociceptive neurons initiate and increase a process known as neurogenic inflammation (Chung et al., 2013; Mizumura and Murase, 2015). Within this context, transient receptor potential (TRP) proteins are known to play an important role in dental nociception. TRP proteins are a large family of cation-permeable ion channels (Latorre et al., 2007; Nilius and Owsianik, 2010) sensitive to electrical, chemical, mechanical, and thermal stimuli, and many of them act as cell receptors involved in environmental detection (Morales-Lázaro et al., 2013). Current research links TRP proteins to hereditary neuropathies, neuronal disorders, and other TRP channel-associated channelopathies (Rosenbaum and Simon, 2007). The group of temperature-activated TRP channels (thermoTRPs) is composed of different sub-families (i.e. vanilloid, melastatin, ankiryn, and canonical) which are important for the detection and integration of peripheral sensory input (Caterina, 2007). Therefore, the understanding of how certain ligands modulate close-open configurations of thermoTRP channels could significantly facilitate the pharmaceutical design and elaboration of TRP drug modulators (Rosenbaum and Simon, 2007; Steinberg et al., 2014; Yang and Zheng, 2017).

The discovery of TRPV1 channels and their role in the signaling process and temperature-mediated nociception has enabled progress in strategies for managing and treating pain (Caterina et al., 1997; Rosenbaum and Simon, 2007; Julius, 2013). TRPV 1-4 thermal detectors are expressed in sensory nerves, transducing both proprioceptive and nociceptive information and providing information about environmental as well as body temperature to the central nervous system (Baez et al., 2014). Agonist-induced activation of TRPV1 channels has been critical to their identification and functional description (Matta and Ahern, 2007; Gao et al., 2016). At the molecular level, the canonical agonist capsaicin binds to the channel with high affinity (< 0.7 interaction score) and enables discrimination of TRPV1 from other vanilloid subtype channels (Yang and Zheng, 2017). Several vanilloid-related compounds are also able to induce transition to the open state with high affinity; interestingly, most of them are compounds of natural origin or minor derivatives of the natural chemical structure (Norman et al., 2007; Cui et al., 2016). Moreover, the gating mechanism in TRP channels as well as traffic to the plasma membrane is strongly regulated by phosphoinositides and their sub-products [diacylglycerol (DAG), inositol triphosphate (IP3)], polyunsaturated fatty acids (PUFAs) as well as metabolites generated by COX, LOX and CYP enzymes (Rosenbaum and Simon, 2007). To date, many compounds capable of inhibiting TRPV1 have been discovered such as resiniferatoxin (agonist), GRC6211 (antagonist), and QX-314 (TRPV1 permeable Na⁺ channel blocker); however, none of them is available yet for clinical therapeutics (Meotti et al., 2014). As allosteric control by lipid binding adds pharmacological properties to TRPV1, here the intention is to discuss the modulation of TRPV1 cellular activity by antagonists in the context of the pharmacological basis of oral pain sensation caused by inflammatory conditions (Steinberg et al., 2014).

The study and development of composites capable of acting on TRPV1 receptors have evolved greatly over time and capsaicin has been identified as the selective agonist ligand (Cheung et al., 2008; Gregorio-Teruel et al., 2014). The mechanism through which capsaicin exerts its function is known as tachyphylaxis, a rapid desensitization phenomenon associated to pore opening of cation channels permeable to sodium and potassium (Gavva et al., 2004; Ohbuchi et al., 2016). It is dependent on the exposure of the ligand to the channel, i.e., that the influx of Ca²⁺ ions has a protective feedback mechanism that presents a reduced response to repeated exposure (Brauchi and Orio, 2011; Yang and Zheng, 2017). The reversible effect of channel desensitization was also observed in situations of overdose and epidermal degeneration of nerve fibers (Rosenbaum and Simon, 2007; Cheung et al., 2008).

In clinics, capsaicin-derived formulations to treat severe and chronic pain are available in a drug delivery system known as transdermal patches (Carnevale and Rohacs, 2016). Formulations with a stable release of 8% capsaicin (80 mg per gram of adhesive) as a dermal patch have been approved for use on neuropathic pain in the US and Europe, and can be used at multiple sites according to patient needs (Trevisani and Szallasi, 2010). Randomized clinical trials in diseases like diabetes, traumatic tissue injuries, herpes zoster infections, and cancer among others, have proven

the clinical safety and efficacy of the formulation in terms of improving patient quality of life (Kulkankorn et al., 2013).

Capsaicin has also been tested in association with other drugs such as local anesthetics (Gerner et al., 2008). Among these is the quaternary lidocaine derivate QX-314, which displays the therapeutic advantage of prolonging the anesthetic effect 10-times when compared to lidocaine hydrochloride (Nakagawa and Hiura, 2013). Ries et al. (2009) utilized capsaicin as a V1 agonist, enabling QX-314 to reach its intracellular binding site in the sodium channel. This scenario suggests that QX-314 in the presence of capsaicin can produce selective blocking mediated by TRPV1 (Binshtok et al., 2007). This pharmacological association was tested *in vivo* for an inferior alveolar nerve block, where a selective, long-lasting effect on sensory perception was noted without affecting motor areas (Lim et al., 2007). The above-described selective and long-lasting activity could be ideal in future formulations for the treatment of odontalgias caused by inflammatory processes, as well as in painful episodes caused by temporomandibular joint disorders (Kim et al., 2010a). Also, it demonstrates the potential of capsaicin and vanilloid-1 derivatives in the clinical management of pain associated with frequent episodes of pulpal inflammation in dentistry (Hossain et al., 2019). Due to the strong involvement of TRPV1 in the transmission of pain processes, structure-activity studies allowed for modifications in the structure of capsaicin and the development of analogous compounds in the attempt to eliminate the undesired irritation effect and pungency (Baskaran et al., 2018).

In this context, numerous specific TRPV1 antagonists have been developed with the aim of inhibiting nociceptive sensitization and the resulting transduction of the pain signaling process (Norman et al., 2007). Capsazepine was described as the first competitive antagonist of the channel, but its poorly selective binding capacity made it fail in clinical trials (Bevan et al., 1992). In models applied to the treatment of dental pain, examples of clinical records of compounds and their respective laboratories include AMG 517 (Amgen), AZD1386 (AstraZeneca), MK2295 (Merck/Neurogen) and SB705498 (GSK) (Trevisani and Szallasi, 2010). The clinical trial coordinated for SB705498 described it primarily as a powerful *in vitro* TRPV1 antagonist with the ability to reverse inflammation and pain *in vivo* (Gunthorpe et al., 2007). The controlled clinical trial was a multicenter (Korea, Italy, and UK), single-blind and placebo randomized study with a total of 145 patients, and the evaluated outcome corresponded to postoperative pain during third molar extraction surgery (Gunthorpe et al., 2007; De Petrocellis and Moriello, 2013). The experimental groups consisted of the treatments: SB705498 (400 mg and 1,000 mg), placebo and co-codamol (2 capsules of 500 mg paracetamol, 12.8 mg phosphatized codeine and 2 placebo capsules) (Cairns, 2009). In January 2019, the phase II results concluded in 2008 were published and no serious adverse effects were found; however, reported effects of SB705498 included headaches (n = 5) and heat sensation (n = 2) for the highest dose of 1,000 mg (Gunthorpe et al., 2007). The pharmaceutical industry has made a concerted effort to generate potent and selective compounds for TRPV1, in particular due to its attractive mechanism of action, which differs from the nonsteroidal


anti-inflammatories, and also in the attempt to find a drug with a wide therapeutic window (Cheung et al., 2008; Trevisani and Szallasi, 2010).

NOVEL APPROACHES FOR DENTAL PAIN TREATMENT AND BEYOND: REGENERATIVE MEDICINE AND OTHERS

Regenerative medicine, and more specifically cell and tissue engineering approaches, have become increasingly popular in dental research over the last years. The key concept underlying these approaches is to develop a causative treatment aiming for long-term regeneration rather than immediate pain control. Due to its complex structure, regeneration of dentin and dental pulp has been proven a difficult and versatile task, which can be firstly categorized according to the primary clinical problem tackled (antibacterial treatment or pulp restoration) and subsequently into the following strategies: a) bio(active)materials, b) stem cells and cell-based approaches, c) exosomes, and d) physical stimuli, as well as combinations of the above approaches (Figure 2).

Bioactive Materials

The use of active materials to control inflammation, infection and pain has been described since ancient times. Ancient Egyptians utilized pastes of ochre, honey, and willow extract (containing salicin) to treat loosening teeth as well as dental pain (Leek, 1967; Forshaw, 2009). The activity attributed to the materials used is mostly antibacterial, with salicin also displaying anti-inflammatory properties. Utilizing bioactive materials is a strategy pursued until today, currently aiming to ensure bio- and cyto-compatibility as well as reliable mechanic properties. Bioactive materials extensively used for pulp capping have been *calcium hydroxide* (causing mineralization by reduction of pyrophosphate) (Farhad and Mohammadi, 2005), *mineral trioxide aggregate* (MTA) (Torabinejad and Chivian, 1999) (a Portland cement-based substrate stimulating proliferation, migration, and differentiation of dental pulp stem cells into odontoblast-like cells to produce a collagen matrix) (Parirokh and Torabinejad, 2010; Paranjpe et al., 2011) and *biodentine* (a tricalcium-silicate based substrate with similar properties as mineral trioxide aggregate in terms of dental pulp stem cell differentiation) (Luo et al., 2014; Malkondu et al., 2014). Newer materials based on bioglass derivatives are aiming to improve overall results of pulp capping, however, so far have only demonstrated decreased setting times and increased compressive strength as main favourable characteristics (Long et al., 2018; Hanada et al., 2019). Addressing the issue of biofilm formation, Yang et al. added the antimicrobial quaternary ammonium salt monomer 2-methacryloxyethyl dodecyl methyl ammonium bromide to Portland cement, and could demonstrate a *Streptococcus mutans*-inhibiting effect for up to 6 months (Yang et al., 2014). Kwon et al. (2019) used 2-methacryloxyethyl phosphorylcholine in combination with mineral trioxide aggregate and found a two-fold effect: on the one hand the composite demonstrated protein-repellent properties and prevented adhesion of *Enterococcus*



faecalis, and on the other hand increased calcium ion deposition on the surface compared to mineral trioxide aggregate alone.

With recent advances in nanoscale manufacturing, an increased number of approaches have utilized the opportunities given by nano-structured materials to effectively deliver drugs into the dental pulp. Cuppini et al. (2019) developed a tricalcium-phosphate based paste containing amoxicillin microspheres, calcium tungstate and indomethacin nanocapsules to deliver antimicrobial and anti-inflammatory compounds to the defect site. Zhu et al. (2017) designed calcium silicate nanoparticles in combination with silver and zinc as a nano-disinfectant for dentin tubules and root canals. However, Ji et al. (2018) pointed out a limitation of this approach as insufficient amounts of nanoparticles are transported reliably through dentinal tubules; thus, they designed magnetic nanoparticles for directed transport through the dental pulp. Iron nanoparticles coated with prednisolone were applied onto the dental pulp and directed through dentinal tubules using external magnets, and authors hypothesized that this approach can serve to transport medications to reduce dental pulp inflammation,

as well as enhance bond strength of composite resin to dentin (Ji et al., 2018).

Another potential way of controlling pain related to dentinal hypersensitivity is to develop materials that promote dentinal regeneration. As type-I collagen is the most abundant structural protein in the extracellular matrix, collagen-derived scaffolds have been of interest to promote homing and differentiation of cells for dental pulp regenerative purposes (Prescott et al., 2008). To induce reparative dentinogenesis, Rutherford et al. (1993) utilized osteogenic protein 1 and bone morphogenic factor 7 in a collagen matrix, as a chemo-attractive layer on dental pulp sealed with temporary dental cement. In the case of osteogenic protein 1, a higher amount of dentin deposition was observed while bone morphogenic factor 7 did not induce reparative dentinogenesis (Rutherford and Gu, 2000). Using a similar approach involving a combination of growth factors (bFGF, VEGF, or PDGF with a basal set of NGF and BMP7), Kim et al. (2010b) highlighted a beneficial effect of applying growth factor cocktails to stimulate cell homing and subsequent expression of von Willebrand factor, dentin sialoprotein, and NGF. Also, Rosa et al. (2013) found that

an injectable type I collagen scaffold in combination with stem cells from human exfoliated deciduous teeth (SHED) was able to express odontoblastic differentiation markers when implanted into root canals. The resulting pulp-like tissue presented similar vascularization and cellularity compared to control dental pulps, and formation of new dentin was observed throughout the root. Furthermore, Jiang et al. (2017) induced root canal regeneration of immature teeth by using a resorbable collagen membrane paired with MTA, and observed complete resolution of signs and symptoms in all patients. However, no statistically significant improvement in regeneration was observed compared to the MTA-only control group. Besides collagen, other materials have also been of interest for promoting pulp regeneration, such as platelet rich derivatives such as platelet rich plasma (PRP) or platelet rich fibrin (PRF). These materials are known for their richness in growth factors, which makes them a popular biomaterial in several areas of regenerative medicine (De Pascale et al., 2015). Regarding dental pulp regeneration, several groups have utilized the strategy of autologous PRP/PRF with varying results. While Torabinejad et al. (2015) and Rodríguez-Benítez et al. (2015) reported improved rates of bone formation and re-vascularization, Bezgin et al. (2015) found no advantages in using PRP for pulpal repair. Overall, further research is necessary in order to clarify the potential effectiveness of PRP and PRF for dental pulp regeneration.

It is important to note that these material-based approaches rely mainly on the regenerative capacity of the dental pulp, augmenting the formation of new dentin by stimulating tissue-resident dental pulp stem cells and controlling inflammation as well as infection. In aging teeth however, dentinal regeneration with material-based approaches has significantly lower success rates (Lipski et al., 2018), creating a demand for combined cell-material approaches.

Cell-Based Approaches

The dental pulp is a tissue rich in vasculature and different cell types such as fibroblasts, dental pulp stem cells (DPSC), and odontoblasts. The latter are capable of forming dentin by production of collagenous and non-collagenous mineralizing proteins (known as secondary dentin). Furthermore, the odontoblast layer is located at the interface between dental pulp and dentin, protruding their processes into the dentinal tubules (**Figure 1B**). The dental pulp is crucial for tooth homeostasis and presents stimulus-dependent strategies for self-repair known as reactive dentinogenesis, where stimuli such as dental caries can induce odontoblasts to deposit tertiary dentin (Smith et al., 1995). Reparative dentinogenesis involves the dental pulp and its cellular components in response to a severe insult. When the odontoblast layer is disrupted, DPSC migrate into the site of injury and differentiate into odontoblast-like cells, forming new but irregular and less tubular dentin (Mitsiadis and Rahiotis, 2004; Tziafas, 2010). The presence of DPSC therefore is crucial for reparative dentinogenesis to occur. However, with age the dentin layer becomes thicker and the pulp chamber decreases in size, limiting the amount of DPSC available for regeneration. Furthermore, the resident DPSC are subjected to senescence

through the p16INK4a/Rb pathway, driving the otherwise highly proliferative cells into cell cycle arrest (Feng et al., 2014). Hence, the main goals in pulpal cell therapy are to deliver DPSC to the damaged, resected or retracted pulp chamber, and/or to transplant odontoblasts to obtain dentin restoration.

DPSC display common features to mesenchymal stem cells such as ability to differentiate into several mesenchymal lineages, as well as a low immunogenicity. The latter makes them suitable for allogeneic stem cell therapies (Pierdomenico et al., 2005). Several studies speculate whether odontoblast-like cells differentiated from DPSC display the same characteristics as native odontoblasts (Shi et al., 2005; Huang et al., 2009). In the early 2000s, proof-of-concept studies demonstrated a high potential of dental-derived stem cells to differentiate into a functional pulp-like complex, using hydroxyapatite in ectopic rat models (Gronthos et al., 2002; Miura et al., 2003). Interestingly, in a similar model using 3D printed hydroxyapatite and DPSC or apical papilla stem cells, Hilken et al. (2017) could not find significant differences between the stem cell- and cell free-groups in terms of angiogenesis. However, it was shown by Huang et al. (2010) that apical papilla stem cells and DPSC are capable of producing pulp-like tissue *de novo*, differentiating into odontoblast-like cells producing dentin-like tissue on existing dentinal walls.

One of the key questions in cell therapy is the route of delivery through which cells are administered to the site of interest, with approaches ranging from free cell transplantation to scaffold-free cell sheets and natural or synthetic scaffolds. Collagen and atelocollagen served as scaffolds to deliver mesenchymal stem cells in several studies (Iohara et al., 2011; Iohara et al., 2013; Hayashi et al., 2015), improving the formation of pulp-like tissue and secondary dentin in dog models (Iohara et al., 2011; Iohara et al., 2013). In a pilot clinical study, functional dentin formation was observed with no adverse effects (Nakashima et al., 2017). Interestingly, DPSC transplantation with platelet-rich fibrin did not show superior results compared to controls and was found to produce mainly periodontal tissue and not pulp (Hausner et al., 2007). Chitosan-hydrogel scaffolds seeded with DPSC in combination with growth factors formed pulp-like tissue and a layer of regenerative dentin in a necrotic-tooth dog model (El Ashiry et al., 2018). Decellularized swine dental pulp seeded with DPSC was found to induce the formation of pulp-like tissue and a layer of odontoblast-like cells in a subcutaneous mouse model (Hu et al., 2017). Also, Zhang et al. (2017) analysed decellularized tooth bud scaffolds seeded with porcine dental epithelial cells, human dental pulp cells, and human umbilical vein endothelial cells in combination with BMP-2, and discovered formation of organized dentin in a mini-pig model. Furthermore, in a scaffold-free approach, Itoh et al. (2018) designed a dual-celltype sheet containing undifferentiated DPSCs and differentiated odontoblast-like cells, reporting pulp regeneration in an ectopic mouse model. Xuan et al. (2018) demonstrated in a minipig-pulpectomy-model that pelleted DPSC are capable of regenerating whole pulp tissue, including the odontoblast layer. In a subsequent clinical trial, they observed similar effects in patients and furthermore sensory nerve regeneration.

Exosome Therapies

One of the main recurrent issues in cell therapy is promoting the survival of transplanted cells in the recipient tissue over time. Factors like mechanical and nutritional stress, hypoxia, or detachment from the extracellular matrix as well as immune responses limit cell survival during and after the transplantation process. Many attempts have been made to increase cell homing and design long-term active cells therapies including biomaterial- or growth factor co-administration, reactive oxygen species protected hydrogels or preconditioning of the cells, among others (reviewed by Baldari et al., 2017). Within this context, exosomes have been proposed as an interesting alternative to cell transplantation and have gained popularity over the last decade, avoiding the challenge of cellular homing. Exosomes were first described in mammalian cell culture in 1981 as the cargo of microvesicles being shed from cells (Trams et al., 1981). They were believed to be a cellular “waste” system, mainly involved in the removal of obsolete molecules from the cell and cell wall (Pan and Johnstone, 1983). Due to key findings such as their role in expression of antigens (Raposo, 1996) and ability to transfer mRNA/miRNA (Valadi et al., 2007), they are now established as one of the main factors in inter-cellular communication. By definition, exosomes are vesicles between 30 and 150 nm in size, and their cell surface markers and cargo can be associated to the phenotype, metabolic status, and biological role of their cell of origin (Ratajczak et al., 2006; Valadi et al., 2007; Alcayaga-Miranda, 2016).

Exosomes have been harvested from several types mesenchymal stem cells and have demonstrated immune-modulatory and pro-angiogenic properties, amongst others (Liang et al., 2016; Cosenza et al., 2018). Within the analysed sources, DPSC have been demonstrated to produce exosomes with the capacity of regulating acute inflammation by suppressing activities of cathepsin B and matrix metalloproteinases in a carrageenan-induced inflammation model (Pivoraite et al., 2015). As mentioned above, exosomes are a tool for cellular communication, representing the cell's current state. Huang et al. (2016) utilized this characteristic and hypothesized that cell-type specific exosomes can trigger lineage-specific differentiation of stem cells. They differentiated DPSCs into odontoblast-like cells to subsequently isolate the exosomes and could verify a superior odontoblast differentiation *in vitro* as well as regeneration of pulp-like tissue in an *in vivo* tooth slice model. Hu et al. (2019) confirmed these findings and analysed underlying mechanisms, finding an activation of the TGF β 1/smads signalling pathway *via* transfer of microRNAs.

Aside from active native cargo, exosomes have shown to be promising system for bio-inspired and targeted drug delivery (reviewed by Zhang et al., 2019). Despite a lack of studies reporting the use of drug-loaded exosomes for the treatment of dental pain, there are several likely possibilities originating from experiences in other tissues. One potential avenue could be the local delivery of anti-inflammatory drugs into the pulp, as curcumin-loaded stem cell or macrophage exosomes have shown to display anti-inflammatory properties (Sun et al., 2010; Kalani et al., 2016). Also, delivery of liposome-encapsulated superoxide dismutase and catalase has demonstrated to locally suppress periodontal inflammation in a dog model (Petelin et al., 2000). Alternative

approaches, originating from advances in nano-engineering, have demonstrated that exosomes can also be directed to specific tissues, making them an attractive option for targeted therapy (Gomari et al., 2018; Tian et al., 2018). Furthermore, several groups have designed artificial ion channels using DNA origami (reviewed by Shen et al., 2014). These channels could potentially resemble the biologic function of channels (i.e. TRP channel) and could be inserted as “mock-channels” into cell membranes. Overall, exosomes appear to be a promising alternative to cell therapy for dental pulp pain treatment; however, their efficacy relies strongly on recipient cells, which could be an issue in cases where the dental pulp is necrotic or a pulpectomy has been performed.

Physical Therapies

Aside from pharmacological pain treatment or causative tissue engineering approaches such as biomaterials and cell-based therapies, physical therapies such as photo-biomodulation or shockwave treatment are an upcoming field in dentistry. Some studies have tried to modulate pain and inflammation with these approaches, including treatment of dental pain.

Low level laser therapy (LLLT) was invented in the 1960s, and in an attempt to identify cancer risks of laser it was found to induce hair growth in mice (Mester et al., 1968). It utilizes wavelengths between 600 nm and 1,000 nm and has been found to trigger many effects at the cellular and tissue levels. For example, cytochrome C oxidase absorbs near-infrared light, triggering downstream events such as changes in ATP, reactive oxygen species and nitric oxide (reviewed by Farivar et al., 2014). Regarding dental pain management, LLLT has been reported to reduce pain in several situations such as postoperatively after root canal retreatment (Arslan et al., 2017) or in cases of dentinal hypersensitivity (Gerschman et al., 1994; Orhan et al., 2011). However, several studies also demonstrate a regenerative effect of LLLT. In an *in vitro* tooth-slice model, LLLT increased odontogenic and angiogenic gene expression (El Nawam et al., 2019). Kim et al. (2018) also reported increased dentinogenic differentiation of DPSC *in vitro* using pulsing LLLT, proposing high production of reactive oxygen species and activation of the TGF- β 1 signalling pathway as potential mechanisms. In rat models it was shown that LLLT increased the density of odontoblasts and induced formation of more regular dentin after mechanical pulp damage (de Santana et al., 2017) and furthermore, stimulated cell proliferation as well as dentin formation after pulpal apoptosis (Shigetani et al., 2011). These results suggest LLLT as a promising non-invasive approach for dental pain management in diverse clinical situations, although further studies are needed to confirm its clinical reproducibility.

Extracorporeal shockwave treatment (ESWT) has demonstrated beneficial effects in several areas of regenerative medicine [e.g. bone regeneration (Schaden et al., 2001) or ischemia-induced tissue necrosis (Mittermayr et al., 2011)]. Shockwaves are sonic pulses characterized by an initial rise in pressure, reaching a positive peak of up to 100 MPa within 10 ns, followed by a negative amplitude of up to 10 MPa and a total life cycle of less than 10 μ s (Ogden et al., 2001). The underlying effect on cells and tissues has been attributed to increased ATP secretion and subsequent activation of the extracellular-signal-regulated kinase pathway (ERK) (Weihs et al., 2014). For DPSC it has been demonstrated

that extracellular ATP activates P2 receptors and downstream signaling events that induce odontogenic differentiation, with ESWT suggested as one of the triggering factors (Wang et al., 2016). Furthermore, it has been shown that ESWT increases efficacy of desensitizing agents for dentine hypersensitivity (Datey et al., 2016). Despite these promising initial studies, there remains a lack of animal and clinical studies associating ESWT to dental pain control, and future research is needed to strengthen these initial observations.

CONCLUSIONS AND FUTURE PERSPECTIVES

Although significant improvements have been made in dental care, dental pain remains a frequent source for patient poor quality of life. Current treatment is mainly centered around a combination of the pharmacological management of symptoms and restorative treatment, nonetheless drugs such as NSAIDs or opioids can potentially present significant side effects and addiction in limited patients. Thus, there are many novel approaches for dental pain management being explored in hopes of improving treatment outcomes in the long-term. Approaches such as TRPV1-targeting drugs have shown promising results so far, although further clinical research is needed in order to demonstrate their efficacy in dental pain management. Furthermore, causative treatment approaches originating from regenerative medicine have the potential to significantly improve the resolution of dental pain. Cell-based and exosome therapies are already being used in other fields, and introduction into clinics could provide long-term solutions for the modulation of local pulp inflammation and nociception. Material-based therapies that promote tooth remineralization have been used in the past, and current advances improving bioactivity and antibacterial effect of materials could provide clinicians with novel approaches for the treatment of dental caries and subsequent dental pain.

Interestingly, physical therapies such as LLLT and ESWT have been widely used for the treatment of inflammation and pain in a range of tissues in humans (Cotler et al., 2015; Lee et al.,

2017; Moya et al., 2018). One of the main advantages of these approaches is their non-invasiveness, generating a biological effect in deep tissues without the need of surgery of restorative treatment. Initial studies regarding use of LLLT and ESWT for dentinal hypersensitivity showed promising results (Gerschman et al., 1994; Datey et al., 2016), as well as the use of ESWT for controlling inflammation in other areas of the mouth. However, there is still a lack of clarity regarding the biological mechanism of ESWT in tissues. Further studies elucidating the mechanisms underlying ESWT as well as effectiveness to resolve dental pulp inflammation and pain are necessary.

As discussed throughout this review, teeth are highly specialized organs with a quite particular nociception mechanism. Nerve fibers within the dental pulp are triggered either by environmental factors (dentinal hypersensitivity) or by the release of pro-inflammatory and bacterial molecules. Thus, the ultimate goal of any therapy should be to reduce local inflammation while restoring protection to dentin and the dental pulp, in order to provide causative treatment and long-term success to patients. Current research should therefore focus on developing novel and causative treatments against dental pain, with limited adverse effects and clinically reproducible results.

AUTHOR CONTRIBUTIONS

CS and BB contributed equally to this manuscript. CS, BB and SA participated in the conception of the review. SA drafted the initial full manuscript. CS, BB and SA wrote and edited the manuscript.

FUNDING

This work was sponsored by Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT) grant #11180101. CS is supported by FONDECYT grant #11180406. BB is supported by FONDECYT grant #3170518. SA is supported by FONDECYT grant #11180101. MiNICAD is supported by Iniciativa Científica Milenio, Ministry of Economy, Development and Tourism, Chile.

REFERENCES

- Alcayaga-Miranda, F., Varas-Godoy, M., and Khoury, M. (2016). Harnessing the Angiogenic potential of stem cell-derived exosomes for vascular regeneration. *Stem Cells Int.* 2016, 3409169. doi: 10.1155/2016/3409169
- Alonso-Ezpeleta, L. O., Gasco-García, C., Castellanos-Cosano, L., Martín-González, J., López-Frías, F. J., and Segura-Egea, J. J. (2012). Postoperative pain after one-visit root-canal treatment on teeth with vital pulps: comparison of three different obturation techniques. *Med. Oral Patol. Oral Cir. Bucal.* 17 (4), e721–e727. doi: 10.4317/medoral.17898
- Andrew, D., and Matthews, B. (2000). Displacement of the contents of dentinal tubules and sensory transduction in intradental nerves of the cat. *J. Physiol.* 529, 791–802. doi: 10.1111/j.1469-7793.2000.00791.x
- Arana-Chavez, V. E., and Massa, L. F. (2004). Odontoblasts: the cells forming and maintaining dentine. *Int. J. Biochem. Cell Biol.* 36, 1367–1373. doi: 10.1016/j.biocel.2004.01.006
- Arslan, H., Doğanay, E., Karataş, E., Ünlü, M. A., and Ahmed, H. M. A. (2017). Effect of low-level laser therapy on postoperative pain after root canal retreatment: a preliminary placebo-controlled, triple-blind, randomized clinical trial. *J. Endod.* 43, 1765–1769. doi: 10.1016/j.joen.2017.06.028
- Ashley, P. F., Parekh, S., Moles, D. R., Anand, P., and MacDonald, L. C. (2016). Preoperative analgesics for additional pain relief in children and adolescents having dental treatment. *Cochrane Database Syst. Rev.* CD008392. doi: 10.1002/14651858.CD008392.pub3
- Baez, D., Raddatz, N., Ferreira, G., Gonzalez, C., and Latorre, R. (2014). "Gating of thermally activated channels," in Current topics in membranes (Elsevier), 51–87. doi: 10.1016/B978-0-12-800181-3.00003-8
- Bailey, E., Worthington, H. V., van Wijk, A., Yates, J. M., Coulthard, P., and Afzal, Z. (2013). Ibuprofen and/or paracetamol (acetaminophen) for pain relief after surgical removal of lower wisdom teeth. *Cochrane Database Syst. Rev.* (12), CD004624. doi: 10.1002/14651858.CD004624.pub2
- Baldari, S., Di Rocco, G., Piccoli, M., Pozzobon, M., Muraca, M., and Toietta, G. (2017). Challenges and strategies for improving the regenerative effects of mesenchymal stromal cell-based therapies. *Int. J. Mol. Sci.* 18, E2087. doi: 10.3390/ijms18102087
- Baskaran, P., Covington, K., Bennis, J., Mohandass, A., Lehmann, T., and Thyagarajan, B. (2018). Binding efficacy and thermogenic efficiency of pungent

- and nonpungent analogs of capsaicin. *Molecules* 23, E3198. doi: 10.3390/molecules23123198
- Bastos, J. L. D., Nomura, L. H., and Peres, M. A. (2006). Dental pain, socioeconomic status, and dental caries in young male adults from southern Brazil. *Cad. Saude Publica* 21, 1416–1423. doi: 10.1590/S0102-311X2005000500014
- Becker, D. E. (2010). Pain management: Part I: managing acute and postoperative dental pain. *Anesth. Prog.* 57, 67–79. doi: 10.2344/0003-3006-57.2.67
- Becker, D. E., and Phero, J. C. (2005). Drug therapy in dental practice: nonopioid and opioid analgesics. *Anesth. Prog.* 52, 140–149. doi: 10.2344/0003-3006(2005)52[140:DTD]2.0.CO;2
- Beighton, D., Al-Haboubi, M., Mantzourani, M., Gilbert, S. C., Clark, D., Zoiopoulos, L., et al. (2010). Oral bifidobacteria: caries-associated bacteria in older adults. *J. Dent. Res.* 89, 970–974. doi: 10.1177/0022034510369319
- Benyamin, R., Trescot, A. M., Datta, S., Buenaventura, R., Adlaka, R., Sehgal, N., et al. (2008). Opioid complications and side effects. *Pain Physician* 11, 105–120.
- Berlin, H., List, T., Ridell, K., and Klingberg, G. (2018). Dentists' attitudes towards acute pharmacological pain management in children and adolescents. *Int. J. Paediatr. Dent.* 28, 152–160. doi: 10.1111/ipd.12316
- Bertinetti, L., Masci, A., Schuetz, R., Barbetta, A., Seidt, B., Wagermaier, W., et al. (2015). Osmotically driven tensile stress in collagen-based mineralized tissues. *J. Mech. Behav. Biomed. Mater.* 52, 14–21. doi: 10.1016/j.jmbbm.2015.03.010
- Bertolini, A., Ferrari, A., Ottani, A., Guerzoni, S., Tacchi, R., and Leone, S. (2006). Paracetamol: new vistas of an old drug. *CNS Drug Rev.* 12, 250–275. doi: 10.1111/j.1527-3458.2006.00250.x
- Bevan, S., Hothi, S., Hughes, G., James, I. F., Rang, H. P., Shah, K., et al. (1992). Capsazepine: a competitive antagonist of the sensory neurone excitant capsaicin. *Br. J. Pharmacol.* 107, 544–552. doi: 10.1111/j.1476-5381.1992.tb12781.x
- Bezgin, T., Yilmaz, A. D., Celik, B. N., Kolsuz, M. E., and Sonmez, H. (2015). Efficacy of platelet-rich plasma as a scaffold in regenerative endodontic treatment. *J. Endod.* 41, 36–44. doi: 10.1016/j.joen.2014.10.004
- Binshtok, A. M., Bean, B. P., and Woolf, C. J. (2007). Inhibition of nociceptors by TRPV1-mediated entry of impermeant sodium channel blockers. *Nature* 449, 607–610. doi: 10.1038/nature06191
- Brauchi, S., and Orio, P. (2011). Voltage sensing in thermo-TRP channels. *Adv. Exp. Med. Biol.* 704, 517–530. doi: 10.1007/978-94-007-0265-3_28
- Brooks, E., Freter, S. H., Bowles, S. K., and Amirault, D. (2017). Multimodal pain management in older elective arthroplasty patients. *Geriatr. Orthop. Surg. Rehabil.* 8, 151–154. doi: 10.1177/2151458517720297
- Brune, K., and Patrignani, P. (2015). New insights into the use of currently available non-steroidal anti-inflammatory drugs. *J. Pain Res.* 8, 105–118. doi: 10.2147/JPR.S75160
- Butler, W. T., and Ritchie, H. (1995). The nature and functional significance of dentin extracellular matrix proteins. *Int. J. Dev. Biol.* 39 (1), 169–179.
- Cairns, B. E. (2009). *Peripheral receptor targets for analgesia: novel approaches to pain treatment*. Hoboken, New Jersey: John Wiley & Sons. doi: 10.1002/9780470522226
- Carnevale, V., and Rohacs, T. (2016). TRPV1: a target for rational drug design. *Pharmacoeut. (Basel)*. 9, E52. doi: 10.3390/ph9030052
- Caterina, M. J. (2007). Transient receptor potential ion channels as participants in thermosensation and thermoregulation. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 292, R64–R76. doi: 10.1152/ajpregu.00446.2006
- Caterina, M. J., Schumacher, M. A., Tominaga, M., Rosen, T. A., Levine, J. D., and Julius, D. (1997). The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* 389, 816–824. doi: 10.1038/39807
- Caviedes-Bucheli, J., Azuero-Holguin, M. M., Correa-Ortiz, J. A., Aguilar-Mora, M. V., Pedroza-Flores, J. D., Ulate, E., et al. (2011). Effect of experimentally induced occlusal trauma on substance P expression in human dental pulp and periodontal ligament. *J. Endod.* 37, 627–630. doi: 10.1016/j.joen.2011.02.013
- Caviedes-Bucheli, J., Muñoz, H. R., Azuero-Holguin, M. M., and Ulate, E. (2008). Neuropeptides in dental pulp: the silent protagonists. *J. Endod.* 34, 773–788. doi: 10.1016/j.joen.2008.03.010
- Chang, D. J., Fricke, J. R., Bird, S. R., Bohidar, N. R., Dobbins, T. W., and Geba, G. P. (2001). Rofecoxib versus codeine/acetaminophen in postoperative dental pain: a double-blind, randomized, placebo- and active comparator-controlled clinical trial. *Clin. Ther.* 23, 1446–1455. doi: 10.1016/S0149-2918(01)80119-3
- Cheung, W. S., Calvo, R. R., Tounge, B. A., Zhang, S.-P., Stone, D. R., Brandt, M. R., et al. (2008). Discovery of piperidine carboxamide TRPV1 antagonists. *Bioorg. Med. Chem. Lett.* 18, 4569–4572. doi: 10.1016/j.bmcl.2008.07.035
- Chiu, I. M., Heesters, B. A., Ghasemlou, N., Von Hehn, C. A., Zhao, F., Tran, J., et al. (2013). Bacteria activate sensory neurons that modulate pain and inflammation. *Nature* 501, 52–57. doi: 10.1038/nature12479
- Chou, R., and Huffman, L. H. (2007). Medications for acute and chronic low back pain: a review of the evidence for an American Pain Society/American College of Physicians Clinical Practice Guideline. *Ann. Intern. Med.* 147, 505. doi: 10.7326/0003-4819-147-7-200710020-00008
- Chung, G., Jung, S. J., and Oh, S. B. (2013). Cellular and molecular mechanisms of dental nociception. *J. Dent. Res.* 92 (11), 948–955. doi: 10.1177/0022034513501877
- Cosenza, S., Toupet, K., Maumus, M., Luz-Crawford, P., Blanc-Brude, O., Jorgensen, C., et al. (2018). Mesenchymal stem cells-derived exosomes are more immunosuppressive than microparticles in inflammatory arthritis. *Theranostics* 8, 1399–1410. doi: 10.7150/thno.21072
- Cotler, H. B., Chow, R. T., Hamblin, M. R., and Carroll, J. (2015). The use of low level laser therapy (LLLT) for musculoskeletal pain. *MOJ Orthop. Rheumatol.* 2, 68. doi: 10.15406/mojor.2015.02.00068
- Cui, M., Gosu, V., Basith, S., Hong, S., and Choi, S. (2016). “Chapter three – polymodal transient receptor potential vanilloid type 1 nociceptor: structure, modulators, and therapeutic applications,” in *Advances in Protein Chemistry and Structural Biology*, (Netherlands: Elsevier), 81–125. doi: 10.1016/b.s.apcsb.2015.11.005
- Cuppini, M., Zatta, K. C., Mestieri, L. B., Grecca, F. S., Leitune, V. C. B., Guterres, S. S., et al. (2019). Antimicrobial and anti-inflammatory drug-delivery systems at endodontic reparative material: synthesis and characterization. *Dent. Mater.* 35, 457–467. doi: 10.1016/j.dental.2019.01.002
- Dal Secco, D., Moreira, A. P., Freitas, A., Silva, J. S., Rossi, M. A., Ferreira, S. H., et al. (2006). Nitric oxide inhibits neutrophil migration by a mechanism dependent on ICAM-1: role of soluble guanylate cyclase. *Nitric Oxide* 15, 77–86. doi: 10.1016/j.niox.2006.02.004
- Datey, A., Adeeb Thaha, C. S., Patil, S. R., Gopalan, J., and Chakravorty, D. (2016). Enhancing the efficiency of desensitizing agents with shockwave treatment—a new paradigm in dentinal hypersensitivity management. *RSC Adv.* 6, 68973–68978. doi: 10.1039/C6RA12342B
- De Pascale, M. R., Sommese, L., Casamassimi, A., and Napoli, C. (2015). Platelet derivatives in regenerative medicine: an update. *Transfus. Med. Rev.* 29, 52–61. doi: 10.1016/j.tmr.2014.11.001
- De Petrocellis, L., and Moriello, A. S. (2013). Modulation of the TRPV1 channel: current clinical trials and recent patents with focus on neurological conditions. *Recent Pat. CNS Drug Discov.* 8, 180–204. doi: 10.2174/1574889808666131209124012
- de Santana, D. A., Fonseca, G. F., Ramalho, L. M. P., Rodriguez, T. T., and Aguiar, M. C. (2017). Effect of low-level laser therapy (λ780 nm) on the mechanically damaged dentin-pulp complex in a model of extrusive luxation in rat incisors. *Lasers Med. Sci.* 32, 1995–2004. doi: 10.1007/s10103-017-2295-6
- Devin, C. J., and McGirt, M. J. (2015). Best evidence in multimodal pain management in spine surgery and means of assessing postoperative pain and functional outcomes. *J. Clin. Neurosci.* 22, 930–938. doi: 10.1016/j.jocn.2015.01.003
- Dubois, R. N., Abramson, S. B., Crofford, L., Gupta, R. A., Simon, L. S., A. Van De Putte, L. B., et al. (1998). Cyclooxygenase in biology and disease. *FASEB J.* 12, 1063–1073. doi: 10.1096/fasebj.12.12.1063
- Durand, S. H., Flacher, V., Romeas, A., Carrouel, F., Colomb, E., Vincent, C., et al. (2006). Lipoteichoic acid increases TLR and functional chemokine expression while reducing dentin formation in in vitro differentiated human odontoblasts. *J. Immunol.* 176, 2880–2887. doi: 10.4049/jimmunol.176.5.2880
- Dye, B. A., Thornton-Evans, G., Li, X., and Iafolla, T. J. (2015). Dental caries and tooth loss in adults in the United States, 2011–2012. *NCHS Data Brief*, (197), 1–8. doi: 10.1080/713604647
- Edwards, J. E., McQuay, H. J., and Moore, R. A. (2002). Combination analgesic efficacy: individual patient data meta-analysis of single-dose oral tramadol plus acetaminophen in acute postoperative pain. *J. Pain Symptom Manage.* 23, 121–130. doi: 10.1016/S0885-3924(01)00404-3
- El Ashiry, E. A., Alamoudi, N. M., El Ashiry, M. K., Bastawy, H. A., El Derwi, D. A., and Atta, H. M. (2018). Tissue engineering of necrotic dental pulp of immature teeth with apical periodontitis in dogs: radiographic and histological evaluation. *J. Clin. Pediatr. Dent.* 42, 373–382. doi: 10.17796/1053-4625-42.5.9

- El Karim, I. A., Lamey, P. J., Linden, G. J., Awawdeh, L. A., and Lundy, F. T. (2006). Caries-induced changes in the expression of pulpal neuropeptide Y. *Eur. J. Oral Sci.* 114 (2), 133–137. doi: 10.1111/j.1600-0722.2006.00343.x
- El Nawam, H., El Backly, R., Zaky, A., and Abdallah, A. (2019). Low-level laser therapy affects dentinogenesis and angiogenesis of in vitro 3D cultures of dentin-pulp complex. *Lasers Med. Sci.* doi: 10.1007/s10103-019-02804-6
- ElSalhy, M., Azizieh, F., and Raghupathy, R. (2013). Cytokines as diagnostic markers of pulpal inflammation. *Int. Endod. J.* 46, 573–580. doi: 10.1111/iej.12030
- Estrela, C., Guedes, O. A., Silva, J. A., Leles, C. R., Estrela, C. R., de, A., et al. (2011). Diagnostic and clinical factors associated with pulpal and periapical pain. *Braz. Dent. J.* 22, 306–311. doi: 10.1590/S0103-64402011000400008
- Farges, J. C., Alliot-Licht, B., Renard, E., Ducret, M., Gaudin, A., Smith, A. J., et al. (2015). Dental pulp defence and repair mechanisms in dental caries. *Mediators Inflamm.* 2015, 230251. doi: 10.1155/2015/230251
- Farges, J. C., Carrouel, F., Keller, J. F., Baudouin, C., Msika, P., Bleicher, F., et al. (2011). Cytokine production by human odontoblast-like cells upon Toll-like receptor-2 engagement. *Immunobiology* 216, 513–517. doi: 10.1016/j.imbio.2010.08.006
- Farhad, A., and Mohammadi, Z. (2005). Calcium hydroxide: a review. *Int. Dent. J.* 55, 293–301. doi: 10.1111/j.1875-595X.2005.tb00326.x
- Farivar, S., Malekshahi, T., and Shiari, R. (2014). Biological effects of low level laser therapy. *J. Lasers Med. Sci.* 5, 58–62.
- Feng, X., Xing, J., Feng, G., Huang, D., Lu, X., Liu, S., et al. (2014). P16 INK4A mediates age-related changes in mesenchymal stem cells derived from human dental pulp through the DNA damage and stress response. *Mech. Ageing Dev.* 141–142, 46–55. doi: 10.1016/j.mad.2014.09.004
- Forshaw, R. J. (2009). The practice of dentistry in ancient Egypt. *Br. Dent. J.* 206, 481–486. doi: 10.1038/sj.bdj.2009.355
- Ganda, K. M. (2008). Dentist's guide to medical conditions, medications, and complications.
- Ganzberg, S. (2018). Concerns about NSAIDs. *J. Am. Dent. Assoc.* 149, 667. doi: 10.1016/j.adaj.2018.06.008
- Gao, Y., Cao, E., Julius, D., and Cheng, Y. (2016). TRPV1 structures in nanodiscs reveal mechanisms of ligand and lipid action. *Nature* 534, 347–351. doi: 10.1038/nature17964
- Gavva, N. R., Klionsky, L., Qu, Y., Shi, L., Tamir, R., Edenson, S., et al. (2004). Molecular determinants of vanilloid sensitivity in TRPV1. *J. Biol. Chem.* 279, 20283–20295. doi: 10.1074/jbc.M312577200
- Gebhart, G. F. (2000). "Scientific issues of pain and distress," in *National Research Council (US) Committee on Regulatory Issues in Animal Care and Use. Definition of pain and distress and reporting requirements for laboratory animals: Proceedings of the workshop held June 22, 2000*. Washington (DC): National Academies Press (US), (2000). doi: 10.17226/10035
- Gerner, P., Pinshtok, A. M., Wang, C.-F., Hevelone, N. D., Bean, B. P., Woolf, C. J., et al. (2008). Capsaicin combined with local anesthetics preferentially prolongs sensory/nociceptive block in rat sciatic nerve. *Anesthesiology* 109, 872–878. doi: 10.1097/ALN.0b013e31818958f7
- Gerschman, J. A., Ruben, J., and Gebart-Eaglemon, J. (1994). Low level laser therapy for dentinal tooth hypersensitivity. *Aust. Dent. J.* 39, 353–357. doi: 10.1111/j.1834-7819.1994.tb03105.x
- Global Burden of Disease Study, C. (2013). Global Burden of Disease: 2013. *Lancet* 2015 386, 743–800. doi: 10.1016/S0140-6736(15)60692-4
- Goldberg, M. (2011). Dentin structure composition and mineralization. *Front. Biosci.* E3, 281. doi: 10.2741/e281
- Gomari, H., Moghadam, M. F., and Soleimani, M. (2018). Targeted cancer therapy using engineered exosome as a natural drug delivery vehicle. *Oncotargets Ther.* 11, 5753–5762. doi: 10.2147/OTT.S173110
- Graham, G. G., and Scott, K. F. (2005). Mechanism of action of paracetamol. *Am. J. Ther.* 12, 46–55. doi: 10.1097/00045391-200501000-00008
- Gregorio-Teruel, L., Valente, P., González-Ros, J. M., Fernández-Ballester, G., and Ferrer-Montiel, A. (2014). Mutation of I696 and W697 in the TRP box of vanilloid receptor subtype I modulates allosteric channel activation. *J. Gen. Physiol.* 143, 361–375. doi: 10.1085/jgp.201311070
- Gronthos, S., Brahimi, J., Li, W., Fisher, L. W., Cherman, N., Boyde, A., et al. (2002). Stem cell properties of human dental pulp stem cells. *J. Dent. Res.* 81, 531–535. doi: 10.1177/154405910208100806
- Gunthorpe, M. J., Hannan, S. L., Smart, D., Jerman, J. C., Arpino, S., Smith, G. D., et al. (2007). Characterization of SB-705498, a potent and selective vanilloid receptor-1 (VR1/TRPV1) antagonist that inhibits the capsaicin-, acid-, and heat-mediated activation of the receptor. *J. Pharmacol. Exp. Ther.* 321, 1183–1192. doi: 10.1124/jpet.106.116657
- Hall-Stoodley, L., Costerton, J., and Stoodley, P. (2004). Bacterial biofilms: from the natural environment to infectious diseases. *Nat. Rev. Microbiol.* 2, 95–108. doi: 10.1038/nrmicro821
- Hall, B. E., Zhang, L., Sun, Z. J., Utreras, E., Prochazkova, M., Cho, A., et al. (2016). Conditional TNF- α overexpression in the tooth and alveolar bone results in painful pulpitis and osteitis. *J. Dent. Res.* 95, 188–195. doi: 10.1177/0022034515612022
- Hanada, K., Morotomi, T., Washio, A., Yada, N., Matsuo, K., Teshima, H., et al. (2019). In vitro and in vivo effects of a novel bioactive glass-based cement used as a direct pulp capping agent. *J. Biomed. Mater. Res. - Part B Appl. Biomater.* 107, 161–168. doi: 10.1002/jbm.b.34107
- Hargreaves, K., and Abbott, P. V. (2005). Drugs for pain management in dentistry. *Aust. Dent. J.* 50, S14–S22. doi: 10.1111/j.1834-7819.2005.tb00378.x
- Hausner, T., Schmidhammer, R., Zandieh, S., Hopf, R., Schultz, A., Gogolewski, S., et al. (2007). Nerve regeneration using tubular scaffolds from biodegradable polyurethane. *Acta Neurochir. Suppl.* 100, 69–72. doi: 10.1007/978-3-211-72958-8_15
- Hayashi, Y., Murakami, M., Kawamura, R., Ishizaka, R., Fukuta, O., and Nakashima, M. (2015). CXCL14 and MCP1 are potent trophic factors associated with cell migration and angiogenesis leading to higher regenerative potential of dental pulp side population cells. *Stem Cell Res. Ther.* 6, 111. doi: 10.1186/s13287-015-0088-z
- Hilkens, P., Bronckaers, A., Ratajczak, J., Gervois, P., Wolfs, E., and Lambrechts, I. (2017). The angiogenic potential of DPSCs and SCAPs in an in vivo model of dental pulp regeneration. *Stem Cells Int.* 2017, 2582080. doi: 10.1155/2017/2582080
- Hossain, M. Z., Bakri, M. M., Yahya, F., Ando, H., Unno, S., and Kitagawa, J. (2019). The role of transient receptor potential (TRP) channels in the transduction of dental pain. *Int. J. Mol. Sci.* 20 (3), 526. doi: 10.3390/ijms20030526
- Hu, L., Gao, Z., Xu, J., Zhu, Z., Fan, Z., Zhang, C., et al. (2017). Decellularized swine dental pulp as a bioscaffold for pulp regeneration. *Biomed Res. Int.* 2017, 9342714. doi: 10.1155/2017/9342714
- Hu, X., Zhong, Y., Kong, Y., Chen, Y., Feng, J., and Zheng, J. (2019). Lineage-specific exosomes promote the odontogenic differentiation of human dental pulp stem cells (DPSCs) through TGF β 1/smads signaling pathway via transfer of microRNAs. *Stem Cell Res. Ther.* 10, 170. doi: 10.1186/s13287-019-1278-x
- Huang, C. C., Narayanan, R., Alapati, S., and Ravindran, S. (2016). Exosomes as biomimetic tools for stem cell differentiation: applications in dental pulp tissue regeneration. *Biomaterials* 111, 103–115. doi: 10.1016/j.biomaterials.2016.09.029
- Huang, G. T.-J., Gronthos, S., and Shi, S. (2009). Mesenchymal stem cells derived from dental tissues vs. those from other sources: their biology and role in regenerative medicine. *J. Dent. Res.* 88, 792–806. doi: 10.1177/0022034509340867
- Huang, G. T.-J., Yamaza, T., Shea, L. D., Djouad, F., Kuhn, N. Z., Tuan, R. S., et al. (2010). Stem/progenitor cell-mediated de novo regeneration of dental pulp with newly deposited continuous layer of dentin in an in vivo model. *Tissue Eng. Part A* 16, 605–615. doi: 10.1089/ten.tea.2009.0518
- Iohara, K., Imabayashi, K., Ishizaka, R., Watanabe, A., Nabekura, J., Ito, M., et al. (2011). Complete pulp regeneration after pulpectomy by transplantation of CD105 + stem cells with stromal cell-derived factor-1. *Tissue Eng. Part A* 17, 1911–1920. doi: 10.1089/ten.tea.2010.0615
- Iohara, K., Murakami, M., Takeuchi, N., Osako, Y., Ito, M., Ishizaka, R., et al. (2013). A novel combinatorial therapy with pulp stem cells and granulocyte colony-stimulating factor for total pulp regeneration. *Stem Cells Transl. Med.* 2, 818. doi: 10.5966/sctm.2012-0132
- Itoh, Y., Sasaki, J. I., Hashimoto, M., Katata, C., Hayashi, M., and Imazato, S. (2018). Pulp regeneration by 3-dimensional dental pulp stem cell constructs. *J. Dent. Res.* 97, 1137–1143. doi: 10.1177/0022034518772260
- Jain, N., Gupta, A., and Meena, N. (2013). An insight into neurophysiology of pulpal pain: facts and hypotheses. *Korean J. Pain.* 26 (4), 347–355. doi: 10.3344/kjp.2013.26.4.347
- Ji, Y., Choi, S. K., Sultan, A. S., Chuncai, K., Lin, X., Dashtimoghdam, E., et al. (2018). Nanomagnetic-mediated drug delivery for the treatment of dental disease. *Nanomedicine* 14, 919–927. doi: 10.1016/j.nano.2018.01.013

- Jiang, X., Liu, H., and Peng, C. (2017). Clinical and radiographic assessment of the efficacy of a collagen membrane in regenerative endodontics: a randomized, controlled clinical trial. *J. Endod.* 43, 1465–1471. doi: 10.1016/j.joen.2017.04.011
- Jones, M. R., Viswanath, O., Peck, J., Kaye, A. D., Gill, J. S., and Simopoulos, T. T. (2018). A brief history of the opioid epidemic and strategies for pain medicine. *Pain Ther.* 7, 13–21. doi: 10.1007/s40122-018-0097-6
- Julius, D. (2013). TRP channels and pain. *Annu. Rev. Cell Dev. Biol.* 29, 355–384. doi: 10.1146/annurev-cellbio-101011-155833
- Kalani, A., Chaturvedi, P., Kamat, P. K., Maldonado, C., Bauer, P., Joshua, I. G., et al. (2016). Curcumin-loaded embryonic stem cell exosomes restored neurovascular unit following ischemia-reperfusion injury. *Int. J. Biochem. Cell Biol.* 79, 360–369. doi: 10.1016/j.biocel.2016.09.002
- Kassebaum, N. J., Bernabé, E., Dahiya, M., Bhandari, B., Murray, C. J. L., and Marcenes, W. (2014). Global burden of severe tooth loss. *J. Dent. Res.* 93, 20S–28S. doi: 10.1177/0022034514537828
- Khan, J., Zusman, T., Wang, Q., and Eliav, E. (2019). Acute and chronic pain in orofacial trauma patients. *Dent. Traumatol.* 2019. doi: 10.1111/edt.12493
- Kim, H. B., Baik, K. Y., Seonwoo, H., Jang, K. J., Lee, M. C., Choung, P. H., et al. (2018). Effects of pulsing of light on the dentinogenesis of dental pulp stem cells in vitro. *Sci. Rep.* 8, 2057. doi: 10.1038/s41598-018-19395-x
- Kim, H. Y., Kim, K., Li, H. Y., Chung, G., Park, C.-K., Kim, J. S., et al. (2010a). Selectively targeting pain in the trigeminal system. *Pain* 150, 29–40. doi: 10.1016/j.pain.2010.02.016
- Kim, J. Y., Xin, X., Moio, E. K., Chung, J., Lee, C. H., Chen, M., et al. (2010b). Regeneration of dental-pulp-like tissue by chemotaxis-induced cell homing. *Tissue Eng. Part A* 16, 3023–3031. doi: 10.1089/ten.tea.2010.0181
- Korn, S., Vassil, T. C., Kotey, P. N.-A., and Fricke, J. R. (2004). Comparison of rofecoxib and oxycodone plus acetaminophen in the treatment of acute pain: a randomized, double-blind, placebo-controlled study in patients with moderate to severe postoperative pain in the third molar extraction model. *Clin. Ther.* 26, 769–778. doi: 10.1016/S0149-2918(04)90076-8
- Krzyściak, W., Jurczak, A., Kościelniak, D., Bystrowska, B., and Skalniak, A. (2014). The virulence of *Streptococcus mutans* and the ability to form biofilms. *Eur. J. Clin. Microbiol. Infect. Dis.* 33, 499–515. doi: 10.1007/s10096-013-1993-7
- Kulkarnakorn, K., Lorsuwansiri, C., and Meesawatson, P. (2013). 0.025% Capsaicin gel for the treatment of painful diabetic neuropathy: a randomized, double-blind, crossover, placebo-controlled trial. *Pain Pract.* 13, 497–503. doi: 10.1111/papr.12013
- Kwon, J.-S., Lee, M.-J., Kim, J.-Y., Kim, D., Ryu, J.-H., Jang, S., et al. (2019). Novel anti-biofouling bioactive calcium silicate-based cement containing 2-methacryloyloxyethyl phosphorylcholine. *PLoS One* 14, e0211007. doi: 10.1371/journal.pone.0211007
- Latorre, R., Brauchi, S., Orta, G., Zaelzer, C., and Vargas, G. (2007). ThermoTRP channels as modular proteins with allosteric gating. *Cell Calcium* 42, 427–438. doi: 10.1016/j.ceca.2007.04.004
- Lee, S., Lee, S., Jeong, M., Oh, H., and Lee, K. (2017). The effects of extracorporeal shock wave therapy on pain and range of motion in patients with adhesive capsulitis. *J. Phys. Ther. Sci.* 29, 1907–1909. doi: 10.1589/jpts.29.1907
- Leek, F. F. (1967). The practice of dentistry in ancient Egypt. *J. Egypt. Archaeol.* 53, 51–58. doi: 10.1177/030751336705300109
- Liang, X., Zhang, L., Wang, S., Han, Q., and Zhao, R. C. (2016). Exosomes secreted by mesenchymal stem cells promote endothelial cell angiogenesis by transferring miR-125a. *J. Cell Sci.* 129, 2182–2189. doi: 10.1242/jcs.170373
- Lim, T. K. Y., MacLeod, B. A., Ries, C. R., and Schwarz, S. K. W. (2007). The quaternary lidocaine derivative, QX-314, produces long-lasting local anesthesia in animal models in vivo. *Anesthesiology* 107, 305–311. doi: 10.1097/01.anes.0000207058.77314.b4
- Lipski, M., Nowicka, A., Kot, K., Postek-Stefańska, L., Wysoczańska-Jankowicz, I., Borkowski, L., et al. (2018). Factors affecting the outcomes of direct pulp capping using Biodentine. *Clin. Oral Investig.* 22, 2021–2029. doi: 10.1007/s00784-017-2296-7
- Logan, N., and Brett, P. (2013). The control of mesenchymal stromal cell osteogenic differentiation through modified surfaces. *Stem Cells Int.* 2013, 10. doi: 10.1155/2013/361637
- Long, Y. Z., Liu, S. Y., Li, W., and Dong, Y. M. (2018). [Physical and chemical properties of pulp capping materials based on bioactive glass]. *Beijing Da Xue Xue Bao.* 50, 887–891. doi: 10.19723/j.issn.1671-167X.2018.05.021
- Luo, Z., Kohli, M. R., Yu, Q., Kim, S., Qu, T., and He, W. X. (2014). Biodentine induces human dental pulp stem cell differentiation through mitogen-activated protein kinase and calcium-/calmodulin-dependent protein kinase II pathways. *J. Endod.* 40, 937–942. doi: 10.1016/j.joen.2013.11.022
- Magloire, H., Maurin, J. C., Couble, M. L., Shibukawa, Y., Tsumura, M., Thivichon-Prince, B., et al. (2010). Topical review. Dental pain and odontoblasts: facts and hypotheses. *J. Orofac. Pain* 24, 335–349.
- Malkondu, Ö., Kazandağ, M. K., and Kazazoğlu, E. (2014). A review on biodentine, a contemporary dentine replacement and repair material. *Biomed Res. Int.* 2014, 1160951. doi: 10.1155/2014/160951
- Matta, J. A., and Ahern, G. P. (2007). Voltage is a partial activator of rat thermosensitive TRP channels. *J. Physiol.* 585, 469–482. doi: 10.1113/jphysiol.2007.144287
- Matute Crespo, M., and Montero Matamala, A. (2017). Avances farmacológicos en el manejo multimodal de la analgesia perioperatoria. *Rev. Esp. Anestesiol. Reanim.* 64, 467–471. doi: 10.1016/j.redar.2017.03.006
- Mehrvarzfar, P., Abbott, P. V., Saghir, M. A., Delvarani, A., Asgar, K., Lotfi, M., et al. (2012). Effects of three oral analgesics on postoperative pain following root canal preparation: a controlled clinical trial. *Int. Endod. J.* 45, 76–82. doi: 10.1111/j.1365-2591.2011.01950.x
- Meotti, F. C., Lemos de Andrade, E., and Calixto, J. B. (2014). “TRP modulation by 1553 natural compounds,” in *Handbook of experimental pharmacology*, (GER Elsevier) 1177–1238. doi: 10.1007/978-3-319-05161-1_19
- Mester, E., Szende, B., and Gärtner, P. (1968). The effect of laser beams on the growth of hair in mice. *Radiobiol. Radiother. (Berl.)* 9, 621–626.
- Miglani, S., Aggarwal, V., and Ahuja, B. (2010). Dentin hypersensitivity: recent trends in management. *J. Conserv. Dent.* 13, 218–224. doi: 10.4103/0972-0707.73385
- Mitsiadis, T. A., and Rahiotis, C. (2004). Parallels between tooth development and repair: conserved molecular mechanisms following carious and dental injury. *J. Dent. Res.* 83, 896–902. doi: 10.1177/154405910408301202
- Mittermayr, R., Hartinger, J., Antonic, V., Meinel, A., Pfeifer, S., Stojadinovic, A., et al. (2011). Extracorporeal shock wave therapy (ESWT) minimizes ischemic tissue necrosis irrespective of application time and promotes tissue revascularization by stimulating angiogenesis. *Ann. Surg.* 253, 1024–1032. doi: 10.1097/SLA.0b013e3182121d6e
- Miura, M., Gronthos, S., Zhao, M., Lu, B., Fisher, L. W., Robey, P. G., et al. (2003). SHED: stem cells from human exfoliated deciduous teeth. *Proc. Natl. Acad. Sci.* 100, 5807–5812. doi: 10.1073/pnas.0937635100
- Mizumura, K., and Murase, S. (2015). “Role of nerve growth factor in pain,” in *Handbook of experimental pharmacology*, (GER Elsevier). 57–77. doi: 10.1007/978-3-662-46450-2_4
- Moore, P. A., and Hersh, E. V. (2014). Combining ibuprofen and acetaminophen for acute pain management after third-molar extractions. *J. Am. Dent. Assoc.* 144, 898–908. doi: 10.14219/jada.archive.2013.0207
- Moore, P. A., Ziegler, K. M., Lipman, R. D., Aminoshariae, A., Carrasco-Labra, A., and Mariotti, A. (2018). Benefits and harms associated with analgesic medications used in the management of acute dental pain. *J. Am. Dent. Assoc.* 149, 256–265.e3. doi: 10.1016/j.adaj.2018.02.012
- Morales-Lázaro, S. L., Simon, S. A., and Rosenbaum, T. (2013). The role of endogenous molecules in modulating pain through transient receptor potential vanilloid 1 (TRPV1). *J. Physiol.* 591, 3109–3121. doi: 10.1113/jphysiol.2013.251751
- Morgan, C. R., Rodd, H. D., Clayton, N., Davis, J. B., and Boissonade, F. M. (2005). Vanilloid receptor 1 expression in human tooth pulp in relation to caries and pain. *J. Orofac. Pain* 19, 248–260.
- Moya, D., Ramón, S., Schaden, W., Wang, C. J., Guiloff, L., and Cheng, J. H. (2018). The role of extracorporeal shockwave treatment in musculoskeletal disorders. *J. Bone Jt. Surg. – Am.* 100, 251–263. doi: 10.2106/JBJS.17.00661
- Murray, H., Locker, D., Mock, D., and Tenenbaum, H. C. (1996). Pain and the quality of life in patients referred to a craniofacial pain unit. *J. Orofac. Pain* 10, 316–323.
- Mutoh, N., Tani-Ishii, N., Tsukinoki, K., Chieda, K., and Watanabe, K. (2007). Expression of toll-like receptor 2 and 4 in dental pulp. *J. Endod.* 33, 1183–1186. doi: 10.1016/j.joen.2007.05.018
- Nack, B., Haas, S. E., and Portnof, J. (2017). Opioid use disorder in dental patients: the latest on how to identify, treat, refer and apply laws and regulations in your practice. *Anesth. Prog.* 64, 178–187. doi: 10.2344/anpr-64-03-09

- Nagi, R., Yashoda Devi, B. K., Rakesh, N., Reddy, S. S., and Patil, D. J. (2015). Clinical implications of prescribing nonsteroidal anti-inflammatory drugs in oral health care — A review. *Oral Surg. Oral Med. Oral Pathol. Oral Radiol.* 119, 264–271. doi: 10.1016/j.oooo.2014.12.002
- Nakagawa, H., and Hiura, A. (2013). Comparison of the transport of QX-314 through TRPA1, TRPM8, and TRPV1 channels. *J. Pain Res.* 6, 223–230. doi: 10.2147/JPR.S41614
- Nakanishi, T., Takegawa, D., Hirao, K., Takahashi, K., Yumoto, H., and Matsuo, T. (2011). Roles of dental pulp fibroblasts in the recognition of bacterium-related factors and subsequent development of pulpitis. *Jpn. Dent. Sci. Rev.* 47, 161–166. doi: 10.1016/j.jdsr.2011.02.001
- Nakashima, M., Iohara, K., Murakami, M., Nakamura, H., Sato, Y., Ariji, Y., et al. (2017). Pulp regeneration by transplantation of dental pulp stem cells in pulpitis: a pilot clinical study. *Stem Cell Res. Ther.* 8, 61. doi: 10.1186/s13287-017-0506-5
- Nilius, B., and Owsianik, G. (2010). Transient receptor potential channelopathies. *Pflügers Arch. - Eur. J. Physiol.* 460, 437–450. doi: 10.1007/s00424-010-0788-2
- Noble, M., Treadwell, J. R., Tregear, S. J., Coates, V. H., Wiffen, P. J., Akafomo, C., et al. (2010). Long-term opioid management for chronic noncancer pain. *Cochrane Database Syst. Rev.* 2010 (1), CD006605. doi: 10.1002/14651858.CD006605.pub2
- Nomura, L. H., Bastos, J. L. D., and Peres, M. A. (2005). Dental pain prevalence and association with dental caries and socioeconomic status in schoolchildren, Southern Brazil, 2002. *Braz. Oral Res.* 18, 134–140. doi: 10.1590/S1806-83242004000200008
- Norman, M. H., Zhu, J., Fotsch, C., Bo, Y., Chen, N., Chakrabarti, P., et al. (2007). Novel vanilloid receptor-1 antagonists: 1. Conformationally restricted analogues of trans-cinnamides †. *J. Med. Chem.* 50, 3497–3514. doi: 10.1021/jm070189q
- Ogden, J. A., Tóth-Kischkat, A., and Schultheiss, R. (2001). Principles of shock wave therapy. *Clin. Orthop. Relat. Res.* (387), 8–17. doi: 10.1097/00003086-200106000-00003
- Ohbuchi, K., Mori, Y., Ogawa, K., Warabi, E., Yamamoto, M., and Hirokawa, T. (2016). Detailed analysis of the binding mode of vanilloids to transient receptor potential vanilloid type I (TRPV1) by a mutational and computational study. *PLoS One* 11, e0162543. doi: 10.1371/journal.pone.0162543
- Ong, C. K.-S., Lirk, P., Seymour, R. A., and Jenkins, B. J. (2005). The efficacy of preemptive analgesia for acute postoperative pain management: a meta-analysis. *Anesth. Analg.* 100, 757–773. doi: 10.1213/01.ANE.0000144428.98767.0E
- Ong, C. K. S., Lirk, P., Tan, C. H., and Seymour, R. A. (2007). An evidence-based update on nonsteroidal anti-inflammatory drugs. *Clin. Med. Res.* 5, 19–34. doi: 10.3121/cm.2007.698
- Orhan, K., Aksoy, U., Can-Karabulut, D. C., and Kalender, A. (2011). Low-level laser therapy of dentin hypersensitivity: a short-term clinical trial. *Lasers Med. Sci.* 26, 591–598. doi: 10.1007/s10103-010-0794-9
- Pan, B. T., and Johnstone, R. M. (1983). Fate of the transferrin receptor during maturation of sheep reticulocytes in vitro: selective externalization of the receptor. *Cell* 33, 967–978. doi: 10.1016/0092-8674(83)90040-5
- Paranjpe, A., Smoot, T., Zhang, H., and Johnson, J. D. (2011). Direct contact with mineral trioxide aggregate activates and differentiates human dental pulp cells. *J. Endod.* 37, 1691–1695. doi: 10.1016/j.joen.2011.09.012
- Parirokh, M., and Torabinejad, M. (2010). Mineral trioxide aggregate: a comprehensive literature review-part III: clinical applications, drawbacks, and mechanism of action. *J. Endod.* 36, 400–413. doi: 10.1016/j.joen.2009.09.009
- Park, S. H., Ye, L., Love, R. M., Farges, J.-C., and Yumoto, H. (2015). Inflammation of the dental pulp. *Mediators Inflamm.* 2015, 980196. doi: 10.1155/2015/980196
- Petelin, M., Pavlica, Z., Ivanuša, T., Šentjurc, M., and Skalerič, U. (2000). Local delivery of liposome-encapsulated superoxide dismutase and catalase suppress periodontal inflammation in beagles. *J. Clin. Periodontol.* 27, 918–925. doi: 10.1034/j.1600-051x.2000.027012918.x
- Petersen, P. E., Bourgeois, D., Ogawa, H., Estupinan-Day, S., and Ndiaye, C. (2005). The global burden of oral diseases and risks to oral health. *Bull. World Health Organ.* 83, 661–669. doi: /S0042-96862005000900011
- Petrini, M., Ferrante, M., Ciavarelli, L., Ceccarini, A., Brunetti, L., Vacca, M., et al. (2012). Prostaglandin E2 to diagnose between reversible and irreversible pulpitis. *Int. J. Immunopathol. Pharmacol.* 25, 157–163. doi: 10.1177/039463201202500118
- Pierdomenico, L., Bonsi, L., Calvitti, M., Rondelli, D., Arpinati, M., Chirumbolo, G., et al. (2005). Multipotent mesenchymal stem cells with immunosuppressive activity can be easily isolated from dental pulp. *Transplantation* 80, 836–842. doi: 10.1097/01.tp.0000173794.72151.88
- Pitts, N. B., Zero, D. T., Marsh, P. D., Ekstrand, K., Weintraub, J. A., Ramos-Gomez, F., et al. (2017). Dental caries. *Nat. Rev. Dis. Prim.* 3. doi: 10.1038/nrdp.2017.30
- Pivoraitė, U., Jarmalavičiūtė, A., Tunaitis, V., Ramanauskaitė, G., Vaitkuvienė, A., Kašėta, V., et al. (2015). Exosomes from human dental pulp stem cells suppress carrageenan-induced acute inflammation in mice. *Inflammation* 38, 1933–1941. doi: 10.1007/s10753-015-0173-6
- Prescott, R. S., Alsanea, R., Fayad, M. I., Johnson, B. R., Wenckus, C. S., Hao, J., et al. (2008). In vivo generation of dental pulp-like tissue by using dental pulp stem cells, a collagen scaffold, and dentin matrix protein 1 after subcutaneous transplantation in mice. *J. Endod.* 34, 421–426. doi: 10.1016/j.joen.2008.02.005
- Primosch, R. E., Antony, S. J., and Courts, F. J. (1993). The efficacy of preoperative analgesic administration for postoperative pain management of pediatric dental patients. *Anesth. Pain Control Dent.* 2, 102–106.
- Raposo, G. (1996). B lymphocytes secrete antigen-presenting vesicles. *J. Exp. Med.* 183, 1161–1172. doi: 10.1084/jem.183.3.1161
- Ratajczak, J., Miekus, K., Kucia, M., Zhang, J., Reca, R., Dvorak, P., et al. (2006). Embryonic stem cell-derived microvesicles reprogram hematopoietic progenitors: evidence for horizontal transfer of mRNA and protein delivery. *Leukemia* 20, 847–856. doi: 10.1038/sj.leu.2404132
- Rechenberg, D. K., Galicia, J. C., and Peters, O. A. (2016). Biological markers for pulpal inflammation: a systematic review. *PLoS One* 11, e0167289. doi: 10.1371/journal.pone.0167289
- Reid, B. C., Chenette, R., and Macek, M. D. (2003). Prevalence and predictors of untreated caries and oral pain among special olympic athletes. *Spec. Care Dent.* 23 (4), 139–142. doi: 10.1111/j.1754-4505.2003.tb00300.x
- Ricciotti, E., and FitzGerald, G. A. (2011). Prostaglandins and inflammation. *Arterioscler. Thromb. Vasc. Biol.* 31, 986–1000. doi: 10.1161/ATVBAHA.110.207449
- Rice, A. S. C., Smith, B. H., and Blyth, F. M. (2016). Pain and the global burden of disease. *Pain* 157 (4), 791–796. doi: 10.1097/j.pain.0000000000000454
- Ries, C. R., Pillai, R., Chung, C. C. W., Wang, J. T. C., MacLeod, B. A., and Schwarz, S. K. W. (2009). QX-314 produces long-lasting local anesthesia modulated by transient receptor potential vanilloid receptors in mice. *Anesthesiology* 111, 122–126. doi: 10.1097/ALN.0b013e3181a9160e
- Rodd, H. D., and Boissonade, F. M. (2000). Substance P expression in human tooth pulp in relation to caries and pain experience. *Eur. J. Oral Sci.* 108, 467–474. doi: 10.1034/j.1600-0722.2000.00924.x
- Rodríguez-Benítez, S., Stambolsky, C., Gutiérrez-Pérez, J. L., Torres Lagares, D., and Segura-Egea, J. J. (2015). Pulp revascularization of immature dog teeth with apical periodontitis using triantibiotic paste and platelet-rich plasma: a radiographic study. *J. Endod.* 41, 1299–1304. doi: 10.1016/j.joen.2015.05.002
- Römling, U., Kjelleberg, S., Normark, S., Nyman, L., Uhlin, B. E., and Åkerlund, B. (2014). Microbial biofilm formation: a need to act. *J. Intern. Med.* 276, 98–110. doi: 10.1111/joim.12242
- Rosa, V., Zhang, Z., Grande, R. H. M., and Nör, J. E. (2013). Dental pulp tissue engineering in full-length human root canals. *J. Dent. Res.* 92, 970–975. doi: 10.1177/0022034513505772
- Rosenbaum, T., and Simon, S. A. (2007). *TRPV1 Receptors and Signal Transduction*. Boca Raton (FL): CRC Press/Taylor & Francis. doi: 10.1201/9781420005844.ch5
- Rutherford, R. B., and Gu, K. (2000). Treatment of inflamed ferret dental pulps with recombinant bone morphogenetic protein-7. *Eur. J. Oral Sci.* 108, 202–206. doi: 10.1034/j.1600-0722.2000.108003202.x
- Rutherford, R. B., Wahle, J., Tucker, M., Rueger, D., and Charette, M. (1993). Induction of reparative dentine formation in monkeys by recombinant human osteogenic Protein-1. *Arch. Oral Biol.* 38, 571–576. doi: 10.1016/0003-9969(93)90121-2
- Schaden, W., Fischer, A., and Sailer, A. (2001). Extracorporeal shock wave therapy of nonunion or delayed osseous union. *Clin. Orthop. Relat. Res.* (387): 90–4. doi: 10.1097/00003086-200106000-00012
- Sessle, B. J. (1986). Recent developments in pain research: central mechanisms of orofacial pain and its control. *J. Endod.* 12, 435–444. doi: 10.1016/S0099-2399(86)80196-0

- Sessle, B. J. (2011). "Peripheral and central mechanisms of orofacial inflammatory pain," in *International Review of Neurobiology*. 97, 179–206. doi: 10.1016/B978-0-12-385198-7.00007-2
- Shahmoradi, M., Bertassoni, L. E., Elfallah, H. M., and Swain, M. (2014). "Fundamental structure and properties of enamel, dentin and cementum BT — Advances in calcium phosphate biomaterials," Ed. B. Ben-Nissan (Berlin, Heidelberg: Springer Berlin Heidelberg), 511–547. doi: 10.1007/978-3-642-53980-0_17
- Sharma, C. V., and Mehta, V. (2013). Paracetamol: mechanisms and updates. *BJA Educ.* 14, 153–158. doi: 10.1093/bjaceaccp/mkt049
- Sheiham, A. (2006). Dental caries affects body weight, growth and quality of life in pre-school children. *Br. Dent. J.* 201, 625–626. doi: 10.1038/sj.bdj.4814259
- Shen, Y., Saboe, P. O., Sines, I. T., Erbakan, M., and Kumar, M. (2014). Biomimetic membranes: a review. *J. Memb. Sci.* 454, 359–381. doi: 10.1016/j.memsci.2013.12.019
- Shi, S., Miura, M., Seo, B. M., Robey, P. G., Bartold, P. M., and Gronthos, S. (2005). The efficacy of mesenchymal stem cells to regenerate and repair dental structures. *Orthod. Craniofacial Res.* 8, 191–199. doi: 10.1111/j.1601-6343.2005.00331.x
- Shibukawa, Y., Sato, M., Kimura, M., Sobhan, U., Shimada, M., Nishiyama, A., et al. (2015). Odontoblasts as sensory receptors: transient receptor potential channels, pannexin-1, and ionotropic ATP receptors mediate intercellular odontoblast-neuron signal transduction. *Pflugers Arch. Eur. J. Physiol.* 467, 843–863. doi: 10.1007/s00424-014-1551-x
- Shigetani, Y., Sasa, N., Suzuki, H., Okiji, T., and Ohshima, H. (2011). GaAlAs laser irradiation induces active tertiary dentin formation after pulpal apoptosis and cell proliferation in rat molars. *J. Endod.* 37, 1086–1091. doi: 10.1016/j.joen.2011.05.020
- Smith, A. J., Cassidy, N., Perry, H., Begue-Kirn, C., Ruch, J. V., and Lesot, H. (1995). Reactionary dentinogenesis. *Int. J. Dev. Biol.* 39, 273–280.
- Solé-Magdalena, A., Martínez-Alonso, M., Coronado, C. A., Junquera, L. M., Cobo, J., and Vega, J. A. (2018). Molecular basis of dental sensitivity: the odontoblasts are multisensory cells and express multifunctional ion channels. *Ann. Anat. - Anat. Anzeiger* 215, 20–29. doi: 10.1016/j.aanat.2017.09.006
- Steinberg, X., Lespay-Rebolledo, C., and Brauchi, S. (2014). A structural view of ligand-dependent activation in thermoTRP channels. *Front. Physiol.* 5, 171. doi: 10.3389/fphys.2014.00171
- Sun, D., Zhuang, X., Xiang, X., Liu, Y., Zhang, S., Liu, C., et al. (2010). A novel nanoparticle drug delivery system: the anti-inflammatory activity of curcumin is enhanced when encapsulated in exosomes. *Mol. Ther.* 18, 1606–1614. doi: 10.1038/mt.2010.105
- Sunshine, A., Laska, E. M., Olson, N. Z., Colon, A., Gonzalez, L., and Tirado, S. (1983). Analgesic effects of oral propiram fumarate, codeine sulfate and placebo in postoperative pain. *Pharmacother. J. Hum. Pharmacol. Drug Ther.* 3, 299–303. doi: 10.1002/j.1875-9114.1983.tb03281.x
- Tian, T., Zhang, H. X., He, C. P., Fan, S., Zhu, Y. L., Qi, C., et al. (2018). Surface functionalized exosomes as targeted drug delivery vehicles for cerebral ischemia therapy. *Biomaterials* 150, 137–149. doi: 10.1016/j.biomaterials.2017.10.012
- Tikhonova, S., Booi, L., D'Souza, V., Crosara, K. T. B., Siqueira, W. L., and Emami, E. (2018). Investigating the association between stress, saliva and dental caries: a scoping review. *BMC Oral Health* 18, 41. doi: 10.1186/s12903-018-0500-z
- Tonetti, M. S., Jepsen, S., Jin, L., and Otomo-Corgel, J. (2017). Impact of the global burden of periodontal diseases on health, nutrition and wellbeing of mankind: a call for global action. *J. Clin. Periodontol.* 44, 456–462. doi: 10.1111/jcpe.12732
- Torabinejad, M., and Chivian, N. (1999). Clinical applications of mineral trioxide aggregate. *J. Endod.* 25, 197–205. doi: 10.1016/S0099-2399(99)80142-3
- Torabinejad, M., Milan, M., Shabahang, S., Wright, K. R., and Faras, H. (2015). Histologic examination of teeth with necrotic pulps and periapical lesions treated with 2 scaffolds: an animal investigation. *J. Endod.* 41, 846–852. doi: 10.1016/j.joen.2015.01.026
- Trams, E. G., Lauter, C. J., Norman Salem, J., and Heine, U. (1981). Exfoliation of membrane ecto-enzymes in the form of micro-vesicles. *BBA - Biomembr.* 645, 63–70. doi: 10.1016/0005-2736(81)90512-5
- Trevisani, M., and Szallasi, A. (2010). Targeting TRPV1: Challenges and issues in pain management. *Open Drug Discov. J.* 2, 37–49. Ferrara Italy. Bentham Open. doi: 10.2174/1877381801002010037
- Tziafas, D. (2010). Dentinogenic potential of the dental pulp: facts and hypotheses. *Endod. Top.* 17, 42–64. doi: 10.1111/j.1601-1546.2010.00248.x
- Valadi, H., Ekström, K., Bossios, A., Sjöstrand, M., Lee, J. J., and Lötvall, J. O. (2007). Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic exchange between cells. *Nat. Cell Biol.* 9, 654–659. doi: 10.1038/ncb1596
- Vanegas, H., Vazquez, E., and Tortorici, V. (2010). NSAIDs, opioids, cannabinoids and the control of pain by the central nervous system. *Pharmaceut. (Basel).* 3, 1335–1347. doi: 10.3390/ph3051335
- Vos, T., Abajobir, A. A., Abbafati, C., Abbas, K. M., Abate, K. H., Abd-Allah, F., et al. (2017). Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 390, 1211–1259. doi: 10.1016/S0140-6736(17)32154-2
- Wang, W., Yi, X., Ren, Y., and Xie, Q. (2016). Effects of adenosine triphosphate on proliferation and odontoblastic differentiation of human dental pulp cells. *J. Endod.* 42, 1483–1489. doi: 10.1016/j.joen.2016.07.013
- Ward, P. A. (1974). The inflammatory mediators. *Ann. N. Y. Acad. Sci.* 221, 290–298. doi: 10.1111/j.1749-6632.1974.tb28228.x
- Wassel, M. O., and Khattab, M. A. (2017). Antibacterial activity against *Streptococcus mutans* and inhibition of bacterial induced enamel demineralization of propolis, miswak, and chitosan nanoparticles based dental varnishes. *J. Adv. Res.* 8, 387–392. doi: 10.1016/j.jare.2017.05.006
- Weihls, A., Fuchs, C., Teuschl, A., Hartinger, J., Slezak, P., Mittermayr, R., et al. (2014). Shock wave treatment enhances cell proliferation and improves wound healing by ATP release-coupled extracellular signal-regulated kinase (ERK) activation. *J. Biol. Chem. Sep* 26, 27090–27104. doi: 10.1074/jbc.M114.580936
- Wierichs, R. J., and Meyer-Lueckel, H. (2015). Systematic review on noninvasive treatment of root caries lesions. *J. Dent. Res.* 94, 261–271. doi: 10.1177/0022034514557330
- Won, J., Vang, H., Kim, J. H., Lee, P. R., Kang, Y., and Oh, S. B. (2018). TRPM7 mediates mechanosensitivity in adult rat odontoblasts. *J. Dent. Res.* 97, 1039–1046. doi: 10.1177/0022034518759947
- Wongrakpanich, S., Wongrakpanich, A., Melhado, K., and Rangaswami, J. (2018). A comprehensive review of non-steroidal anti-inflammatory drug use in the elderly. *Aging Dis.* 9, 143–150. doi: 10.14336/AD.2017.0306
- Xuan, K., Li, B., Guo, H., Sun, W., Kou, X., He, X., et al. (2018). Deciduous autologous tooth stem cells regenerate dental pulp after implantation into injured teeth. *Sci. Transl. Med.* 10, eaf3227. doi: 10.1126/scitranslmed.aaf3227
- Yang, F., and Zheng, J. (2017). Understand spiciness: mechanism of TRPV1 channel activation by capsaicin. *Protein Cell* 8, 169–177. doi: 10.1007/s13238-016-0353-7
- Yang, Y., Huang, L., Dong, Y., Zhang, H., Zhou, W., Ban, J., et al. (2014). In vitro antibacterial activity of a novel resin-based pulp capping material containing the quaternary ammonium salt MAE-DB and Portland cement. *PLoS One* 9, e112549. doi: 10.1371/journal.pone.0112549
- Zhang, M., Zang, X., Wang, M., Li, Z., Qiao, M., Hu, H., et al. (2019). Exosome-based nanocarriers as bio-inspired and versatile vehicles for drug delivery: recent advances and challenges. *J. Mater. Chem. B* 7, 2421–2433. doi: 10.1039/C9TB00170K
- Zhang, W., Vazquez, B., Oreadi, D., and Yelick, P. C. (2017). Decellularized tooth bud scaffolds for tooth regeneration. *J. Dent. Res.* 96, 516–523. doi: 10.1177/0022034516689082
- Zhu, J., Liang, R., Sun, C., Xie, L., Wang, J., Leng, D., et al. (2017). Effects of nanosilver and nanozinc incorporated mesoporous calcium-silicate nanoparticles on the mechanical properties of dentin. *PLoS One* 12, e0182583. doi: 10.1371/journal.pone.0182583

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Schuh, Benso and Aguayo. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.