

Case report

Open Access

***Stenotrophomonas maltophilia* prosthetic valve endocarditis: a case report**

Sophie Bayle¹, Clarisse Roverly¹, Pascal Sbragia², Didier Raoult³ and Philippe Brouqui*¹

Address: ¹Service des maladies infectieuses et tropicales, Hôpital Nord, Chemin des Bourrelys, 13915 Marseille cedex 20, France, ²Service de cardiologie, Hôpital Nord, Chemin des Bourrelys, 13915 Marseille cedex 20, France and ³Service de Microbiologie, Hôpital de la Timone, Marseille, France

Email: Sophie Bayle - sophiebayle@voila.fr; Clarisse Roverly - clarisse.roverly@medecine.univ-mrs.fr; Pascal Sbragia - pascal.sbragia@mail.ap-hm.fr; Didier Raoult - didier.raoult@medecine.univ-mrs.fr; Philippe Brouqui* - philippe.brouqui@medecine.univ-mrs.fr

* Corresponding author

Published: 23 May 2008

Received: 5 September 2007

Journal of Medical Case Reports 2008, **2**:174 doi:10.1186/1752-1947-2-174

Accepted: 23 May 2008

This article is available from: <http://www.jmedicalcasereports.com/content/2/1/174>

© 2008 Bayle et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Introduction: *Stenotrophomonas maltophilia* is an environmental bacterium increasingly involved in nosocomial infections and resistant to most antibiotics. It is important to recognize and efficiently treat infections with this bacterium as soon as possible.

Case presentation: We present a case of *Stenotrophomonas maltophilia* prosthetic valve endocarditis secondary to an indwelling catheter infection. The patient was cured without surgery. We review other cases of *S. maltophilia* endocarditis from the literature and describe the peculiarities of this case.

Conclusion: *S. maltophilia* endocarditis is a rare disease that is often hospital-acquired and related to an indwelling catheter infection. The high lethality is likely related to the intrinsic resistance of nosocomial bloodstream infections to commonly prescribed antibiotics.

Introduction

Stenotrophomonas maltophilia is a Gram-negative aerobic bacillus widely distributed in natural and human environments and has become increasingly responsible for nosocomial infections such as bacteremia, pneumonia, urinary tract infections, skin and soft tissue infections, ocular infections and meningitis [1]. It is generally considered as an opportunist pathogen with potential risk factors being: malignancies, debilitating illnesses, prior therapy with broad-spectrum antibiotics, chronic respiratory diseases, especially cystic fibrosis, prolonged endotracheal intubation and indwelling vascular catheters. *S. maltophilia* infective endocarditis (IE) is a rare and poorly described

disease [2-5]. Here, we present the 29th reported case of *S. maltophilia* endocarditis along with a literature review.

Case presentation

In November 2005, a 34-year-old Algerian woman was admitted to the infectious disease department for a suspected prosthetic mitral valve IE. She had a medical history of acute rheumatic fever when she was 10 years old. She had a mitral valve replacement with a prosthetic mechanic valve eight years ago. She was hospitalized at the end of a recent pregnancy because she needed intravenous heparin therapy in relay of oral anticoagulant for prevention of thrombosis of the mechanical valve prosthesis. Three days after delivery, she presented with a fever

and peripheral catheter-related infection. The catheter was in place for six days; it was inserted at the admission of the patient to the high-risk pregnancy unit. The site of the catheter insertion appeared inflammatory and indurate. All three blood cultures (automated blood culture BACTEC 9240 system, Becton Dickinson, Le pont de Claix, France) grew on day 2 with *S. maltophilia*. Transoesophageal echocardiography (TE) performed one day after the appearance of fever showed two small vegetations on the mitral mechanic valve without dehiscence. Based on the criteria established by Duke, the diagnosis of prosthetic mitral valve IE caused by *S. maltophilia* was confirmed. She was first treated empirically with vancomycin and gentamicin for 24 h. Antibiotics were then modified when we discovered Gram-negative bacillus growing in the blood cultures: imipenem and gentamicin were administered. She was treated with trimethoprim-sulfamethoxazole (SXT, trimethoprim (TMP) 4800 mg/day and sulfamethoxazole (SMZ) 960 mg/day) and gentamicin 3 mg/kg/day intravenously in accordance with antimicrobial susceptibility. In fact, effective treatment was started two days after the onset of disease and she had no fever on day 2 of this treatment. No cardiac complications occurred. Gentamicin was discontinued after seven days. SXT was then given *per os* (TMP 4800 mg/day and SMZ 960 mg/day). On day 8, a TE showed no vegetation. Antibiotics were continued for a total duration of six weeks. She remained well and showed no signs of fever and had a normal TE at her three-month follow-up visit.

Discussion

Here we present the 29th reported case of *S. maltophilia* endocarditis. *S. maltophilia* IE is a rare disease [1-5]. In 2002, Crum et al reported one case of *S. maltophilia* IE and reviewed 24 additional cases [3]. A Medline search revealed three additional cases of *S. maltophilia* IE [2,4,5]. The characteristics of reported cases of *S. maltophilia* IE are summarized in additional files 1 [see additional file 1]. Underlying risk factors were found in all patients (27/27) including 8/27 (29.6%) of intravenous drug users (IVDUs). Among *S. maltophilia* IE, 14/27 (51.94%) were of nosocomial origins (4/14, 28.6% being catheter-related infections and 10/14, 71.4% related to recent valve replacement (< 9 months) whereas 13/27 (48.1%) were community-acquired (8/13, 61.5% being acquired in IVDUs). *S. maltophilia* is intrinsically resistant to many common antibiotics and broad-spectrum β -lactams, such as imipenem. Combination antibiotherapy is indicated with SXT as the agent of choice [1-6]. Therapy for *S. maltophilia* infections is problematic because of the broad resistance to antibiotics that is characteristic of this organism. Antibiotic treatments currently used for nosocomial primary bloodstream infections [7], such as vancomycin and imipenem, have no effect on *S. maltophilia*. This may explain the high mortality reported (9/28, 32%) and the

frequency of complications (17/25, 68%) in *S. maltophilia* IE. Surgery is necessary in 9/17 (52.9%) cases involving prosthetic valves. Indeed, only 6/17 (35.3%) of *S. maltophilia* prosthetic valve IE are cured without surgery.

S. maltophilia endocarditis is a rare and severe disease, often nosocomial and related to central indwelling catheter infection. Our case is unusual as IE was secondary to peripheral catheter-related infection. As *S. Maltophilia* is resistant to the first line of antibiotic treatment used in these cases, empiric antibiotic treatment is often ineffective. Delay in effective antibiotic treatment is a major risk factor for mortality in *S. Maltophilia* bacteremia [8]. As a consequence, it is important to identify this microorganism as quickly as possible. Standard bacterial identification and susceptibility testing is normally a 48-h process. Polymerase chain reaction (PCR) can be used to decrease the identification time for Gram-negative bacteria isolated from blood culture, which has also been suggested for *Pseudomonas aeruginosa* infections [9]. In the light of this case report, it is also important to recall that guidelines on the prevention of intravascular device-related bloodstream infection have to be generalized and respected [10]. In fact, the catheter was left in place for too long in this previously healthy woman, and nosocomial infection could have easily been prevented in this patient.

Conclusion

S. maltophilia is a rare and severe disease, and is usually hospital-acquired. It is important to identify this microorganism as quickly as possible since *S. maltophilia* is resistant to first line antibiotic therapy used in case of nosocomial infections. This case report also emphasizes that prevention of intravascular device-related bloodstream infection is a major measure to prevent this very severe disease.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

SB, CR, PB and DR contributed to the clinical work-up of the patient and writing-up of the manuscript. PS was in charge of the patient in the department of cardiology and performed the echocardiography. All authors read and approved the final manuscript.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

Additional material

Additional file 1

Stenotrophomonas maltophilia infectious endocarditis case reports in the literature. The data provided represent the *Stenotrophomonas maltophilia* infectious endocarditis cases reported in the literature.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1752-1947-2-174-S1.doc>]

References

1. Candel FJ, Lopez R, Valdivia A, Nunez MJ, Roca-Arbones V, JJ PG: **[Endocarditis due to *Stenotrophomonas maltophilia*].** *Enferm Infecc Microbiol Clin* 2002, **20**:477-478.
2. Crum NF, Utz GC, Wallace MR: **Stenotrophomonas maltophilia endocarditis.** *Scand J Infect Dis* 2002, **34**:925-927.
3. Lopez RR, Lado Lado FL, Sanchez A, Rodriguez L I, Gamallo TR, Lorenzo Z V, Rodriguez-Otero L: **[Endocarditis caused by *Stenotrophomonas maltophilia*: report of a case and review of literature].** *An Med Interna* 2003, **20**:312-316.
4. Meimoun P, Mainardi JL, Berrebi A, Choucair J, Chevalier P, Buu-Hoi A, Gutmann L, Carpentier A: **[Stenotrophomonas maltophilia endocarditis following mitral valve prosthesis implantation. Report of a case].** *Arch Mal Coeur Vaiss* 1999, **92**:1389-1392.
5. Senol E: **Stenotrophomonas maltophilia: the significance and role as a nosocomial pathogen.** *J Hosp Infect* 2004, **57**:1-7.
6. Muder RR, Harris AP, Muller S, Edmond M, Chow JW, Papadakis K, Wagener MW, Bodey GP, Steckelberg JM: **Bacteremia due to *Stenotrophomonas (Xanthomonas) maltophilia*: a prospective, multicenter study of 91 episodes.** *Clin Infect Dis* 1996, **22**:508-512.
7. Mermel LA, Farr BM, Sherertz RJ, Raad II, O'Grady N, Harris JS, Craven DE: **Guidelines for the Management of Intravascular Catheter-Related Infections.** *Clin Infect Dis* 2001, **32**:1249-1272.
8. Wu PS, Lu CY, Chang LY, Hsueh PR, Lee PI, Chen JM, Lee CY, Chan PC, Chang PY, Yang TT, Huang LM: **Stenotrophomonas maltophilia bacteremia in pediatric patients-- a 10-year analysis.** *J Microbiol Immunol Infect* 2006, **39**:144-149.
9. Jaffe RI, Lane JD, Bates CW: **Real-time identification of *Pseudomonas aeruginosa* direct from clinical samples using a rapid extraction method and polymerase chain reaction (PCR).** *J Clin Lab Anal* 2001, **15**:131-137.
10. N.P. OG, M A, E.P. D, D.G. M, H. M, R.D MC, LA M, M.L P, II R, A R, R.A W: **Guidelines for the prevention of intravascular catheter-related infections.** *Clin Infect Dis* 2002, **35**:1281-1307.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

