

## Research Article

# Purification and the Secondary Structure of Fucoidanase from *Fusarium* sp. LD8

Wu Qianqian, Ma Shuang, Xiao Hourong, Zhang Min, and Cai Jingmin

Department of Biological and Environmental Sciences, Hefei University, Hefei 230022, China

Correspondence should be addressed to Cai Jingmin, caijingmin@hfu.edu.cn

Received 15 January 2011; Revised 25 April 2011; Accepted 25 July 2011

Copyright © 2011 Wu Qianqian et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The fucoidanase from *Fusarium* sp. (LD8) was obtained by solid-state fermentation. The fermented solid medium was extracted by citric acid buffer, and the extracts were precipitated by acetone and purified by Sephadex G-100 successively. The results showed that the specific fucoidanase activity of purified enzyme was 22.7-fold than that of the crude enzyme. The recovery of the enzyme was 23.9%. The purified enzyme gave a single band on SDS-PAGE gel, and the molecular weight of fucoidanase was about 64 kDa. The isoelectric point of the enzyme was 4.5. The enzyme properties were also studied. The results showed that the optimum temperature and pH were 60°C and 6.0, respectively; the temperature of half inactivation was 50°C, and the most stable pH for the enzyme was 6.0.  $K_M$ , and the  $V_{max}$  of the enzyme was 8.9 mg·L<sup>-1</sup> and 2.02 mg·min<sup>-1</sup>·mL<sup>-1</sup> by using fucoidan from *Fucus vesiculosus* as substrate. The compositions of the secondary structure of fucoidanase were estimated by FTIR, the second derivative spectra, and the curve-fitting analysis of the amide I bands in their spectra. The results showed that  $\beta$ -sheet was the dominant component (58.6%) and  $\alpha$ -helix was the least (12%); the content of  $\beta$ -turn and random coil were 15.39% and 14.5%, respectively.

## 1. Introduction

Sulfated polysaccharides, named fucoidan, could be extracted from many marine brown algae. The main differences of structural characterization in algal fucoidan originate from their species; the backbone of fucoidan isolated from *Laminaria cichorioides* [1] was consisting of  $\alpha$  (1–3) linked fucopyranose residues, and that from *L. japonica* was primarily  $\alpha$  (1–3) linked fucopyranose residues (75%) and a few  $\alpha$  (1–4) fucopyranose linkages (25%) [2]; The fucoidan from *Fucus serratus* [3], *F. evanescens* [4], and *F. distichus* [5] was shown to contain a backbone consisting of alternating (1–3) and (1–4) linked  $\alpha$ -L-fucopyranose residues.

Investigators were interested in screening fucoidan which was isolated from different brown algal species; fucoidan was attributed several different bioactivities including anticoagulant [6–8], antithrombotic/antithrombin activity [9], antiviral [10], and other activities [11]. The biological effects of fucoidan depended on the molecular mass, sulfate content, and sugar constituents [12, 13]. The low molecular weight fucoidan (LMWF) could be obtained by fucoidanase (E.C.3.2.1.44) enzymolysis, and it could hydrolyze fucoidan to sulfated LMWF without removal of its side substitute groups.

Fucoidanase could be extracted from hepatopancreas of invertebrates [14, 15], marine bacteria [16–25], and marine fungi [26–28].

There were some papers focusing on fermentation conditions of fucoidanase from marine fungus [26–28] and marine bacteria [23, 29], a few papers concerning purifications of fucoidanase from hepatopancreas [14] and marine bacteria reported [17, 20–22]. In order to elucidate the structure of fungal fucoidanase, we investigate fucoidanases isolated from different species of fungi. The present work is devoted to purification and the structure analysis of a fucoidanase from marine fungus *Fusarium* sp. LD8.

## 2. Materials and Methods

**2.1. Chemicals.** Fucoidan from *Fucus vesiculosus* was obtained from Sigma Chemical Co., Sephadex; G-100 was purchased from Pharmacia Biotech Corporation (Sweden). Fucoidan from *Laminaria* sp. was purchased from Rizhao Jiejing Ocean Biotechnology Development Co., Ltd (PRC). All of the other chemical reagents were analytical pure grade made in China.

**2.2. Microorganism.** The marine fungus LD8 was isolated from sand of North Sea in German which was identified as *Fusarium* sp. (LD8), and it could be used for the synthesis of fucoidanase. Solid-state medium for LD8 consisted of 7.5 g wheat bran, 0.5 g kelp powder, 0.1 g glucose, 0.04 g NaNO<sub>3</sub>, and 7.5 mL artificial sea water in 250 mL flasks [27].

**2.3. Chemical Analysis.** The protein concentration was determined by the Bradford method [30].

#### 2.4. Enzyme Assay

**2.4.1. Fucoïdanase Activity.** Fucoïdanase activity was measured by the dinitrosalicylic acid technique [31] to estimate the release of reducing sugars at 540 nm as follows: a mixture consisting of 1 mL substrate solution (1% fucoidan (w/v) from *Fucus vesiculosus* dissolved with 0.02 mol·L<sup>-1</sup> citric acid-sodium citric buffer, pH 6.0) and 0.1 mL enzyme solution (crude extract or pure enzyme) was incubated at 50°C for 10 min, using inactivated enzyme solution as blank CK. Using fucose as standard, the calibration curve function was  $y = 1.611x$  ( $x$ : the quantity of fucose, mg;  $y$ : the absorbance at 540 nm,  $R^2 = 0.9983$ ).

One unit (IU) of fucoïdanase activity is defined as the amount of enzyme that releases 1 μmol of fucose per minute under the assay conditions.

**2.4.2. Fucosidase Activity.** Fucosidase activity was measured under the following conditions: the reaction mixture contained 1 mL substrate solution (1% *p*-nitrophenyl- $\alpha$ -L-fucoside (w/v) dissolved with 0.02 mol·L<sup>-1</sup> citric acid-sodium citric buffer, pH 6.0) and 0.1 mL enzyme solution (purified enzyme) was incubated at 40°C for 2 h. One unit (IU) of fucosidase activity is defined as the amount of enzyme that releases 1 μmol of *p*-nitrophenyl per minute under the assay conditions.

**2.4.3. Amylase Assay.** The reaction mixture containing 1 mL of 2% soluble starch (w/v) in acetate buffer (pH 5.5) and 1 mL of enzyme solution was incubated with shaking at 40°C for 30 min. The reaction was stopped by boiling water for 5 min, and after centrifugation the released reducing sugar was measured by the dinitrosalicylic acid method [31]. One unit (IU) of amylase activity is defined as the amount of enzyme that liberated 1 μmol reducing sugar (as glucose) per min under the assay conditions.

**2.5. Extraction, Purification, and Purity Identification of Fucoïdanase.** The LD8 was cultivated at 28°C for 48 h on the solid-state medium in 250 mL flask.

All the following steps were accomplished at 4°C except being indicated specifically.

50 g fermented culture medium was extracted with citric acid-citric sodium buffer (pH 6.0) for 0.5 h. After being filtrated through six-layer carbasus, the filtrate was centrifuged at 15,000 g for 0.5 h, and then ice-cold acetone was added to the supernatant to a final concentration of 66.7% (v/v)

with gentle stirring. Insoluble material was obtained by centrifugation at 15,000 g for 0.5 h. The precipitate was dissolved in pH 6.0, 0.02 mol·L<sup>-1</sup> citric acid-citric sodium buffer and was centrifuged at 15,000 g for 0.5 h, and the clear solution was collected for use. The enzyme solution was concentrated to about 10 mL by low-temperature vacuum concentration and then loaded to Sephadex G-100 column (2.5 × 100 cm), which has been balanced well with 0.1 mol·L<sup>-1</sup> citric acid-citric sodium buffer. The enzyme was eluted at room temperature at the flow rate of 0.33 mL·min<sup>-1</sup>; fractions were collected at 20 min intervals. The fractions with the highest enzymatic activity were pooled, concentrated by low-temperature vacuum concentration to 10 mL, dialyzed in deionized water, lyophilized, and stored at -18°C.

SDS-PAGE was performed to identify the purity of the purified enzyme and calculate the  $M_w$  of the proteins in the gel. SDS-PAGE was 7% (w/v) polyacrylamide gel containing 10% (w/v) SDS; molecular markers were myosin (220 kDa),  $\alpha$ -2 macroglobulin (170 kDa),  $\beta$ -galactosidase (116 kDa), transferrin (76 kDa), and glutamic dehydrogenase (53 kDa), respectively. The protein was stained by Coomassie bright blue R-250.

**2.6. Characteristics of the Purified Enzyme.** Isoelectrofocusing was performed on gel rods. The electrode solutions were 2% (w/v) sodium hydroxide solution at the cathode and 5% (v/v) phosphoric acid solution at the anode. Each gel was focused in 100 V, 2 mA for 2 h, followed by 150–160 V, 4 mA for 5 h, which allowed for a sharp focalisation of fucoïdanase. Following isoelectric focusing, the gels were cut into 0.5 cm for measuring pH and enzyme activity. The protein was stained by 0.5% amino black (w/v).

The pH relative activity of the fucoïdanase was determined by detecting the fucoïdanase activity over a pH range of 3–8 with three kinds of buffer solutions (50 mmol·L<sup>-1</sup> sodium citrate buffer, 50 mmol·L<sup>-1</sup> sodium phosphate buffer, 50 mmol·L<sup>-1</sup> sodium carbonate buffer) at 50°C. The pH stability of the enzyme was studied by detecting the residual activity after the enzyme being incubated for 3 h at room temperature under different pH value. To obtain optimal reactive temperature, a mixture consisting of 1 mL substrate solution (1% fucoidan (w/v) from *Fucus vesiculosus* dissolved with 0.02 mol·L<sup>-1</sup> citric acid-sodium citric buffer, pH 6.0) and 0.1 mL enzyme solution was incubated at different temperatures for 10 min. For thermal stability study, 0.1 mL of enzyme solution was incubated at temperature range from 30 to 80°C for 1 h, then cooled rapidly in ice bath for 5 min, and then removed to 25°C. The residual enzyme activity was detected at 50°C for 10 min.

The substrate specificity of the enzyme was examined by using different fucoidan from *Fucus vesiculosus* and *Laminaria* sp. as substrates. Michaelis constants ( $K_M$ ) and maximum reaction velocities ( $V_{max}$ ) were calculated by the double-reciprocal plot method of Lineweaver and Burk [32].

**2.7. FTIR Assessment of the Secondary Structure of Fucoïdanase.** Infrared spectra were recorded on a MAGNA-IR750 Fourier transform spectrometer (INCOLET




FIGURE 1: Gel filtration column chromatography. Chromatography of fucoidanase obtained from the column of Sephadex G-100 ( $2.5 \times 100$  cm). After the column had been washed well with  $0.1 \text{ mol}\cdot\text{L}^{-1}$  citric acid-citric sodium buffer (pH 6.0), it was eluted with 2000 mL of the same buffer solution at a flow rate of  $0.33 \text{ mL}\cdot\text{min}^{-1}$  with 6.6 mL elute in each fraction.

INSTRUMENTW, USA). For the spectrum range from  $1500\sim 2200 \text{ cm}^{-1}$ , 32 scans were collected at a spectral resolution of  $4 \text{ cm}^{-1}$ . Pure fucoidanase (10 mg) was mixed with 100 mg of dried potassium bromide (KBr). Water vapor was purged from sample room. The spectrum of the amide I band of fucoidanase was obtained, and self-deconvolution and curve-fitting methods were used to analyze the secondary structure of the fucoidanase.

### 3. Result

**3.1. Purification of Fucoidanase.** Two protein peaks (E1 and E2) were observed. E2 peak showed fucoidanase activities (Figure 1) whose purity had been detected by SDS-PAGE (see below). The crude enzyme extraction was purified by 66.7% acetone precipitation and Sephadex G-100 gel chromatography (Table 1). The purification fold of fucoidanase activity of 1 mg protein was enhanced from 1-fold to 23.9-folds while recovery rate was decreased from 100% to 22.7%. The fractions with higher fucoidanase activity (tube no. 65 to 87) were pooled and concentrated to 10 mL. To exclude the reducing carbohydrates from carbon sources used for cultivation (starch, kelp polysaccharides), fucoidanase-related enzymes such as fucosidase and amylase activity of purified protein were determined, but there is no fucosidase or amylase activity (Table 2).

**3.2. Determination of Isoelectric Point and Molecular Mass of Fucoidanase.** The result exhibited that the purified

fucoidanase gave a single band on isoelectric electrophoresis, and the isoelectric point of the enzyme was pH 4.5 (Figure 2).

The purified fucoidanase gave a single band on SDS-PAGE gel, which suggested that relatively pure fucoidanase had been obtained. The molecular mass of the fucoidanase was about 64 kDa by SDS-PAGE (Figure 3) which was different from that of *Dendryphiella arenaria* TM94 (180 kDa). Molecular markers used were myosin (220 kDa),  $\alpha$ -2 macroglobulin (170 kDa),  $\beta$ -galactosidase (116 kDa), transferrin (76 kDa), and glutamic dehydrogenase (53 kDa), respectively.

**3.3. Effect of pH on Fucoidanase Activity and Stability.** Effect of pH on the activity of fucoidanase obtained from *Fusarium* sp. LD8 was shown in Figure 4. The maximum enzyme activity was at pH 6. The optimal pH of this enzyme was very close to that from marine fungus *Dendryphiella arenaria* TM94 and *Vibrio* sp. N-5, while the optimal pH of fucoidanase from Hepatopancreas of *Patinopecten yessoensis* was 5.5.

The effect of pH on stability was also determined. The results showed that the enzyme displayed stability at pH 6.0, whereas at pH 5.0 and 8.0, an activity loss of about 50% occurred after 6 h incubation at room temperature ( $25^\circ\text{C}$ ), respectively.

**3.4. Effect of Temperature on Fucoidanase Activity and Stability.** The optimum temperature for maximal activity of the fucoidanase was  $60^\circ\text{C}$  at pH 6.0 (Figure 5(a)). At  $30^\circ\text{C}$  the activity of fucoidanase decreased to 12.5%, while at  $80^\circ\text{C}$  to 18.75%. The result showed that the optimal temperature of fucoidanase from TM94 was higher than that of the fucoidanase in *Vibrio* sp. N-5, whose optimum temperature was  $40^\circ\text{C}$  [33].

The residual activity of fucoidanase was examined after preincubating it at different temperatures for 1 hr, and the temperature at which enzyme lost half activity was  $50^\circ\text{C}$  (Figure 5(b)). It was completely inactivated at above  $80^\circ\text{C}$ . The temperature of lost half activity is different from those of *Vibrio* sp. N-5 and *Dendryphiella arenaria* TM94. The enzyme of *Dendryphiella arenaria* TM94 and *Vibrio* sp. N-5 showed that their optimal temperature of half lost inactivation was at  $56^\circ\text{C}$  and  $40^\circ\text{C}$ , respectively [33].

**3.5. Determination of  $K_M$ ,  $V_{\max}$  and Affinity for Fucoidanase.** The kinetic parameters of fucoidanase were examined using *Fucus vesiculosus* fucoidan and *Laminaria* sp. fucoidan as substrates. The  $K_M$  values of the enzyme determined by Lineweaver-Burk method were  $8.9 \text{ mg}\cdot\text{L}^{-1}$  and  $10.9 \text{ mg}\cdot\text{mL}^{-1}$ , respectively. At the same time, the  $V_{\max}$  values for both substrates were  $2.02 \text{ mg}\cdot\text{min}^{-1}\cdot\text{mL}^{-1}$  and  $2.06 \text{ mg}\cdot\text{min}^{-1}\cdot\text{mL}^{-1}$ , respectively. The enzyme showed higher affinity to fucoidan from *Fucus vesiculosus*.

**3.6. The Second Structure of Fucoidanase.** The original IR spectrum for the fucoidanase was showed in Figure 6. The

TABLE 1: Summary of Purification of Fucoidanase.

Purification step	Total protein (mg)	Total activity (IU)	Specific activity (IU·mg <sup>-1</sup> )	Purification (fold)	Yield (100%)
Crude enzyme	11668.43	127.96	0.011	1	100
Acetone precipitation	509.23	80.43	0.16	14.5	62.8
Sephadex G-100	120.43	30.64	0.25	22.7	23.9




FIGURE 2: Determination of isoelectric point of the fucoidanase (a)  $R_f$  value of the pH; (b)  $R_f$  of the purified fucoidanase on original isoelectric focusing gel isoelectrofocusing was performed on gel rods. The electrode solutions were 2% (w/v) sodium hydroxide solution at the cathode and 5% (v/v) phosphoric acid solution at the anode. Each gel was focused in 100 V, 2 mA for 2 h, followed by 150–160 V, 4 mA for 5 h. Following isoelectric focusing, the gels were cut into 0.5 cm for measuring pH and enzyme activity. The protein was stained by 0.5% amino black.

TABLE 2: Fucoidanase and fucoidanase-related activity of purified protein.

Enzyme activity	Fucoidanase	E2 Fucosidase	Amylase
IU·mL <sup>-1</sup>	1.88	—	—

—: no enzyme activity.

bands at about 1620~1700 cm<sup>-1</sup> can be mainly attributed to the ( $\nu$ (C=O)) and is called amide I. The shape of the amide I band is representative of fucoidanase secondary structure. The bands of the 1650~1658 cm<sup>-1</sup>, 1600~1640 cm<sup>-1</sup>, 1640~1650 cm<sup>-1</sup>, 1660~1695 cm<sup>-1</sup> regions were, respectively, assigned to  $\alpha$ -helix,  $\beta$ -sheet, random coil, and  $\beta$ -turn structure.

The second derivative and a curve-fitting treatment can be carried out to estimate quantitatively the relative proportion of each component representing a type of secondary structure. The fourth derivative function was calculated by the PeakFit 4.12 software to determine the number of components in the amide I region for the second derivative spectra and the curve-fitting process (Table 3). According to this band composition, the amide I profile of fucoidanase contains four major components that can be linked with  $\alpha$ -helix,  $\beta$ -sheet, random coil, and  $\beta$ -turn structure where  $\beta$ -sheet is evidently the most intense component. The results showed that  $\beta$ -sheet was the dominant component (58.6%),  $\alpha$ -helix was the least (12%), and the content of  $\beta$ -turn and random coil were 15.39% and 14.5%.

## 4. Discussion

Fucoidanase could be isolated from marine invertebrates as *Haliotis* sp., sea cucumber, sponges, and molluscs [34] and was produced by marine bacteria as *Pseudomonas atlantica*, *P. carrageenovora*, *P. alteromonas*, *Vibrio* sp., *Bacillus* sp. HTP2, *Pseudoalteromonas citrea*, Family flavobacteriaceae [19, 24] and marine fungi *Dendryphiella arenaria* TM94 [26], *Fusarium* sp. LD8 [27], *Aspergillus niger*, *Penicillium purpurogenum*, *Mucor* sp. [28].

Fucoidanase had different molecular weight in different organisms. The molecular weight ( $M_w$ ) of LD8 fucoidanase was 64 kDa; it is close to that of fucoidanase E1, E2, and E3 of *Vibrio* sp. N-5 (39 kDa, 68 kDa, and 68 kDa, resp.) [33], while the  $M_w$  of LD8 fucoidanase was lower than that of fucoidanase from hepatopancreas of *Patinopecten yessoensis* (100–200 kDa) [14] and *Dendryphiella arenaria* TM94 (180 kDa) [35].

Fucoidanase from *Fusarium* sp. LD8 was more sensitive to pH and temperature. The catalytic activity of the fucoidanase of LD8 reached maximum at pH 6.0, which is very close to that of the fucoidanases from bacteria *Vibrio* sp. N-5 and *Dendryphiella arenaria* TM94. The enzyme activity decreased rapidly in LD8 when pH is below or above 6.0. At pH 5.0 and 7.0, the enzyme retained 68.2% and 86.4% of the enzyme activity at pH 6, respectively. The fucoidanase from LD8 had a relatively higher optimal temperature (60°C) compared with that of bacteria *Vibrio* sp. N-5 (37°C), Flavobacteriaceae SW5 (room temperature),






FIGURE 3: Determination of the  $M_w$  of the fucoidanase by SDS-PAGE. Picture of electrophoresis of fucoidanase: lane 1 is  $M_w$  marker; lane 2 is the fucoidanase; (b) standard curve and  $M_w$  function of the  $M_w$  marker SDS-PAGE was conducted using discontinuous electrophoresis method with Phastsystem, 5.5% (w/v) polyacrylamide, pH 8.3, 250 V, 10 mA, in the concentration gel with 7.0% polyacrylamide (w/v), 250 V, 30 mA, in the separation gel. Protein was stained with Coomassie brilliant blue R-250. Molecular markers used were (220 kDa),  $\alpha$ -2 macroglobulin (170 kDa),  $\beta$ -galactosidase (116 kDa), transferrin (76 kDa), and glutamic dehydrogenase (53 kDa).




FIGURE 4: Effects of pH on fucoidanase activity (a) and stability. (b) Effects of pH on fucoidanase activity was determined under the following method: 1.0 mL substrate solution (pH 3–8) in buffer solutions ( $50 \text{ mmol}\cdot\text{L}^{-1}$  sodium citrate buffer,  $50 \text{ mmol}\cdot\text{L}^{-1}$  sodium phosphate buffer,  $50 \text{ mmol}\cdot\text{L}^{-1}$  sodium carbonate buffer) was added to 0.1 mL enzyme solution and then was incubated at  $50^\circ\text{C}$  for 10 min. The pH stability of the enzyme was determined by assaying the residual activity after incubating enzyme for 3 h at the room temperature at different pH levels ranging from 3 to 9. Residual fucoidanase activity was determined by adding 0.1 mL enzyme solution to 1.0 mL of substrate solution (citric acid-citric sodium buffer, pH 6.0) and then was incubated at  $50^\circ\text{C}$  for 10 min.

and marine fungi *Dendryphiella arenaria* TM94 ( $50^\circ\text{C}$ ). The temperature of half inactivation of LD8 fucoidanase was  $50^\circ\text{C}$ , while that of bacteria *Vibrio* sp. N-5 was  $65^\circ\text{C}$  [33].

Effects of temperature on the secondary structure of LD8 fucoidanase were studied by Gaussian fitting to the deconvoluted spectra of fucoidanase at amide I region [36, 37].

A decrease of the  $\beta$ -turn structure and an increase of  $\alpha$ -helix of amide I region had been observed when treated temperature was below  $60^\circ\text{C}$ . While treated temperature was above  $60^\circ\text{C}$ , the contents of  $\alpha$ -helix,  $\beta$ -sheet, random coil, and  $\beta$ -turn had no changes. The above result was consistent with our conclusion that the optimal enzyme reaction tem-




FIGURE 5: Effects of temperature on fucoidanase activity (a) and stability. (b) Effects of temperature on fucoidanase activity was obtained by the following method: adding 0.1 mL of enzyme solution to 1.0 mL of substrate solution (citric acid-citric sodium buffer, pH 6.0), then incubating for 10 min at different temperatures from 30°C to 80°C, respectively. Temperature on stability of the fucoidanase was determined by assaying the residual activity under the following method: 0.1 mL of enzyme solution was incubated at 30 to 80°C for 1 h, rapidly cooled in an ice bath for 5 min, and then removed to 25°C. When the sample reached 25°C, 1.0 mL of substrate solution was added, and the residual enzyme activity was determined and expressed relative to the maximum activity.

TABLE 3: The results of Gaussian fitting of fucoidanase in amide I region.

Position	Area	Content (%)	Components (assignments)
1606	7.89140	21.1	$\beta$ -sheet
1618	7.40320	19.8	$\beta$ -sheet
1631	2.36538	17.7	$\beta$ -sheet
1645	5.40532	14.5	random
1658	4.31620	11.5	$\alpha$ -helix
1672	3.38760	9.06	$\beta$ -turn
1685	6.60977	6.33	$\beta$ -turn

perature was 60°C; The enzyme activity decreased rapidly in LD8 below or above pH 6.0. It was suggested that the enzyme activity of fucoidanase was closely related to the proportion of  $\alpha$ -helix and  $\beta$ -turn structure with no direct relation to  $\beta$ -sheet and random coil structure.

## Abbreviations

LMWF: The low molecular weight fucoidan  
 $K_M$ : Michaelis constants  
 $V_{max}$ : Maximum reaction velocities  
 $M_w$ : Molecular weight.

## Acknowledgments

This work was supported by funds from Anhui Provincial Nature Science Foundation (no. 03043104), Anhui Provin-




FIGURE 6: The FTIR spectra of fucoidanase. Fucoidanase (10 mg) was mixed with 100 mg of dried potassium bromide (KBr). Water vapor was purged from sample room. A baseline drawn between the spectra data points at 1500 and 2200 cm<sup>-1</sup>; 32 scans were collected at a spectral resolution of 4 cm<sup>-1</sup>.

cial Excellent Youth Science and Technology Foundation (no. 04043051), Anhui Advanced School Science and Research Fund Project for Young Teacher, Hefei University Scientific Research and Development Fund.

## References

- [1] S. D. Anastuyk, N. M. Shevchenko, E. L. Nazarenko et al., "Structural analysis of a highly sulfated fucan from the brown alga *Laminaria cichorioides* by tandem MALDI and ESI mass

- spectrometry," *Carbohydrate Research*, vol. 345, no. 15, pp. 2206–2207, 2010.
- [2] J. Wang, Q. Zhang, Z. Zhang, H. Zhang, and X. Niu, "Structural studies on a novel fucogalactan sulfate extracted from the brown seaweed *Laminaria japonica*," *International Journal of Biological Macromolecules*, vol. 47, no. 2, pp. 126–131, 2010.
- [3] M. I. Bilan, A. A. Grachev, A. S. Shashkov, N. E. Nifantiev, and A. I. Usov, "Structure of a fucoidan from the brown seaweed *Fucus serratus* L.," *Carbohydrate Research*, vol. 341, no. 2, pp. 238–245, 2006.
- [4] S. D. Anastyuk, N. M. Shevchenko, E. L. Nazarenko, P. S. Dmitrenok, and T. N. Zvyagintseva, "Structural analysis of a fucoidan from the brown alga *Fucus evanescens* by MALDI-TOF and tandem ESI mass spectrometry," *Carbohydrate Research*, vol. 344, no. 6, pp. 779–787, 2009.
- [5] M. I. Bilan, A. A. Grachev, N. E. Ustuzhanina, A. S. Shashkov, N. E. Nifantiev, and A. I. Usov, "A highly regular fraction of a fucoidan from the brown seaweed *Fucus distichus* L.," *Carbohydrate Research*, vol. 339, no. 3, pp. 511–517, 2004.
- [6] I. R. L. Albuquerque, K. C. S. Queiroz, L. G. Alves, E. A. Santos, E. L. Leite, and H. A. O. Rocha, "Heterofucans from Dictyota menstrualis have anticoagulant activity," *Brazilian Journal of Medical and Biological Research*, vol. 37, no. 2, pp. 167–171, 2004.
- [7] P. A. S. Mourão, "Use of sulfated fucans as anticoagulant and antithrombotic agents: future perspectives," *Current Pharmaceutical Design*, vol. 10, no. 9, pp. 967–981, 2004.
- [8] W. J. Kim, Y. K. Koo, M. K. Jung et al., "Anticoagulating activities of low-molecular weight fuco-oligosaccharides prepared by enzymatic digestion of fucoidan from the sporophyll of Korean *Undaria pinnatifida*," *Archives of Pharmacal Research*, vol. 33, no. 1, pp. 125–131, 2010.
- [9] T. Nishino, T. Yamauchi, M. Horie, T. Nagumo, and H. Suzuki, "Effects of a fucoidan on the activation of plasminogen by u-PA and t-PA," *Thrombosis Research*, vol. 99, no. 6, pp. 623–634, 2000.
- [10] D. J. Schaeffer and V. S. Krylov, "Anti-HIV activity of extracts and compounds from algae and cyanobacteria," *Ecotoxicology and Environmental Safety*, vol. 45, no. 3, pp. 208–227, 2000.
- [11] H. Maruyama, H. Tamauchi, M. Hashimoto, and T. Nakano, "Antitumor activity and immune response of Mekabu fucoidan extracted from Sporophyll of *Undaria pinnatifida*," *In Vivo*, vol. 17, no. 3, pp. 245–249, 2003.
- [12] T. Nishino, C. Nishioka, H. Ura, and T. Nagumo, "Isolation and partial characterization of a novel amino sugar-containing fucan sulfate from commercial *Fucus vesiculosus* fucoidan," *Carbohydrate Research*, vol. 255, pp. 213–224, 1994.
- [13] L. Chevolot, A. Foucault, F. Chaubet et al., "Further data on the structure of brown seaweed fucans: relationships with anticoagulant activity," *Carbohydrate Research*, vol. 319, no. 1–4, pp. 154–165, 1999.
- [14] T. Fujikawa, K. Koyabu, and M. Wada, "Enzymes in hepatopancreas of abalone active on fucoidan(1), Crude enzyme and unabsorbed fraction on CM-cellulose," *Nippon Nogeikagaku Kaishi*, vol. 53, pp. 87–95, 1979 (Japanese).
- [15] K. Kitamura, M. Masaru, and T. Yasui, "Enzymic degradation of fucoidan by fucoidanase from the hepatopancreas of *Patinopecten yessoensis*," *Bioscience, Biotechnology, and Biochemistry*, vol. 56, pp. 490–494, 1992.
- [16] W. Yaphe and K. Morgan, "Enzymic hydrolysis of fucoidin by *Pseudomonas atlantica* and *pseudomonas carrageenovora*," *Nature*, vol. 183, no. 4663, pp. 761–762, 1959.
- [17] S. Furukawa, T. Fujikawa, D. Koga, and A. Ide, "Production of fucoidan-degrading enzymes, fucoidanase, and fucoidan sulfatase by *Vibrio* sp. N-5," *Nippon Suisan Gakkaishi*, vol. 58, pp. 1499–1503, 1992 (Japanese).
- [18] I. Y. Bakunina, O. I. Nedashkovskaya, S. A. Alekseeva et al., "Degradation of fucoidan by the marine proteobacterium *Pseudoalteromonas citrea*," *Mikrobiologiya*, vol. 71, pp. 41–47, 2002.
- [19] T. Sakai, H. Kimura, and I. Kato, "A marine strain of Flavobacteriaceae utilizes brown seaweed fucoidan," *Marine Biotechnology*, vol. 4, no. 4, pp. 399–405, 2002.
- [20] T. Sakai, K. Ishizuka, and I. Kato, "Isolation and characterization of a fucoidan-degrading marine bacterium," *Marine Biotechnology*, vol. 5, no. 5, pp. 409–416, 2003.
- [21] T. Sakai, K. Ishizuka, K. Shimanaka, K. Ikai, and I. Kato, "Structures of oligosaccharides derived from *Cladosiphon okamuranus* fucoidan by digestion with marine bacterial enzymes," *Marine Biotechnology*, vol. 5, no. 6, pp. 536–544, 2003.
- [22] T. Sakai, H. Kimura, and I. Kato, "Purification of sulfated fucoglucuronomannan lyase from bacterial strain of *Fucobacter marina* and study of appropriate conditions for its enzyme digestion," *Marine Biotechnology*, vol. 5, no. 4, pp. 380–387, 2003.
- [23] P. Wang, J. Cai, S. Qin et al., "Fermentation of marine bacterium *Bacillus* sp. H-TP2 for fucoidanase and enzyme properties," *Food and Fermentation Industry*, vol. 30, pp. 13–15, 2004 (Chinese).
- [24] A. M. Urvantseva, I. Y. Bakunina, O. I. Nedashkovskaya, S. B. Kim, and T. N. Zvyagintseva, "Distribution of intracellular fucoidan hydrolases among marine bacteria of the family Flavobacteriaceae," *Applied Biochemistry and Microbiology*, vol. 42, no. 5, pp. 484–491, 2006.
- [25] V. Descamps, S. Colin, M. Lahaye et al., "Isolation and culture of a marine bacterium degrading the sulfated fucans from marine brown algae," *Marine Biotechnology*, vol. 8, no. 1, pp. 27–39, 2006.
- [26] K. Wu, Q. Wu, J. Yan et al., "Production of fucoidanase from the marine fungus *Dendryphiella arenaria* TM94 by solid substrate fermentation," *ESMB*, vol. 5, pp. 12–14, 2002.
- [27] Q. Wu, J. Cai, K. Wu, and S. Qin, "Solid state fermentation of fucoidanase of marine fungus LD8. Marine Sciences," *Marine Sciences*, vol. 28, pp. 23–27, 2004 (Chinese).
- [28] R. M. R. Jasso, C. N. A. Gonzalez, L. Pastrana, and J. A. Teixeira, "Identification and evaluation of fungal strains with fucoidan degradation potential," in *Proceedings of the 10th International Chemical and Biological Engineering Conference (CHEMPOR '08)*, pp. 2106–2109, Braga, Portugal, September 2008.
- [29] T. Sakai, T. Kawai, and I. Kato, "Isolation and characterization of a fucoidan-degrading marine bacterial strain and its fucoidanase," *Marine Biotechnology*, vol. 6, no. 4, pp. 335–346, 2004.
- [30] M. M. Bradford, "A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding," *Analytical Biochemistry*, vol. 72, no. 1–2, pp. 248–254, 1976.
- [31] G. L. Miller, "Use of dinitrosalicylic acid reagent for determination of reducing sugar," *Analytical Chemistry*, vol. 31, no. 3, pp. 426–428, 1959.
- [32] H. Lineweaver and D. Burk, "The determination of enzyme dissociation constants," *Journal of the American Chemical Society*, vol. 56, no. 3, pp. 658–666, 1934.

- [33] S. Furukawa, T. Fujikawa, D. Koga, and A. Ide, "Purification and some properties of exo-type fucoidanase from *Vibrio* sp. N-5," *Bioscience, Biotechnology, and Biochemistry*, vol. 56, pp. 1829–1834, 1992.
- [34] R. Daniel, O. Berteau, J. Jozefonvicz, and N. Goasdoue, "Degradation of algal (*Ascophyllum nodosum*) fucoidan by an enzymatic activity contained in digestive glands of the marine mollusc *Pecten maximus*," *Carbohydrate Research*, vol. 322, no. 3-4, pp. 291–297, 1999.
- [35] Q. Wu, M. Zhang, K. Wu, B. Liu, J. Cai, and R. Pan, "Purification and characteristics of fucoidanase obtained from *Dendryphiella arenaria* TM94," *Journal of Applied Phycology*, vol. 23, no. 2, pp. 197–203, 2011.
- [36] S. Ma, *The studies on the preparation, enzymatic properties and characterization of fucoidanase from marine Fusarium*, M.S. thesis, Anhui Agriculture University, China, 2006.
- [37] S. Ma, H. Xiao, Q. Wu, R. Pan, and J. Cai, "Investigation of fucoidanase by FTIR spectra," *Spectroscopy and Spectral Analysis*, vol. 28, pp. 590–593, 2010 (Chinese).