

RESEARCH ARTICLE

Open Access



# Occurrence, virulence genes, and antimicrobial profiles of *Escherichia coli* O157 isolated from ruminants slaughtered in Al Ain, United Arab Emirates

Dawood Al-Ajmi<sup>\*</sup> , Shafeeq Rahman and Sharmila Banu

## Abstract

**Background:** Shiga toxin-producing *Escherichia coli* (STEC) is a major source of food-borne illness around the world. *E. coli* O157 has been widely reported as the most common STEC serogroup and has emerged as an important enteric pathogen. Cattle, in particular have been identified as a major *E. coli* O157:H7 reservoir of human infections; however, the prevalence of this organism in camels, sheep, and goats is less understood. The aim of this study was to evaluate the occurrence and concentration of *E. coli* serotype O157 in the feces of healthy camels ( $n = 140$ ), cattle ( $n = 137$ ), sheep ( $n = 141$ ) and goats ( $n = 150$ ) slaughtered in United Arab Emirates (UAE) for meat consumption between September 2017 and August 2018. We used immunomagnetic separation coupled with a culture-plating method to detect *E. coli* O157. Non-sorbitol fermenting colonies were assessed via latex-agglutination testing, and positive cultures were analyzed by performing polymerase chain reactions to detect genes encoding attaching and effacing protein (*eaeA*), hemolysin A (*hlyA*, also known as *ehxA*) and Shiga toxin (*stx1* and *stx2*), and *E. coli* O157:H7 specific genes (*rfb* O157, *uidA*, and *fliC*). All *E. coli* O157 isolates were analyzed for their susceptibility to 20 selected antimicrobials.

**Results:** *E. coli* O157 was observed in camels, goats, and cattle fecal samples at abundances of 4.3, 2, and 1.46%, respectively, but it was undetectable in sheep feces. The most prevalent *E. coli* O157 gene in all STEC isolates was *stx2*; whereas, *stx1* was not detected in any of the samples. The fecal samples from camels, goats, and cattle harbored *E. coli* O157 isolates that were 100% susceptible to cefotaxime, chloramphenicol, ciprofloxacin, norfloxacin, and polymyxin B.

**Conclusion:** To our knowledge, this is the first report on the occurrence of *E. coli* O157 in slaughter animals in the UAE. Our results clearly demonstrate the presence of *E. coli* O157 in slaughtered animals, which could possibly contaminate meat products intended for human consumption.

**Keywords:** *Escherichia coli* O157, Antimicrobial resistance, Food pathogen testing

\* Correspondence: [dawoods@uaeu.ac.ae](mailto:dawoods@uaeu.ac.ae)

Department of Integrative Agriculture, College of Food and Agriculture, United Arab Emirates University (UAEU), Al Ain, United Arab Emirates



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Background

Access to safe food is a basic human need but remains a major health concern across the world, with food-borne infections considered as a major global public health issue. These infections not only affect the health and well-being of consumers, but also adversely impact economies in food-exporting countries. Consequently, the frequency with which food items are rejected at border crossings due to microbial and chemical contaminants has increased dramatically over the past few years, resulting in huge economic losses and food wastage for both importing and exporting countries [1, 2]. Specifically, the lack of efficient food-safety programs in Arab countries prevented the export of food products into the international market in 2003 [3]. Raw meat, salads, and unpasteurized dairy products pose important food-contamination risks and increase the burden of food-borne diseases [4]. However, a few Arab countries have carried out microbial-contamination profiling regarding food-borne outbreaks [3]. As a result, 1926 cases of food-borne illness were reported in Lebanon in 2003 and 779 cases were reported in Libya in 2004, while 112, 904 cases of acute gastroenteritis and diarrhea were reported in Oman in 2002 [3].

Recently, enterohemorrhagic strains of *Escherichia coli* have emerged as significant enteric pathogens. During the early 1980s, various serotypes were implicated in human disease, and *E. coli* O157 was the most prevalent. This serotype is responsible for severe abdominal illness, specifically enterohemorrhagic colitis (HC) and hemolytic uremic syndrome (HUS), and generally causes severe diarrhea [5, 6]. *E. coli* O157 belongs to the larger category of Shiga toxin-producing *E. coli* (STEC), which can produce Shiga toxin type 1 (*Stx*<sub>1</sub>), Shiga toxin type 2 (*Stx*<sub>2</sub>), or both, along with other variants [7]. *E. coli* O157 can infect humans via various routes; however, a large proportion of infections and human outbreaks have occurred following the consumption of contaminated food products, such as meat and dairy products [8]. Although the prevalence of STEC infections is often low, major outbreaks resulting in serious medical conditions have occurred throughout the world [9].

Typically, healthy cattle serve as the primary reservoir for STEC, but they are also carried by sheep and other animals [10]. Several host and farm practices have been associated with an increasing *E. coli* O157:H7 prevalence, including the animal age, season, and herd type [11, 12]. Currently, another major concern for human health is the rise in antimicrobial resistance due to the overuse of antibiotics in livestock production and human diseases treatment in developing countries [13, 14]. Various studies have been conducted worldwide to better understand the antimicrobial-resistance profiles of food-borne pathogens. However, no studies have been

conducted to study the burden and drug-sensitivity profile of *E. coli* O157 in the United Arab Emirates (UAE). An abattoir or slaughterhouse is a place that has been approved by an authority to practice hygienic slaughtering and processing of meat products for human consumption [15]. Due to increases in the global population and urbanization, improper slaughterhouse practices can facilitate the growth and spread of pathogenic organisms if the slaughtered animals harbor pathogens in their feces. Thus, it is necessary to determine the prevalence and concentration of *E. coli* O157 in fecal samples in order to evaluate the extent of the risk for contamination. Such assessment would necessitate the development of cost-effective and safe interventions to control *E. coli* O157 shedding at the farm level to prevent further food-chain contamination.

Thus, this study aimed to evaluate the occurrence, concentration, virulence genes, and antimicrobial-resistance profiles of *E. coli* O157 in feces of healthy camels, cattle, sheep, and goats slaughtered for meat consumption. This is anticipated to be the first report of the occurrence and abundance of *E. coli* O157 in these animals within the UAE.

## Results

Examination of fecal samples collected from a slaughterhouse in Al Ain, UAE over ten months revealed that the occurrences of *E. coli* serotype O157 in camels, goats, and cattle were 4.3, 2, and 1.46%, respectively. However, *E. coli* O157 was not detected in the sheep fecal samples. Furthermore, no significant differences were observed in the occurrence of *E. coli* O157 between male and female animals or between different breeds. Given that no other groups have reported the prevalence of *E. coli* O157 in slaughtered animals in the UAE, we performed microbiological and molecular tests to confirm the presence of *E. coli* O157 strains.

Biochemical tests confirmed the presence of *E. coli* O157 strains. The strains were sorbitol and indole-negative, but positive for citrate utilization. Bacterial enumeration was carried out by plating the samples and manually counting the colonies to quantify the colony-forming units (CFUs). The CFU concentrations (CFU g<sup>-1</sup> feces) are presented in Table 1. The results indicate that the fecal concentrations of *E. coli* O157 were sufficiently high to promote the spread of infection. We also sought to understand the seasonal prevalence of *E. coli* O157 and found that during the ten-month study, *E. coli* O157 isolates were only obtained from the samples in February, March, and April. According to the National Center of Meteorology, the temperature range during these three months was 37 °C–45 °C. There was no statistically significant difference ( $p > 0.05$ ) in the prevalence of *E. coli* O157 between sex, origin, or breed.

**Table 1** *E. coli* O157-positive cases and the *E. coli* O157 concentrations in animal feces

Species	Number of animals	<i>E. coli</i> O157-positive animals	Percentage (%)	<i>E. coli</i> O157 Concentration (mean CFU per g feces)
Camel	140	6	4.3	$4 \times 10^4$
Goat	150	3	2.0	$4 \times 10^4$
Cattle	137	2	1.46	$2 \times 10^3$
Sheep	141	0	0.0	0

CFU colony-forming unit, used to estimate the number of viable bacteria cells in a sample

However, due to the low number of positive samples and the prevalence, it was difficult to perform a comprehensive statistical analysis.

#### Polymerase chain reaction (PCR) analysis

Out of the 12 positive camel fecal samples, five (42%) lacked the *fliCH7* gene, and the virulence gene, *hlyA*, was present in nine samples (75%). In addition, the *eaeA* gene was present in 11 of the 12 camel fecal samples. Similarly, of the two positive cattle samples, one contained the *flicH7* gene. Further, the *hlyA* gene was present in only one of the cattle samples, whereas the *eaeA* gene was present in both samples. All ten *E. coli* O157-positive goat samples (100%) lacked the *flicH7* gene. Alternatively, the *hlyA* and *eaeA* genes were present in 6/10 (60%) of the goat samples. The most prevalent gene in the *E. coli* O157 isolates from all species was *stx<sub>2</sub>*, whereas *uidA* and *stx<sub>1</sub>* was absent in all samples (Table 2). For some positive culture samples, we were unable to confirm the presence of *E. coli* O157 by PCR amplification, and those samples were considered negative for *E. coli* O157.

#### Antimicrobial susceptibility of isolates

All samples isolated from goats, cattle, and camels were subjected to antimicrobial susceptibility testing. We found that all (100%) of the *E. coli* O157 isolates were susceptible to cefotaxime, chloramphenicol, ciprofloxacin, norfloxacin, and polymyxin B. Kanamycin showed growth-inhibition zones with 20 isolates out of 24 positive samples. Other antimicrobials showed irregular susceptibility and resistance patterns that could not be considered effective (Table 3).

#### Discussion

Human *E. coli* O157:H7 infections primarily originate from animal food sources [16]. Specifically, domestic ruminants, including cattle, sheep, and goats, have been reported to be major natural reservoirs for *E. coli* O157 and contribute significantly to the epidemiology of human infections [17]. In this study, we surveyed the

**Table 2** Distribution of virulent genes detected by PCR

Sample number	Species	Genes						
		<i>rfbE</i>	<i>flicH7</i>	<i>hlyA</i>	<i>uidA</i>	<i>eaeA</i>	<i>stx<sub>2</sub></i>	<i>stx<sub>1</sub></i>
1	Camel	+	+	-	-	+	+	-
2	Camel	+	+	+	-	+	+	-
3	Camel	+	-	-	-	+	+	-
4	Camel	+	+	-	-	-	+	-
5	Camel	+	+	+	-	+	+	-
6	Camel	+	-	+	-	+	+	-
7	Camel	+	-	+	-	+	+	-
8	Camel	+	+	+	-	+	+	-
9	Camel	+	+	+	-	+	+	-
10	Camel	+	+	+	-	+	+	-
11	Camel	+	-	+	-	+	+	-
12	Camel	+	-	+	-	+	+	-
13	Cattle	+	-	-	-	+	+	-
14	Cattle	+	+	+	-	+	+	-
15	Goat	+	-	-	-	-	+	-
16	Goat	+	-	+	-	+	+	-
17	Goat	+	-	+	-	+	+	-
18	Goat	+	-	-	-	-	+	-
19	Goat	+	-	-	-	-	+	-
20	Goat	+	-	+	-	+	+	-
21	Goat	+	-	+	-	+	+	-
22	Goat	+	-	+	-	+	+	-
23	Goat	+	-	+	-	+	+	-
24	Goat	+	-	-	-	-	+	-

occurrence and concentration of *E. coli* O157 (CFU g<sup>-1</sup> feces) in fecal samples from animals in a slaughterhouse in Al Ain, UAE. We successfully identified *E. coli* O157 in 4.3, 2, and 1.46% of the camel, goat and cattle samples, respectively, with concentrations ranging from TFTC (too few to count) to  $4 \times 10^4$  CFU g<sup>-1</sup> feces. This observed *E. coli* O157:H7 prevalence in the UAE slaughterhouse is in agreement with previous studies carried out in countries such as Ethiopia, South Africa, the United Kingdom, and Ireland, where the prevalences of *E. coli* O157:H7 in abattoirs were reported as 2.7, 2.8, 2.9 and 3.2% respectively [18–22]. Furthermore, in this study, the highest prevalence of *E. coli* O157 (4.3%) was found in camel feces. This result was consistent with the reports from neighboring countries, including Saudi Arabia (2.4%) [23], but slightly lower than those found in Ethiopia (8%) and Iran (6.4–9.6%) [24, 25]. Moreover, reports from Qatar showed that *E. coli* O157 was ten times more abundant in camel fecal samples than in cattle fecal samples [26]. Interestingly, Al-Gburi (2016) [27] reported that in Iraq, camels harbored the *E. coli* O157

**Table 3** Antimicrobials used, their symbols, and corresponding zones of inhibition for gram-negative enteric bacteria

Antimicrobial used	Symbol	Diameter of zone of inhibition (mm)		
		Resistant	Intermediate	Susceptible
Amoxicillin (25 µg)	AML	≤13	14–16	≥17
Ampicillin (10 µg)	AMP	≤13	14–17	≥18
Bacitracin (10 µg)	B	≤14	15–16	≥17
Cefotaxime (30 µg)	CTX	≤14	15–22	≥23
Cefoxitin (30 µg)	FOX	≤14	15–17	≥18
Ceftazidime (10 µg)	CAZ	≤12	13–17	≥18
Cefuroxime sodium (5 µg)	CXM	≤14	15–22	≥23
Chloramphenicol (30 µg)	CHL	≤14	15–20	≥21
Ciprofloxacin (30 µg)	CIP	≤15	16–20	≥21
Cloxacillin (5 µg)	CLOXA	≤13	14–16	≥17
Doxycycline (30 µg)	DO	≤13	14–16	≥17
Gentamycin (10 µg)	GEN	≤12	13–15	≥16
Kanamycin (30 µg)	K	≤13	14–17	≥18
Nalidixic acid (30 µg)	NAL	≤13	14–18	≥19
Nitrofurantoin (300 µg)	F	≤13	14–17	≥18
Norfloxacin (10 µg)	NOR	≤12	13–16	≥17
Polymyxin B (300 units)	PMB	≤8	9–11	≥12
Streptomycin (10 µg)	STR	≤11	12–14	≥15
Tetracycline (30 µg)	TET	≤12	15–18	≥19
Vancomycin (30 µg)	VAN	≤14	15–16	≥17

most frequently (19%) among all animals tested, and these isolates were found to be multi-drug resistant.

Previously, researchers studying UAE camels were unable to isolate *E. coli* O157 from their fecal samples. A study conducted by Moore et al. [28] on racing camel calves from the UAE failed to show the presence of *E. coli* O157, and similar results were obtained by El-Sayed et al. [29] in a large herds of camels. However, this discrepancy with our results may be due to the absence of a reliable “gold standard” for detecting the presence of STEC *E. coli* O157 in samples, as well as the inherent difficulty associated with directly comparing the results obtained in this study with those obtained in other studies that used different culture methods. In addition, results from above studies were primarily based on the characterization of very few colonies picked from the samples, without specific screening and isolation techniques. In contrast, rather than using a direct culturing method, we employed enrichment in buffered-peptone water (BPW), followed by immunomagnetic separation of *E. coli* O157 (IMS) and culture on cefixime-tellurite sorbitol MacConkey (CT-SMAC) agar, to improve the isolation efficiency [30]. All isolates were confirmed as *E. coli* O157 via latex agglutination testing and PCR.

However, we were unable to identify *E. coli* O157 strains from sheep fecal samples. Similar results were

published by Alhelfi et al. [31], where no *E. coli* O157 was detected in either cattle or sheep fecal samples within the target area studied in the United Kingdom. The failure to detect positive samples may be indicative of several factors, such as the study design, sample size, geographical origin, age and sex of the animals, abattoir conditions, animal husbandry, and/or the diet, which have been shown to impact the prevalence rates of *E. coli* O157 in livestock [32, 33]. In addition, a previous prevalence study [34] on suckling lambs did not detect *E. coli* O157 in all the 265 animals studied. However, it is worth noting that the samples in that study were not collected during the summer months, which might have affected the presence of *E. coli* O157 strains. The relationship between *E. coli* O157 and the gastrointestinal microbiome flora is not well understood. Indeed, factors encoded within the locus of enterocyte effacement (LEE), such as intimin, the translocated intimin receptor (*tir*), the *EspA* translocator, and other candidate bacterial factors have been implicated in the colonization and persistence of the strain in tested experimental ruminants such as sheep and cattle [35, 36]. It is important to note that in those previous experiments, the researchers screened neither for other pathogenic serogroups of *E. coli* that may be prevalent in sheep, nor for other bacterial pathogens. Thus, the absence of these strains in sheep

does not indicate that sheep pose zero risk. Our study did not have access to sufficient information regarding the local sheep population, including animal ages, breeds, husbandry, and diet history. Hence, additional studies on larger herds, using a meta-analysis approach, are recommended to confirm the absence of *E. coli* O157 in local sheep population.

Although cattle were previously reported to be the most common *E. coli* O157 reservoir [37], in our study we only observed a prevalence of approximately 1.46%. Alternatively, data from a study carried out in Riyadh, Saudi Arabia, showed that 10.7% of cattle feces samples contained *E. coli* O157:H7. Moreover, cattle from Jordan had a higher prevalence of *E. coli* O157:H7 (12.22%) than cattle from other countries in Asia [38], and the lowest prevalence was reported in Taiwan (0.13%) [39]. In another report, it was shown that the prevalence rate of *E. coli* O157:H7 in fecal samples from cattle throughout the world ranged from 2.4 to 24% [40].

We also examined whether seasonal variation occurred in the prevalence of *E. coli* O157, as such fluctuations in cattle and meat products reportedly influence human *E. coli* O157 infection rates [41, 42]. Specifically, the prevalence in cattle feces are low in the winter and high in the spring, peaking in summer [43]. The warmer summer months may provide more suitable environments for *E. coli* O157:H7 survive and potentially multiply outside of the host in soil, feed, and water, resulting in a continual source of infection or re-infection of cattle populations. Al Ain, UAE, is located in a tropical dry area where the average temperature during different seasons can enhance bacterial growth. In fact, previous findings have shown that the prevalence of *E. coli* O157 increased during the summer months and declined in the winter [44]. Our results showed that the highest prevalence occurred in spring and the early summer months (February to April). This may likely reflect the adverse effect of the intense heat during the summer months in the UAE, when daytime temperatures may reach as high as 50 °C, causing *E. coli* O157:H7 to decline due to the inability of the organism to persist in the extreme environment. However, our findings are in agreement with previous studies carried out in Riyadh, Saudi Arabia, where the highest prevalence was reported in the spring and early autumn months [23]. In fact, we were unable to detect *E. coli* O157:H7 in any other months, similar to what reported in other parts of the world.

The concentrations of *E. coli* O157 within the animal fecal samples were also quantified. In camels, the CFUs ranged from TFTC to  $4 \times 10^4$  CFU g<sup>-1</sup> feces, whereas the goat and cattle samples contained  $4 \times 10^4$  CFU g<sup>-1</sup> feces and  $2 \times 10^3$  CFU g<sup>-1</sup> feces, respectively. These results demonstrate that animals infected with *E. coli* O157 could be considered super shedders. Such high

fecal *E. coli* O157 levels present the carcasses of these animals as a high risk for contamination if the proper precautions are not taken. Animals in Scottish slaughterhouses reportedly shed bacteria at a level of  $> 10^4$  CFU/g, constituting  $> 96\%$  of the bacteria that were shed by all animals tested [45]. Together, these findings underscore the importance of the presence of high-shedding animals in a herd to the overall prevalence of colonization in the entire cattle population [45].

Shiga toxins are associated with HC and HUS, while intimin is responsible for attaching/effacing (A/E) lesions on intestinal epithelial cells. As such, enterohemolysins have been proposed as potential epidemiological markers for STEC strains [46]. Intimin, encoded by the *eaeA* gene, adheres to intestinal mucosa and causes intestinal lesion formation [47]. Enterohemolysin, encoded by *hlyA*, leads to erythrocyte lysis, which may contribute to iron intake by intestinal bacteria [48]. Previous reports showed that *E. coli* O157:H7/H<sup>-</sup> strains isolated from the feces of slaughtered ruminants exhibited *stx2* gene prominence over the *stx1* gene [34, 49, 50]. In our study, *stx1* was absent from all strains irrespective of the species. Data from various studies have shown that *stx2* and *eaeA* are clinically important virulence genes, and the carriage of these genes was associated with the severity of human disease, especially HUS [51, 52]. Hence, our present data suggest that *E. coli* O157 strains harboring the major virulence genes may be more virulent to humans than strains lacking these genes. Nevertheless, it has been reported that the production of major virulence genes are not essential for pathogenesis, as several sporadic cases of HUS were induced by *Stx* and *eaeA*-negative strains [53].

Antimicrobial resistance has been recognized as a global health issue for many decades. Food animals are considered key reservoirs of antibiotic-resistant bacteria because certain antibiotic-resistance genes identified in the bacteria of animal food products have also been identified in humans [54]. However, effectively treating *E. coli* O157:H7/H<sup>-</sup> infections is challenging, due to differing opinions presented by various investigators [55]. Many studies have reported an increasing incidence of multi-drug resistant *E. coli* O157:H7/H<sup>-</sup> strains isolated from the feces of slaughtered ruminants [56–58]. Our results show that all isolated strains were susceptible to antibiotics; cefotaxime, chloramphenicol, ciprofloxacin, norfloxacin, and polymyxin B. These findings are in agreement with previous reports of the antimicrobial-resistance patterns of *E. coli* O157:H7 isolates from animal and human sources [59, 60]. Further, previous reports on antibiotic-resistant bacteria in the feces of slaughtered ruminants demonstrated that the isolates showed high or intermediate resistance against multiple antimicrobials [56–58]. In fact, one study found that

gentamycin resistance was the most common (56.0%), followed by ampicillin (48.0%), erythromycin (40.0%), amoxicillin (16.0%), tetracycline (12.0%), chloramphenicol (8.0%), nalidixic acid (8.0%), and streptomycin (4.0%) [57]. The high levels of resistance to the antibiotics evaluated in that study perhaps reflect their wide use in the veterinary sector for treating various infections. Antimicrobial resistance emerges due to widespread use of antibiotics in animals and humans, which subsequently transfers the resistance genes and bacteria between animals and products, humans, and the environment [61]. It is worth considering that antibiotic resistance in food-borne pathogens have increased virulence and can increase the burden on human health by increasing the risk of contracting an infection in immunocompromised persons and reducing the treatment options for illnesses. In this study, most isolates displayed an intermediate resistance profile, indicating the potential for resistance to occur in the future. Attention must also be given to the elevated intermediate susceptibility profiles of such antibiotics. Some of the drugs we studied such as amoxicillin and nalidixic acid are commonly prescribed for *E. coli* O157 treatment in adults but showed resistance in our study. There have been limited studies on antibiotic resistance in *E. coli* O157 strains isolated from ruminant feces in the UAE. However, testing of only a small study population for antibiotic susceptibility, variability in specific resistance genes within isolates, differences in preferred antibiotics, or origins of strains, may impact the observed resistance or even lead to false conclusion of higher resistance rates.

## Conclusions

In this study, the presence of the *E. coli* O157 serotype, its major virulence genes, and associated antibiotic resistance were investigated in strains isolated from the feces of slaughtered ruminants in the UAE, including, camels, cattle, sheep and goats. The presence of high concentrations of pathogens in some animals, along with seasonal variations, highlights the need for improved risk-mitigation strategies to screen for high-shedding animals prior to slaughter. In this study, we established the presence of high-shedding animals at slaughterhouses in the UAE, which pose increased contamination risks for both the food chain and the environment. Our results also demonstrate that bacterial shedding at a high concentration by a few animals may play a more significant role in pathogen dissemination than the prevalence rate, as it can directly cause carcass contamination. Although using antibiotics to treat *E. coli* O157 infections is controversial, monitoring antibiotic resistance in the strains continues to be useful for epidemiological purposes. Moreover, further studies are needed to accurately determine the level of *E. coli* O157 contamination in the

carcasses and hides of animals in slaughterhouses. This study has some limitations specifically related to the small sample size. The number of positive samples was insufficient to detect monthly variations in the prevalence within statistically acceptable confidence limits; however, it did enable detection of the minimum prevalence rate in the species studied.

This work clearly supports the need for a platform aimed at designing regular screening programs in food safety and processing units, in order to ensure early and rapid detection of food-borne pathogens. The need for suitable control measures for such animals cannot be underestimated, and further studies are needed to devise mitigation strategies that will reduce the risk of large-scale contamination of the food chain or environment. Studies are also needed to characterize the contamination of *E. coli* O157 isolates and other non-*E. coli* O157 pathogens in meat food products.

## Methods

### Animal fecal samples

This study was conducted on healthy camels, cattle, sheep and goats slaughtered in the public abattoir in Al Ain, UAE, during the study period. Most animals were transported to the slaughterhouse and, hence, we studied the animals starting from the lairage. Samples were collected for a period of ten months, from September 2017 to July 2018. As per the project design, the samples were collected and tested on a biweekly basis, and the breed, species, gender, and fecal consistency of the animal were noted. All animals were local, except for the cattle. The mature Holstein dairy cattle were culled from a local dairy farm. A total of 568 fecal samples were collected from healthy animals in the slaughterhouse (sheep: 141, goats: 150, camels: 140, cattle: 137). Approximately 10 cm of the recto-anal junction was excised immediately after the slaughter and the fecal samples were collected, stored on ice, transported to our laboratory, and examined. Microbial testing was performed within 3 h of sample collection.

### IMS technique

Approximately 10 g of each fecal sample was diluted in 90 mL of BPW (Oxoid) and mixed for 30 s. A 15 mL aliquot of each fecal mixture was stored for later use. Next, 1 mL of enriched fecal sample was mixed with 20  $\mu$ L of magnetic beads coated with an anti-O157 antibody and IMS was performed according to the manufacturer's instructions (Oxoid). The bead suspension (100  $\mu$ L) was then plated onto CT-SMAC agar (Oxoid). The plates were incubated at 37 °C for 24 h and presumptive *E. coli* O157 colonies were identified as *E. coli* O157 that did not ferment sorbitol. These colonies were picked and inoculated into nutrient agar slants at 37 °C for 24 h and

stored in a refrigerator for further biochemical analysis. These isolates were further verified using conventional biochemical tests as described by Harrigan [62]. Tests were performed to examine indole production and motility using Sulfide Indole Motility (SIM) medium (Merck); citrate utilization with Simmons citrate agar (Merck); and methyl red and vogues-Proskauer using MR-VP medium (Merck) [63].

#### Latex agglutination test

*E. coli* O157 strains were confirmed by growing non-sorbitol-fermenting colonies (white-gray color) on CT-SMAC medium and agglutinating then with *E. coli* O157 latex reagent (Oxoid), as per the manufacturer's instructions. Isolates showing visible agglutination following reaction with the test latex solution were again subcultured for virulence gene identification.

#### Enumeration of *E. coli* O157

Latex agglutination test-positive samples were directly enumerated. Briefly, 100  $\mu$ L of each enriched fecal sample was serially diluted three times in 900  $\mu$ L of BPW (i.e., diluted by  $10^{-1}$ -,  $10^{-2}$ -, and  $10^{-3}$ -fold). The serially diluted samples were separated by IMS and the number of *E. coli* O157 cells was counted [64]. Finally, 100  $\mu$ L of separated bacteria-bead complexes were plated onto two SMAC plates (50  $\mu$ L/plate) and incubated at 37 °C for 18–24 h. After incubation, the CFUs were counted and recorded. The CFUs ranged from too few to count (TFTC) to too numerous to count (TNTC).

#### DNA extraction

After enumerating the positive samples from each animal in each collection, they were subjected to DNA extraction, as previously described [65]. Pure culture samples were centrifuged at 10,000 rpm for 1 min, and each supernatant was discarded and re-suspended with

sterile water. The tubes were incubated at 100 °C in a dry bath for 10 min, after which they were transferred to ice for 5–10 min. Crude DNA in the supernatant was obtained by centrifugation. The DNA was later analyzed for the presence of virulence genes.

#### Confirmation of *E. coli* O157 by multiplex PCR

All biochemically confirmed *E. coli* isolates that were O157 agglutination-positive were tested for the presence of genes in the *E. coli* O157 antigen gene locus, *rfbEO157:H7*, which codes for GDP perosamine synthetase (*rfbO157*), *uidA*, and the H7 flagellar protein (*fliCH7*) by multiplex PCR analysis, as previously described by Al-Ajmi et al. [65]. Briefly, 1  $\mu$ L of crude DNA extract was amplified in 25  $\mu$ L mixtures containing 10  $\mu$ L PCR master mix (Thermo Scientific), 5  $\mu$ L nuclease free water, and 1  $\mu$ L of each forward and reverse primer (Table 4) for *rfbE*, *flicH7*, and *uidA*. A negative control without template DNA was also included in the analysis. The thermal cycling program was as follows: initial denaturation at 95 °C for 2 min; 30 cycles of denaturation (30 s, 94 °C), annealing (15 s, 58 °C), and extension (1 min 68 °C); and a final extension (10 min, 68 °C).

An additional multiplex PCR analysis was carried out using the *eaeA*, *hlyA*, *Stx1*, and *Stx2* primers to quantify virulence genes (Table 4) [66]. Briefly, 1  $\mu$ L of crude DNA extract was amplified in 20- $\mu$ L reaction mixtures containing 10  $\mu$ L of PCR master mix, 5  $\mu$ L nuclease-free water, and 1  $\mu$ L of each forward and reverse primer. The following thermal cycling program was used: 1 min at 94 °C; 30 cycles of 30 s at 94 °C, 1 min at 55 °C, 1 min at 72 °C and a 5 min extension at 72 °C, with a no-template negative control. The PCR products were separated on 2% agarose gels containing ethidium bromide, and DNA bands were visualized on an ultraviolet light box using a gel-image capture system and software (Bio-Rad).

**Table 4** Primers used to confirm the presence of *E. coli* O157:H7 and virulence genes

Target gene	Primer name	Primer sequence (5'-3')	Amplification size (bp)
<i>rfbE</i>	O157AF	AAGATTGCGCTGAAGCCTTTG	497
	O157AR	CATTGGCATCGTGTGGACAG	
<i>flicC</i>	FLICH7 F	GCGCTGTGAGTTCTATCGAGC	625
	FLICH7 R	CAACGGTGACTTTATCGCCATTCC	
<i>uidA</i>	PT-2	GCGAAAACGTGGAATTGGG	252
	PT-3	TGATGCTCCATAACTTCTG	
<i>hlyA</i>	hlyA F	GCATCATCAAGCGTACGTTCC	534
	hlyA R	AATGAGCCAAGCTGGTTAAGCT	
<i>stx2</i>	Stx2F	GGCACTGTCTGAAACTGCTCC	255
	Stx2R	TCGCCAGTTATCTGACATTCTG	
<i>stx1</i>	Stx1F	ATAAATCGCCATTCTGTTGACTAC	180
	Stx1R	AGAACGCCCACTGAGATCATC	

### Antimicrobial testing

The Kirby–Bauer disk-diffusion method was performed to test nine antimicrobials, which are commonly used for treating human infections. An inoculum from each *E. coli* O157 strain was grown in 5 mL Mueller–Hinton (MH) broth (Himedia) at 37 °C and assessed for turbidity to a McFarland 0.5 standard, according to the Clinical and Laboratory Standards Institute (CLSI) [67] standards. All test discs were obtained from Oxoid. A standard reference strain (*E. coli* ATCC 25922), which is sensitive to all tested antimicrobial agents, was used as the control strain. MH agar plates were swabbed with sterile cotton swabs, and antimicrobial paper disks containing amoxicillin (25 µg), ampicillin (10 µg), bacitracin (10 µg), cefotaxime (30 µg), ceftazidime (10 µg), cefuroxime sodium (5 µg), chloramphenicol (30 µg), ciprofloxacin (30 µg), cloxacillin (5 µg), doxycycline (30 µg), gentamycin (10 µg), kanamycin (30 µg), nalidixic acid (30 µg), nitrofurantoin (300 µg), norfloxacin (10 µg), polymyxin B (300 U), streptomycin (10 µg), tetracycline (30 µg), and vancomycin (30 µg) were impregnated onto the surface of inoculated MH agar plates. The plates were incubated at 37 °C for 16–18 h, and the diameters of each zone of microbial-growth inhibition around the antimicrobial disk were measured. The minimum inhibitory concentration was determined for each antimicrobial agent, and each agent was characterized as being sensitive, having intermediate resistance, or being resistant according to criteria established by the CLSI [67].

### Statistical analysis

Statistical analysis of the breed of animal, sex, season of sampling and antimicrobial susceptibility were analyzed using the SAS software (SAS, V8.02) and comparisons between means were made using least significant difference at a *p* value of 0.05.

### Abbreviations

AML: Amoxicillin; AMP: Ampicillin; B: Bacitracin; BPW: Buffered Peptone Water; CAZ: Ceftazidime; CFUs: colony-forming units; CHL: Chloramphenicol; CIP: Ciprofloxacin; CLOXA: Cloxacillin; CLSI: Clinical and Laboratory Standards Institute; CT-SMAC: cefixime-tellurite sorbitol MacConkey; CTX: Cefotaxime; CXM: Cefuroxime sodium; DO: Doxycycline; F: Nitrofurantoin; FOX: Cefoxitin; GEN: Gentamycin; HC: enterohemorrhagic colitis; HUS: hemolytic uremic syndrome; IMS: Immuno Magnetic Separation; K: Kanamycin; MIC: Minimum Inhibitory Concentration; NAL: Nalidixic acid; NOR: Norfloxacin; PCR: Polymerase Chain Reaction; PMB: Polymyxin B; STEC: Shiga toxin-producing *Escherichia coli*; STR: Streptomycin; TET: Tetracycline; VAN: Vancomycin

### Acknowledgments

We would like to acknowledge the University editorial service for editing the document.

### Authors' contributions

Conceived and designed the study: DA and SR. Conducted the experiments and analyzed the results: DA, SR, and SB. Drafted the manuscript: DA and SR.

All authors reviewed the manuscript. All authors read and approved the final manuscript.

### Funding

This material is based upon work that was supported by funding provided by UAE University, United Arab Emirates, through a research grant (start-up grant number 31F098). The funding body had no role in the design of the study, collection, analysis, or interpretation of data, or writing of the manuscript.

### Availability of data and materials

The datasets used and/or analyzed during this study are available from the corresponding author upon reasonable request.

### Ethics approval and consent to participate

This study was conducted at a slaughterhouse with approval by UAE University and conduct of the experiment was permitted and ethical clearance was waived due to no major involvements of human and animal subjects necessitating ethical issues. The study purpose was explained to respective individuals and verbal agreement was obtained before proceeding to the study.

### Consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

Received: 22 January 2020 Accepted: 12 July 2020

Published online: 16 July 2020

### References

- Fung F, Wang HS, Menon S. Food safety in the 21st century. *Biom J*. 2018; 41:88–95.
- Kamleh R, Jurdi M, Annous BA. Management of microbial food safety in Arab countries. *J Food Prot*. 2012;75:2082–90.
- Food and Agriculture Organization of the United Nations and World Health Organization. FAO/WHO regional meeting on food safety for the Near East, Amman, Jordan. The impact of current food safety systems in the Near East/eastern Mediterranean region on human health. 2005. [ftp://ftp.fao.org/es/esn/food/meetings/NE\\_wp2\\_en.pdf](ftp://ftp.fao.org/es/esn/food/meetings/NE_wp2_en.pdf). Accessed 30 Oct 2019.
- Saleh I, Zouhairi O, Alwan N, Hawi A, Barbour E, Harakeh S. Antimicrobial resistance and pathogenicity of *Escherichia coli* isolated from common dairy products in the Lebanon. *Ann Trop Med Parasitol*. 2009;103:39–52.
- Banatvala N, Griffin PM, Greene KD, Barrett TJ, Bibb WF, Green JH, et al. The United States national prospective hemolytic uremic syndrome study: microbiologic, serologic, clinical, and epidemiologic findings. *J Infect Dis*. 2001;183:1063–70.
- Kim JC, Chui L, Wang Y, Shen J, Jeon B. 2016. Expansion of Shiga toxin-producing *Escherichia coli* by use of bovine antibiotic growth promoters. *Emerg Infect Dis*. 2016;22:802–9.
- Vallance BA, Finlay BB. Exploitation of host cells by enteropathogenic *Escherichia coli*. *Proc Natl Acad Sci U S A*. 2000;97:8799–806.
- Rangel JM, Sparling PH, Crowe C, Griffin PM, Swerdlow DL. Epidemiology of *Escherichia coli* O157:H7 outbreaks, United States, 1982–2002. *Emerg Infect Dis*. 2005;11:603–9.
- Buchholz U, Bernard H, Werber D, Böhmer MM, Renschmidt C, Wilking H, et al. 2011. German outbreak of *Escherichia coli* O104:H4 associated with sprouts. *N Engl J Med*. 2011;365:1763–70.
- Persad AK, LeJeune JT. Animal reservoirs of Shiga toxin-producing *Escherichia coli*. *Microbiol Spectr*. 2014;2:EHCC-0027-2014.
- Widgren S, Söderlund R, Eriksson E, Fasth C, Aspan A, Emanuelson U, et al. Longitudinal observational study over 38 months of verotoxigenic *Escherichia coli* O157:H7 status in 126 cattle herds. *Prev Vet Med*. 2015;121: 343–52.
- Segura A, Auffret P, Bibbal D, Bertoni M, Durand A, Jubelin G, et al. Factors involved in the persistence of a Shiga toxin-producing *Escherichia coli* O157:H7 strain in bovine feces and gastro-intestinal content. *Front Microbiol*. 2018;9:375.
- Helke KL, McCrackin MA, Galloway AM, Poole AZ, Salgado CD, Marriott BP. Effects of antimicrobial use in agricultural animals on drug-resistant



- foodborne salmonellosis in humans: a systematic literature review. *Crit Rev Food Sci Nutr.* 2017;57:472–88.
14. Pornsukarom S, van Vliet AH, Thakur S. Whole genome sequencing analysis of multiple *Salmonella* serovars provides insights into phylogenetic relatedness, antimicrobial resistance, and virulence markers across humans, food animals and agriculture environmental sources. *BMC Genomics.* 2018; 19:801.
  15. Alonge DO. Textbook of meat hygiene in the tropics. Ibadan: Farm Coe Press; 1991.
  16. Jo MY, Kim JH, Lim JH, Kang MY, Koh HB, Park YH, et al. Prevalence and characteristics of *Escherichia coli* O157 from major food animals in Korea. *Int J Food Microbiol.* 2004;95:41–9.
  17. Griffin PM, Tauxe RV. The epidemiology of infections caused by *Escherichia coli* O157: H7, other enterohemorrhagic *E. coli*, and the associated hemolytic uremic syndrome. *Epidemiol Rev.* 1991;13:60–98.
  18. McCluskey BJ, Rice DH, Hancock DD, Hovde CJ, Besser TE, Gray S, et al. Prevalence of *Escherichia coli* O157 and other Shiga-toxin-producing *E. coli* in lambs at slaughter. *J Vet Diagn Investig.* 1999;11:563–5.
  19. De Boer E, Heuvelink AE. Methods for the detection and isolation of Shiga toxin-producing *Escherichia coli*. *Symp Ser Soc Appl Microbiol.* 2000;29: 1335–43S.
  20. Dutta S, Deb A, Chattopadhyay UK, Tsukamoto T. Isolation of Shiga toxin producing *Escherichia coli* including O157:H7 strains from dairy cattle and beef samples marketed in Calcutta, India. *J Med Microbiol.* 2000;49:765–7.
  21. Carney E, O'Brien SB, Sheridan JJ, McDowell DA, Blair IS, Duffy G. Prevalence and level of *Escherichia coli* O157 on beef trimmings, carcasses and boned head meat at a beef slaughter plant. *Food Microbiol.* 2006;23:52–9.
  22. Taye M, Berhanu T, Berhanu Y, Tamiro F, Terefe D. Study on carcass contaminating *Escherichia coli* in apparently healthy slaughtered cattle in Haramaya University slaughter house with special emphasis on *Escherichia coli* O157:H7, Ethiopia. *J Vet Sci Technol.* 2013;4:132.
  23. Bosilevac JM, Gasseem MA, Al Sheddy IA, Almaiman SA, Al-Mohizea IS, Alowaimar A, et al. Prevalence of *Escherichia coli* O157: H7 and *Salmonella* in camels, cattle, goats, and sheep harvested for meat in Riyadh. *J Food Prot.* 2015;78:89–96.
  24. Rahimi E, Hontaz H, Hemmafzadeh F. The prevalence of *Escherichia coli* O157:H7, *Listeria monocytogenes* and *Campylobacter* spp. on bovine carcasses in Isfahan, Iran. *Int J Vet Res.* 2008;4:365–70.
  25. Hashemi M, Khanzadi S, Jamshadi A. Identification of *Escherichia coli* O157: H7 isolated from cattle carcasses in Mashhad abattoir by multiplex PCR. *World Appl Sci J.* 2010;6:703–8.
  26. Mohammed HO, Stipetic K, Salem A, McDonough P, Chang YF, Sultan A. 2015. Risk of *Escherichia coli* O157: H7, non-O157 Shiga toxin-producing *Escherichia coli*, and *Campylobacter* spp. in food animals and their products in Qatar. *J Food Prot.* 2015;78:1812–8.
  27. Al-Gburi NM. Prevalence of *Escherichia coli* O157: H7 in camels fecal samples. *J Genet Environ Resour Conserv.* 2016;4:46–50.
  28. Moore JE, McCalmont M, Xu JR, Nation G, Tinson AH, Cartothers L. Prevalence of fecal pathogens in calves of racing camels (*Camelus dromedarius*) in the United Arab Emirates. *Trop Anim Health Prod.* 2002;4: 283–7.
  29. El-Sayed A, Ahemd S, Awad W. Do camels (*Camelus dromedarius*) play an epidemiological role in the spread of Shiga toxin producing *Escherichia coli* (STEC) infection? *Trop Anim Health Prod.* 2008;40:469–73.
  30. Chapman PA, Siddons CA. A comparison of immunomagnetic separation and direct culture for the isolation of verocytotoxin-producing *Escherichia coli* O157 from cases of bloody diarrhoea, non-bloody diarrhoea and asymptomatic contacts. *J Med Microbiol.* 1996;44:267–71.
  31. Alhelfi NA, Adam H, Jones DL, Williams AP. Absence of *E. coli* O157: H7 in sheep and cattle faeces in North Wales. *Vet Rec.* 2013;173:143.
  32. Chapman PA, Cerdán AT, Ellin M, Ashton R, Harkin MA. *Escherichia coli* O157 in cattle and sheep at slaughter, on beef and lamb carcasses and in raw beef and lamb products in South Yorkshire. *UK Int J Food Microbiol.* 2001; 64:139–50.
  33. Rhoades JR, Duffy G, Koutsoumanis K. Prevalence and concentration of verocytotoxigenic *Escherichia coli*, *Salmonella enterica* and *Listeria monocytogenes* in the beef production chain: a review. *Food Microbiol.* 2009;26:357–76.
  34. Battisti A, Lovari S, Franco A, Diegidio A, Tozzoli R, Caprioli A, et al. Prevalence of *Escherichia coli* O157 in lambs at slaughter in Rome. *Central Italy Epidemiol Infect.* 2006;134:415–9.
  35. Vlisidou I, Marchés O, Dziva F, Mundy R, Frankel G, Stevens MP. Identification and characterization of *EspK*, a type III secreted effector protein of enterohaemorrhagic *Escherichia coli* O157:H7. *FEMS Microbiol Lett.* 2006;263:32–40.
  36. Torres AG, Milflores-Flores L, Garcia-Gallegos JG, Patel SD, Best A, La Ragione RM, et al. Environmental regulation and colonization attributes of the long polar fimbriae (LPF) of *Escherichia coli* O157:H7. *Int J Med Microbiol.* 2007; 297:177–85.
  37. Gansheroff LJ, O'Brien AD. *Escherichia coli* O157: H7 in beef cattle presented for slaughter in the US: higher prevalence rates than previously estimated. *Proc Natl Acad Sci U S A.* 2000;97:2959–61.
  38. Osaili TM, Alaboudi AR, Rahalah M. Prevalence and antimicrobial susceptibility of *Escherichia coli* O157: H7 on beef cattle slaughtered in Amman abattoir. *Meat Sci.* 2013;93:463–8.
  39. Lin YL, Chou C, Pan T. Screening procedure from cattle faeces and the prevalence of *Escherichia coli* O157:H7 in Taiwan dairy cattle. *J Microbiol Immunol Infect.* 2001;34:17–24.
  40. Jacob ME, Almes KM, Shi X, Sargeant JM, Nagaraja TG. *Escherichia coli* O157: H7 genetic diversity in bovine fecal samples. *J Food Prot.* 2011;74:1186–8.
  41. Barkocy-Gallagher GA, Arthur TM, Rivera-Betancourt M, Nou X, Shackelford SD, Wheeler TL, et al. Seasonal prevalence of Shiga toxin-producing *Escherichia coli*, including O157:H7 and non-O157 serotypes, and *Salmonella* in commercial beef processing plants. *J Food Prot.* 2003;66:1978–86.
  42. Seker E, Kus FS. The prevalence, virulence factors and antibiotic resistance of *Escherichia coli* O157 in feces of adult ruminants slaughtered in three provinces of Turkey. *Veterinarski Arhiv.* 2019;89:107–21.
  43. Brichta-Harhay DM, Guerini MN, Arthur TM, Bosilevac JM, Kalchayanand N, Shackelford SD, et al. *Salmonella* and *Escherichia coli* O157:H7 contamination on hides and carcasses of cull cattle presented for slaughter in the United States: an evaluation of prevalence and bacterial loads by immunomagnetic separation and direct plating methods. *Appl Environ Microbiol.* 2008;74:6289–97.
  44. Gautam R, Bani-Yaghoob M, Neill WH, Döpfer D, Kaspar C, Ivanek R. Modeling the effect of seasonal variation in ambient temperature on the transmission dynamics of a pathogen with a free-living stage: example of *Escherichia coli* O157: H7 in a dairy herd. *Prev Vet Med.* 2011;102:10–21.
  45. Omisakin F, MacRae M, Ogden ID, Strachan NJ. Concentration and prevalence of *Escherichia coli* O157 in cattle feces at slaughter. *Appl Environ Microbiol.* 2003;69:2444–7.
  46. Farrokh C, Jordan K, Auvray F, Glass K, Oppegaard H, Raynaud S, et al. Review of Shiga-toxin-producing *Escherichia coli* (STEC) and their significance in dairy production. *Int J Food Microbiol.* 2013;162:190–212.
  47. Cornick NA, Booher SL, Moon HW. 2002. Intimin facilitates colonization by *Escherichia coli* O157: H7 in adult ruminants. *Infect Immun.* 2002;70:2704–7.
  48. Schwidder M, Heinisch L, Schmidt H. Genetics, toxicity, and distribution of Enterohemorrhagic *Escherichia coli* Hemolysin. *Toxins.* 2019;11:502.
  49. Johnsen G, Wasteson Y, Heir E, Berget OI, Herikstad H. *Escherichia coli* O157: H7 in faeces from cattle, sheep and pigs in the southwest part of Norway during 1998 and 1999. *Int J Food Microbiol.* 2001;65:193–200.
  50. Aslantaş Ö, Erdoğan S, Cantekin Z, Gülaçtı İ, Evrendilek GA. Isolation and characterization of verocytotoxin-producing *Escherichia coli* O157 from Turkish cattle. *Int J Food Microbiol.* 2006;106:338–42.
  51. Friedrich AW, Bielaszewska M, Zhang WL, Pulz M, Kuczys T, Ammon A, et al. *Escherichia coli* harboring Shiga toxin 2 gene variants: frequency and association with clinical symptoms. *J Infect Dis.* 2002;185:74–84.
  52. Beutin L, Krause G, Zimmermann S, Kaulfuss S, Gleier K. Characterization of Shiga toxin-producing *Escherichia coli* strains isolated from human patients in Germany over a 3-year period. *J Clin Microbiol.* 2004;42:1099–108.
  53. Ko H, Maymani H, Rojas-Hernandez C. Hemolytic uremic syndrome associated with *Escherichia coli* O157: H7 infection in older adults: a case report and review of the literature. *J Med Case Rep.* 2016;10:175.
  54. Founou LL, Founou RC, Essack SY. Antibiotic resistance in the food chain: a developing country-perspective. *Front Microbiol.* 2016;7:1881.
  55. Wong A. Epistasis and the evolution of antimicrobial resistance. *Front Microbiol.* 2017;8:246.
  56. Manna SK, Brahmane MP, Manna C, Batabyal K, Das R. Occurrence, virulence characteristics and antimicrobial resistance of *Escherichia coli* O157 in slaughtered cattle and diarrhoeic calves in West Bengal, India. *Lett Appl Microbiol.* 2006;43:405–9.
  57. Rahimi E, Nayeypour F. Antimicrobial resistance of *Escherichia coli* O157: H7/NM isolated from faeces of ruminant animals in Iran. *J Cell Anim Biol.* 2012;6:104–8.

58. Dulo F, Feleke A, Szonyi B, Fries R, Baumann MP, Grace D. Isolation of multidrug-resistant *Escherichia coli* O157 from goats in the Somalia region of Ethiopia: a cross-sectional, abattoir-based study. *PLoS One*. 2015;10: e0142905.
59. Faris G, Mekonen E. Prevalence and antibiotic susceptibility of *E. coli* strains in UTI: Prevalence and antibiotic susceptibility of *E. coli* strains amongst patients with complaints of urinary tract infection. LAP LAMBERT Academic Publishing; 2012.
60. Abreham S, Teklu A, Cox E, Tessema TS. *Escherichia coli* O157: H7: distribution, molecular characterization, antimicrobial resistance patterns and source of contamination of sheep and goat carcasses at an export abattoir, Mojo. *Ethiopia BMC Microbiol*. 2019;19:215.
61. Scott L, McGee P, Minihan D, Sheridan JJ, Earley B, Leonard N. The characterization of *E. coli* O157:H7 isolates from cattle faces and feedlot environment using PFGE. *Vet Microbiol*. 2006;114:331–6.
62. Harrigan WF, McCance ME. *Laboratory methods in food and dairy microbiology*. Academic Press Inc. (London) Ltd.; 1976.
63. NPH (National Public Health Service for Wales) Detection of *Escherichia coli* O157 by Automated Immunomagnetic Separation. Standard Method. Issued by Standards Unit, Evaluations and standards laboratory with the regional food, Water and Environmental Coordinators Forum. Wales: SOPs from the Health Protection Agency; 2006. p. 1–15.
64. Gilbert RA, Tomkins N, Padmanabha J, Gough JM, Krause DO, McSweeney CS. Effect of finishing diets on *Escherichia coli* populations and prevalence of enterohaemorrhagic *E. coli* virulence genes in cattle feces. *J Appl Microbiol*. 2005;99:885–94.
65. Al-Ajmi D, Padmanabha J, Denman SE, Gilbert RA, Al Jassim RAM, McSweeney CS. Evaluation of a PCR detection method for *Escherichia coli* O157:H7/H–bovine faecal samples. *Lett Appl Microbiol*. 2006;42:386–91.
66. Patton AW, Patton JC. Detection and characterization of Shiga toxicogenic *Escherichia coli* using multiplex PCR assays for *stx1*, *stx2* *eaeA*, enterohaemorrhagic *E. coli hlyA*, *rfbO111*, and *rfbO157*. *J Clin Microbiol*. 1998; 36:598–602.
67. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing; 24th informational supplement. Wayne: CLSI Document M100-S24; 2014.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

