

Research article

Open Access

# Bioinformatics analysis suggests base modifications of tRNAs and miRNAs in *Arabidopsis thaliana*

Kei Iida<sup>1,2</sup>, Hailing Jin<sup>3</sup> and Jian-Kang Zhu\*<sup>1</sup>

Address: <sup>1</sup>Department of Botany and Plant Sciences, University of California, Riverside, CA 92521, USA, <sup>2</sup>Bioinformatics and Systems Engineering Division, RIKEN Yokohama Institute, 1-7-22, Suehiro, Tsurumi, Yokohama 230-0045, Japan and <sup>3</sup>Department of Plant Pathology, University of California, Riverside, California 92521, USA

Email: Kei Iida - kiida@base.riken.jp; Hailing Jin - hailing.jin@ucr.edu; Jian-Kang Zhu\* - jian-kang.zhu@ucr.edu

\* Corresponding author

Published: 9 April 2009

Received: 27 October 2008

BMC Genomics 2009, 10:155 doi:10.1186/1471-2164-10-155

Accepted: 9 April 2009

This article is available from: <http://www.biomedcentral.com/1471-2164/10/155>

© 2009 Iida et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

**Background:** Modifications of RNA bases have been found in some mRNAs and non-coding RNAs including rRNAs, tRNAs, and snRNAs, where modified bases are important for RNA function. Little is known about RNA base modifications in *Arabidopsis thaliana*.

**Results:** In the current work, we carried out a bioinformatics analysis of RNA base modifications in tRNAs and miRNAs using large numbers of cDNA sequences of small RNAs (sRNAs) generated with the 454 technology and the massively parallel signature sequencing (MPSS) method. We looked for sRNAs that map to the genome sequence with one-base mismatch (OMM), which indicate candidate modified nucleotides. We obtained 1,187 sites with possible RNA base modifications supported by both 454 and MPSS sequences. Seven hundred and three of these sites were within tRNA loci. Nucleotide substitutions were frequently located in the T arm (substitutions from A to U or G), upstream of the D arm (from G to C, U, or A), and downstream of the D arm (from G to U). The positions of major substitution sites corresponded with the following known RNA base modifications in tRNAs: N1-methyladenosine (m<sup>1</sup>A), N2-methylguanosine (m<sup>2</sup>G), and N2-N2-methylguanosine (m<sup>2</sup><sub>2</sub>G).

**Conclusion:** These results indicate that our bioinformatics method successfully detected modified nucleotides in tRNAs. Using this method, we also found 147 substitution sites in miRNA loci. As with tRNAs, substitutions from A to U or G and from G to C, U, or A were common, suggesting that base modifications might be similar in tRNAs and miRNAs. We suggest that miRNAs contain modified bases and such modifications might be important for miRNA maturation and/or function.

## Background


In rRNAs, tRNAs, snRNAs, and some mRNAs, the bases of nucleotides are often modified [1-5]. Especially in tRNAs, many types of base modifications have been characterized [1,2]. These modifications often involve methylation or acetylation, and may contribute to the stability of tRNA molecules when they form tertiary structures [4,6].

Another well-characterized tRNA modification is RNA editing from adenine (A) to inosine (I) [7,8]. This A to I editing is explained by deamination of A by adenosine deaminase [7]. In the yeast alanine tRNA, adenosine bases on the anticodon are deaminated to I. The edited inosine can form base pairs with uridine (U), cytosine (C), or adenosine in codons of mRNAs, and thereby expands the

**Table 1: Data set and mapping results**

	All		Perfect match		OMM		OMM excluding terminal miss match	
	Unique	Reads	Unique	Reads	Unique	Reads	Unique	Reads
454 data set 1 <sup>a</sup>	96,080	236,245	19,085 (19.9%)	69,071 (29.2%)	14,269 (14.9%)	32,808 (13.9%)	4,889 (5.1%)	11,162 (4.7%)
454 data set 2 <sup>b</sup>	36,562	58,163	18,027 (49.3%)	18,027 (31.0%)	4,709 (12.9%)	6,081 (10.5%)	931 (2.5%)	1,091 (1.9%)
MPSS data set <sup>c</sup>	116,853	3,126,002	84,207 (72.1%)	2,501,573 (80.0%)	13,242 (11.3%)	240,355 (7.7%)	6,505 (5.6%)	112,299 (3.6%)

(a) Jin et al. 2008  
 (b) Lu et al. 2006  
 (c) Lu et al. 2005



**Figure 1**  
**Distribution of positions of observed substitutions in sRNA sequences.** Sequences were obtained with the MPSS method (A), with 454 technology by Lu et al. [22] (B), and our own sequence set (C). For each graph, the left side shows the number of substitutions associated with initial nucleotides (where "1" is the first nucleotide) and right side shows the number of substitutions associated with terminal nucleotides (where "-1" is the last nucleotide). For the entire data set, substitutions were especially abundant in the initial/terminal three nucleotides.

**Table 2: Modified nucleotides found in tRNA**

Region name	Genomic nucleotide	(a)	(b)	TPQ for non-modified bases	# of tRNA	Uniq # of sRNA	Total TPQ of sRNA	Edited nucleotides	TPQ of edited nucleotides	Type(s) of tRNA
T_loop	A	(5)	m <sup>1</sup> A	4454.04	383	4198	12127.1	U,G,C	9005.67,2958.02,163.49	Ala,Arg,Asp,Cys,Gln,Glu,Gly,His,Ile,Leu,Lys,Phe,Pro,Ser,Thr,Trp,Val
D->AntiC	G	(3)	m <sup>2</sup> G	372.31	34	257	894.15	U,A,C	660.14,155.40,78.64	Asn,Met,Tyr
Acc->D	G	(1)	m <sup>2</sup> G	23314.9	138	682	767.54	C,U,A	378.05,212.08,176.87	Arg,Lys,Met,Ser,Trp,Tyr,Val
D_stem_5'	C	(2)	ac <sup>4</sup> C	17339.82	17	51	212.86	A,U,G	197.48,8.42,6.96	Arg,Gly
AntiC_loop	A	(4)	I	0	6	86	98.95	G	98.95	Thr
D_stem_5'	T	N/A	N/A	11624.69	60	180	43.2	C,A	23.40,19.80	Tyr
Acc_stem_5'	G	N/A	N/A	16999.53	17	45	38.64	A,U	32.15,6.38	Arg,Gly
D_loop	G	N/A	N/A	2420.16	6	12	24.9	A	24.9	Ser
D_loop	A	N/A	N/A	1348	1	2	22	U,C	13.90,8.10	Gln
Acc_stem_5'	T	N/A	N/A	2406.42	6	12	11.4	C,G	9.42,1.98	Ser
AntiC_loop	G	N/A	m <sup>1</sup> G	47.12	26	52	9.1	U,C	8.32,0.78	Pro
T_stem_5'	G	N/A	N/A	313.29	3	6	8.61	A,C	7.89,0.69	Asp

(a) Corresponding position in Figure 2.

(b) Corresponding modified base.

decoding capacities [8]. Although RNA modification in tRNAs, especially mitochondrial tRNAs, has been studied in *Arabidopsis* [9], the understanding of RNA modification in *Arabidopsis* tRNAs is still limited.

MicroRNAs (miRNAs) are a recently described class of functional RNAs and were initially found in *C. elegans* [10]. Several editing events in miRNAs have been reported. One example is A to I editing in primary miRNA precursor transcripts (pri-miRNAs). A to I editing has been documented for human pri-miR-22 and pri-miR-142 [11-13]. In the case of human pri-miR-142, A to I editing represses the maturation process carried out by the cleaving enzyme, Droscha [12,13]. Another example of RNA editing in miRNAs is found in primary transcripts of the human miR-376 cluster [14]. In this case, the edited pri-miRNAs are processed into mature miRNAs, and the mature miRNA targets different transcripts than the non-edited miRNA. These examples demonstrated the importance of RNA editing on miRNAs. Like humans, *Arabidopsis thaliana* also has hundreds of miRNA genes [15]. In *Arabidopsis*, the methyltransferase enzyme HEN1 introduces 2'-O methyl groups to the ribose of 3' terminal nucleotide of miRNAs [16,17]. This modification is important for the stability and accumulation of miRNAs. However, there has been no report of RNA modification in the bases of miRNAs in *Arabidopsis* or other plants.

The current study is focused on RNA base modifications in *Arabidopsis*. Because some modified bases are read differently from unmodified bases by reverse transcriptases during cDNA synthesis [18], we expect that the modified nucleotides will be read as different nucleotides from

genomic ones if the bases of RNAs are modified. Our analysis utilized several sets of high throughput cDNA sequences of small RNAs (sRNAs). We used our own sequences generated by the "454" technology [19,20]. During the analysis of our sequences, we observed that many of the sequences could not be mapped to either the nuclear or organellar genomes [19]. We suspected that such non-mapped sequences may contain information of RNA modifications. We found that the sRNAs that were not mapped to the genome, mapped perfectly with the genome sequences except for one base mismatches (OMM). In the current work, we also used public sequences obtained with the 454 technology and the "massively parallel signature sequencing" (MPSS) method [21,22]. The MPSS dataset is especially important here since the sequences were generated by a completely different technology from the 454. We listed substituted sites only if the sites were supported by sequences from both 454 technology and MPSS. This strategy allowed us to avoid detecting simple sequencing errors resulting from one or the other of the sequencing technologies. In this study, we first analyzed RNA base modifications in tRNA molecules. Next, we investigated potential RNA base modifications in miRNAs. Ours is the first report of large scale analysis of RNA base modifications of tRNAs in *Arabidopsis*. More importantly, our bioinformatics analysis suggests similar base modifications (e.g. methylation and acetylation) in miRNAs.

## Results and discussion

### Mapping sRNAs to the genome

We mapped sRNA sequences to the *Arabidopsis* nuclear genome with the alignment program SOAP [23]. Out of



**Table 3: Substitutions found in miRNAs**

Name of microRNA	Summary of substitutions <sup>a</sup>
MIR159a, MIR159b	5:G(27900.95)>CA(21), 12:G(28142.15)>CA(45.4), 13:G(35092.7)>UA(144.86)
MIR161	6:G(20643.65)>CA(50.2), 9:A(20057.15)>GCU(109.2), 13:C(18208.1)>GA(8.3)
MIR162a, MIR162b	14:G(3040.5)>UA(58.8)
MIR165a, MIR166a, MIR166b, MIR166c, MIR166d, MIR166e, MIR166f, MIR166g	4:G(54554.7)>CUA(134.7), 6:C(54554.7)>UA(209.21), 7:C(44547.6)>U(149.93), 8:A(35038.15)>GU(83.7), 9:G(47675.1)>CUA(328.62), 10:G(54539.9)>A(106.29), 11:C(54539.9)>GUA(278.15), 12:U(54539.9)>GCA(379.38), 13:U(54539.9)>CA(256.11)
MIR167a, MIR167b, MIR167d	4:A(219813.04)>GCU(76.99), 5:G(219813.04)>CUA(325.81), 7:U(158835.06)>GCA(466.4), 8:G(158835.06)>CUA(326.2), 9:C(158835.06)>GUA(602.6), 10:C(158835.06)>GU(431.4), 11:A(219826.94)>GCU(3686.02), 12:G(219826.94)>CU(1018.4), 13:C(158835.06)>GU(336.74)
MIR168a, MIR168b	9:U(15948.36)>CA(36.6), 10:G(15933.26)>A(24.3), 12:A(15933.26)>GC(22.2), 13:G(15933.26)>UA(43.1), 14:G(15933.26)>UA(41.3)
MIR169a, MIR169b, MIR169c, MIR169d, MIR169e, MIR169h, MIR169i, MIR169j, MIR169k, MIR169l, MIR169m, MIR169n	6:C(24472.01)>GU(40.32), 8:A(87930.11)>GCU(77.31), 9:G(58812.37)>CUA(501.61), 10:G(37156.57)>CUA(161.5), 11:A(10397.81)>GC(26.88), 13:G(87930.11)>CUA(1043.6), 14:A(34341.26)>G(44.23), 15:C(38019.92)>GUA(109.1)
MIR170	5:U(50752.3)>GC(244.7), 8:G(50752.3)>CA(23.8)
MIR171a	7:A(2397.55)>GU(54.5)
MIR172a, MIR172b, MIR172c	8:U(211955.28)>GC(157.21), 11:U(83771.26)>C(90.98)
MIR319a, MIR319b	9:A(1866.44)>CU(37)
MIR390a, MIR390b	PRE74:A(6807.15)>G(21.5), PRE76:C(6807.15)>U(78.6), 6:C(41119.9)>UA(27.6), 7:A(20563.45)>G(48.5)
MIR858	5:G(12416.25)>CU(13.15), 12:G(12416.25)>CA(8.55)


(a) Format here is: position in mature miRNA:un-substituted nucleotide at this position (TPQ of miRNAs without substitution) > substituted nucleotide(s) (TPQ of miRNAs with substitution). In MIR390, two substitutions were found in pre-miRNA regions; the positions and frequencies of these substitutions are indicated following "PRE": position from 5' end of pre-miRNA:unsubstituted nucleotide at this position (TPQ of sRNAs without substitution)>substituted nucleotide (TPQ of the sRNAs with substitution).

### Base modifications in tRNAs

In the current analysis, the largest fraction of observed substitution sites corresponded with tRNAs. Because we used sRNAs of up to only 40 nt in length, we inferred that the tRNA sequences came from fragmented tRNAs. Such fragmented tRNAs would be consistent with recent reports of detection of cellular tRNA fragments [25,26]. Given that most substitution sites corresponded with tRNAs, these were the first substitution sites that we analyzed. We predicted secondary structures of tRNAs with tRNAscan-SE [27] and then mapped the substitution sites on the second-

ary structures. The substitution sites were clustered into five major regions and several minor regions (Table 2). The most abundant type of substitution was found in the T loop (Table 2, Fig. 2). In these regions, A to U or to G substitutions were common. Out of 639 tRNAs encoded in the *Arabidopsis* genome, 383 have substitutions of this type. We found this type of substitution on tRNAs for all amino acids except asparagine, methionine, and tyrosine. This position corresponds with the position of a known RNA modification, N1-methyladenosine (m<sup>1</sup>A) [5,28,29]. We concluded that the inferred modification was a methylation of ade-





**Figure 4**

**Illustration of substitution patterns found in sRNAs corresponding to tRNAs.** Changes of A to T or G and G to T, C, or A are abundant. Similar substitution patterns were found in miRNAs.


ref. [34]]. Despite the small number of tRNA fragments containing the anticodon loop, we found some substitutions which correspond to the well-known modified base, 1-methylguanosine ( $m^1G$ ) at position 37 of the anticodon loop (Table 2, substitution of G to U/C).

The locations of predicted RNA modifications based on OMMs corresponded well with locations of known modifications reported in the literature. Out of 703 substitution sites found in tRNAs, 604 sites (86%) can be accounted for by the 6 known base modifications discussed above. Clearly these bias were not caused by chance, which was supported by binomial test with  $p$ -value less than  $1 \times 10^{-15}$ . Substitution patterns were also consistent with the known chemistry of reverse transcription in the literature. Research on reverse transcription of tRNAs showed that  $m^1A$  nucleotides formed base pairs with A, U, G, or C [18]. In agreement with Steinberg and Cedergren [31], we found that the substituted nucleotides at known  $m^2G$  sites were mainly U and A (Table 2). Therefore, our predictions of base modifications in tRNAs are consistent with previous studies. We suggest that RNA

modification types can be classified based on substitution patterns found in OMM sRNAs.

#### Base modifications in miRNAs

After describing base modifications in tRNAs, we next considered possible base modifications in miRNAs. We found 147 cases of substitutions in miRNAs, which could be classified into 13 families (Fig. 3, Table 3). The substitution pattern of A was similar to that found in tRNAs (Fig 4). Adenine nucleotides were read mainly as thymine (T) and guanine in OMM sRNAs. This similarity of substitution suggests that N1-methyladenosines could be a major modified RNA base in miRNAs, as in tRNAs (Table 2). In the cases of substitutions of G, all of the other three nucleotides (C, A, and T) were observed (Fig 4). This substitution pattern was similar to that of  $m^2G$  rather than of  $m^2G$  in tRNAs (Table 2). We suspect that  $m^2G$  might be the main modified G in miRNAs, although it is possible that miRNAs may have different types of G modifications. We also examined the substitutions on genomic C and T sites, although there are few such substitutions in tRNAs. These RNA modifications may differ from those in tRNAs




**Figure 5**  
**Examples of substitutions found in miRNAs and relationship with their targets.** We show miRNAs that have the most (miR167; A) and the second most abundant (miR169; B) substitutions. In both cases, complementarity between miRNAs and their targets was reduced by the substitutions.

or may be rare in tRNAs. We found that some fraction of the substitutions occurring in miRNAs were similar with those in tRNAs. Moreover, the substitution patterns were not random. Based on these results, we concluded that these substitutions are not simple sequencing errors, but reflect novel RNA base modifications.

A to I editing is found in human miRNAs [14]. Based on the results with tRNAs, A to I editing was expected to be observed as substitution to G (Table 2). Such substitutions were found in miR169 (site 14) and miR319 (site 7) (Table 3). However, A to G substitutions did not represent the major substitution pattern (Fig. 4). Therefore, although A to I editing seems to occur in Arabidopsis miRNAs, such modifications are evidently uncommon.

**Potential biological impacts of base modifications in miRNAs**

Our bioinformatics analysis strongly suggested RNA base modifications in miRNAs. What would be the biological effects of base modifications in miRNAs? We considered



**Figure 6**  
**The rates of substitutions for tRNAs, miRNAs, rRNAs, and all other data.** In tRNAs, substitution rates were more than 0.2 at 438 sites. In miRNAs, however, substitution rate was only 0.02 at the site with maximal substitution rate.

two possibilities. The first one is that the modification may enhance or prohibit the recognition of mRNA targets or cause recognition of novel targets, as is the case for RNA editing in human miR-376 [14]. The other possibility is that RNA base modifications may affect the maturation of miRNA, as with RNA editing in human pri-miR-142 [12,13].

With respect to the first hypothesis, we examined how putative RNA modification sites may affect the relationship between miRNAs and their targets, assuming that the pairing between a modified nucleotide and its target nucleotide can be mimicked by pairing of the substituted nucleotide and the target (Fig. 5). Although the RNA modifications occurred in sites that could alter target recognition, the modifications did not seem to enhance target recognition. We also tried to predict some novel target transcripts with modified miRNA sequences but we were unable to find novel targets with miRNAs containing substitutions. These results suggest that the base modifications may not enhance target recognition or create new targets. Consistent with this notion, putative modification sites are spread widely within the length of a miRNA molecule (Fig. 3).

The second hypothesis was difficult to test because our data came from mature miRNAs rather than pri-miRNA sequences. However, several results in this study may be



consistent with the second hypothesis. First, we found that nucleotides outside of mature miRNAs were also modified in miR390a (Fig. 3). Second, we surveyed EST sequences corresponding with pre-miRNAs. They also had several substitutions from genomic nucleotides that were located outside of regions of mature miRNAs (see Additional files 2 and 3). These results suggest that pre-miRNA already have modified bases. Third, it is known that several RNA modification enzymes affecting tRNAs recognize secondary and/or tertiary structures of tRNA molecules [35-37]. It is possible that tRNA modification enzymes might also modify pri-/pre miRNAs. In the case of miRNAs, it is unknown whether the base modifications may reinforce the fold-back structures of pre-miRNAs. However, we observed that many substitutions were in the positions where Watson-Crick or wobble base pairs were initially observed (Fig. 3). In such cases, the modifications may prevent the formation of the fold-back structures. Thus, it is possible that modified pri-/pre-miRNA are less efficiently recognized than non-modified ones by the processing enzyme, DCL1. The low modification rate (Table 3, Fig. 6) in the current data sets is also consistent with the second hypothesis, which proposes that modified pri/pre miRNAs are more prone to degradation and thus less likely to yield mature miRNAs [12,13]. Our analysis suggests that base modifications in miRNAs may prevent the maturation of miRNAs, although it is also possible that some base modifications in miRNAs may inhibit or enhance the targeting of miRNAs.

## Conclusion

Our data on tRNAs clearly show that substitutions in cDNAs can indicate RNA base modifications. This research thus found a new analytical strategy for predicting RNA base modifications and provided the first detailed bioinformatics analysis on RNA modifications in *Arabidopsis* tRNAs.

More importantly, this is the first report of the possibility of RNA base modifications in miRNAs with methyl or acetyl groups. Our results, which are based on carefully chosen sRNA sequences, strongly suggest that some RNA bases might be modified in miRNAs. We showed that one-base mismatches in sRNAs were not random and much more abundant than expected, and that many substitution sites were supported by both MPSS and 454 technologies. It follows that these substitutions cannot be explained by sequencing errors. More importantly, substitution patterns in adenine and guanine of miRNAs were similar to those of tRNAs. This similarity suggests common base modifications in miRNAs and tRNAs, although the nature of base modifications in miRNAs can only be determined experimentally in the future. Base modifications likely provide another regulatory mechanism for the biogenesis and/or functioning of miRNAs.

## Methods

### Data sets

We used 238,538 sRNA sequences (of which 98,035 were unique) sequenced by the 454 technology (Table 1) [19,20]. The small RNA libraries were constructed from plants treated with abiotic stresses (cold, drought, salt, copper, UV and ABA, respectively) or infected with bacterial or fungal pathogens [38]. We also used public sRNA sequences obtained by Lu et al. [21,22] with 454 and MPSS technologies, which have 58,178 reads and 3,126,002 reads respectively. The read numbers are used with normalization in TPQ (Transcripts Per Quarter million), according to ref. [21]. When a single type of sRNA had multiple loci on the genome, the TPQ value was divided by the number of the loci.

### Mapping sRNAs to the genome

We mapped sRNA sequences to the genome sequences with the alignment program, SOAP [23]. We used the program with the "-v 1" option, enabling the program to find one mismatch when the sRNA is mapped to the genome. We added a further criterion for the sRNAs that mapped to the genome with one mismatch. If a single sRNA had more than two OMM loci, with substitutions in different positions or with different nucleotides, we discarded it. After the analysis of the substitution sites (Table 1, Fig 1), we selected OMM sRNAs only if the substitution sites were not in the initial or terminal three bases. Then, we compared the results of sRNAs from 454 and MPSS technologies. If a substitution of a genomic locus was supported with sRNAs from both 454 and MPSS technologies, we selected it for further analysis. These criteria decreased technology-dependent errors.

### Statistical tests

We used probabilities of OMM sRNAs caused by sequencing errors based on reported sequencing error rates [24,21]: 0.25% for MPSS sequences and 0.004% for 454 sequences. We calculated the expected numbers of OMM sRNAs caused by sequencing errors for each substitution site. Then we performed chi-square tests to assess whether the read numbers of OMM sRNAs were significantly larger than expected from sequencing errors.

### Analysis of tRNAs and miRNAs

In the analysis of RNA modification in tRNAs and miRNAs, we had to determine the clover leaf structure of tRNAs and fold-back structures of miRNAs. For this purpose, we used tRNAscan-SE [27] and information about miRNAs published at miRBase [39]. Perl scripts were used to summarize the results.

### Authors' contributions

KI performed all bioinformatics analyses. HJ contributed sRNA sequences. JKZ conceived of the study. KI and JKZ

wrote the paper. All authors read and approved the final manuscript.

## Additional material

### Additional File 1

*Distribution of numbers of OMM sRNAs supporting substitutions. 249 of total 1,187 sites (21.0%) were supported by the minimal number of unique OMM sRNAs (one OMM sRNA from 454 data set and one from MPSS data set). This graph has a local peak where the number of unique OMM sRNA is six. 245 substitution sites were supported more than 10 unique OMM sRNAs. This graph shows the numbers of unique OMM sRNAs supporting each substitution site.*

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-10-155-S1.ppt>]

### Additional File 2

*Substitutions found in EST sequences for pre-miRNAs. Panels A, B, and C show multiple alignments of non-substituted sequences and EST sequences for ath-MIR398c, ath-MIR835, and ath-MIR860, respectively. In the multiple alignments lines labeled with miRNA IDs represent non-substituted sequences. In these lines, nucleotides included in mature miRNAs are shown as capital letters. Bottom lines of each alignment block indicate the positions of substitutions. Note that substitution sites were observed not only within mature miRNA regions, but also outside of the mature miRNAs. Pound symbols in panel C indicate the region outside of EST.*

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-10-155-S2.pdf>]

### Additional File 3

*Detail of substitutions. Raw information of all observed base modifications was described in the file. The information was summarized according to categories of corresponding genes (tRNA, miRNA, rRNA, and the others). Genomic positions of the observed base modifications, bases of nucleotides in the cases of unsubstituted and substituted, TPQs supporting each bases, definitions for corresponding genes, and results of chi-squared tests are described. Details of the chi-squared test were described in methods.*

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-10-155-S3.xls>]

## Acknowledgements

This work was supported by National Institutes of Health grants R01GM070795 and R01GM059138 (J.-K. Zhu) and a long-time fellowship from TOYOBO bio-technology foundation (K. Iida). We greatly appreciate Mr. Satoshi Takahasi for his initial efforts on analysis of sRNA data.

## References

- Limbach PA, Crain PF, McCloskey JA: **Summary: the modified nucleosides of RNA.** *Nucleic Acids Res* 1994, **22**:2183-2196.
- Rozenski J, Crain PF, McCloskey JA: **The RNA Modification Database: 1999 update.** *Nucleic Acids Res* 1999, **27**:196-197.
- Ofengand J: **Ribosomal RNA pseudouridines and pseudouridine synthases.** *FEBS Lett* 2002, **514**:17-25.
- Johansson MJ, Byström AS: **The Saccharomyces cerevisiae TANI gene is required for N4-acetylcytidine formation in tRNA.** *RNA* 2004, **10**:712-719.
- Grosjean H, Auxilien S, Constantinesco F, Simon C, Corda Y, Becker HF, Foiret D, Morin A, Jin YX, Fournier M, Fourrey JL: **Enzymatic conversion of adenosine to inosine and to N1-methylinosine in transfer RNAs: a review.** *Biochimie* 1996, **78**:488-501.
- Rife JP, Cheng CS, Moore PB, Strobel SA: **N 2-methylguanosine is iso-energetic with guanosine in RNA duplexes and GNRA tetraloops.** *Nucleic Acids Res* 1998, **26**:3640-3644.
- Schaub M, Keller W: **RNA editing by adenosine deaminases generates RNA and protein diversity.** *Biochimie* 2002, **84**:791-803.
- Keller W, Wolf J, Gerber A: **Editing of messenger RNA precursors and of tRNAs by adenosine to inosine conversion.** *FEBS Lett* 1999, **4**:71-76.
- Dietrich A, Small I, Cosset A, Weil JH, Maréchal-Drouard L: **Editing and import: strategies for providing plant mitochondria with a complete set of functional transfer RNAs.** *Biochimie* 1996, **78**:518-29.
- Lee RC, Feinbaum RL, Ambros V: **The C. elegans heterochronic gene lin-4 encodes small RNAs with antisense complementarity to lin-14.** *Cell* 1993, **75**:843-854.
- Luciano DJ, Mirsky H, Vendetti NJ, Maas S: **RNA editing of a miRNA precursor.** *RNA* 2004, **10**:1174-1177.
- Yang W, Chendrimada TP, Wang Q, Higuchi M, Seeburg PH, Shiekhattar R, Nishikura K: **Modulation of microRNA processing and expression through RNA editing by ADAR deaminases.** *Nat Struct Mol Biol* 2006, **13**:13-21.
- O'Connell MA, Keegan LP: **Drosha versus ADAR: wrangling over pri-miRNA.** *Nat Struct Mol Biol* 2006, **13**:3-4.
- Kawahara Y, Zinshteyn B, Sethupathy P, Iizasa H, Hatzigeorgiou AG, Nishikura K: **Redirection of silencing targets by adenosine-to-inosine editing of miRNAs.** *Science* 2007, **315**:1137-1140.
- Fahlgren N, Howell MD, Kasschau KD, Chapman EJ, Sullivan CM, Cumble JS, Givan SA, Law TF, Grant SR, Dangl JL, Carrington JC: **High-throughput sequencing of Arabidopsis microRNAs: evidence for frequent birth and death of MIRNA genes.** *PLoS ONE* 2007, **2**:e219.
- Yu B, Yang Z, Li J, Minakhina S, Yang M, Padgett RW, Steward R, Chen X: **Methylation as a crucial step in plant microRNA biogenesis.** *Science* 2005, **307**:932-935.
- Ramachandran V, Chen X: **Small RNA metabolism in Arabidopsis.** *Trends Plant Sci* 2008, **13**:368-374.
- Wittig B, Wittig S: **Reverse transcription of tRNA.** *Nucleic Acids Res* 1978, **5**:1165-1178.
- Jin H, Vacic V, Girke T, Lonardi S, Zhu JK: **Small RNAs and the regulation of cis-natural antisense transcripts in Arabidopsis.** *BMC Mol Biol* 2008, **9**:6.
- Margulies M, Egholm M, Altman WE, Attiya S, Bader JS, Bemben LA, Berka J, Braverman MS, Chen YJ, Chen Z, Dewell SB, Du L, Fierro JM, Gomes XV, Godwin BC, He W, Helgesen S, Ho CH, Irzyk GP, Jando SC, Alenquer ML, Jarvie TP, Jirage KB, Kim JB, Knight JR, Lanza JR, Leamon JH, Lefkowitz SM, Lei M, Li J, et al.: **Genome sequencing in microfabricated high-density picolitre reactors.** *Nature* 2005, **437**:376-380.
- Lu C, Tej SS, Luo S, Haudenschild CD, Meyers BC, Green PJ: **Elucidation of the small RNA component of the transcriptome.** *Science* 2005, **309**:1567-1569.
- Lu C, Kulkarni K, Souret FF, MuthuVallippan R, Tej SS, Poethig RS, Henderson IR, Jacobsen SE, Wang W, Green PJ, Meyers BC: **MicroRNAs and other small RNAs enriched in the Arabidopsis RNA-dependent RNA polymerase-2 mutant.** *Genome Res* 2006, **16**:1276-1288.
- Li R, Li Y, Kristiansen K, Wang J: **SOAP: short oligonucleotide alignment program.** *Bioinformatics* 2008, **24**:713-714.
- Meyers BC, Tej SS, Vu TH, Haudenschild CD, Agrawal V, Edberg SB, Ghazal H, Decola S: **The use of MPSS for whole-genome transcriptional analysis in Arabidopsis.** *Genome Res* 2004, **14**:1641-1653.
- Kawaji H, Nakamura M, Takahashi Y, Sandelin A, Katayama S, Fukuda S, Daub CO, Kai C, Kawai J, Yasuda J, Carninci P, Hayashizaki Y: **Hidden layers of human small RNAs.** *BMC Genomics* 2008, **9**:157.
- Lee SR, Collins K: **Starvation-induced cleavage of the tRNA anticodon loop in Tetrahymena thermophila.** *J Biol Chem* 2005, **280**:42744-42749.
- Schattnr P, Brooks AN, Lowe TM: **The tRNAscan-SE, snscan and snoGPS web servers for the detection of tRNAs and snoRNAs.** *Nucleic Acids Res* 2005, **33**:V686-V689.

28. Roovers M, Wouters J, Bujnicki JM, Tricot C, Stalon V, Grosjean H, Droogmans L: **A primordial RNA modification enzyme: the case of tRNA (m1A) methyltransferase.** *Nucleic Acids Res* 2004, **32**:465-476.
29. Ozanick SG, Bujnicki JM, Sem DS, Anderson JT: **Conserved amino acids in each subunit of the heterologomeric tRNA m1A58 Mtase from *Saccharomyces cerevisiae* contribute to tRNA binding.** *Nucleic Acids Res* 2007, **35**:6808-6819.
30. Edqvist J, Stråby KB, Grosjean H: **Enzymatic formation of N2,N2-dimethylguanosine in eukaryotic tRNA: importance of the tRNA architecture.** *Biochimie* 1995, **77**:54-61.
31. Steinberg S, Cedergren R: **A correlation between N2-dimethylguanosine presence and alternate tRNA conformers.** *RNA* 1995, **1**:886-891.
32. Niederberger C, Gräub R, Costa A, Desgrès J, Schweingruber ME: **The tRNA N2,N2-dimethylguanosine-26 methyltransferase encoded by gene *trm1* increases efficiency of suppression of an ochre codon in *Schizosaccharomyces pombe*.** *FEBS Lett* 1999, **464**:67-70.
33. Kim SH, Sussman JL, Suddath FL, Quigley GJ, McPherson A, Wang AH, Seeman NC, Rich A: **The general structure of transfer RNA molecules.** *Proc Natl Acad Sci USA* 1974, **71**:4970-4974.
34. Agris PF, Vendeix FA, Graham WD: **tRNA's wobble decoding of the genome: 40 years of modification.** *J Mol Biol* 2007, **366**:1-13.
35. Barraud P, Golinelli-Pimpaneau B, Atmanene C, Sanglier S, Van Dorselaer A, Droogmans L, Dardel F, Tisné C: **Crystal structure of *Thermus thermophilus* tRNA m1A58 methyltransferase and biophysical characterization of its interaction with tRNA.** *J Mol Biol* 2008, **377**:535-550.
36. Urbonavicius J, Armengaud J, Grosjean H: **Identity elements required for enzymatic formation of N2,N2-dimethylguanosine from N2-monomethylated derivative and its possible role in avoiding alternative conformations in archaeal tRNA.** *J Mol Biol* 2006, **357**:387-399.
37. Ishitani R, Yokoyama S, Nureki O: **Structure, dynamics, and function of RNA modification enzymes.** *Curr Opin Struct Biol* 2008, **18**:330-339.
38. Katiyar-Agarwal S, Gao S, Vivian-Smith A, Jin H: **A novel class of bacteria-induced small RNAs in *Arabidopsis*.** *Genes Dev* 2007, **21**:3123-3134.
39. Griffiths-Jones S, Saini HK, van Dongen S, Enright AJ: **miRBase: tools for microRNA genomics.** *Nucleic Acids Res* 2008, **36**:D154-D158.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

