

Original Article

Prevalence and Genetic Analysis of α - and β -Thalassemia and Sickle Cell Anemia in Southwest Iran

Forozan H. Nezhad¹, Khojasteh H. Nezhad^{1*}, Parastoo M. Choghakabodi^{1,2}, Bijan Keikhaei¹

¹Research Center of Thalassemia and Hemoglobinopathies, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

²Department of Microbiology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

ARTICLE INFO

Article History

Received 11 March 2018

Accepted 25 August 2018

Keywords

Deletion mutants
Khuzestan
sickle cell anemia
thalassemia

ABSTRACT

This prospective study assessed the prevalence and genetic analysis of α - and β -thalassemia and sickle cell anemia (SCA) in Southwest Iran. Hematological indices were measured in 17,581 couples living in Khuzestan Province, Southwest Iran. Individuals with mean corpuscular volume <80 , mean corpuscular hemoglobin <27 , hemoglobin A2 $\geq 3/5$ were considered as β -thalassemia traits. Prevalence of minor β -thalassemia, α -thalassemia, SCA, iron deficiency anemia, and silent thalassemia were respectively identified in 995 (5.6%), 1169 (6.65%), 1240 (7.05%), 911 (5.18%), and 1134 (6.45%) individuals using a multiplex amplification refractory mutation system, and direct DNA sequencing of globin genes. Three codons IVS-II-1 (G \rightarrow A; 26%; $n = 13$), IVS-I-1 (G \rightarrow T; 16%; $n = 8$), and IVS-I-110 (G \rightarrow A; 14%; $n = 7$) were the most frequent mutants and IVS-II-1 was the most common β -thalassemia mutation. Also, based on a gap-polymerase chain reaction assay, genotype frequencies of α -globin mutations were $-\alpha^{3.7\text{ kb}}$ (50%; $n = 25$), Med/ $\alpha\alpha^{\text{thal}}$ (12%; $n = 6$), and $-\alpha 4.2/\alpha\alpha$ (10%; $n = 5$), which were the most frequent deletion mutants (72% in total). The most common deletion (50%) was $-\alpha^{3.7\text{ kb}}$. Our data suggest that the population of Southwest Iran is at high risk of α - and β -thalassemia caused by these deletion mutants and SCA. Our findings will be useful for developing an efficient control program and genetic counseling.

© 2018 Atlantis Press International B.V.

This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).

1. INTRODUCTION

Thalassemia is a genetic abnormality involving mutations of the genes responsible for hemoglobin production in the blood. There are two broad types of thalassemia, α - and β -thalassemia; each of which has a different prevalence among certain ethnicities or population groups. Approximately 4.5 of every 10,000 live births throughout the world are affected by thalassemia (<http://www.ironhealthalliance.com/disease-states/thalassemia/epidemiology-and-pathophysiology.jsp>). α -Thalassemia is more frequent in Southeast Asia than in other areas of the world, and up to 40% of genetic traits have been found in thalassemia traits (TTs) (1–30%). People living in the Mediterranean, African, and South Asian areas are more likely to be affected by β -thalassemia. Genetic prevalence of β -thalassemia throughout the world is 2–18% (affected by a gene mutation) in the Eastern Mediterranean and 0–11% in Southeast Asia [1]. Approximately 5% of the global population have a variation in the α or β part of the hemoglobin molecule, although some of these are asymptomatic and known as silent traits. In fact, only 1.7% of the global population have signs as a result of the gene mutations, known as α -TT. However, tribal or ethnic groups are more likely to be affected and 5–30% of the population may be symptomatic among these groups [1].

Iran has an area of 1,648,000 km², and like many other countries in the region, has a large number of patients with major thalassemia. α -Thalassemia is infrequent in Iran. Gene frequency of β -thalassemia is high, and it varies considerably from area to area. Its highest rate ($>10\%$) is found around the Caspian Sea and Persian Gulf. Prevalence of β -thalassemia in other areas is between 4% and 8% [2]. In Southern Iran, the IVS-II-1 (G \rightarrow A) mutation is the most frequent (31%) mutation for β -thalassemia. In Khuzestan Province, in Southern Iran, the frequency of β -thalassemia minor is also high and reaches 10% [2].

According to a study conducted by Zandian et al. [3] in Khuzestan, from 152 volunteer couples (342 individuals) from Arab ethnic groups in Dashte-Azadegan and Khorramshahr cities, 3.63% and 10.57%, respectively, had sickle cell trait; Of these, 84.21% had normal hematological indices [mean corpuscular volume (MCV) and mean corpuscular hemoglobin (MCH)]. A few studies on genetic diversity of β -thalassemia mutations and sickle cell anemia (SCA) have been conducted in Iran, particularly in Khuzestan Province. Therefore we decided to carry out this cross-sectional and prospective study to assess the genetic diversity and prevalence of α - and β -thalassemia and their relationship with each other, detection of genetic diversity of α - and β -thalassemia minor, and detection of genetic diversity of SCA among 17,581 volunteer couples living in Khuzestan Province, Southwest Iran. Eventually data for the analysis of mutated genes, mutational pattern, as well as their prevalence were submitted to the Research Center of Thalassemia and Hemoglobinopathies, Jundishapur University of Medical Sciences, Ahvaz, Iran. Data were then submitted to Khuzestan

*Corresponding author. Email: khojasteh.h32@yahoo.com

Healthcare and Treatment Institute, Khuzestan, Iran to develop a control program in order to reduce the prevalence of thalassemia.

2. MATERIALS AND METHODS

2.1. Study Design

This was a prospective study that was carried out in the Department of Pathology and Thalassemia Control Cell, Shafa Hospital in Ahvaz, the capital city of Khuzestan Province, Southern Iran (Latitude: 31.436015, Longitude: 49.041312) between February 2014 and December 2017. A total of 17,581 prospective couples from Ahvaz (Latitude: 31.3183272, Longitude: 48.67061869999998) and Abadan (Latitude: 30.347296, Longitude: 48.2934) cities of Khuzestan Province were screened for the presence of thalassemia or any structural variant. All participants and their family members with hematological indices were questioned about their medical history. A 5-mL intravenous blood sample was collected in ethylenediamine tetraacetic acid (EDTA) anticoagulant. Red cell indices were measured on an automated hematology analyzer (Sysmex KX 21; Sysmex Corporation, Kobe, Japan). Hemoglobin (Hb)A2 and HbF were studied by high-performance liquid chromatography (HPLC) used for chromatographic separation of human Hb [4,5].

2.2. Sample Collection and Preparation

Five milliliters of intravenous blood was collected in a vacuum collection tube containing EDTA, which was stored at 3–8°C for a maximum 7 days if processing were delayed. HbA2 calibrators and normal and abnormal controls were analyzed at the beginning of each run.

2.3. Thalassemia Screening

At the first, the men's red cell indices were checked by complete blood count. If they had microcytosis (MCH < 27 pg or MCV < 80,

the women's red cell indices were tested too. When both were microcytic, their HbA2 concentrations were measured by HPLC (Model D10; Bio-Rad, France, using an ELITech Kit; ELITech Group, Puteaux, France). Tris-glycine buffer was used in order to measure HbA2 by HPLC. Electrophoresis was performed on cellogel at pH 8.5 in Tris-glycine buffer for 90 minutes, as described previously [4,5]. A concentration >3.5% was indicative of β -TT. Microcytic individuals with HbA2 concentration in the normal range (1.5–3.5%) were treated with iron and their indices rechecked. Patients who had TT (MCV < 80 fL, MCH < 27 pg/L, and HbA2 \geq 3.5%) were examined using multiplex amplification refractory mutation system (M-ARMS) to detect α - and β -thalassemia mutations, direct DNA sequencing of α - and β -globin genes, and gap-polymerase chain reaction (PCR) for globin gene deletions, respectively.

Inclusion criteria for the iron deficiency anemia (IDA) group were Hb <13 g/dL for men and <12 g/dL for women, MCV < 80 fL and MCH < 27 pg for both sexes, and ferritin < 28 ng/mL for men and < 6 ng/mL for women [6,7]. Exclusion criteria for the IDA group included the presence of mutations associated with α -TTs and/or β -TTs. For inclusion in the β -TT group, individuals had MCV < 80 fL, MCH < 27 pg, and HbA2 > 3.5% [4,5]. α -TT was confirmed by the presence of mutations (Fig. 1).

Silent α/β -thalassemia carriers have no signs or symptoms of the disease, but are able to pass it on to their children. In our study, silent thalassemia carriers showed a normal hematological picture or slight alterations in some hematological parameters (mild changes in erythrocyte morphology, or MCV below the normal mean), and normal Hb status or slight alterations in HbA2 level, while α/β -globin synthesis ratio was abnormal (>1 or <1). Genotype and some hematological parameters in some individuals with silent or α - or β -thalassemia are shown (see Table 1).

2.4. Thalassemia Mutation Analysis

Genomic DNA was extracted from peripheral blood lymphocytes using a QIAamp DNA Blood Mini Kit (QIAGEN, Valencia, CA, USA).




Figure 1 | Inclusion and exclusion criteria. Hb, hemoglobin; IDA, iron deficiency anemia; MCH, mean corpuscular hemoglobin; MCV, mean corpuscular volume; TT, thalassemia trait

Table 1 | Distribution of some hematological parameters in subjects with silent α - or β -thalassemia

Silent thalassemia	Hb (g/dL)	MCV	HbA2	α/β ratio	Genotype	
					β	α
Silent β^+ thalassemia (-101 C \rightarrow T mutation)	14.2 \pm 1.6	85 \pm 1.2	3.3 \pm 0.3	1.1 \pm 0.14	-101/b _A	$\alpha\alpha/\alpha\alpha$
Silent β^+ thalassemia (IVS II 844 C \rightarrow G mutation)	14.4 \pm 1.8	86 \pm 1.5	3.4 \pm 0.23	1.10 \pm 0.03	AA ♂ CP ♀	$\alpha\alpha/\alpha\alpha$ $\alpha\alpha/\alpha\alpha$
Silent α^+ thalassemia (-a ₂ ^{NcoI} mutation)	13.6 \pm 0.2	75 \pm 0.2	2.3 \pm 0.23	0.88 \pm 0.08	β^A/β^A	$\alpha^{\text{NcoI}}\alpha/\alpha\alpha$
Silent α^+ thalassemia (a ₂ ^{HphI} mutation)	13.5 \pm 1.1	78 \pm 1.2	2.4 \pm 0.04	0.85 \pm 0.12	β^A/β^A	$\alpha^{\text{HphI}}\alpha/\alpha\alpha$

Data are presented as mean \pm SD; Hb, hemoglobin; MCV, mean corpuscular volume.

Table 2 | Multiplex gap-PCR protocol for the diagnosis of 3.7- and 4.2-kb α^+ -thalassemia deletions

Multiplex gap-PCR protocol for the diagnosis of - $\alpha^{3.7}$ and - $\alpha^{4.2}$ deletions

Primer sequences [1]

Primer	Description	Sequence	Annealing T(°C)
1	$\alpha 2/3.7$ -F	CCCCTCGCCAAGTCCACCC	64
2	3.7/20.5-R	AAAGCACTCTAGGGTCCAGCG	64
3	$\alpha 2$ -R	AGACCAGGAAGGGCCGGTG	64
4	4.2-R	CCCGTTGGATCTTCTCATTTC	64
5	4.2-F	GGTTACCCATGTGGTGCCTC	64

PCR mix

Component	μL
$\alpha 2/3.7$ -F (10 μM)	1.0
$\alpha 2$ -R (10 μM)	0.25
$\alpha 2/20.5$ -R (10 μM)	1.0
4.2-F (10 μM)	1.0
4.2-R (10 μM)	1.5
10 \times buffer (750 mM Tris-HCl pH 8.8, 200 mM (NH ₄) ₂ SO ₄ , 0.1% Tween 20)	2.5
25 mM MgCl ₂	1.5
dNTPs (1 mM)	5.0
Betaine (5 M)	3.75
DMSO (10%)	1.25
Platinum Taq (5 U/mL)	0.1
DNA template (100 ng/mL)	
Water	

Gel electrophoresis conditions

Run PCR products out on 1.5% (1:1 Nusieve: agarose) gel for 2–3 h

Interpretation of results

PCR fragment size (bp)	Genotype	Product of primers
2020	α^+ -thalassaemia: - $\alpha^{3.7}$	1 + 2
1800	Normal ($\alpha\alpha$)	1 + 3
1628	α^+ -thalassaemia: - $\alpha^{4.2}$	4 + 5

DMSO, dimethyl sulfoxide; PCR, polymerase chain reaction; T, temperature.

β -Globin gene mutations were first characterized using an M-ARMS to detect common mutations in Southwest Iranian populations, including IVS-II-1 (G \rightarrow A), IVS-I-1 (G \rightarrow T), IVS-I-110 (G \rightarrow A), CDs 36/37 (-T), IVS-I-5 (G \rightarrow C), cd5, IVS-I-6 (T \rightarrow C), and cd39 (C + T), as previously described [8]. Uncommon β -thalassemia genes were further characterized by direct DNA sequencing of all coding regions and exon-intron boundaries to detect uncommon point mutations, as described previously [9]. α - and β -Thalassemia mutations were subsequently screened by gap-PCR to detect entire globin gene deletions, as previously described [10–13].

2.5. Gap-PCR

Gap-PCR techniques (amplification using oligoprimers flanking deletion breakpoints) are used to detect different types of globin gene deletions, such as common α -thalassemia deletion mutations and α -gene duplication [11]. A typical gap-PCR test is illustrated for the diagnosis of α -thalassemia, and the primers can be multiplexed [14,15], as shown in Tables 2 and 3. The 3.7- and 4.2-kb α^+ -thalassemia deletions can be detected in one assay [11,12], and the Mediterranean thalassemia deletion (-^{MED}) in the

Table 3 | Multiplex gap-PCR protocol for the diagnosis of α -MED thalassemia deletion

Multiplex gap-PCR protocol for the diagnosis of α -MED deletions			
Primer sequences [2]			
Primer	Name	Sequence	Annealing T (°C)
1	MED(F)	CGATGAGAACATAGTGAGCAGAATTGCAGG	60
2	MED(R)	ACGCCGACGTTGCTGCCAGCTTCTTCCAC	60
3	α 2-R	AGACCAGGAAGGGCCGGTG	64
4	4.2-R	CCCGTTGGATCTTCTCATTCC	64

PCR mix

Component	μ L
MED(F) (10 μ M)	0.4
MED(R) (10 μ M)	0.4
10 \times buffer (750 mM Tris-HCl pH 8.8, 200 mM (NH ₄) ₂ SO ₄ , 0.1% Tween 20)	2.5
25 mM MgCl ₂	1.5
dNTPs (1 mM)	4.0
Betaine (5 M)	3.75
DMSO (10%)	1.25
Platinum Taq (5 U/mL)	0.1
DNA template (100 ng/mL)	1.0
Water	6.2

Gel electrophoresis conditions

Run PCR products out on 2% (1:1 Nusieve:agarose) gel for 1–1.5 h

Interpretation of results

PCR fragment size (bp)	Genotype	Product of primers
1010	Normal ($\alpha\alpha$)	3 + 4
875	α^+ -thalassaemia: α -MED	1 + 2

PCR, polymerase chain reaction; T, temperature.

second assay [12]. Specific primer details are listed in Tables 2 and 3 for the multiplex diagnosis of the common α -thalassemia deletion genotypes.

2.6. Statistical Analysis

Statistical analysis of hematologic indices and the results of molecular tests and mutation types among carrier couples was performed by one-way analysis of variance with SPSS software (SPSS Inc., Chicago, IL, USA).

3. RESULTS

Among all volunteers, 995 (5.6%) had β -thalassemia minor (435 female and 560 male). Of these, 695 cases (69.85%) belonged to the city of Abadan and 300 (30.15%) to Ahvaz. Average value of HbA2 among female and male populations in the city of Abadan was 4.97% and 5.23%, respectively. In Ahvaz, the average value of HbA2 was 4.3% in women and 5.01% in men. The mean hematological indices among TTs are shown in Table 4.

In a molecular study of 50 individuals (5%) randomly selected using M-ARMS, eight codons from the β -globin defects accounted for 90% of the total β -thalassemia mutations: IVS-II-1 (G \rightarrow A; 26%; n = 13), IVS-I-1 (G \rightarrow T; 16%; n = 8), IVS-I-

110 (G \rightarrow A; 14%; n = 7), CDs 36/37 (–T; 10%; n = 5), IVS-I-5 (G \rightarrow C; 8%; n = 4), IVSI-6 (T \rightarrow C; 6%; n = 3), cd5 (6%; n = 3), and cd39 (C + T; 4%; n = 2); the most common mutation was IVS-II-1 (Fig. 1). Prevalence of α -TTs was 6.65% (n = 1169). Also, 50 individuals were tested for α -thalassemia gene mutations based on a gap-PCR assay. Genotype frequencies of α -globin mutations were $-\alpha^{3.7\text{ kb}}$ (50%; n = 25), Med/ $\alpha\alpha^{\text{thal}}$ (12%; n = 6), and $-\alpha 4.2/\alpha\alpha$ (10%; n = 5), which were the most frequent deletion mutants (72% in total). The $-\alpha^{3.7\text{ kb}}$ deletion was the most common deletion (Fig. 2).

Prevalence of SCA, IDA, and silent thalassemia was 1240 (7.05%), 911 (5.18%), and 1134 (6.45%) respectively, as shown in Table 5 and Fig. 2. Distribution of SCA haplotypes among people in Southwest Iran is shown in Table 6. Genotype and hematological characteristics of subjects with silent α - or β -thalassemia are presented in Table 1.

4. DISCUSSION

Thalassemia is common in Iran due to a tradition of inbreeding, a conservative religious culture, and a large number of ethnic groups in different areas of the country. Genetic prevention programs based on hospital-based screening and prenatal diagnosis were started in 1997, and Iranian laws were modified between 1998 and 2005 to permit abortion of affected fetuses [16]. Some research on the clinical and laboratory presentations of β -thalassemia

Table 4 | Mean of hematological indices among thalassemia traits^a

Blood indices		Type of mutation				p
		No mutation	α	β	α and β	
MCV (fL)	Female	77.60 ± 6.09	74.65 ± 5.86	62.72 ± 5.13	67.5 ± 4.75	0.001
	Male	77.14 ± 6.07	75.11 ± 6.10	64.02 ± 5.99	68.39 ± 7.25	0.001
	Total	77.40 ± 6.03	75.40 ± 6.04	63.64 ± 5.72	68.36 ± 6.22	>0.001
MCH (pg)	Female	24.99 ± 2.38	23.7 ± 2.32	20.01 ± 2.42	21.51 ± 1.71	0.001
	Male	24.83 ± 2.37	24.34 ± 2.36	20.56 ± 2.70	22.57 ± 2.34	0.001
	Total	24.92 ± 2.36	24.3 ± 2.34	20.29 ± 2.57	22.56 ± 2.04	>0.001
RBC	Female	4.77 ± 0.37	5.04 ± 0.14	5.43 ± 0.60	5.22 ± 0.65	0.001
	Male	5.65 ± 0.51	5.79 ± 0.47	6.29 ± 0.57	6.19 ± 0.70	0.001
	Total	5.14 ± 0.61	5.40 ± 0.57	5.89 ± 0.73	5.76 ± 0.82	>0.001
HbA2 (%)	Female	2.57 ± 0.37	2.60 ± 0.53	4.97 ± 0.89	5.39 ± 0.93	0.001
	Male	2.91 ± 0.96	2.95 ± 1.08	5.23 ± 1.23	4.87 ± 1.30	0.001
	Total	2.72 ± 0.711	2.77 ± 0.86	5.11 ± 1.08	5.08 ± 1.1	>0.001

^aMean MCH and MCV among β -TTs were less than those of α -thalassemia traits (mean difference: -4.01 and -11.76, respectively, $p = 0.001$). Mean HbA2 and RBC increased among the TTs (mean difference: 2.36 and 0.62, respectively, $p = 0.001$). Mean MCH and MCV among α - and β -TTs were less than in individuals with normal indices (mean difference: 2.36 and 9.04, respectively, $p = 0.001$). Mean MCH, MCV, HbA2, and RBC among α -TTs were not significantly different ($p > 0.001$). Comparisons are expressed according to one-way analysis of variance; Data are presented as mean ± SD; HbA2, hemoglobin A2; MCH, mean corpuscular hemoglobin; MCV, mean corpuscular volume; RBC, red blood cell; SD, standard deviation; TT, thalassemia trait.




Figure 2 | Genotype frequencies of β -thalassemia mutations

syndromes has been conducted in Iran, and unfortunately, no systematic large-scale investigation has been performed to clarify the status of thalassemia in Iran [17,18].

There are numerous gene mutations responsible for β -thalassemia in Iran [19]. These mutations are originally of Iranian, Mediterranean, Turkish, Kurdish, Egyptian, Indian, Asian Indian, Tunisian, Chinese, and Afro-American origin [20]. This notable heterogeneity may be due to Islamic education, which emphasizes fraternity and hospitality, and urges Muslims to accept newcomers with open arms. Iran hosts the largest number of refugees in the world; mostly from Iraq and Afghanistan during the past two decades. This massive ethnic/genetic commixture has led to an unexpectedly high number of different mutations of β -thalassemia in this country. Iran, particularly Khuzestan Province, is a thalassemia hot zone [20].

Yao et al. [21] demonstrated that the Li people in Hainan Province have a high incidence of $-\alpha^{4.2}$ and $-\alpha^{3.7}$ thalassemia. Our study showed that Khozestan Province has high genotype frequencies of α -globin mutations, including $-\alpha^{3.7}/\alpha\alpha$ ($-\alpha^{3.7\text{ kb}}$; 50%; $n = 25$), Med/ $\alpha\alpha^{\text{thal}}$ (12%; $n = 6$), and $-\alpha^{4.2}/\alpha\alpha$ ($-\alpha^{4.2\text{ kb}}$; 10%; $n = 5$), which were the most frequent deletion mutants (72% in total). The $-\alpha^{3.7\text{ kb}}$ deletion was the most common α -thalassemia deletion among

the α -TTs. This can be explained by the fact that this group of α -thalassemia deletions is common in Asia, where a high prevalence of α^+ -thalassemia has been observed.

Recent studies have revealed the presence of >47 different β -globin gene mutations responsible for β -thalassemia in Iran. IVS-II-1 (G → A) mutation followed by IVS-I-5 (G → C), codons 8/9 (+G), IVS-I-110 (G → A), IVS-I-1 (G → A), 25-bp deletion, IVS-I-6 (T → C), codon 5 (-CT), and codon 39 (C → T) mutations were the most frequent mutations, which accounted for 85% of the total β -thalassemia defects in Iran [19,22].

Rezaee et al. [23] attempted to study the origin of β -thalassemia mutations in different parts of Iran. They demonstrated that β -thalassemia mutations in different regions have different distribution patterns. These β -thalassemia mutations are indicative of the ancestral origin of the people who migrated to Iran from other regions of the world. In this study [23], in the southwest near the Arabian Peninsula, codons IVS-I-1 (G → A), 8/9 (+G), IVS-I-110 (G → A), IVS-II-1 (G → A), IVS-I-6 (T → C) and IVS-I-5 (G → C) were common mutations that caused 86% of cases of β -thalassemia. Codon IVS-I-1 is the most frequent mutation in Southwest Iran.

Karimi et al. [24] carried out a prospective study demonstrating that the IVS-II-1, IVS-I-110, IVS-I-1, and FSC 8/9 mutations were the most prevalent in the country, and IVS-II-1 was the most frequent in Southern Iran (having the highest rate of 24%).

In our study, the thalassemia syndromes (α - and β -thalassemia mutations, sickle cell disease) were assessed using M-ARMS, direct DNA sequencing of α - and β -globin genes, and gap-PCR among the 17,581 volunteer couples living in Khuzestan Province. Eight codons out of the β -globin defects accounted for 90% of the total β -thalassemia mutations: IVS-II-1 (G → A; 26%; $n = 13$), IVS-I-1 (G → T; 16%; $n = 8$), IVS-I-110 (G → A; 14%; $n = 7$), CD36/37 (-T; 10%; $n = 5$), IVS-I-5 (G → C; 8%; $n = 4$), IVS-I-6 (T → C; 6%; $n = 3$), cd5 (6%; $n = 3$), and cd39 (C + T; 4%; $n = 2$). IVS-II-1 (G → A)


Table 5 | Frequency distribution of SCA, β -thalassemia minor, IDA, and silent β - or α -thalassemia among thalassemia carriers in Southwest Iran

City	SCA	β -Thalassemia minor	IDA	Silent β^+ -thalassemia	Silent α^+ -thalassemia
Abadan	888 (76)	695 (69.8)	530 (3)	711 (4)	112 (0.63)
Ahvaz	280 (24)	300 (30.2)	381 (2.16)	224 (1.27)	87 (0.49)

Data are presented as n (%); IDA, iron deficiency anemia; SCA, sickle cell anemia.

Table 6 | Distribution of sickle-cell anemia haplotypes among different ethnic groups in Southwest Iran

Ethnic groups	Frequency (%)
Arab-Indian	0.37
Benin	0.17
Bantu	0.113
Senegal	0.18

**Figure 3** | Genotype frequencies of α -thalassemia mutations

mutation was the most frequent (26%) mutation for β -thalassemia in Khuzestan Province. Our study has confirmed the previous studies.

Prevalence of α -TTs was 6.65% ($n = 1169$) and $-\alpha^{3.7kb}$ was the most common α -thalassemia deletion (50%; $n = 25$), which was consistent with Zandian et al. [25]. It indicates the importance of identification of this gene in couples to prevent the occurrence of hemoglobin H (HbH) disease and hydrops fetalis.

Our study demonstrated the distribution of SCA haplotypes among different ethnic groups in Southwest Iran. The Arab ethnic group has the highest frequency of SCA traits among all ethnic groups, confirming the results of Zandian et al. [26].

5. CONCLUSION

The genetic basis and clinical severity of α - and β -thalassemia are heterogeneous among Iranians due to the presence of multiple ethnic groups in the country. The β -thalassemic IVSII-1 (G \rightarrow A) mutation had the highest frequency in Southwest Iran. As a Mediterranean mutation, it might reflect its independent origin, genetic admixture, and/or gene flow from neighboring countries. A broad spectrum of α -thalassemia alleles has been detected among Iranians and $-\alpha^{3.7kb}$ was the most prevalent thalassemia mutation. The results of this study could be useful to upgrade the program for prevention of neonatal thalassemia in Iran.

CONFLICTS OF INTEREST

Dr. Forozan H. Nezhad, First author, has received research grants from Ahvaz Jundishapur University of Medical Sciences. Other co-authors report no conflicts of interest relevant to this article.

REFERENCES

- [1] Xu XM, Zhou YQ, Luo GX, Liao C, Zhou M, Chen PY, et al. The prevalence and spectrum of alpha and beta thalassaemia in Guangdong Province: implications for the future health burden and population screening. *J Clin Pathol* 2004;57: 517–22.
- [2] Abolghasemi H, Amid A, Zeinali S, Radfar MH, Eshghi P, Rahiminejad MS, et al. Thalassemia in Iran: epidemiology, prevention, and management. *J Pediatr Hematol Oncol* 2007; 29:233–8.
- [3] Zandian K, Nateghi J, Kaikhahi B, Najmabadi H. Elucidation of α -thalassemia mutations in Khuzestan, Iran. *Genetics in the 3rd Millennium* 2007;5:1120–5.
- [4] Greer JP, Foerster J, Rodgers GM, Paraskevas F, Glader B, Arber DA, et al. *Wintrobe's clinical hematology*. 12th ed. Philadelphia: Lippincott Williams and Wilkins; 2009.
- [5] Schleider CT, Mayson SM, Huisman TH. Further modification of the microchromatographic determination of hemoglobin A2. *Hemoglobin* 1977;1:503–4.
- [6] World Health Organization. *Guidelines for the use of iron supplements to prevent and treat iron deficiency anemia*. Geneva: World Health Organization; 1998.
- [7] World Health Organization. *Iron deficiency anaemia: assessment, prevention and control. A guide for programme managers*. Geneva: World Health Organization; 2001.
- [8] Bhardwaj U, Zhang YH, Lorey F, McCabe LL, McCabe ER. Molecular genetic confirmatory testing from newborn screening samples for the common African-American, Asian Indian, Southeast Asian, and Chinese beta-thalassemia mutations. *Am J Hematol* 2007;78:249–55.
- [9] Sirichotiyakul S, Saetung R, Sanguansermisri T. Analysis of beta-thalassemia mutations in northern Thailand using an automated fluorescence DNA sequencing technique. *Hemoglobin* 2003;27:89–95.
- [10] Old J, Hartevelde CL, Traeger-Synodinos J, Petrou M, Angastiniotis M, Galanello R. *Prevention of thalassaemias and other haemoglobin disorders: Volume 2: Laboratory protocols*. 2nd edition. Nicosia, Cyprus: Thalassaemia International Federation; 2012.
- [11] Dodé C, Krishnamoorthy R, Lamb J, Rochette J. Rapid analysis of $-\alpha^{3.7}$ thalassaemia and alpha alpha alpha anti 3.7 triplication by enzymatic amplification analysis. *Br J Haematol* 1993;83:105–11.

- [12] Bowden DK, Vickers MA, Higgs DR. A PCR-based strategy to detect the common severe determinants of alpha thalassaemia. *Br J Haematol* 1992;81:104–8.
- [13] Liu YT, Old JM, Miles K, Fisher CA, Weatherall DJ, Clegg JB. Rapid detection of alpha-thalassaemia deletions and alpha-globin gene triplication by multiplex polymerase chain reactions. *Br J Haematol* 2000;108:295–9.
- [14] Embury SH. Advances in the prenatal and molecular diagnosis of the hemoglobinopathies and thalassemsias. *Hemoglobin* 1995;19:237–61.
- [15] Old J. Haemoglobinopathies. *Prenat Diagn* 1996;16:1181–86.
- [16] Strauss BS. Genetic counseling for thalassaemia in the Islamic Republic of Iran. *Perspect Biol Med* 2009;52:364–76.
- [17] Nasab AH. Clinical and laboratory findings in the initial diagnosis of homozygous beta thalassaemia in Fars Province, Iran. *Br J Haematol* 1979;43:57–61.
- [18] Gharib R, Ayazi S. Electrocardiographic findings in Iranian children with severe chronic anemia. Observations in iron deficiency, thalassaemia major, and miscellaneous other anemic states. *Clin Pediatr* 1972;11:630–33.
- [19] Merat A, Haghshenas M, Pour ZM, Plonczynski MW, Harrell AN, Coleman MB, et al. β -thalassaemia in southwestern Iran. *Hemoglobin* 1993;17:427–37.
- [20] Akhavan-Niaki H, Derakhshandeh-Peykar P, Banihashemi A, Mostafazadeh A, Asghari B, Ahmadifard MR, et al. A comprehensive molecular characterization of beta thalassaemia in a highly heterogeneous population. *Blood Cells Mol Dis* 2011;47:29–32.
- [21] Yao H, Chen X, Lin L, Wu C, Fu X, Wang H, et al. The spectrum of α - and β -thalassaemia mutations of the Li people in Hainan Province of China. *Blood Cells Mol Dis* 2014;53:16–20.
- [22] Najmabadi H, Karimi-Nejad R, Sahebjam S, Pourfarzad F, Teimourian S, Sahebjam F, et al. The β -thalassaemia mutation spectrum in the Iranian population. *Hemoglobin* 2001;25:285–96.
- [23] Rezaee AR, Banoei MM, Khalili E, Houshmand M. Beta-Thalassaemia in Iran: new insight into the role of genetic admixture and migration. *Scientific World J* 2012;2012:635183.
- [24] Karimi M, Yarmohammadi H, Farjadian S, Giordano PC. Beta-thalassaemia intermedia from southern Iran: IVS-II-1 (G \rightarrow A) is the prevalent thalassaemia intermedia allele. *Hemoglobin* 2002;26:147–54.
- [25] Zandian K, Nateghi J, Keikhaie B, Pedram M, Hafezi-Nejad N, Hadavi V, et al. alpha-thalassaemia mutations in Khuzestan Province, Southwest Iran. *Hemoglobin* 2008;32:546–52.
- [26] Zandian KM, Pedram M, Ghahfarokhi FK. Pre-marriage sickle cell screening program in south region of Iran, a pilot study on 50 cases of sickle trait. *Iran J Blood Cancer* 2009;2:55–7.