

Coronary Artery Bypass Graft Patients' Perception about the Risk Factors of Illness: Educational Necessities of Second Prevention

Abstract

Background: Patients' beliefs about the cause of cardiac disease (perceived risk factors) as part of the global psychological presentation are influenced by patients' health knowledge. Hence, the present study aimed to assess the relationship between actual and perceived risk factors, identification of underestimated risk factors, and indication of underestimation of every risk factor. **Materials and Methods:** In this cross-sectional study, data of 313 coronary artery bypass graft (CABG) patients admitted to one hospital in the west of Iran were collected through a demographic interview, actual risk factors' checklist, open single item of perceived risk factors, and a life stressful events scale. Data were analyzed by means of Spearman's correlation coefficients and one-sample Z-test for proportions. **Results:** Although there are significant relations between actual and perceived risk factors related to hypertension, family history, diabetes, smoking, and substance abuse ($P < 0.05$), there is no relation between the actual and perceived risk factors, and patients underestimate the role of actual risk factors in disease ($P < 0.001$). The patients underestimated the role of aging (98.8%), substance abuse (95.2%), overweight and obesity (94.9%), hyperlipidemia (93.1%), family history (90.3%), and hypertension (90%) more than diabetes (86.1%), smoking (72.5%), and stress (54.7%). **Conclusion:** Cardiac patients seem to underestimate the role of aging, substance abuse, obesity and overweight, hyperlipidemia, family history, and hypertension more than other actual risk factors. Therefore, these factors should be highlighted to patients to help them to (i) increase the awareness of actual risk factors and (ii) promote an appropriate lifestyle after CABG surgery.

Keywords: Cardiovascular disease, patients' perception, risk factors, second prevention

Introduction

Cardiovascular diseases (CVDs) are one of the most common chronic diseases and one of the main causes of mortality worldwide.^[1] In the past years, approximately 15 million persons in Iran were affected by CVDs, while a large part of the general population is at risk of developing cardiovascular pathologies in the next decades.^[2] Cardiac risk factors include biological, environmental, physiological, behavioral, and psychological factors.^[3] For example, a recent study by Compare *et al.* suggested that personality traits such as social inhibition and anger are associated with the presence and severity of CVDs.^[4] Moreover, the link between emotional regulation and depression seems to represent a potential mediator between heart and mind.^[5] All these cardiac risk factors have a direct effect on the incidence and persistence of CVDs.^[6] In addition, they can affect the patients' health attitudes.^[7] Indeed, according to the health

belief model, patients' attitudes, cognitive beliefs, and their emotional reactions to disease and treatment can predict their health behaviors independently.^[8-10] For example, poor adherence to treatment represents a major limitation in the management of patients at a high risk of cardiovascular events.^[11] However, the mismatch between actual and perceived risk factors may affect the patient's cognitions^[12] and lead to increased anxiety.^[13,14] Patients' beliefs about the cause of cardiac disease (perceived risk factors) are part of the global psychological presentation of disease which is affected by patients' health literacy.^[7] It has been suggested that wrong thoughts about cardiac disease and its risk factors can increase the probability of negative outcomes and of further hospitalization.^[15] In addition, perceived risk, or personal beliefs about the probability of facing health threats, and the true perception of risk factors of disease could promote a healthy lifestyle.^[16] Indeed, according to Byrne *et al.*,^[10] secondary

Ali Soroush^{1,2},
Saeid Komasi^{1,3},
Mozhgan Saeidi¹,
Behzad
Heydarpour¹,
Danilo Carrozzino⁴,
Mario Fulcheri⁴,
Paolo Marchettini⁵,
Massimo Rabboni⁶,
Angelo Compare⁷

¹Cardiac Rehabilitation Center, Imam Ali Hospital, Kermanshah University of Medical Sciences, ²Lifestyle Modification Research Center, Imam Reza Hospital, Kermanshah University of Medical Sciences, ³Clinical Research Development Center, Imam Reza Hospital, Kermanshah University of Medical Sciences, Kermanshah, Iran, ⁴Department of Psychological, Health and Territorial Sciences, University "G. d'Annunzio" of Chieti-Pescara, Chieti, ⁵Department of Neurology, San Raffaele Hospital Milano and Pain center, Centro Diagnostico Italiano, Milano, ⁶Department of Psychiatry, Papa Giovanni XXIII Hospital, Bergamo, ⁷Department of Human and Social Sciences, University of Bergamo, Bergamo, Italy

Address for correspondence:
Dr. Mozhgan Saeidi,
Cardiac Rehabilitation Center,
Imam Ali Hospital, Kermanshah
University of Medical Sciences,
Shahid Beheshti Boulevard,
Kermanshah, Iran.
E-mail: m_saeidi20@yahoo.com

Access this article online

Website: www.annals.in

DOI: 10.4103/aca.ACA_19_17

Quick Response Code:



How to cite this article: Soroush A, Komasi S, Saeidi M, Heydarpour B, Carrozzino D, Fulcheri M, *et al.* Coronary artery bypass graft patients' perception about the risk factors of illness: Educational necessities of second prevention. *Ann Card Anaesth* 2017;20:303-8.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

prevention needs longtime management of risk factors by patients with the persistent cardiac disease. Although, failure of rehabilitation and secondary prevention programs and the inability of patients in the lifestyle modification is a result of their dysfunctional beliefs about disease.

Although Perkins-Porras *et al.*^[7] found that there is a relation between actual and perceived risk factors, other studies indicated that there is no proportion between actual and perceived risk factors among patients.^[12,16] This lack of proportion could be explained by patients' poor knowledge and awareness of risk factors. A recent study, for example, reported that 25% of people have no correct perception about unhealthy nutrition, poor exercise, and smoking in the incidence of CVDs.^[17] There is more worry about some of these risk factors, and it has been suggested that 83% of cardiac patients with diabetes and 69% of cardiac patients who smoke have a poor perception about the role of these risk factors in their cardiac problems.^[16] Based on different studies, patients consider stress (21%–64.4%), smoking (11.3%–58.8%), hypertension (3.2%–53.9%), hereditary and genetics (4.3%–48.8%), unhealthy nutrition (13.3%–46.6%), overweight and obesity (0.5%–39.4%), and other factors as the main causes of their disease.^[7,18-21] Approximately 10%–65% of patients are unable to identify the cause of their disease,^[18,21] while about 60% of cardiac patients have two or more risk factors for a further cardiac event.^[16] Thus, cardiac patients usually underestimate the risk of disease and its risk factors. These cognitions may increase the risk of illness.^[16]

It is clear that cardiac patients underestimate the risk factors, but it is not obvious which factors are underestimated more than others. Therefore, the present study aimed to respond to three questions: (1) Is there any relation between actual and perceived risk factors? (2) Which risk factors are underestimated significantly? (3) What is the level of underestimation of each risk factor? It is possible that the responses are effective in the design of secondary preventions and future training in cardiac rehabilitation.

Materials and Methods

Design and context

In this cross-sectional study, a number of cardiovascular patients who were admitted to Imam Ali hospital in Kermanshah city, Iran, for coronary artery bypass graft (CABG), volunteered to participate before their discharge. This hospital is a specialized cardiac center located in western Iran, with a capacity of 214 beds.

Inclusion and exclusion criteria

Inclusion criteria included: (1) 30–80 years of age, (2) appropriate perception and emotional–physical abilities, (3) being able to understand the language and fluent speech. Exclusion criteria included: (1) fatigue and lack of tendency for participation and (2) having risk factors out

of the assessed risk factors of this study. The study was conducted following the guidelines of the Declaration of Helsinki.

Participants

Among 368 CABG patients who were admitted in the department of cardiac surgery from May to August 2015, 333 patients met the inclusion criteria and therefore were included in this study. Overall, 35 cases were excluded (21 cases: aging, 3 cases: cognitive dysfunction, and 11 cases: refused to participate) from the study. In addition, twenty cases were excluded from the study due to different risk factors than those assessed in this study. Thus, the final sample was of 313 patients. Furthermore, *a priori* sample size analysis (level of error 0.05) suggested that a minimum of 188 participants were required to obtain significant results. Data were collected after obtaining written consent form.

Data collection

At first, researchers performed a short interview with patients to check the inclusion criteria. Then, demographic checklist was collected. The clinical psychologist of the team administered the life stressful events scale.^[22] The psychologist read the questions and registered answers for patients with an age higher than 50 years and with illiterate patients. In addition, the team referred to the medical records to obtain more information. Perceived risk factors were assessed through a single-item question,^[2,3,20,21] in addition, patients' viewpoints were recorded. Then, actual risk factors were assessed and recorded by a cardiologist. Finally, data were entered into SPSS software and analyzed.

Instruments

Demographics and actual risk factors' checklist

Actual risk factors' checklist was fulfilled through cardiologist's and clinical psychologist's interviews with patients. It included information about demographic variables (age, gender, education level, and occupational status) and actual risk factors (diabetes, hypertension, hyperlipidemia, body mass index, history of smoking, opiate abuse, and familial history of cardiac disease).^[1,23] Body mass index was measured by a nutritionist.

Perceived cardiac risk factors

According to the new category about the perceived risk factors for CVDs,^[2,3,13,20,21] we used an open single-item approach, asking all participants "Which factor do you know as the main factor for your disease?" Patients' responses were recorded accurately.

Holmes and Rahe Scale of Stress of Life Events

Holmes and Rahe Scale of Stress of Life Events questionnaire was made by Holmes and Rahe in 1963 to evaluate the presence of 41 stressful events ("list of recent events"). Based on this scale, the life changes during the

recent 6–12 months were evaluated and the total score was obtained through a sum of the scores. The item value is between 11 and 100. A total score ranging from 150 to 200 is associated with an increased probability (equal to 37%) to develop a future disease, while this probability increases to 50%–80% if scores are in ranges between 200–300 or higher than 300.^[22] The reliability of this scale was measured by Vafaii,^[24] which reported an acceptable internal validity (Cronbach's $\alpha = 0.68$). We used a cutoff score of >150 to identify stressed participants.

Statistical analyses

Spearman's correlation coefficient was used for assessment of the relations between variables. In addition, one-sample Z-test for proportions was used to compare actual and perceived risk factors. The percentages of actual and perceived risk factors and the mean of the stress score were reported. All statistical analyses were performed using Statistical Package for the Social Sciences version 20.0 (IBM SPSS, Armonk, NY, USA). All statistical tests were two sided; $P \leq 0.05$ was considered statistically significant. Before analyses, statistical assumptions were checked;^[25] results evidenced that all variables were normally distributed.

Results

The mean age of participants was 61 years and 63.6% of them were male. The demographic data and stress scores (computed from Holmes and Rahe Scale of Stress of Life Events questionnaire) of the entire sample are shown in Table 1.

Table 2 shows the correlation between actual and perceived risk factors. Results evidenced statistically

Table 1: The sample demographics and stress

| Variable | Frequency (%) |
|---------------------|---------------|
| Sex | |
| Female | 114 (36.4) |
| Male | 199 (63.6) |
| Education | |
| Illiterate | 159 (50.8) |
| Under diploma | 95 (30.3) |
| Diploma | 34 (10.9) |
| Academic | 25 (8.0) |
| Marital status | |
| Married | 271 (86.6) |
| Divorced or widowed | 42 (13.4) |
| Occupation | |
| Employee | 21 (6.7) |
| Self-employed | 106 (33.9) |
| Homemakers | 109 (34.8) |
| Retired | 50 (16.0) |
| Unemployed | 27 (8.6) |
| Age (mean±SD) | 61.39±10.9 |
| Stress (mean±SD) | 158.3±111.5 |
| BMI (mean±SD) | 26.67±3.86 |

SD: Standard deviation, BMI: Body mass index

significant relations between some actual and perceived risk factors, such as hypertension ($P < 0.001$), familial history ($P = 0.011$), diabetes ($P < 0.001$), smoking ($P < 0.001$), and substance abuse ($P < 0.001$). Results in women showed statistically significant correlations between a number of actual and perceived risk factors, such as hypertension ($P = 0.019$), diabetes ($P = 0.011$), and smoking ($P < 0.001$). Finally, results in men showed statistically significant correlations between actual and perceived risk factors which include hypertension ($P = 0.022$), familial history ($P = 0.023$), diabetes ($P < 0.001$), smoking ($P < 0.001$), and substance abuse ($P = 0.002$). There were no significant relations between other actual and perceived risk factors ($P > 0.05$).

Table 3 shows the proportionality between actual and perceived risk factors. The results of this table indicate that there is no proportion between any of the actual and perceived risk factors. Results evidenced that patients generally have a poor perception about actual risk factors and they significantly underestimate their role ($P < 0.001$). Based on the results of this table, overweight and obesity, hypertension, stress, genetics and familial history of cardiac disease, hyperlipidemia, diabetes, smoking, aging, and substance abuse are the prevalent actual risk factors. However, the prevalent perceived risk factors were stress, smoking, hypertension, diabetes, genetic and familial history, obesity, hyperlipidemia, substance abuse, and aging. Regardless of the ranking of perceived risk factors, the results indicate that patients underestimated the role of aging (98.8%), substance abuse (95.2%), overweight and obesity (94.9%), hyperlipidemia (93.1%), familial history (90.3%), hypertension (90%) more than diabetes (86.1%), smoking (72.5%), and stress (54.7%).

Discussion

This study was carried out to assess the patients' causal beliefs, identification of underestimated risk factors, and indication of underestimation of every risk factor. One of the main aspects of CVDs is the patients' beliefs about the causes of the disease itself (causal reference or perceived risk factor). According to the patients' view, this risk factor is the main cause of illness.^[26] In accordance with a previous study,^[7] our findings showed that there are significant relations between hypertension, familial history, diabetes, smoking, and substance abuse with perceived risk factors. Indeed, Perkins-Porras *et al.* reported that 90% of patients with smoking and hypertension, 72% of patients with familial history, 85% of patients with overweight and obesity, and 49% of patients with low physical exercise consider these factors as causes of their disease.^[7]

In addition, according to other studies,^[20,21,27] patients seem to have a poor perception about their actual risk factors and they significantly underestimate their role. Jensen and Moser^[18] suggested that, compared to men,

Table 2: The correlation between the actual and perceived risk factors

| Actual/perceived risk factor | Total (n=313) | | Female (n=114) | | Male (n=199) | |
|------------------------------|---------------|-------|----------------|-------|--------------|-------|
| | r | P | r | P | r | P |
| Obesity | 0.091 | 0.106 | 0.087 | 0.357 | 0.098 | 0.168 |
| Hypertension | 0.220 | 0.001 | 0.219 | 0.019 | 0.163 | 0.022 |
| Stress | 0.067 | 0.235 | 0.093 | 0.323 | 0.051 | 0.473 |
| Family history | 0.144 | 0.011 | 0.122 | 0.197 | 0.162 | 0.023 |
| Hyperlipidemia | 0.079 | 0.162 | 0.151 | 0.108 | 0.051 | 0.477 |
| Diabetes | 0.313 | 0.001 | 0.238 | 0.011 | 0.376 | 0.001 |
| Smoking | 0.406 | 0.001 | 0.321 | 0.001 | 0.371 | 0.001 |
| Aging | 0.094 | 0.096 | - | - | 0.126 | 0.076 |
| Drug abuse | 0.204 | 0.001 | - | - | 0.218 | 0.002 |

Table 3: The proportionality between actual and perceived risk factors, as assessed through binomial test

| Risk factor | Actual risk | | Perceived risk | | Underestimation (%) | P |
|----------------|-------------|---------|----------------|---------|---------------------|-------|
| | n (%) | Ranking | n (%) | Ranking | | |
| Obesity | 215 (68.7) | 1 | 11 (3.5) | 6 | 94.9 | 0.001 |
| Hypertension | 170 (54.3) | 2 | 17 (5.4) | 3 | 90.0 | 0.001 |
| Stress | 148 (47.3) | 3 | 67 (21.4) | 1 | 54.7 | 0.001 |
| Family history | 124 (39.6) | 4 | 12 (3.8) | 5 | 90.3 | 0.001 |
| Hyperlipidemia | 102 (32.6) | 5 | 7 (2.2) | 7 | 93.1 | 0.001 |
| Diabetes | 101 (32.3) | 6 | 14 (4.5) | 4 | 86.1 | 0.001 |
| Smoking | 98 (31.3) | 7 | 27 (8.6) | 2 | 72.5 | 0.001 |
| Aging | 83 (26.5) | 8 | 1 (0.3) | 9 | 98.8 | 0.001 |
| Drug abuse | 42 (13.4) | 9 | 2 (0.6) | 8 | 95.2 | 0.001 |

women have more awareness of cardiac risk factors, but less than one-third of them can identify the actual risk factors. Age and socioeconomic demographics are the main factors related to underestimation of risk factors and risk of CVDs in general population.^[28] The role of fasting blood sugar, obesity, high cholesterol, and hypertension is underestimated by cardiac patients in the development of further outcome of disease.^[29-34] Based on another result, cardiac patients are aware of most of the risk factors as well, but they emphasize on one factor as the main cause of their disease severely. This is probably due to a sense of more control on disease.^[35]

According to the results, although patients have better perception about stress, smoking, and diabetes, the causal role of these risk factors is underestimated in more than 50% of cases whereas the factors of aging, substance abuse, obesity and overweight, hyperlipidemia, familial history, and hypertension in more than 90% of cases. It seems that the type of attitude about the cause of disease and perceived risk factors are influenced by cultural differences.^[26,36] Hence, each of the risk factors has a "special" ranking in different cultures. For example, one systematic review showed that patients know stress as a cause in 41% of cases and they refer to an unhealthy lifestyle only in 31% of cases.^[37] After stress as the main cause of CVDs,^[19,26] patients refer to the role of smoking, hypertension, misfortune, familial history, and physiologic risk factors such as diabetes as the major causes of CVDs.^[7,17,18,20,21]

Our findings indicated that patients underestimate the role of aging more than the other factors. Indeed, aging is a risk factor for atherosclerosis.^[38] In addition, aging as one nonspecific parameter and biomarker of CVDs has the greatest effect in the course of disease.^[2] In line with one study,^[34] it appears that obesity and hyperlipidemia are severely underestimated risk factors. Indeed, our results evidenced that overweight and obesity are in 11th rank of perceived risk factors (39.4% of estimation).^[7] Therefore, cardiac patients are aware of the role of inappropriate nutrition and lack of exercise as well, but they tend to (i) emphasize on the role of stress^[35] and (ii) underestimate overweight and obesity. Sanderson *et al.*^[39] reported that only 35% of individuals know familial history as a cause of CVDs, but hypertension, despite underestimation, has a better situation compared to the others.^[34] Hence, according to the role of correction of thoughts and patients' inefficient cognitions about risk factors in reducing anxiety and depression and medical costs and improving the quality of life, it is recommended that the correction of patients' wrong cognitions is concerned as a part of secondary prevention programs.^[2,13]

Strengths and limitations

The assessment of nine risk factors of CVDs which include stress, smoking, hypertension, diabetes, genetic and familial history, obesity, hyperlipidemia, substance abuse, and aging, over than the past studies,^[29-34] and applying the recommended short instrument for the assessment

of patients' perceived risk factors^[2,3,13,20,21] are the positive points of our study. On the other hand, illiterate cases (51%), sampling from one hospital in the west of Iran, and lack of assessment of the other risk factors such as unhealthy nutrition and physical activity are concerned as limitations.

Conclusion

Cardiac patients seem to underestimate the role of aging, substance abuse, obesity and overweight, hyperlipidemia, family history, and hypertension more than other actual risk factors. Therefore, these factors should be highlighted to patients to help them to (i) increase the awareness of actual risk factors and (ii) promote an appropriate lifestyle after CABG surgery.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

- Heydarpour B, Saeidi M, Ezzati P, Soroush A, Komasi S. Sociodemographic predictors in failure to complete outpatient cardiac rehabilitation. *Ann Rehabil Med* 2015;39:863-71.
- Komasi S, Saeidi M. Aging is an important cause for a lack of understanding of the main risk factor in cardiac rehabilitation patients. *Thrita* 2015;4:e32751.
- Komasi S, Saeidi M. Screening for depressive symptoms at the beginning of outpatient. Cardiac rehabilitation by assessed perceived risk factors by patients. *Clin Med Rev Case Rep* 2015;2:1-5.
- Compare A, Mommersteeg PM, Faletra F, Grossi E, Pasotti E, Moccetti T, *et al.* Personality traits, cardiac risk factors, and their association with presence and severity of coronary artery plaque in people with no history of cardiovascular disease. *J Cardiovasc Med (Hagerstown)* 2014;15:423-30.
- Compare A, Zarbo C, Shonin E, Van Gordon W, Marconi C. Emotional regulation and depression: A potential mediator between heart and mind. *Cardiovasc Psychiatry Neurol* 2014;2014:324374.
- Karami J, Komasi S, Maesoomi M, Saeidi M. Comparing the effects of two methods of relaxation and interpersonal cognitive problem solving (ICPS) on decreasing anxiety and depression in cardiac rehabilitation patients. *Urmia Med J* 2014;25:298-308.
- Perkins-Porras L, Whitehead DL, Steptoe A. Patients' beliefs about the causes of heart disease: Relationships with risk factors, sex and socio-economic status. *Eur J Cardiovasc Prev Rehabil* 2006;13:724-30.
- van der Wal MH, Jaarsma T, Moser DK, Veeger NJ, van Gilst WH, van Veldhuisen DJ. Compliance in heart failure patients: The importance of knowledge and beliefs. *Eur Heart J* 2006;27:434-40.
- Phillips LA, Diefenbach MA, Abrams J, Horowitz CR. Stroke and TIA survivors' cognitive beliefs and affective responses regarding treatment and future stroke risk differentially predict medication adherence and categorised stroke risk. *Psychol Health* 2015;30:218-32.
- Byrne M, Walsh J, Murphy AW. Secondary prevention of coronary heart disease: Patient beliefs and health-related behaviour. *J Psychosom Res* 2005;58:403-15.
- Parati G, Omboni S, Compare A, Grossi E, Callus E, Venco A, *et al.* Blood pressure control and treatment adherence in hypertensive patients with metabolic syndrome: Protocol of a randomized controlled study based on home blood pressure telemonitoring vs. conventional management and assessment of psychological determinants of adherence (TELEBPMET Study). *Trials* 2013;14:22.
- Murphy B, Worcester M, Higgins R, Le Grande M, Larritt P, Goble A. Causal attributions for coronary heart disease among female cardiac patients. *J Cardiopulm Rehabil* 2005;25:135-43.
- Saeidi M, Komasi S, Heydarpour B, Momeni K, Zakiei A. Those who perceive their disease as a physiological or psychological risk factor experience more anxiety at the beginning of the cardiac rehabilitation program. *Res Cardiovasc Med* 2016;5:e29291.
- Saeidi M, Komasi S, Heydarpour B, Karim H, Nalini M, Ezzati P. Predictors of clinical anxiety aggravation at the end of a cardiac rehabilitation program. *Res Cardiovasc Med* 2015;5:e30091.
- Furze G, Lewin RJ, Murberg T, Bull P, Thompson DR. Does it matter what patients think? The relationship between changes in patients' beliefs about angina and their psychological and functional status. *J Psychosom Res* 2005;59:323-9.
- Abed MA, Khalil AA, Moser DK. Awareness of modifiable acute myocardial infarction risk factors has little impact on risk perception for heart attack among vulnerable patients. *Heart Lung* 2015;44:183-8.
- Panagiotakos DB, Georgousopoulou EN, Polychronopoulos E, Vassilakou T, Chrysohoou C, Pitsavos C, *et al.* Beliefs and attitudes regarding cardiovascular disease risk factors: A health survey in 10,141 Greek men and women (2006-2012). *Int J Cardiol* 2013;168:4847-9.
- Jensen LA, Moser DK. Gender differences in knowledge, attitudes, and beliefs about heart disease. *Nurs Clin North Am* 2008;43:77-104; vi-vii.
- Kelly-Irving M, Mulot S, Inamo J, Ruidavets JB, Atallah A, Lang T. Improving stroke prevention in the French West Indies: Limits to lay knowledge of risk factors. *Stroke* 2010;41:2637-44.
- Saeidi M, Komasi S, Soroush A, Zakiei A, Shakeri J. Gender differences in patients' beliefs about biological, environmental, behavioral, and psychological risk factors in a cardiac rehabilitation program. *J Cardiothorac Med* 2014;2:215-20.
- Saeidi M, Soroush A, Komasi S, Moemeni K, Heydarpour B. Attitudes toward cardiovascular disease risk factors among patients referred to a cardiac rehabilitation center: Importance of psychological attitudes. *Shiraz Emed J* 2015;16:e22281.
- Holmes TH, Rahe RH. The social readjustment rating scale. *J Psychosom Res* 1967;11:213-8.
- Komasi S, Saeidi M, Montazeri N, Masoumi M, Soroush A, Ezzati P. Which factors unexpectedly increase depressive symptom severity in patients at the end of a cardiac rehabilitation program? *Ann Rehabil Med* 2015;39:872-9.
- Vafaii B. Evaluation of the relation between life stresses and blood neoplastic diseases in males and females aged between 30 and 50. *Tabriz Med J* 2000;34:47-54.
- Pallant J. *SPSS Survival Manual: A Step by Step Guide to Data Analysis using SPSS for Windows*. 2nd ed., Ver. 12. Australia: Allen & Unwin; 2005.
- Darr A, Astin F, Atkin K. Causal attributions, lifestyle change, and coronary heart disease: Illness beliefs of patients of South Asian and European origin living in the United Kingdom. *Heart Lung* 2008;37:91-104.
- Patel M, Phillips-Caesar E, Boutin-Foster C. Attitudes and beliefs regarding cardiovascular risk factors among Bangladeshi immigrants in the US. *J Immigr Minor Health* 2014;16:994-1000.
- Oertelt-Prigione S, Seeland U, Kendel F, Rucke M, Flöel A, Gaismaier W, *et al.* Cardiovascular risk factor distribution and subjective risk estimation in urban women – The BEFRI study: A randomized cross-sectional study. *BMC Med* 2015;13:52.
- Bjørnholt JV, Erikssen G, Aaser E, Sandvik L, Nitter-Hauge S, Jervell J, *et al.* Fasting blood glucose: An underestimated risk factor for cardiovascular death. Results from a 22-year follow-up of healthy

- nondiabetic men. *Diabetes Care* 1999;22:45-9.
30. Huang PY, Buring JE, Ridker PM, Glynn RJ. Awareness, accuracy, and predictive validity of self-reported cholesterol in women. *J Gen Intern Med* 2007;22:606-13.
 31. Stommel M, Schoenborn CA. Accuracy and usefulness of BMI measures based on self-reported weight and height: Findings from the NHANES and NHIS 2001-2006. *BMC Public Health* 2009;9:421.
 32. Schenker N, Raghunathan TE, Bondarenko I. Improving on analyses of self-reported data in a large-scale health survey by using information from an examination-based survey. *Stat Med* 2010;29:533-45.
 33. Taylor A, Dal Grande E, Gill T, Pickering S, Grant J, Adams R, *et al.* Comparing self-reported and measured high blood pressure and high cholesterol status using data from a large representative cohort study. *Aust N Z J Public Health* 2010;34:394-400.
 34. Van Eenwyk J, Bensley L, Ossiander EM, Krueger K. Comparison of examination-based and self-reported risk factors for cardiovascular disease, Washington state, 2006-2007. *Prev Chronic Dis* 2012;9:E117.
 35. French D, Maissi E, Marteau TM. The purpose of attributing cause: Beliefs about the causes of myocardial infarction. *Soc Sci Med* 2005;60:1411-21.
 36. Farooqi A, Nagra D, Edgar T, Khunti K. Attitudes to lifestyle risk factors for coronary heart disease amongst South Asians in Leicester: A focus group study. *Fam Pract* 2000;17:293-7.
 37. Cherrington CC, Moser DK, Lennie TA, Kennedy CW. Illness representation after acute myocardial infarction: Impact on in-hospital recovery. *Am J Crit Care* 2004;13:136-45.
 38. Tabei SM, Senemar S, Saffari B, Ahmadi Z, Haqparast S. Non-modifiable factors of coronary artery stenosis in late onset patients with coronary artery disease in Southern Iranian population. *J Cardiovasc Thorac Res* 2014;6:51-5.
 39. Sanderson SC, Waller J, Humphries SE, Wardle J. Public awareness of genetic influence on chronic disease risk: Are genetic and lifestyle causal beliefs compatible? *Public Health Genomics* 2011;14:290-7.