



Corrigendum: The Influence of Radio-Frequency Transmit Field Inhomogeneities on the Accuracy of G-ratio Weighted Imaging

Tim M. Emmenegger^{1,2}, Gergely David^{1,2}, Mohammad Ashtarayeh², Francisco J. Fritz², Isabel Ellerbrock^{2,3}, Gunther Helms⁴, Evelyne Balteau⁵, Patrick Freund^{1,6,7} and Siawoosh Mohammadi^{2,6*}

¹ Spinal Cord Injury Center Balgrist, University Hospital Zurich, University of Zurich, Zurich, Switzerland, ² Department of Systems Neuroscience, University Medical Center Hamburg-Eppendorf, Hamburg, Germany, ³ Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden, ⁴ Medical Radiation Physics, Clinical Sciences Lund (IKVL), Lund University, Lund, Sweden, ⁵ GIGA Institute, University of Liège, Liège, Belgium, ⁶ Department of Neurophysics, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany, ⁷ Wellcome Trust Centre for Neuroimaging, University College London, London, United Kingdom

Keywords: myelin volume fraction, axon volume fraction, radio-frequency transmit field inhomogeneities, B₁+ correction, multi-parameter mapping, diffusion MRI, magnetization transfer saturation, MR g-ratio

OPEN ACCESS

Edited and reviewed by:

Tim B. Dyrby, Technical University of Denmark, Denmark

*Correspondence:

Siawoosh Mohammadi s.mohammadi@uke.de

Specialty section:

This article was submitted to Brain Imaging Methods, a section of the journal Frontiers in Neuroscience

Received: 08 September 2021 Accepted: 20 September 2021 Published: 07 October 2021

Citation:

Emmenegger TM, David G,
Ashtarayeh M, Fritz FJ, Ellerbrock I,
Helms G, Balteau E, Freund P and
Mohammadi S (2021) Corrigendum:
The Influence of Radio-Frequency
Transmit Field Inhomogeneities on the
Accuracy of G-ratio Weighted
Imaging. Front. Neurosci. 15:772745.
doi: 10.3389/fnins.2021.772745

A Corrigendum on

The Influence of Radio-Frequency Transmit Field Inhomogeneities on the Accuracy of G-ratio Weighted Imaging

by Emmenegger, T. M., David, G., Ashtarayeh, M., Fritz, F. J., Ellerbrock, I., Helms, G., Balteau, E., Freund, P., and Mohammadi, S. (2021). Front. Neurosci. 15:674719. doi: 10.3389/fnins.2021.674719

In the original article, there was a mistake in **Figure 5** as published. The MVF map was shown twice in the original figure (middle and bottom rows), instead the AVF map should have been depicted in the middle row. The corrected **Figure 5** that appears below.

In the published article, there was an error regarding the affiliation for Isabel Ellerbrock. As well as having affiliation 3, she should also have 2. There was an error regarding the affiliation for Gergely David. As well as having affiliation 1, he should also have 2.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Emmenegger, David, Ashtarayeh, Fritz, Ellerbrock, Helms, Balteau, Freund and Mohammadi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

1




FIGURE 5 | Voxel-wise maps of group-averaged g^{B1}_{MR}, AVF^{B1}_{MR}, and MVF^{B1}_{MR}, restricted to the group WM mask (cf. section "Definition of White Matter Masks"). Depicted are a single sagittal (x = 100), coronal (y = 91), and axial (z = 85) slice.