

Impaired polymorphonuclear neutrophils in the oral cavity of edentulous individuals

Patrick Rijkschroeff , Bruno G. Loos, Elena A. Nicu

Department of Periodontology, Academic Centre for Dentistry Amsterdam, Amsterdam, the Netherlands

Rijkschroeff P, Loos BG, Nicu EA. Impaired polymorphonuclear neutrophils in the oral cavity of edentulous individuals.

Eur J Oral Sci 2017; 125: 371–378. © 2017 The Authors. *Eur J Oral Sci* published by John Wiley & Sons Ltd

Oral health is characterized by functional oral polymorphonuclear neutrophils (oPMNs). Edentulism might be associated with a loss of oPMNs because these cells enter the oral cavity primarily through the gingival crevices. The main aim of this study was to investigate the numbers of oPMNs in rinse samples obtained from edentulous ($n = 21$) and dentate ($n = 20$) subjects. A second study aim was to investigate possible differences between oPMNs and peripheral blood polymorphonuclear neutrophils (cPMNs). Apoptosis/necrosis and cell-activation markers (CD11b, CD63 and CD66b) were analyzed using flow cytometry. Reactive oxygen species (ROS) production was determined either without stimulation (constitutive) or in response to 10 μ M phorbol myristate acetate or *Fusobacterium nucleatum*. The edentulous subjects presented with lower oPMN counts and higher percentages of apoptotic/necrotic oPMNs compared with dentate subjects. Furthermore, oPMNs from edentulous donors expressed low levels of all three activation markers and low constitutive ROS. In contrast, oPMNs from dentate subjects expressed high levels of all three activation markers and a higher level of constitutive ROS than cPMNs. When challenged, oPMNs from edentulous subjects showed no upregulation in ROS production, whereas oPMNs from dentate subjects retained their ability to respond to stimulation. The functional characteristics of cPMNs were comparable between edentulous and dentate subjects. This study demonstrates that despite having functional cPMNs, edentulous subjects have low oPMN numbers that are functionally impaired.

Elena A. Nicu, Department of Periodontology, Academic Centre for Dentistry Amsterdam, Gustav Mahlerlaan 3004, 1081 LA, Amsterdam, the Netherlands

E-mail: e.nicu@acta.nl

Key words: apoptosis; granulocytes; innate immunity; oral health; reactive oxygen species

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Accepted for publication June 2017

Over the past century, the number of elderly people has significantly increased (1). Edentulism is prevalent among elderly people worldwide and is highly associated with socio-economic status (1–3). The World Health Organization reported that within the European region, over 30% of people 65–74 yr of age have lost all their natural teeth (4). Denture use is common among the elderly in the Netherlands. In 2009, approximately 41% of the Dutch population over 65 yr of age wore full dentures and 34% wore partial dentures, according to Statistics Netherlands (CBS) (5). Several studies reported that the patients' quality of life decreases with tooth loss. Edentulous subjects wearing full dentures experience more psychological discomfort and social disability compared with patients with partial dentures or with dentate subjects, in addition to oral discomfort, such as functional limitations and physical pain (6–8).

Currently, oral health is becoming increasingly recognized as imperative for maintaining systemic health (9, 10). Innate immune responses play a crucial role in the equilibrium within the oral cavity (11, 12). Important innate immune cells are the polymorphonuclear

neutrophils (PMNs). They are the predominant leukocyte effector cells in the blood circulation, which can migrate to tissues in response to chemoattractant molecules such as lipopolysaccharide or lipoteichoic acid (released by microorganisms) or to products released from the host cells following tissue damage (13).

In a recent study, oral PMNs (oPMNs) were investigated in a group of orally and systemically healthy dentate individuals (14). Viable PMNs were found to be capable of performing innate immune functions, such as active degranulation and production of reactive oxygen species (ROS). A few early studies investigated leukocyte counts in the oral cavity of edentulous subjects; however, limited knowledge exists concerning PMN functionality in the edentulous oral cavity (15–17).

The first aim of the current study was to investigate the presence of oPMNs in edentulous subjects and to determine their functionality. To this end, we analyzed the numbers, membrane integrity, cell-activation status, and reactive oxygen species (ROS) production capabilities of the oPMNs from a group of edentulous subjects and a comparable group of dentate subjects. The second aim was to investigate possible differences between

the oPMNs and the PMNs from peripheral blood [i.e. the circulatory system PMNs (cPMNs)] in both groups.

Material and methods

Study population

For this case-control study, volunteers were recruited from the Academic Centre for Dentistry Amsterdam (ACTA), the Netherlands. Participants were either patients at the dental clinics or non-professional personnel. All subjects presented without any history of pathological conditions related to altered PMN levels or function (such as hematological disorders, diabetes mellitus, antibiotic use within the last 6 months, recent history of illness or fever, allergies, alcoholism, and pregnancy). Furthermore, individuals with removable partial dentures, night guards, orthodontic banding, (peri)oral piercings, or apparent oral lesions were also excluded. The study was conducted in accordance with the Declaration of Helsinki (2008) of the World Medical Association and approximating Good Clinical Practice guidelines. Subjects were informed about the purpose of this study, received written information, and had given written consent before the start of the study. This study was approved by the Medical Ethics Committee of the VU University Medical Center (reference 2012-210#B2012406) and was registered at the ISRCTN registry (ISRCTN 13234871).

Edentulous subjects could be included if they had been wearing a full denture for a minimum of 1 yr. Dentate subjects were screened for their periodontal condition using the Dutch Periodontal Screening Index (18, 19). The latter subjects were included if they had a minimum of 24 teeth and only if they showed a maximum probing depth of 4–5 mm in the absence of gingival recession. Additionally, bitewing radiographs were taken in order to confirm the absence of alveolar bone loss.

Clinical collection procedure for oPMNs and cPMNs

The oPMNs and cPMNs were isolated and analyzed on the same day. The collection procedure was based on a previously described protocol (14). Briefly, in order to avoid contamination, oral rinse samples were taken before any other intra-oral procedures. Edentulous subjects were asked to remove their dentures before supervised sample collection. All subjects were instructed not to gargle or to clear their throat during the sampling procedure. The oPMNs were obtained by four serial rinses of the oral cavity with 10 ml of sterile sodium chloride solution (0.9% NaCl) for 30 s, with an intermission of 4 min 30 s. The pooled sample was kept on ice in a 50-ml centrifuge tube (Sigma-Aldrich Chemie, Zwijndrecht, the Netherlands) until the end of the collection procedure. Venous blood samples were drawn from the antecubital fossa of all subjects into 9-ml sodium heparin blood-collection tubes (BD Vacutainer, Breda, the Netherlands) and were maintained at room temperature until processed using the PMN isolation procedure. The subsequent isolation procedures of the oPMNs and the cPMNs were carried out as previously described (14).

Cell counts

The oPMN and the cPMN counts were made using a Muse Cell Analyzer (Merck Millipore, Darmstadt,

Germany) with the Muse Count & Viability Kit (Merck Millipore), according to the manufacturer's instructions.

Membrane integrity assessment

Morphological changes associated with apoptosis are a late manifestation of the programmed cell death pathway. During cell death, PMNs lose their membrane integrity. Propidium iodide (PI) is a fluorescent dye that can diffuse into cells and bind to DNA when membrane integrity is affected. Propidium iodide-positive (PI⁺) cells are regarded as late apoptotic or necrotic (20, 21). In brief, oPMNs and cPMNs were incubated for 15 min in the dark at room temperature according to the manufacturer's instructions (BD Pharmingen FITC Annexin V Apoptosis Detection Kit; BD Biosciences, San Diego, CA, USA). Flow cytometric analysis was performed within 1 h using a BD Accuri C6 flow cytometer (BD Biosciences). The percentages of PI⁺ PMNs were calculated.

Cell activation status

Low-affinity immunoglobulin-Fc γ receptor IIIb (CD16b) can be found on the surface membrane of mature PMNs and can be used as a PMN identification marker. Expression of receptors on the surface of PMNs was measured by flow cytometry. The PMNs were gated according to CD16 expression and the sideward scatter profile. To study PMN activation, samples were measured for the expression of phycoerythrin (PE)-conjugated monoclonal antibodies (mAbs) (at a final concentration of 10 $\mu\text{g ml}^{-1}$) against CD11b and CD63 and for expression of fluorescein isothiocyanate (FITC)-conjugated mAb against CD66b (all from BD Biosciences). Samples were incubated for 30 min on ice in the dark. Expression of cell-surface markers was measured on the FL-1 (Green) and FL-2 (Red) channels and calculated as mean fluorescence intensity (MFI). Results were corrected for non-specific binding, by subtracting the MFI values of the corresponding IgG1 isotype-control antibodies (BD Biosciences).

Bacterial culture

Fusobacterium nucleatum strain ATCC10953 was obtained commercially (DSMZ, Braunschweig, Germany) and grown anaerobically (80% nitrogen, 10% carbon dioxide, and 10% hydrogen) in a brain-heart infusion broth supplemented with 5 $\mu\text{g ml}^{-1}$ of hemin (Sigma-Aldrich) and 1 $\mu\text{g ml}^{-1}$ of menadione (Sigma-Aldrich). The bacteria were isolated from the broth cultures by centrifugation, washed twice in sterile PBS, then diluted with sterile PBS, to yield a final suspension of 4×10^8 cells ml^{-1} , which was stored at -20°C .

ROS production assay

Dihydrorhodamine 123 (DHR) (Sigma-Aldrich) is a non-fluorescent probe that can diffuse passively across cell membranes; DHR can be oxidized by hydrogen peroxide and superoxide, and broken down into the green fluorescent rhodamine 123, which is detectable in the FL-1 (Green) channel of a flow cytometer (22, 23). Unstimulated samples were incubated for 30 min with 2 mM DHR, in a shaking (50 r.p.m.) water bath at 37°C .

Additionally, PMNs were stimulated with 10 μM phorbol myristate acetate (PMA; Sigma-Aldrich) or non-opsonized *F. nucleatum* at a ratio of PMN:Fn of 1:20. Results were expressed as the MFI.

Statistical analysis

Data analyses were performed using IBM SPSS 21.0 software (IBM, Chicago, IL, USA). Boxplots with Tukey whiskers were generated using Prism version 5.02 (Graphpad Software, La Jolla, CA, USA). Medians and ranges were calculated. The distribution of our data was non-normal. For this reason, comparisons of continuous variables between edentulous and dentate subjects were analyzed using the Mann–Whitney *U*-test and comparisons of categorical data were analyzed using the chi-square test. Within groups (edentulous or dentate), comparisons between unstimulated and stimulated samples (*F. nucleatum* or PMA), and oPMNs vs. cPMNs, were performed using the Wilcoxon signed-rank test. Values of *P* < 0.05 were considered statistically significant.

Results

Characteristics of subjects

The study population consisted of 21 edentulous participants and 20 dentate participants (Table 1) with a median age of 66 yr for the edentulous group and 63 yr for the dentate group (*P* = 0.043). The majority of participants were male (13 were edentulous and 15 were dentate, *P* = 0.505). Among the edentulous participants, four were current smokers, while one participant from the dentate group was a current smoker

(*P* = 0.343). Twelve edentulous individuals reported medication use compared with two dentate participants (*P* = 0.003).

PMN counts

The results of the oPMN and the cPMN counts are shown in Table 1. After four serial rinses, the median number of oPMNs isolated from edentulous participants (0.3×10^6 oPMNs) was lower than the median number isolated from dentate participants (0.5×10^6 oPMNs) (*P* = 0.030). The yield of cPMNs from a 9-ml tube of blood was higher for edentulous participants (median: 13.0×10^6 cPMNs) than for dentate participants (median: 9.7×10^6 cPMNs) (*P* = 0.022). One blood sample from an edentulous subject was discarded because of technical failure.

Assessment of membrane integrity

The oPMNs and the results of the flow cytometric membrane integrity assessment are presented in Fig. 1 and Table S1. A higher percentage of late apoptotic/necrotic oPMNs was found for edentulous participants (median: 64.5% oPMNs) than for dentate participants (median: 43.1% oPMNs) (*P* = 0.002) (Fig. 1C). In contrast, uptake of PI by cPMNs was low in all samples, and did not differ significantly between edentulous and dentate participants (*P* = 0.301) (Fig. 1D). Consequently, the percentage of late apoptotic/necrotic cells was higher among oPMNs than among cPMNs for both the edentulous and the dentate participants (both *P* < 0.001).

Table 1

Summary of participants' background characteristics, dental features and counts of oral polymorphonuclear neutrophils (oPMNs) and circulatory system polymorphonuclear neutrophils (cPMNs)

Variable	Edentulous subjects (<i>n</i> = 21)	Dentate subjects (<i>n</i> = 20)	<i>P</i> -value*
Age (yr)	66 (44–82)	63 (49–77)	0.043
Males	13 (62)	15 (75)	0.505
Smokers	4 (19)	1 (5)	0.343
Medication users	12 (57) [†]	2 (10) [‡]	0.003
Last tooth extracted			
<5 yr ago	1 (5)	NA	
>5 yr ago	20 (95)	NA	
Last modification denture			
<5 yr ago	11 (52)	NA	
>5 yr ago	10 (48)	NA	
Oral discomfort [§]	2 (10)	NA	
Counts			
oPMNs [¶]	0.3×10^6 (0.1–1.0)	0.5×10^6 (0.1–4.5)	0.030
cPMNs ^{**}	13.0×10^6 (3.7–33.4)	9.7×10^6 (0.7–29.4)	0.022

Values represent median (range) or *n* (%). NA, not applicable.

*Continuous variables were analyzed using the Mann–Whitney *U*-test and categorical data were analyzed using the χ^2 test. Values of *P* < 0.05 were considered statistically significant and are highlighted in bold.

[†]Self-reported medication use: anti-arrhythmic drug (*n* = 1), antidepressant (*n* = 1), beta blocker (*n* = 1), calcium-channel blocker (*n* = 1), diuretics (*n* = 3), gastric acid inhibitor/antacid (*n* = 5), cholesterol-lowering drugs (*n* = 5), calcium supplements (*n* = 1).

[‡]Self-reported medication use: anti-hypertensive drugs (*n* = 1), cholesterol-lowering drugs (*n* = 2).

[§]Self-reported denture-related discomfort; mouth dryness (*n* = 2).

[¶]Total number of oPMNs collected after 4 × 30 s rinsing with 4 min 30 s intermission, after the isolation and purification steps.

**Total cPMNs obtained from a 9-ml tube of blood, after the isolation and purification steps.




Fig. 1. Membrane integrity assessment. Polymorphonuclear neutrophils (PMNs) isolated from the oral cavity of dentate (A) and edentulous (B) participants, and the results of the flow-cytometric analysis of propidium iodide (PI) uptake for oral PMNs (oPMNs) (C) and circulatory system PMNs (cPMNs) (D). Fewer oPMNs were present in the samples originating from the edentulous participants. Black arrows indicate oPMNs, white arrows represent corresponding propidium iodide-positive (PI⁺) cells, indicating late apoptotic/necrotic cells. Boxplots show medians, interquartile ranges, and Tukey whiskers. * $P = 0.002$ (Mann-Whitney U -test). Scale bar = 50 μm .

Cell activation status

CD16⁺ cells, indicating mature PMNs, were analyzed for the expression of cell-activation markers CD11b, CD63, and CD66b using flow cytometry. The results of the immunohistochemistry analyses are presented in Fig. 2 and Table S1. In dentate participants, the median MFI of oPMNs was 9.4×10^4 for CD11b, 4.1×10^4 for CD63, and 15.6×10^4 for CD66b. A different pattern of expression was observed in edentulous participants, with all activation markers being present at very low levels on oPMNs, namely median MFI of 0.8×10^4 for CD11b, 0.5×10^4 for CD63, and 1.4×10^4 for CD66b. Seven of 21 of the edentulous individuals showed no expression of the activation markers on oPMNs (i.e. the binding of the specific mAbs against CD11b, CD63, and CD66b was equal to or lower than the background binding of the isotype-control antibodies). Consequently, expression of the CD markers was lower on oPMNs from edentulous participants than on oPMNs from dentate participants (all $P < 0.001$) (Fig. 2A). All three markers were expressed at low levels on cPMNs, with no statistical differences between edentulous and dentate participants (Fig. 2B). In dentate participants, oPMNs were activated to a higher degree compared with cPMNs (CD11b and CD63, both $P < 0.001$; CD66b, $P = 0.001$) (Table S1). In contrast, in edentulous participants, the expression of CD11b and CD66b did not differ between oPMNs and cPMNs, whereas CD63 was expressed at a higher level on oPMNs than on cPMNs ($P = 0.011$) (Table S1).

ROS production assay

The results for the expression of the ROS production are shown in Fig. 3 and Table S1. The unstimulated oPMNs produced comparable median levels of ROS in the edentulous participants (median MFI: 32.0×10^4) and the dentate participants (median MFI: 54.9×10^4) (Fig. 3A). In the dentate group, the oPMNs were capable of responding to both PMA (median MFI: 179.9×10^4) and *F. nucleatum* (median MFI: 122.1×10^4), which resulted in higher levels of ROS compared with unstimulated oPMNs (PMA and *F. nucleatum* both $P = 0.001$) (Fig. 3A). In contrast, no upregulation was observed in oPMNs from edentulous participants, in response to either PMA (median MFI: 16.2×10^4) or *F. nucleatum* (median MFI: 28.3×10^4) (Fig. 3A). As a result, the stimulated ROS levels were lower for oPMNs from edentulous participants compared with the levels of ROS produced by the oPMNs from dentate participants (PMA, $P < 0.001$; *F. nucleatum*, $P = 0.002$) (Fig. 3A).

The unstimulated cPMNs showed low levels of ROS expression, with no differences between the edentulous participants (median MFI: 7.7×10^4) and the dentate participants (median MFI: 11.4×10^4) (Fig. 3B). Stimulation with PMA and *F. nucleatum* induced robust upregulation compared with unstimulated cPMNs in both groups (PMA and *F. nucleatum*, both $P < 0.001$) (Fig. 3B and Table S1). The levels of ROS produced by cPMNs in response to stimulation were comparable in the edentulous (PMA median MFI: 4.4×10^6 ;




Fig. 2. Cell activation status. Immunohistochemistry of oral polymorphonuclear neutrophil (oPMN) and circulatory system polymorphonuclear neutrophil (cPMN) populations. Polymorphonuclear neutrophils (PMNs) were gated based on CD16 expression and sideward scatter. Representative examples show a distinct subset of CD16⁺ cells, indicating mature oPMNs (A) and cPMNs (B). Flow-cytometric analysis of the cell-activation markers CD11b, CD63, and CD66b on oPMNs (C) and cPMNs (D) are presented as mean fluorescence intensity (MFI). For each CD marker, corresponding isotype-control antibodies were used to determine autofluorescent signals, which were subtracted from the MFI. Boxplots show medians, interquartile ranges, and Tukey whiskers, and black dots represent outliers. All **P* < 0.05 (Mann–Whitney *U*-test).

F. nucleatum median MFI: 5.7×10^6) and the dentate participants (PMA median MFI: 5.2×10^6 , *P* = 0.277; *F. nucleatum* median MFI: 4.9×10^6 , *P* = 0.988).

The unstimulated oPMNs expressed higher levels of ROS compared with the cPMNs, both in the edentulous group (*P* = 0.001) and the dentate group (Table S1, *P* = 0.007). However, the ROS levels observed upon stimulation were lower in oPMNs than in cPMNs (PMA and *F. nucleatum*, both *P* < 0.001) (Table S1).

Discussion

The edentulous participants had fewer oPMNs compared with the dentate participants. The oPMNs from the edentulous participants presented with compromised

membranes and no functional respiratory burst capabilities, whereas the dentate participants presented with functional oPMNs. Of importance was that the functional status of the cPMNs from both the edentulous and dentate individuals was comparable. Our findings suggest that oPMNs are functionally impaired in edentulous participants.

One plausible explanation for the low numbers and altered phenotype of the oPMNs from the edentulous group comes from an anatomical perspective. The site of origin of the oPMN cannot be identical in dentate and edentulous participants. The gingival crevice has long been proposed and accepted as the main port of entry for oPMNs into the oral cavity. In dentate participants, substantial numbers of PMNs ($\pm 3.0 \times 10^4$ PMNs min⁻¹) have been reported to migrate constantly from the blood circulation through crevices around the




Fig. 3. Reactive oxygen species (ROS) production. Results for the flow-cytometric analysis of ROS production from unstimulated and stimulated oral polymorphonuclear neutrophils (oPMNs) (A) and circulatory system polymorphonuclear neutrophils (cPMNs) (B). Stimulated polymorphonuclear neutrophils (PMNs) were incubated with either phorbol myristate acetate (PMA) or *Fusobacterium nucleatum* (*F.n.*). Boxplots show medians, interquartile ranges, and Tukey whiskers, and black dots represent outliers. * $P < 0.05$ (Mann–Whitney U -test), # $P < 0.05$ (Wilcoxon signed-rank test) representing comparison with the corresponding unstimulated samples.

teeth (24–28). This migration is facilitated by the presence of the lymphatic vessels and the structure of the junctional epithelium, whose number of epithelial layers decreases in the apical part, together with fewer tight junctions observed between the cells (29–31). In response to oral microorganisms, proinflammatory molecules are generated within the gingival tissues, leading to increased chemotaxis for PMNs (32, 33). In edentulous participants, both the gingival crevices and the junctional epithelia are lost, which affects the access of PMNs to the oral cavity. A second explanation for the low numbers and the altered oPMN phenotype is that in general, denture wearers harbor a less diverse microbiome and the large majority (i.e. >70%) is heavily colonized with *Candida* species (34, 35). Polymorphonuclear neutrophils have been shown to exhibit diminished defensive activity against *Candida* biofilms, including poor recruitment, impaired ROS production and reduced killing (36, 37). In the current study, edentulous participants also presented with fewer numbers of oPMNs and diminished functionality compared with dentate participants.

It can be discussed whether oPMNs can be found at all in the oral cavity of edentulous participants. However, despite the absence of teeth and gingival crevices, PMNs probably have a surveillance function in the edentulous oral cavity similar to that described for the nasal, lung, or vaginal mucosae, or the intestinal lumen (13, 38–41). A previous study characterized the immune cells present at various oral sites in healthy participants. These authors demonstrated the presence of migrating PMNs in biopsies from buccal mucosa (42). Another study demonstrated the possibility of active transepithelial migration of PMNs in an in vitro model of the multilayered oral epithelium (43). Combined, these observations strongly suggest that oPMNs serve a surveillance function also in the edentulous oral mucosa. On the other hand, the results in the current study show that the oPMNs in edentulous participants have very limited activation potential, as the levels of primary and secondary granules (CD63 and CD66b) were low and no respiratory burst could be further induced. We speculate that the oPMNs emerging at the interface between the epithelial barrier and the mucosal

biofilm have fulfilled their patrolling function, are exhausted or necrotic, and then are shed and consequently appear in the sample rinse fluid. Interestingly, in the present study, the numbers of oPMNs observed in edentulous participants were higher than reported previously (15–17) and may be related to the different sample collection method. In the present study, all participants were asked to rinse vigorously with a saline solution, in order to dislodge the PMNs attached to the mucosa throughout the oral cavity, whereas various reports isolated oPMNs from unstimulated saliva or by drooling (15–17). Also, in the present study a calibrated cell-counting device was used instead of visual identification with a light microscope. The cell-counting device is not prone to human error and is capable of identifying PMNs in low-abundance cell suspensions (44). Nevertheless, to end the discussion on the existence of oPMN in edentulous participants, we cannot exclude the possibility that non-visible transient wounds are an additional source of PMNs in the oral cavity, even though the absence of intra-oral wounds was confirmed in this study by careful clinical examination.

Analysis of cPMNs was included to account for possible differences already occurring in the blood circulation. Importantly, our data showed comparable cPMNs parameters amongst the participants; only the cPMN numbers were higher in the edentulous participants. We cannot exclude a possible effect of the general health status on the cPMN counts (45). As tooth loss has been proposed as a marker for systemic changes associated with a higher risk for developing chronic disease, including changes in lifestyle and socio-economic factors, infection, and inflammation (46–51), it is not surprising that smoking and the use of medication were reported more frequently in edentulous participants.

Altogether, edentulous participants display impaired oral PMNs, despite having functional PMNs in the blood circulation. These oPMNs are present in lower numbers and for the most part are non-functional compared with those from dentate participants. We suggest that this severely hampered aspect of the ‘normal’ innate immunity in the oral cavity of edentulous participants can make them less resistant to perturbations and more prone to dysbiotic conditions, such as oral

mucositis, *Candida* infections, and even oral cancer. These findings underline the importance of the presence and maintenance of the natural teeth where the gingival crevices act as a gateway for preserving the innate defense mechanisms within the oral cavity.

Acknowledgements – The authors wish to thank Britt van den Berg, Jeanine Busropan, Koen Hermes, and Madeline Kosho for their contributions in patient recruitment and collection of the samples.

Conflicts of interest – The authors declare that they have no conflict of interest. The Department of Periodontology is supported by a grant from the University of Amsterdam for research into the focal point ‘Oral Infections and Inflammation’.

References

- PETERSEN PE. The World Oral Health Report 2003: continuous improvement of oral health in the 21st century—the approach of the WHO Global Oral Health Programme. *Community Dent Oral Epidemiol* 2003; **31**: 3–23.
- SALOMON JA, WANG H, FREEMAN MK. Healthy life expectancy for 187 countries, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2013; **380**: 2144–2162.
- McMICHAEL AJ, McKEE M, SHKOLNIKOV V, VALKONEN T. Mortality trends and setbacks: global convergence or divergence? *Lancet* 2004; **363**: 1155–1159.
- PETERSEN PE, KANDELMAN D, ARPIN S, OGAWA H. Global oral health of older people-call for public health action. *Community Dent Health* 2010; **27**: 257–267.
- Gezondheid, Leefstijl, Zorggebruik; t/m 2009. Statistics Netherlands (CBS), 2010.
- ALBAKER AM. The oral health-related quality of life in edentulous patients treated with conventional complete dentures. *Gerodontology* 2013; **30**: 61–66.
- SWELEM AA, GUREVICH KG, FABRIKANT EG, HASSAN MH, AQOU S. Oral health-related quality of life in partially edentulous patients treated with removable, fixed, fixed-removable, and implant-supported prostheses. *Int J Prosthodont* 2014; **27**: 338–347.
- HEYDECKE G, TEDESCO LA, KOWALSKI C, INGLEHART MR. Complete dentures and oral health-related quality of life - do coping styles matter? *Community Dent Oral Epidemiol* 2004; **32**: 297–306.
- VEDIN O, HAGSTROM E, GALLUP D, NEELY ML, STEWART R, KOENIG W, BUDAJ A, SRITARA P, WALLENTIN L, WHITE HD, HELD C. Periodontal disease in patients with chronic coronary heart disease: Prevalence and association with cardiovascular risk factors. *Eur J Prev Cardiol* 2015; **22**: 771–778.
- LOOS BG. Systemic markers of inflammation in periodontitis. *J Periodontol* 2005; **76**: 2106–2115.
- EBENFELT A, LUNDQVIST H, DAHLGREN C, LUNDBERG C. Neutrophils in mucosal secretion are functionally active. *J Clin Exp Immunol* 1996; **106**: 404–409.
- LANDZBERG M, DOERING H, ABOODI GM, TENENBAUM HC, GLOGAUER M. Quantifying oral inflammatory load: oral neutrophil counts in periodontal health and disease. *J Periodontol Res* 2015; **50**: 330–336.
- KOLACZKOWSKA E, KUBES P. Neutrophil recruitment and function in health and inflammation. *Nat Rev Immunol* 2013; **13**: 159–175.
- RIJKSCHROEFF P, JANSSEN ID, VAN DER WEIJDEN FA, KEIJSER BJ, LOOS BG, NICU EA. Oral polymorphonuclear neutrophil characteristics in relation to oral health: a cross-sectional, observational clinical study. *Int J Oral Sci* 2016; **8**: 191–198.
- LANTZMAN E, MICHMAN J. Leukocyte counts in the saliva of adults before and after extraction of teeth. *Oral Surg Oral Med Oral Pathol* 1970; **30**: 766–773.
- WRIGHT DE. Leucocytes in the saliva of edentulous and caries-free subjects. *Arch Oral Biol* 1962; **7**: 581–585.
- CALONIUS PE. The leukocyte count in saliva. *Oral Surg Oral Med Oral Pathol* 1958; **11**: 43–46.
- VAN DER VELDEN U. The Dutch periodontal screening index validation and its application in The Netherlands. *J Clin Periodontol* 2009; **36**: 1018–1024.
- MANTILLA GOMEZ S, DANSER MM, SIPOS PM, ROWSHANI B, VAN DER VELDEN U, VAN DER WEIJDEN GA. Tongue coating and salivary bacterial counts in healthy/gingivitis subjects and periodontitis patients. *J Clin Periodontol* 2001; **28**: 970–978.
- TURINA M, MILLER FN, MCHUGH PP, CHEADLE WG, POLK HC Jr. Endotoxin inhibits apoptosis but induces primary necrosis in neutrophils. *Inflamm* 2005; **29**: 55–63.
- VERMES I, HAANEN C, STEFFENS-NAKKEN H, REUTELINGSBERGER C. A novel assay for apoptosis. Flow cytometric detection of phosphatidylserine expression on early apoptotic cells using fluorescein labelled Annexin V. *J Immunol Methods* 1995; **184**: 39–51.
- ROTHER G, OSER A, VALET G. Dihydrorhodamine 123: a new flow cytometric indicator for respiratory burst activity in neutrophil granulocytes. *Naturwiss* 1988; **75**: 354–355.
- RICHARDSON MP, AYLIFFE MJ, HELBERT M, DAVIES EG. A simple flow cytometry assay using dihydrorhodamine for the measurement of the neutrophil respiratory burst in whole blood: comparison with the quantitative nitrobluetetrazolium test. *J Immunol Methods* 1998; **219**: 187–193.
- ATTSTROM R, EGELBERG J. Emigration of blood neutrophil polymorphonuclear leukocytes and monocytes into the gingival crevice. *J Periodontol Res* 1970; **5**: 48–55.
- SCHIOTT CR, LOE H. The origin and variation in number of leukocytes in the human saliva. *J Periodontol Res* 1970; **5**: 36–41.
- ASHKENAZI M, DENNISON DK. A new method for isolation of salivary neutrophils and determination of their functional activity. *J Dent Res* 1989; **68**: 1256–1261.
- LANDZBERG M, DOERING H, ABOODI GM, TENENBAUM HC, GLOGAUER M. Quantifying oral inflammatory load: oral neutrophil counts in periodontal health and disease. *J Periodontol Res* 2014; **50**: 330–336.
- DELIMA AJ, VAN DYKE TE. Origin and function of the cellular components in gingival crevice fluid. *Periodontol* 2000; **31**: 55–76.
- SCHROEDER HE. Transmigration and infiltration of leucocytes in human junctional epithelium. *Helv Odontol Acta* 1973; **17**: 6–18.
- SCHROEDER HE, ROSSINSKY K, LISTGARTEN MA. Human Junctional Epithelium as a Pathway for Inflammatory Exudation. *J Biol Buccale* 1989; **17**: 147–157.
- PAPADAKOU P, BLETSA A, YASSIN MA, KARLSEN TV, WIIG H, BERGGREEN E. Role of Hyperplasia of Gingival Lymphatics in Periodontal Inflammation. *J Dent Res* 2017; **96**: 467–476.
- HIRSCHFELD J. Dynamic interactions of neutrophils and biofilms. *J Oral Microbiol* 2014; **6**: 26102.
- WONG CH, HEIT B, KUBES P. Molecular regulators of leucocyte chemotaxis during inflammation. *Cardiovasc Res* 2010; **86**: 183–191.
- SACHDEO A, HAFFAJEE AD, SOCRANSKY SS. Biofilms in the edentulous oral cavity. *J Prosthodont* 2008; **17**: 348–356.
- O'DONNELL LE, ROBERTSON D, NILE CJ, CROSS LJ, RIGGIO M, SHERRIFF A, BRADSHAW D, LAMBERT M, MALCOLM J, BUIJS MJ, ZAURA E, CRIELAARD W, BRANDT BW, RAMAGE G. The oral microbiome of denture wearers is influenced by levels of natural dentition. *PLoS ONE* 2015; **10**: e0137717.
- NETT JE. The Host's Reply to *Candida* Biofilm. *Pathogens* 2016; **5**. <https://doi.org/10.3390/pathogens5010033>
- KATRAGKOU A, KRULAK MJ, SIMITSOPOULOU M, CHATZIMOSCHOU A, TAPARKOU A, COTTEN CJ, PALIOGIANNI F, DIZAMATAFTSI E, TSANTALI C, WALSH TJ, ROILIDES E. Interactions between human phagocytes and *Candida albicans* biofilms alone and in combination with antifungal agents. *J Infect Dis* 2010; **201**: 1941–1949.
- FOURNIER BM, PARKOS CA. The role of neutrophils during intestinal inflammation. *Mucosal Immunol* 2012; **5**: 354–366.

39. SUZUKI M, WATANABE T, SUKO T, MOGI G. Comparison of sinusitis with and without allergic rhinitis: characteristics of paranasal sinus effusion and mucosa. *Am J Otolaryngol* 1999; **20**: 143–150.
40. DURHAM SR. Mechanisms of mucosal inflammation in the nose and lungs. *Clin Exp Allergy* 1998; **28**: 11–16.
41. SZABADY RL, McCORMICK BA. Control of neutrophil inflammation at mucosal surfaces by secreted epithelial products. *Front Immunol* 2013; **4**: 1–6.
42. DUTZAN N, KONKEL JE, GREENWELL-WILD T, MOUTSOPOULOS NM. Characterization of the human immune cell network at the gingival barrier. *Mucosal Immunol* 2016; **9**: 1163–1172.
43. WEINDL G, NAGLIK JR, KAESLER S, BIEDERMANN T, HUBE B, KORTING HC, SCHALLER M. Human epithelial cells establish direct antifungal defense through TLR4-mediated signaling. *J Clin Invest* 2007; **117**: 3664–3672.
44. MERCK MILLIPORE MANUAL. Precise and accurate counts and viability measurements across multiple cell lines using the muse (TM) cell count & viability assay. *Biotechniques* 2012; **52**: 200–203.
45. LAKSHMI S, LAKSHMANAN A, GANESH P, SARAVANAN A. Effect of intensity of cigarette smoking on haematological and lipid parameters. *J Clin Diagn Res* 2014; **8**: 11–13.
46. TAGUCHI A, SANADA M, SUEI Y, OHTSUKA M, LEE K, TANIMOTO K, TSUDA M, OHAMA K, YOSHIZUMI M, HIGASHI Y. Tooth loss is associated with an increased risk of hypertension in postmenopausal women. *Hypertens* 2004; **43**: 1297–1300.
47. HUNG HC, JOSHIPURA KJ, COLDITZ G, MANSON JE, RIMM EB, SPEIZER FE, WILLETT WC. The association between tooth loss and coronary heart disease in men and women. *J Public Health Dent* 2004; **64**: 209–215.
48. OKORO CA, BALLUZ LS, EKE PI, AJANI UA, STRINE TW, TOWN M, MENSAH GA, MOKDAD AH. Tooth loss and heart disease: findings from the Behavioral Risk Factor Surveillance System. *Am J Prev Med* 2005; **29**: 50–56.
49. ZANELLA SM, PEREIRA SS, BARBISAN JN, VIEIRA L, SABACHUJFI E, HAAS AN, ROSING CK. Periodontal disease, tooth loss and coronary heart disease assessed by coronary angiography: a cross-sectional observational study. *J Periodontal Res* 2016; **51**: 221–227.
50. BROWN DW. Complete edentulism prior to the age of 65 years is associated with all-cause mortality. *J Public Health Dent* 2009; **69**: 260–266.
51. VEDIN O, HAGSTROM E, BUDAJ A, DENCHEV S, HARRINGTON RA, KOENIG W, SOFFER J, SRITARA P, STEBBINS A, STEWART RH, SWART HP, VIIGIMAA M, VINEREANU D, WALLENTIN L, WHITE HD, HELD C. Tooth loss is independently associated with poor outcomes in stable coronary heart disease. *Eur J Prev Cardiol* 2015; **22**: 771–778.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Table S1. Overview of results of analysis of oPMNs and cPMNs.