

Research

Open Access

Insulin sensitivity and Lp(α) concentrations in normoglycemic offspring of type 2 diabetic parents

Agathoklis Psyrogiannis*, Ioannis Habeos and Venetsana Kyriazopoulou

Address: Division of Endocrinology and Diabetes, Department of Medicine, University of Patras Medical School, Patras, Greece

Email: Agathoklis Psyrogiannis* - apsirog@med.upatras.gr; Ioannis Habeos - ihabeos@med.upatras.gr;

Venetsana Kyriazopoulou - vkyriazo@med.upatras.gr

* Corresponding author

Published: 29 October 2003

Received: 24 September 2003

Lipids in Health and Disease 2003, **2**:8

Accepted: 29 October 2003

This article is available from: <http://www.Lipidworld.com/content/2/1/8>

© 2003 Psyrogiannis et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: Offspring of at least 1 parent with type 2 diabetes are more resistant to the insulin action, exhibit higher incidence of dyslipidemia and are more prone to cardiovascular diseases. The association between Lp(α) and coronary heart disease is well established. An association between Lp(α) concentration and insulin sensitivity was examined in this study.

We investigated the serum LP(α) in 41 offspring of 41 families of type 2 diabetic subjects (group I) with normal glucose tolerance, compared to 49 offspring who their parents had no history of type 2 diabetes, matched for sex, age, BMI, WHR and blood pressure (group II). Serum Lp(α), triglycerides, insulin resistant index, HDL, LDL-cholesterol and insulin were measured.

Results: The offspring of type 2 diabetic subjects had higher fasting serum triglycerides (mean \pm SD 199.3 ± 184.2 vs. 147.1 ± 67.9 ng/dl, $p < 0.05$) lower HDL-cholesterol (37.3 ± 9.0 vs. 44.6 ± 7.8 , $p < 0.001$) and particularly higher Insulin resistance Index (HOMA-IR) (2.84 ± 1.39 vs. 1.67 ± 0.77 , $p < 0.001$).

They also had higher serum LP(α) concentration (15.4 ± 6.7 vs. 8.6 ± 6.0 , $p < 0.001$). By simple linear analysis in the offspring of type 2 diabetic parents there was no correlation of Lp(α) concentration with insulin resistance index Homa-IR ($r = 0,016$ $p = NS$).

Conclusions: We conclude that serum LP(α) is significantly increased in offspring of type 2 diabetic subjects but was not related to insulin sensitivity.

Background

Patients with type 2 diabetes mellitus are characterized by resistance to insulin-stimulated glucose uptake [1,2], dyslipidemia (especially increased triglyceride and decreased high-density lipoprotein (HDL) levels [3,4], hypertension [5] and coronary heart disease [6,7].

Lipoprotein (α) [Lp(α)], first described by Berg [8] in 1963, is a low density lipoprotein (LDL) - like substance

with a specific apoprotein, apoprotein (α) [apo (α)], bound to apo- β 100 by disulfide bridges [9,10].

Numerous studies have suggested that lipoprotein (α) concentrations may be an independent risk factor for coronary heart disease [11-15]. However, the molecular mechanism by which Lp(α) might promote atherosclerosis has not been clarified.

Table 1: Clinical characteristics of the offspring of type 2 diabetes parents (Group 1) and offspring of normal parents (Group 2).

Parameter	Group 1	Group 2	P value
Number (M/F)	41 (28/13)	49 (31/18)	NS
Age (Yr)	41.4 ± 6.8	43.5 ± 5.5	NS
BMI (Kg.m ⁻²)	28.1 ± 4.8	27.6 ± 3.5	NS
WHR	0.87 ± 0.1	0.8 ± 0.08	NS
Systolic BP (mmHg)	119.4 ± 15.5	120.8 ± 10.8	NS
Diastolic BP (mmHg)	72.1 ± 12.8	73.9 ± 10.1	NS

Table 2: Laboratory profile of the offspring of type 2 diabetes parents (Group 1) and offspring of normal parents (Group 2).

Parameter	Group 1	Group 2	P value
Triglyceride (mg/dl)	199.3 ± 184.2	147.1 ± 67.9	< 0.05
Cholesterol (mg/dl)	215.4 ± 41.6	199.2 ± 29.8	< 0.05
HDL-cholesterol (mg/dl)	37.3 ± 9.0	44.6 ± 7.8	< 0.001
LDL-cholesterol (mg/dl)	143.7 ± 31.7	124.6 ± 25.1	< 0.01
Lp(α) (mg/dl)	15.4 ± 6.7	8.6 ± 6.0	< 0.001

The relationship of Lp(α) concentrations with insulin resistance remains controversial. Some studies have shown increase Lp(α) concentration in subjects with NIDDM [16–18] whereas other studies have showed similar Lp(α) concentrations in type 2 diabetic subjects compared with normoglycemic subjects[19]. In other studies of nondiabetic subjects, insulin concentrations have not been associated with Lp(α) concentrations [20,21]. One study showed no elevated Lp(α) concentrations in normoglycemic insulin-resistant subjects [22].

In this report, we examine the association of Lp(α) concentrations to insulin sensitivity, calculated using the Homeostasis model Assessment (HOMA-IRI) in normoglycemic offspring of at least 1 parent with type 2 diabetes.

Results

The clinical characteristics of the study population are shown in Table 1 and the laboratory profile of the study population are shown in Table 2.

The group of offspring with type 2 diabetes parents as well as the control group had a similar distribution of age, sex, BMI, WHR systolic and diastolic BP. As expected the mean fasting serum triglycerides, cholesterol, and LDL-cholesterol were significantly higher in group 1 compared to the control group. HDL-cholesterol was significantly lower in group 1 compared to the control group.

The glucose and insulin concentrations in the fasting state and after an oral glucose challenge are shown in Table 3.

The mean fasting glucose and insulin concentrations were significantly higher in group 1 compared to group 2:

fasting glucose: 4.5 ± 0.6 vs. 4.3 ± 0.6 mmol/l (p < 0.05)

fasting insulin: 14.0 ± 6.4 vs. 8.6 ± 3.2 μU/ml (p < 0.005)

Serum insulin concentration 30 min post glucose challenge was significantly higher in the offspring group 1 (122.2 ± 94.3 vs. 81.7 ± 51.6 μU/ml, p < 0.005) as well at 60 min post glucose challenge (134.4 ± 103.1 vs. 99.9 ± 48.9 μU/ml, p < 0.05) whereas there was no difference at 120 min.

In the offspring of the diabetic parents the insulin resistance index was significantly higher compared to control group (2.84 ± 1.39 vs. 1.67 ± 0.77, p < 0.001). 22 (53.6%) of the 41 subjects of group 1, had HOMA-IR values > 2.5 indicating that they were insulin resistant [23–29], compared to only 6 (12.2%) of the 49 subjects of group 2.

Lp(α) concentration was significantly higher in group 1 compared to group 2 (15.4 ± 6.7 vs. 8.6 ± 6.0 ng/dl, p < 0.001).

In the offspring of the diabetic parents the insulin resistance index was positively correlated with serum triglycer-

Table 3: Serum insulin and glucose and insulin resistance index in offspring and control subjects in the fasting state and after an oral glucose challenge (120 min)

	Group 1 no 41	Group 2 no 49	P value
Glucose (mmol/l)			
Fasting	4.53 ± 0.62	4.26 ± 0.56	0.036
30 min	8.4 ± 2.0	8.3 ± 1.5	NS
60 min	8.4 ± 3.0	8.9 ± 2.5	NS
120 min	5.4 ± 1.3	5.6 ± 1.2	NS
Insulin (μU/ml)			
Fasting	14.0 ± 6.4	8.6 ± 3.2	< 0.001
30 min	122.2 ± 94.3	81.7 ± 50.6	0.013
60 min	134.4 ± 103.1	99.9 ± 48.9	0.042
120 min	76.9 ± 8.1	70.5 ± 40.7	NS
Insulin Resistance Index (HOMA-IR)	2.84 ± 1.39	1.67 ± 0.77	< 0.001

Table 4: Correlation (r) and significance (P) of HOMA-IR with serum concentration of Lipids and Lp(α).

Parameter	Group 1		Group 2	
	r	P value	r	P value
Cholesterol	0.177	NS	0.231	NS
Triglycerides	0.319	p < 0.01	0.099	NS
HDL	0.304	p < 0.05	0.055	NS
LDL	0.342	p < 0.02	0.161	NS
Lp(α)	0.016	NS		

ides and LDL. There was no correlation of HOMA-IR with any of the above parameter in the control group (Table 4).

There was no correlation of HOMA-IR with Lp(α) concentration in the offspring of the diabetic parents.

Discussion

Resistance to insulin-stimulated glucose uptake is a characteristic finding in patients with type 2 diabetes [1,2]. The observation that concordance for type 2 diabetes approaches 100% when one identical twin has the disease, strongly suggests that the decisive component of this syndrome is genetic in nature [30]. Several studies have shown that offspring of at least one parent with type 2 diabetes display hyperinsulinism and are more resistant to insulin action than offspring of parents whose glucose tolerance was normal [31–34].

The result of our study supports and expands those findings. We found that offspring of patients with type 2 diabetes when compared with normal individuals whom their parents had no history of type 2 diabetes matched for sex, age and BMI, were relative hyperinsulinemic, had

higher fasting serum triglycerides, lower HDL-cholesterol and higher insulin resistance index.

Moreover, the serum Lp(α) concentration was significantly higher compared with control group. The overall impression from the large number of publications on diabetes mellitus and Lp(α), is that neither insulin dependent nor non-insulin dependent diabetes has a direct effect on serum Lp(α) [16–19,31–34]. Some reports of increased levels in non-insulin dependent diabetes may result from the inclusion of patients with established CHD in the population studied. However, there does appear to be an increase in serum Lp(α) in insulin-dependent diabetes when nephropathy is present and this may be the case even when this is at the stage of microalbuminuria [35–42]. Thus the rise seems to occur with much less marked proteinuria than in primary renal disease. Possibly this is because there is some specific feature of the nephropathy in diabetes, which triggers the Lp(α) response early in the disease or perhaps microalbuminuria in diabetes is a marker for a more extensive vasculopathy which is the major stimulus for Lp(α) to increase.

There has been little published data on insulin sensitivity and Lp(α) concentrations. Lp(α) concentrations were not significantly related to insulin sensitivity measured by the intravenous glucose tolerance test in nondiabetic subjects [43]. Haffner et al [22] demonstrated that insulin-resistant normoglycemic men did not have elevated Lp(α) concentrations and that Lp(α) concentrations are principally controlled at the locus encoding apo (α) in agreement with the data of other studies [44–47].

Moreover, Haffner et al suggested that the relationship between insulin sensitivity and Lp(α) concentrations might be at least partially dependent on apo (α) phenotype [22]. One possibility for such an association could be that genes for apo (α) genotype and insulin sensitivity could be in linkage disequilibrium.

To our knowledge this is the first report in which the levels of Lp(α) concentration and insulin sensitivity in normoglycemic offspring of type 2 diabetic parents were studied. Our study demonstrates that Lp(α) concentrations were not significantly correlated with HOMA-IR ($r = 0.016$). The impression from this finding and in agreement with previous studies [16,19–21,35–43], is that serum Lp(α) concentration is not related to insulin sensitivity. However, Lp(α) concentration was significantly higher in the group of offspring of diabetic parents compared to the control group ($p < 0.001$).

Conclusions

We conclude that serum LP(α) is significantly increased in offspring of type 2 diabetic subjects but was not related to insulin sensitivity.

Methods

Subjects

We studied 54 healthy offspring of 54 separate families with type 2 diabetes parents. The subjects were Caucasians of Greek origin. They all had a parent who had developed type 2 diabetes mellitus after 50 years of age. Two patients were the offspring whom both parents were diabetic. Maturity-onset diabetes of the young was excluded on the basis of the late onset of diabetes. The 54 subjects (group 1) were matched to 54 subjects with no family history for diabetes and were used 95 controls (group 2). The subjects in both groups were matched for sex, age, Body Mass Index (BMI) and waist-to-hip ratio (WHR).

All subjects had normal physical examination and normal values for routine laboratory parameters including hematology, c-reactive protein, HbA1c (upper limit of normal 6.5%), adrenal, kidney, liver and thyroid function. The subjects were not under any medication known to affect carbohydrate or lipoprotein metabolism and all denied alcohol intake.

After an appropriate carbohydrate diet for one week, an oral Glucose Tolerance test (OGTT) was performed and patients with impaired glucose tolerance or type 2 diabetes were excluded from the study. Forty one subjects remained for group 1 and 49 for group 2. Informed consent was obtained from all participants in the study.

Study design

After 12-h overnight fast, blood samples were drawn for serum glucose, insulin, lipids, lipoprotein and Lp(α). The subjects then ingested 75 g of glucose load over a 2-min period. Blood samples were drawn at 30, 60 and 120 min thereafter for measurement of serum glucose and insulin concentrations. Plasma glucose was measured with an automatic analyzer by the glucose oxidase method. Immunoreactive insulin (IRI) was measured with an insulin radioimmunoassay (RIA) kit.

We used the homeostatic model to assess insulin resistance, where HOMA-IR = fasting plasma insulin ($\mu\text{U/ml}$) \times fasting plasma glucose (mmol/l) / 22.5 [23–25]. HOMA-IR provides a reliable estimate of insulin resistance across the range of glucose tolerance using either the frequently sampled intravenous glucose tolerance test (FSIGTT) with minimal model analysis or the glucose clamp technique as the gold standard measure of insulin sensitivity [26,27].

Total serum cholesterol, serum triglyceride and HDL-cholesterol were assayed with a commercial kit (Boehringer Mannheim, Germany). LDL cholesterol was calculated according to the Friedewald equation [28]. Lp(α) was estimated by enzyme immunoassay (Macra Lp(α), Strategic Diagnostic, Newark, NJ).

The study design was approved by the ethical and scientific committee of the Hospital.

Statistical analysis

Results are expressed as mean \pm SD. Data were analyzed according to the unpaired student's t-test or by the non-parametric Mann-Whitney U test of Wilcoxon's test. Pearson's correlation coefficient or the Spearman's test were used depending on the normal distribution of the variables. A p-value of less than 5% was considered statistically significant.

References

1. Reaven GM: **Insulin resistance in non-insulin dependent diabetes mellitus does it exist and can it be measured?** *Am J Med* 1983, **Suppl 1A**:3-17.
2. De Fronzo RA, Bonadonna RC and Ferrannini E: **Pathogenesis of NIDDM: a balanced overview.** *Diabetes Care* 1999, **15**:318-368.
3. Garg A, Helderan JH, Koffler M, Ayuso R, Rosenstock J and Raskin P: **Relationship between lipoprotein levels and in vivo insulin action in normal white men.** *Metabolism* 1988, **37**:982-987.
4. Mykkanen L, Haffner SM, Ronnema L, Lehto S, Ronnema T, Pyorala K and Laakso M: **Is there a sex difference in the association of**

- insulin levels and insulin sensitivity with lipids and lipoproteins? *Metabolism* 1994, **43**:523-528.
5. Ferrannini E, Buzzigoli G, Bonadonna R, Giorico MA, Oleggini M, Graziadei L, Pedrinelli R, Brandi L and Bevilacqua S: **Insulin resistance in essential hypertension.** *N Engl J Med* 1987, **317**:350-357.
 6. Pyorala K, Savolainen E, Kaukola S and Haapakoski J: **Plasma insulin as coronary heart disease risk factor: relationship to other risk factors and predictive value during 9^{1/2} year follow-up of the Helsinki Policeman population.** *Acta Med Scand* 1985, **701**(Suppl):38-52.
 7. Ducimetiere P, Eschwege E, Papoz L, Richard JL, Claude JR and Rosselin G: **Relationship of plasma insulin levels to the incidence of myocardial infarction and coronary heart disease mortality in middle-aged population.** *Diabetologia* 1980, **19**:205-210.
 8. Berg K: **A new serum type system in man – The Lp system.** *Acta Pathol Microbiol Scand* 1963, **59**:369-382.
 9. Utermann G: **The mysteries of lipoprotein (a).** *Science* 1989, **246**:904-910.
 10. Scann AM and Fress GM: **Heterogeneity and biological relevance.** *J Clin Invest* 1990, **85**:1709-1715.
 11. Berg K, Dahlen G and Frick MH: **Lp(α) lipoprotein and pre-beta lipoprotein in patients with coronary heart disease.** *Clin Genet* 1974, **6**:230-235.
 12. Dahlen GH, Guyton JR, Attar M, Farmer JA, Kautz JA and Gotto AM Jr: **Association of levels of lipoprotein LP(α), plasma lipids and other lipoproteins with coronary artery disease documented by angiography.** *Circulation* 1986, **74**:758-765.
 13. Sandholzer C, Boerwinkle E, Saha N, Tong MC and Utermann G: **Apolipoprotein (α) phenotypes Lp(α) concentration and plasma lipid levels in relation to coronary heart disease in a Chinese population: Evidence for the role of the Apo (α) Gene in Coronary Heart Disease.** *J Clin Invest* 1992, **89**:1040-1046.
 14. Sandholzer C, Saha N, Kark JD, Rees A, Jaross W, Dieplinger H, Hop-pichler F, Boerwinkle E and Utermann G: **Apo(α) isoforms predict risk for Coronary Heart Disease. A study in six populations.** *Arteriosclerosis Thromb* 1992, **12**:1214-1226.
 15. Mohan V, Deepa R, Haranath SP, Premalatha G, Rema M, Sastry NG and Enas EA: **Lipoprotein (α) is an Independent Risk Factor for Coronary Artery Disease in NIDDM Patients in South India.** *Diabetes Care* 1998, **21**:1819-1823.
 16. Ramirez LC, Arauz-Pacheco C, Lackner C, Albright G, Adams BV and Raskin P: **Lipoprotein (α) levels in diabetes mellitus: relationship to metabolic control.** *Ann Intern Med* 1992, **117**:42-47.
 17. Ishida Y, Kazumi T and Yoshida M: **Serum lipoprotein (α) in patients with non-insulin-dependent diabetes mellitus (Abstract).** *Diabetes* 1991, **40**(Suppl 1):188A.
 18. Heller FR, Jamart J, Honore P, Derue G, Novik V, Galanti L, Parfony A, Hondekijn JC and Buysschaert M: **Serum lipoprotein (α) in patients with diabetes mellitus.** *Diabetes Care* 1993, **16**:819-823.
 19. Haffner SM, Morales PA, Stern MP and Gruber MK: **Lp(α) concentrations in NIDDM.** *Diabetes* 1992, **41**:1267-1272.
 20. Sundell IB, Nilsson TK, Hallmans G, Hellsten G and Dahlen GH: **Interrelations between plasma levels of plasminogen activator inhibitor, tissue plasminogen activator, lipoprotein (α), and established cardiovascular risk factors in a North Swedish population.** *Atherosclerosis* 1989, **80**:9-16.
 21. Haffner SM, Gruber KK, Morales PA, Hazuda HP, Valdez RA, Mitchell BD and Stern MP: **Lipoprotein (α) concentrations in Mexican Americans and non-Hispanic whites: The San Antonio Heart Study.** *Am J Epidemiol* 1992, **136**:1060-1068.
 22. Haffner SM, Karhapaa P, Rainwater DL, Mykkanen L, Aldrete G Jr and Laakso M: **Insulin sensitivity and Lp(α) concentrations in Normoglycemic men.** *Diabetes Care* 1995, **18**:193-199.
 23. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF and Turner RC: **Homeostasis model assessment: insulin resistance and b-cell function from fasting plasma glucose and insulin concentrations in man.** *Diabetologia* 1985, **28**:412-419.
 24. Haffner SM, Kennedy E, Gonzalez C, Stern MP and Mietinen H: **A prospective analysis of the HOMA model: the Mexico City Diabetes Study.** *Diabetes Care* 1996, **19**:1138-1146.
 25. Duncan MH, Singh BM, Wise PH, Carter G and Alagband-Zadeh J: **A single measure of insulin resistance in human disease.** *Lancet* 1995, **346**:120-121.
 26. Duncan MH, Singh BM, Wise PH, Carter G and Alagband-Zadeh J: **Ability of alternative indices of insulin sensitivity to predict cardiovascular risk: comparison with the "minimal model". Insulin Resistance Atherosclerosis (IRAS) Investigators.** *Ann Epidemiol* 1998, **8**:358-369.
 27. Bonora E, Targher G, Alberiche M, Bonadonna RC, Saggiani F, Zenere MB, Monauni T and Muggeo M: **Homeostasis model assessment closely mirrors the glucose clamp technique in the assessment of insulin sensitivity.** *Diabetes Care* 2000, **23**:57-63.
 28. Friedewald WT, Lery RI and Fredrickson DS: **Estimation of the concentration of low density lipoprotein cholesterol in plasma without use of the preparative ultracentrifuge.** *Clin Chem* 1972, **18**:499-502.
 29. Taniguchi A, Fukushima M, Sakai M, Miwa K, Makita T, Nagata I, Nagasaka S, Doi K, Okumura T, Fukuda A, Kishimoto H, Fukuda T, Nakai-shi S, Tokuyama K and Nakai Y: **Remnant-like particle cholesterol, triglycerides and insulin resistance in nonobese Japanese type 2 diabetic patients.** *Diabetes Care* 2000, **23**:1766-1769.
 30. Barnett AH, Eff C, Leslie RD and Pyke DA: **Diabetes in identical twins: A study of 200 pairs.** *Diabetologia* 1981, **20**:87-93.
 31. Ho LT, Chang ZY, Wang JT, Li SH, Liu YF, Chen YD and Reaven GM: **Insulin insensitivity in offspring of parents with type 2 diabetes mellitus.** *Diabetic Medicine* 1990, **7**:31-34.
 32. Ganesh RDG, Volkman A and Leslie HP et al.: **Insensitivity to insulin in offspring of non-insulin dependent diabetic patients.** *Diabetes Nutrition Metabolism* 1988, **3**:235-237.
 33. Haffner SM, Stern MP, Hazuda HP, Mitchell BD and Patterson JK: **Increased insulin concentrations in non-diabetic offspring of diabetic parents.** *N Engl J Med* 1988, **319**:1297-1301.
 34. Leslie RD, Volkman HP, Poncher M, Hanning I, Orskov H and Alberti KG: **Metabolic abnormalities in children of non-insulin dependent diabetics.** *Brit Med J* 1986, **293**:840-841.
 35. Haffner SM, Tattle KR and Rainwater DL: **Decrease of lipoprotein (α) with improved glycemic control in IDDM subjects.** *Diabetes Care* 1991, **14**:302-307.
 36. Kapelrud H, Bangstad HJ, Dahl-Jorgensen K, Berg K and Hanssen KF: **Serum Lp(α) lipoprotein concentrations in insulin dependent diabetic patients with microalbuminuria.** *Brit Med J* 1991, **3**:675-678.
 37. Levutsky LL, Scann AM and Gould SH: **Lipoprotein (α) levels in black and white children and adolescents with IDDM.** *Diabetes Care* 1991:283-287.
 38. Nakata H, Horita K and Eto M: **Alteration of lipoprotein (α) concentration with glycaemic control in non-insulin dependent diabetic subjects without complications.** *Metabolism* 1993:1323-1326.
 39. Ritter MM, Loscar M, Richter WO and Schwandt P: **Lipoprotein (α) in diabetes mellitus.** *Clinica Chimica Acta* 1993, **4**:45-54.
 40. Winocour PH, Bhatnagar D, Ishola M, Arrol S and Durrington PN: **Lipoprotein (α) and microvascular disease in type I (insulin dependent) diabetes mellitus.** *Diabetic Medicine* 1991:922-927.
 41. Winocour PH, Durrington PN, Bhatnagar D, Mbewu AD, Ishola M, Mackness M and Arrol S: **A cross-sectional evaluation of cardiovascular risk factors in coronary heart disease associated with type I (insulin dependent) diabetes mellitus.** *Diabetes Research and Clinical Practice* 1992, **8**:173-184.
 42. Wolffenbuttel BH, Leurs PB, Sels JP, Rondas-Colbers GJ, Menheere PP and Nieuwenhuijzen Kruseman AC: **Improved blood glucose control by insulin therapy in type 2 diabetic patients has no effect on lipoprotein (α) levels.** *Diabetic Medicine* 1993, **10**:427-430.
 43. Sidhu M, Crook D, Godsland IF, Walton C, Wynn V and Oliver MF: **Inverse relationship between Lp(α) levels and first-phase insulin secretion.** *Diabetes* 1992, **41**:1341-1345.
 44. Utermann G, Menzel HJ, Kraft HG, Dube HC, Kemmler HG and Seitz C: **Lp(α) glycoprotein phenotypes: inheritance and relation of Lp(α) – lipoprotein concentrations in plasma.** *J Clin Invest* 1987, **80**:458-465.
 45. Rainwater DL, Manis GS and Vandeberg JL: **Hereditary and dietary effects on apolipoprotein (α) isoforms and Lp(α) in baboons.** *J Lipid Res* 1989, **30**:549-558.
 46. Boerwinkle E, Leffert CC, Lin J, Lackner C, Chiesa G and Hobbs HH: **Apolipoprotein (α) gene accounts for greater than 90% of the variation in plasma lipoprotein (α) concentrations.** *J Clin Invest* 1992, **90**:52-60.
 47. Gaubatz JW, Ghanem KI, Guevara J Jr, Nava ML, Patsch W and Morrisett JD: **Polymorphic forms of human apolipoprotein (α):**

inheritance and relationship of their molecular weights to plasma levels of lipoprotein (α). *J Lipid Res* 1990, **31**:603-6.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

