


ORIGINAL ARTICLE

Blood pressure-related differences in brain health between young African Americans and Caucasian Americans

Junyeon Won¹ | Sushant M. Ranadive¹ | Daniel D. Callow^{1,2} | Shuo Chen³ | J. Carson Smith^{1,2} 

¹Department of Kinesiology, University of Maryland, College Park, MD, USA

²Program in Neuroscience and Cognitive Science, University of Maryland, College Park, MD, USA

³Department of Epidemiology and Public Health, University of Maryland School of Medicine, Baltimore, MD, USA

Correspondence

J. Carson Smith, Department of Kinesiology, University of Maryland, School of Public Health, 2351 SPH Bldg #255, College Park, MD 20742, USA.
Email: carson@umd.edu

Abstract

Background: Although there are moderating effects of race on blood pressure (BP) and brain health in older adults, it is currently unknown if these race-related differences in cardiovascular and associated brain function are also present in younger adults. The purpose of this study was to investigate the interaction between race and BP on brain health in younger African (AA) and Caucasian Americans (CA).

Methods: We studied 971 younger adults (29.1 ± 3.5 years; 180 AAs and 791 CAs) who volunteered to participate in the Human Connectome Project. Cognitive composite scores, brain volume, and cortical thickness using MRI were cross-sectionally assessed. ANCOVA was used to examine interactions between race and mean arterial pressure (MAP) on cognitive test scores and brain structure.

Results: After controlling for age, sex, education, and BMI, there were significant Race \times MAP interaction effects on cognitive composite scores and cortical thickness. Among AAs but not CAs, as MAP increased, both global cognitive performance and entorhinal cortex (ERC) thickness decreased.

Conclusions: MAP was an important moderator of racial differences in cognitive performance and ERC thickness. Our findings suggest that young AAs may carry a greater hypertension-associated risk for cognitive brain health deficit. Interventions that address early signs of hypertension in AAs are needed to determine if the racial disparities in BP-related brain health in late adulthood can be reduced.

KEYWORDS

blood pressure, cognitive function, cortical thickness, mean arterial pressure, race, younger adults

1 | BACKGROUND

Beyond the well-documented role of hypertension as a leading risk factor for cardiovascular diseases, the linkage between vascular function and brain health has been extensively studied (Iadecola & Gottesman, 2019). For example, impaired vascular function is associated with cerebral hypoperfusion (Muller et al., 2012), cerebral blood vessel

damage (Bohannon et al., 2002), neurovascular inflammation (Morillas et al., 2012), reduced white matter integrity (Faraco & Iadecola, 2013), and smaller brain tissue volumes in regions that are important to cognitive function and memory, such as the frontal cortex and hippocampus (Beauchet et al., 2013). Due to its deleterious impact on brain function, hypertension has been highlighted as an important biomarker to predict future cognitive impairment (e.g., memory, attention,

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Physiological Reports* published by Wiley Periodicals LLC on behalf of The Physiological Society and the American Physiological Society

and executive function) (Gorelick & Nyenhuis, 2012; Saxby et al., 2003) and development of dementia (Faraco & Iadecola, 2013).

African Americans (AA) have a disproportionately greater incidence and severity of hypertension and hypertension-related cognitive impairments than Caucasian Americans (CA) (Redmond et al., 2011). Middle-aged AAs have been shown to exhibit a faster decline in global cognition, which was associated with higher systolic blood pressure (SBP) and mean arterial pressure (MAP) relative to CAs (Levine et al., 2019). In a neuroimaging study of hypertensive older adults, AAs, compared to their CA counterparts, exhibited impaired network connectivity and cortical thickness of the insula (Chand et al., 2017), a key node of salience network that is important in detecting and filtering salient stimuli and processing information related to emotion, cognition, and reward (Menon & Uddin, 2010). Indeed, higher cumulative blood pressure (BP) over a lifetime, with an onset at a younger age, may be a critical contributor to the faster rate of later-life cognitive decline among AAs relative to CAs (Levine et al., 2020).

Most of what we currently know about interactions between race and BP on brain health outcomes have come from studies of older and hypertensive individuals. Whether or not the moderating effects of BP on indices of brain function are present in normotensive younger AA and CA adults has not been addressed. Understanding these BP-related associations with cognitive performance and brain structure in younger AA and CA adults is an essential step toward identifying modifiable factors and implementing early interventions that may help address racial disparities in late-life cognitive health. Thus, the present study aimed to investigate the moderating role of BP on cognitive function, brain volume, and cortical thickness in younger, normotensive AAs and CAs. Based on evidence showing greater MAP-related decline in global cognition among middle-aged AAs relative to CAs (Levine et al., 2019), we hypothesized that negative associations between MAP and cognitive performance, cortical and hippocampal volumes, and cortical thickness will be greater in younger AAs compared to CAs.

2 | METHODS

2.1 | Participants

The present study used open-access data from the University of Minnesota (Wu-Min) Human Connectome (HC) Project (<https://www.humanconnectome.org/study/hcp-young-adult/document/1200-subjects-data-release>; Van Essen et al., 2013). In short, healthy younger adults were primarily recruited from Missouri. A complete list of eligibility

criteria was previously reported (Van Essen et al., 2013). Briefly, exclusion criteria included: (a) Significant history of psychiatry disorder, substance abuse, neurological, or cardiovascular disease; (b) two or more seizures after age 5 or a diagnosis of epilepsy; (c) any genetic disorder (e.g., cystic fibrosis or sick cell disease); (d) Multiple sclerosis, cerebral palsy, brain tumor, or stroke; and (e) Moderate or severe claustrophobia. Self-reported race was used in this study. Participants provided written consent approved by the Institutional Review Board of Washington University. The entire experimental procedure and data share were performed in compliance with the relevant guidelines and regulations (Glasser et al., 2016). Participants visited Washington University two different days for magnetic resonance imaging (MRI) scanning and BP/cognitive assessments, respectively.

2.2 | Blood Pressure Measurement, Cognitive Tests, and Mental Health

Prior to the in-person visit, participants were asked about (a) whether they had ever been told by a physician that they had high BP, and (b) whether they were currently taking medications for high BP. Each participant's resting BP was measured at their visit using established guidelines. Sitting BP in the left arm was measured using a semi-automatic BP monitor (ADC). A comprehensive battery of neuropsychological tests was administered to evaluate cognitive function. Detailed descriptions about the test administrations and scoring can be found in previous studies (Gur et al., 2010; Weintraub et al., 2013). A comprehensive battery of neuropsychological tests was performed to evaluate cognitive function, including (a) Picture Sequence Memory (to measure episodic memory); (b) Dimensional Change Card Sort (to measure executive function and cognitive flexibility); (c) Flanker (to measure executive function); (d) Oral Reading Recognition Test (to measure reading decoding); (e) picture vocabulary (to measure vocabulary knowledge); (f) Pattern Comparison Test (to measure processing speed); (g) Variable Short Penn Line Orientation Test (VSPLIT; to measure spatial orientation processing); (h) Short Penn Continuous Performance (SPEC; to measure sustained attention); and (i) List Sorting (to measure working memory). Composite scores were calculated as the average of these subtest scores using the NIH Toolbox Cognition Battery Reuben et al., 2013) that included (a) total composite score; (b) crystallized composite score (normalized scores of picture vocabulary and oral reading tests); (c) early cognitive composite score (normalized scores of picture vocabulary, flanker, dimensional change card sort, and picture sequence memory); and (d) cognitive

fluid composite score (normalized scores of flanker, dimensional change card sort, picture sequence memory, and list sorting). For the present analysis, we used these four cognitive composite scores as measurements of cognitive performance. Anxiety/depression score assessed as part of life function questionnaire (Achenbach, 2003) in the HC Project was tested for the between-group difference as a potential covariate affecting BP.

2.3 | MRI data acquisition

Whole-brain MRI was conducted using a customized Siemens 3.0 Tesla MR Scanner at Washington University. A 32-channel head coil was used for radio frequency transmission and reception. A high-resolution T1-weighted anatomical image was acquired with gradient echo sequence: field of view = 224 mm, voxel size = $0.7 \times 0.7 \times 0.7$ mm, slice thickness = 0.9 mm, repetition time = 2400 ms, echo time = 2.14 ms, inversion time = 1000 ms, flip angle = 8° , and duration = 7:40 min (Van Essen et al., 2012).

2.4 | MRI data analysis

MRI data were processed using HC Project's optimal processing pipeline (Glasser et al., 2013) where FreeSurfer's (version 5.3.0) automatic cortical reconstruction process for cortical parcellation and subcortical segmentation (recon-all) (Fischl, 2012) was included. Based on prior evidence documenting the association between hypertension and hippocampal atrophy (Den Heijer et al., 2005; Korf et al., 2004), FreeSurfer-defined bilateral hippocampal volume was chosen *a priori* as region of interest. The interactive effect between race and BP on total gray matter volume was also analyzed as a control to test whether the hypothesized higher BP-related brain atrophy was driven by the total brain volume rather than bilateral hippocampal volume, specifically.

FreeSurfer's segmentation analysis also quantified cortical thickness by assessing the closest distance between the gray/white boundary and gray/cerebrospinal fluid boundary (Fischl & Dale, 2000). Hypertension is associated with decreased cortical thickness (Seo et al., 2012) and affects brain regions that are vulnerable to aging including entorhinal cortex (ERC) (Raz, 2000). ERC is a brain region where early accumulation of amyloid plaques begins and it later proliferates into the hippocampus (Braak & Braak, 1991). Therefore, bilateral ERC thickness was chosen *a priori* as a region of interest. We also selected bilateral insula *a priori* based on previous evidence suggesting differences in insula thickness between hypertensive AA versus CA older individuals (Chand et al., 2017).

2.5 | Statistical analyses

We used the Shapiro–Wilk test to determine normality of the demographic data. Between-group differences (AA vs CA) in demographic characteristics were measured using independent sample *t*-tests (or Wilcoxon rank sum tests) for continuous variables and a chi-squared test for the categorical variable. A Shapiro–Wilk test was also performed to determine normality of the cognitive and brain MRI measurements. Due to significant demographic differences between AAs and CAs, between-group differences in cognitive performance and brain MRI measures were determined using ANCOVA after adjusting age, gender, education, and BMI.

MAP ($[(DBP \times 2) + SBP]/3$) was selected as a measurement of BP as it represents average BP assessed from SBP and DBP. ANCOVA was used to examine the Race \times MAP interaction on the cognition and brain structure variables. To accomplish this, race and MAP were first set as independent variables, and age, education, and BMI were added as covariates. Gender was also included as a covariate based on prior investigations showing significant differences in ICV (Raz, 2000; Won et al., 2019) and cortical thickness (Gautam et al., 2013; Savic & Arver, 2014) between men and women. For the volumetric analyses, total intracranial volume estimated by FreeSurfer was further included as a covariate. Next, a race by MAP interaction term was added to the model. Cognitive test results and cortical volume and thickness, respectively, were included as dependent variables. Lastly, Benjamini–Hochberg false discovery rate (FDR) correction (Benjamini & Hochberg, 1995) was conducted to control the family-wise error rate for multiple comparisons in the interaction results. The statistical significance was determined using a two-tailed $\alpha = 0.05$. All statistical analyses were performed using JASP (version 0.12.2).

3 | RESULTS

3.1 | Participants

Detailed study recruitment protocol and retention have been reported previously (Van Essen et al., 2012, 2013). Among a total of 1207 participants who completed the study protocol, 126 participants whose race were neither the AA nor CA were excluded. Additionally, 85 participants with Hispanic/Latino or unknown ethnicity were also excluded. Next, 21 participants with missing BP data were excluded and 4 outliers (Cook's distance >0.5) in MAP were further excluded. Finally, cognitive test performance for the remaining 971 participants was analyzed. For the brain volume and cortical thickness assessments, 67 individuals were further excluded due to missing MRI data. Thus, data from a total of 904

participants were included in the brain volume and thickness analyses.

The participants had an average age of 29.1 years, BMI of 27.0 kg/m², and 14.8 years of education. 55.2% of the participants were women. AAs had significantly higher BMI ($t[967] = 5.303$, $p < 0.0001$, $d = 0.438$, 95%CI for $d = 0.274$, 0.601), fewer years of education ($t[967] = 8.686$, $p < 0.0001$, $d = 0.717$, 95% CI for $d = 0.552$, 0.882), and greater MAP ($t[968] = 2.602$, $p = 0.009$, $d = 0.214$, 95%CI for $d = 0.052$, 0.377) compared with CAs. As there was no significant between-group difference in anxiety/depression symptom score ($t[965] = 0.100$, $p = 0.920$, $d = 0.008$, 95%CI for $d = -0.153$, 0.170), anxiety/depression symptom score was not included as a covariate in the subsequent analyses (see Table 1). None of the participants was taking BP medication.

3.2 | Between-group differences on cognitive and brain assessments

After controlling for age, gender, education, and BMI, AAs, compared with CAs, exhibited consistently lower cognitive composite scores, including total composite score ($F[1,944] = 109.081$, $p < 0.0001$, $\eta^2_p = 0.104$), crystallized composite score ($F[1,951] = 86.583$, $p < 0.0001$, $\eta^2_p = 0.083$), early cognitive composite score ($F[1,948] = 74.491$, $p < 0.0001$, $\eta^2_p = 0.076$), and cognitive fluid composite score ($F[1,944] = 62.353$, $p < 0.0001$, $\eta^2_p = 0.061$). AAs also had significantly smaller total gray matter volume ($F[1,895] = 22.314$, $p < 0.0001$, $\eta^2_p = 0.024$), bilateral hippocampal volume ($F[1,895] = 4.729$, $p = 0.029$,

$\eta^2_p = 0.005$), bilateral ERC thickness ($F[1,896] = 33.963$, $p < 0.0001$, $\eta^2_p = 0.036$), and bilateral insula thickness ($F[1,896] = 18.170$, $p = 0.0006$, $\eta^2_p = 0.013$) than CAs (see Table 2).

3.3 | Interaction between race and MAP on cognition

Table 3 illustrates the interaction between race and MAP on cognitive composite scores. ANCOVA results revealed significant interactions between race and MAP on total composite score ($F[1,943] = 5.833$, $p = 0.015$, $\eta^2_p = 0.006$) (Figure 1; Panel a), crystallized composite score ($F[1,950] = 4.324$, $p = 0.037$, $\eta^2_p = 0.004$) (Figure 1; Panel b), and early cognitive composite score ($F[1,947] = 3.956$, $p = 0.046$, $\eta^2_p = 0.004$) (Figure 1; Panel c). The interaction between race and MAP on the cognitive fluid composite score was in the same direction but was not statistically significant ($F[1,943] = 3.548$, $p = 0.059$, $\eta^2_p = 0.003$) (Figure 1; Panel d). The p -value of the significant effects was greater than FDR correction threshold (FDR-corrected threshold for total composite score $p = 0.012$, crystallized composite score $p = 0.025$, and early cognitive composite score $p = 0.037$).

3.4 | Interactions between race and MAP on brain volume/cortical thickness

While there was a significant interactive effect between race and MAP on the bilateral ERC thickness

TABLE 1 Demographics and blood pressure data of the participants.

| | Total sample (<i>n</i> = 971) | African Americans (<i>n</i> = 180) | Caucasian Americans (<i>n</i> = 791) | Group differences |
|-------------------------------|-----------------------------------|--|--|-------------------------------------|
| | Mean (SD) | Mean (SD) | Mean (SD) | <i>p</i> -value (Cohen's <i>d</i>) |
| Demographics | | | | |
| Age (years) | 29.1 (3.5) | 28.9 (3.7) | 29.1 (3.4) | 0.439 (0.064) |
| Female (n, %) | 535 (55.2) | 111 (61.6) | 424 (53.6) | 0.056 _F |
| BMI (kg/m ²) | 27.0 (5.5) | 29.1 (6.5) | 26.6 (5.2) | 1.40e−7 (0.217) |
| Education (years) | 4.8 (1.8) | 13.8 (2.0) | 15.0 (1.7) | 1.57e−17 (0.717) |
| Blood pressure (mm Hg) | | | | |
| SBP | 124.6 (14.4) | 126.9 (15.8) | 124.1 (14.0) | 0.019 (0.082) |
| DBP | 77.2 (10.9) | 79.0 (11.7) | 76.8 (10.7) | 0.015 (0.123) |
| MAP | 93.0 (11.1) | 94.9 (11.9) | 92.5 (10.8) | 0.009 (0.103) |
| Mental health | | | | |
| ASR Anxious/Depression Score | 53.8 (5.9) | 53.8 (5.8) | 53.7 (5.9) | 0.920 (0.00001) |

Notes: _F indicates Fisher's exact test; BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; MAP, mean arterial pressure; mmHg, millimeter of mercury; ASR Anxious/Depression Score, measured as part of life function questionnaire (Achenbach, 2003) in the Human Connectome Project; Bold indicates $p < 0.05$.

TABLE 2 Cognitive test, cortical volume, and thickness data of the participants

| | Total sample (<i>n</i> = 971) | African Americans (<i>n</i> = 180) | Caucasian Americans (<i>n</i> = 791) | Group differences |
|---|-----------------------------------|--|--|--------------------------------|
| | Mean (SD) | Mean (SD) | Mean (SD) | <i>p</i> -value (η^2_p) |
| Cognitive tests | | | | |
| Total Cognitive Composite Score | 111.6 (20.6) | 94.0 (20.4) | 115.6 (18.4) | 2.05e−24 (0.104) |
| Crystallized Composite Score | 108.2 (17.1) | 94.2 (18.1) | 111.4 (15.2) | 8.97e−20 (0.083) |
| Early Cognitive Composite Score | 105.6 (16.0) | 93.7 (15.9) | 108.3 (14.8) | 3.89e−18 (0.076) |
| Cognitive Fluid Composite Score | 104.9 (17.1) | 93.9 (16.7) | 107.4 (16.2) | 7.94e−15 (0.061) |
| Regional brain volume (mm³) | | | | |
| Total gray matter | 686755.6 (67984.6) | 639904.3 (60895.5) | 697026.7 (65159.3) | 4.52e−6 (0.023) |
| Bilateral hippocampus | 8913.7 (896.8) | 8415.8 (860.9) | 9021.7 (868.1) | 0.044 (0.004) |
| Cortical thickness (mm) | | | | |
| Entorhinal cortex | 6.7 (0.4) | 6.5 (0.4) | 6.7 (0.4) | 1.82e−7 (0.030) |
| Insula | 6.0 (0.2) | 5.9 (0.2) | 6.0 (0.2) | 8.11e−6 (0.022) |

Notes: η^2_p , partial eta-squared effect size; The group difference analyses were adjusted for age, gender, education, and BMI. Bold indicates $p < 0.05$.

TABLE 3 Interactions between race and MAP on cognitive test performance, gray matter and hippocampal volume, and entorhinal and insula cortex thickness

| | <i>F</i> (df) | <i>p</i> -value (η^2_p) |
|---------------------------------|------------------|--------------------------------|
| Cognitive tests | | |
| Total Cognitive Composite Score | 5.833 (1,943) | 0.015 (0.006) |
| Crystallized Composite Score | 4.324 (1,950) | 0.037 (0.004) |
| Early Cognitive Composite Score | 3.956 (1,947) | 0.046 (0.004) |
| Cognitive Fluid Composite Score | 3.548 (1,943) | 0.059 (0.003) |
| Brain volume | | |
| Total Gray Matter | 0.00006 (1, 894) | 0.993 (0.000) |
| Bilateral Hippocampus | 1.838 (1, 894) | 0.175 (0.001) |
| Cortical thickness | | |
| Entorhinal Cortex | 13.669 (1, 894) | 0.0002 (0.015) |
| Insula | 1.290 (1, 894) | 0.256 (0.001) |

Notes: df, degrees of freedom, η^2_p , partial eta-squared effect size; The interaction analyses were adjusted for age, gender, education, and BMI. Bold indicates $p < 0.05$.

($F[1,892] = 12.587$, $p = 0.0004$, $\eta^2_p = 0.013$) (Figure 2), no significant interactions were observed for total gray matter volume ($F[1,891] = 0.00002$, $p = 0.987$, $\eta^2_p = 2.92e-7$), bilateral hippocampal volume ($F[1,891] = 1.887$, $p = 0.169$, $\eta^2_p = 0.002$), and bilateral insula thickness ($F[1,892] = 0.982$, $p = 0.321$, $\eta^2_p = 0.001$). The p -value of the interaction for ERC thickness was lower than the FDR-corrected threshold ($p = 0.012$).

3.5 | Association between ERC thickness and cognitive test results

As the similar Race \times MAP interaction patterns were identified in cognitive test results and ERC thickness, we further explored the association between ERC thickness and cognitive test results using multivariable linear regression (adjusted for age, gender, and education). We performed this post-hoc analysis only in AAs because of the consistent negative associations between MAP and cognitive test outcomes and ERC thickness, which were not observed in CAs (Figures 1 and 2). These analyses in AAs showed that ERC thickness was not significantly associated with total cognitive composite score ($\beta = 0.119$, $p = 0.063$, 95% CI -0.324 , 12.335), crystallized composite score ($\beta = 0.057$, $p = 0.344$, 95% CI -2.784 , 7.930), or early cognitive composite score ($\beta = 0.131$, $p = 0.057$, 95% CI -0.146 , 10.341).

4 | DISCUSSION

After controlling for age, gender, education, and BMI, our results demonstrate that the markers of brain health between AA and CA younger adults were moderated by BP. As hypothesized, there was a significant Race \times MAP interaction on cognitive composite scores and ERC thickness, such that greater MAP was associated with lower cognitive performance and smaller ERC thickness, which was specific to AAs. The current analyses are consistent with, and expand on, previous evidence documenting that MAP-related associations with lower global cognitive function are stronger in older AAs than their older CA counterparts (Levine




FIGURE 1 Significant interaction between race and MAP on cognitive task performances (adjusted for age, gender, education, and BMI). The r , r^2 and p values indicate the correlation of MAP and cognitive test results for each racial group. The interaction statistical results are presented in Table 3. Notes: MAP, mean arterial pressure; mmHg, millimeters of mercury




FIGURE 2 Significant interaction between race and MAP on entorhinal cortex thickness (adjusted for age, gender, education, and BMI). The r , r^2 , and p values indicate the correlation of MAP and cognitive test results for each racial group. The interaction statistical result is presented in Table 3. Notes: ERC, entorhinal cortex; MAP, mean arterial pressure; mmHg, millimeter of mercury

et al., 2019). The prior finding utilized the Six-item Screener Scores (Callahan et al., 2002) as a primary cognitive outcome and three other cognitive tests including Alzheimer's disease word list learning, word list delayed recall, and animal fluency as secondary cognitive outcomes. The present results extend the previous study (Levine et al., 2019) by investigating the significant race by MAP interactions on cognitive composite scores measured using a comprehensive neuropsychological test battery. Although the association between cognitive performance and hypertension in younger adults has been documented (Suhr et al., 2004), the present findings provide novel insight to suggest BP may moderate cognitive performance differently in younger AA compared to CA. Given long-term exposure to higher BP from young to mid-adulthood accelerates cognitive decline in midlife (Mahinrad et al., 2020), our findings indicate that the racial disparity in higher BP-associated cognitive function may begin in early adulthood. While our results should be interpreted with caution as the significant interactions on cognitive outcomes did

not survive FDR correction, the consistency of the magnitude and direction of these effects gives confidence that these findings are not merely spurious, and that accepting the null hypothesis would result in a Type II error. Notably, despite the small effect size for these associations, the fact that they were observed in a younger and normotensive cohort has implications for long-term brain health, as it is plausible that disparity between the AAs and CAs may only get stronger with age.

It has been suggested that the differences in hypertension between the race groups are relevant to a higher prevalence of obesity and cardiovascular diseases in AAs (Pan et al., 2009). In support of these prior findings, we observed AAs had a significantly greater average body mass index (BMI) than CAs. Furthermore, the average BMI ($29.1 \pm 6.5 \text{ kg/m}^2$) of AAs was just below the typical cut-off for the obese category ($>30 \text{ kg/m}^2$). Despite BMI's significant limitations as an accurate index for body composition (Frankenfield et al., 2001), it provides epidemiological insights on this critical

demographic difference between race groups. Another possible physiological contributor to our findings may be genetic differences between race groups related to kidney function. AAs are more likely to experience kidney failure and disease compared to CAs (Harding et al., 2017). Indeed, a recent genome-wide association meta-analysis suggested a genetic link between kidney system function and risk of Alzheimer's disease in AAs (Kunkle et al., 2021). Given the strong relationship between kidney function and BP (Young et al., 2002), genetic differences linked with kidney function between the race groups might have been associated with the interaction between race and MAP in the present study. Unfortunately, these genetic and kidney-related biomarkers are not available in the HC database we analyzed. More importantly, the socio-economic and/or educational disadvantages (e.g., less access to healthcare, educational and cognitively enriching opportunities) in AAs are also associated and potentially contribute to the racial disparity in development of chronic diseases (Frieden, 2013). Although we have statistically accounted for education, we were unable to adequately address these important socio-economic determinants in the present study.

Lower insula thickness in older hypertensive in AAs relative to their age-matched CA counterparts has been documented in a previous study (Chand et al., 2017). In partial agreement with this finding, we have observed lower insula cortex thickness in younger AAs compared to CAs. However, there was no significant interaction between MAP and race on the insula thickness. We also tested the interactive effects on the hippocampus, a brain area crucial for memory, because higher BP is linked with lower hippocampal perfusion (Zola-Morgan et al., 1986) and reduced hippocampal volume (Beauchet et al., 2013). Higher BP-related lower hippocampal volume, however, was not detected in the present study. Although there were no significant effects on hippocampal and insula thickness, we discovered a Race \times MAP interaction on ERC thickness. One of the earliest changes in neurodegenerative diseases occur within the ERC, and its shrinkage serves as a predictor of future cognitive decline (Du et al., 2001). Age-related dysfunction in ERC has also been shown to be affected by hypertension (Raz, 2000). Yet, no previous work has corroborated BP-related differences in ERC thickness patterns between AAs and CAs in older adults. Taken together, these results indicate that BP-related brain changes may differ by age, such that while the hippocampal and insula may not be impacted by BP in younger adults, higher MAP-related lower ERC cortical thickness may begin during younger adulthood in AAs. Moreover, although only at borderline statistical significance, there were moderate associations between lower ERC thickness and lower cognitive test scores in AAs, further suggesting that ERC cortical thinning may play a role in the relationship between BP and cognitive performance in younger AA individuals.

4.1 | Strengths and Limitations

Prior evidence studying older adults used a limited number of cognitive assessments (Levine et al., 2019) or subject cognitive deficit questionnaires (Hajjar et al., 2016) to determine the BP-related racial disparity on cognitive function. Conversely, the present study employed composite scores from a comprehensive neuropsychological battery to robustly test a wide range of cognitive domains from memory to executive function. We also incorporated the HC Project's high-resolution neuroimaging data into this analysis to elucidate the possible mechanism deriving BP-related disparity in cognitive performance between the race groups.

Despite these strengths, there are several limitations to this study that warrant caution when interpreting the results. There was an unequal number of participants between the race groups. However, a bootstrapping analysis provided nearly identical results, suggesting the sample sizes were not an issue. To gain mechanistic insights, future studies need to utilize another physiological measurement such as systemic inflammation, neurotrophic factors/biomarkers, cerebral blood flow regulation, and consider potential genetic influences. Furthermore, beyond BP, future studies need to examine aortic pulse wave velocity which is the gold-standard technique to investigate arterial stiffness of the large central arteries. The aortic stiffening and associated augmented blood flow pulsatility are closely related to impairment in brain structure and cognitive function (Barnes & Corkery, 2018). Thus, compared to using BP, analyzing aortic pulse wave velocity will advance our understanding of the relationship between vascular structure and brain health. The current study is also subject to the limitations of testing a healthy and normotensive population; thus, younger clinical populations with chronic hypertensive conditions may not demonstrate the significant interactions between race and BP on cognitive and brain measurements found in the present study. Also, since this cross-sectional design limits interpreting the directionality of the current results, further longitudinal studies are necessary to fully understand the moderating effects of BP on long-term brain health. Last but not least, we acknowledge that the present investigation is constrained by the HC Project methods and database. Unfortunately, data were not available to explore the social-cultural factors of race (e.g., income, occupation, zip code, perceived stress, parental support, quality and quantity of education, and exposure to racial discrimination or other traumatic events). Indeed, previous evidence suggests that racial discrimination (Calvin et al., 2003; Coogan et al., 2020) and traumatic events (Edmondson & Känel, 2017; SIVERS GHBAH., 1998) significantly influence the cardiovascular and cognitive health. However, we were not able to account for these factors in our analyses. Nevertheless, we did find that

there were no significant differences in depression or anxiety symptoms between AAs and CAs in the present study, suggesting these mental health-related outcomes may not be associated with the interactions we observed.

5 | CONCLUSIONS

In conclusion, our findings indicate that a higher MAP is associated with lower cognitive performance in younger AAs, but not in CAs. This suggests that AAs, from early adulthood, may carry a greater risk of BP-related late-life cognitive decline. Furthermore, the structural integrity of the ERC, a key region in medial temporal lobe networks related to memory, may be an important mediator or moderator of blood pressure-related cognitive function in younger AA adults. Although the effects we observed were relatively small, the BP-related differences in brain health between AAs and CAs were apparent in younger healthy cohort which may accumulate over time and most likely to have long-term consequences in older people. The present study is limited in its ability to elucidate the underlying social and physiological mechanisms underlying this BP-related racial disparity. Further efforts are needed to identify these underlying mechanisms and thereby develop practical and effective interventions to mitigate racial differences in cardiovascular and brain health.

ACKNOWLEDGMENTS

Data were provided by the Human Connectome Project, WU-Minn Consortium (Principal Investigators: Drs. Van Essen and Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at Washington University. We thank Human Connectome Project Team for sharing the data, and the participants for their time as volunteers.

CONFLICT OF INTEREST

The authors have no potential conflicts of interest to report.

AUTHOR CONTRIBUTIONS

JW, SMR, and JCS conceived and designed the study. JW, SC, and DDC conducted the analyses. JW wrote the paper, and all authors edited the paper.

ORCID

J. Carson Smith  <https://orcid.org/0000-0001-6142-0920>

REFERENCES

Achenbach, T. M., & Rescorla, L. A. (2003) *Manual for the ASEBA adult forms & profiles.* : University of Vermont, Research Center for Children, Youth.

- Barnes, J. N., & Corkery, A. T. (2018). Exercise improves vascular function, but does this translate to the brain? *Brain Plasticity*, 4(1), 65–79.
- Beauchet, O., Celle, S., Roche, F., Bartha, R., Montero-Odasso, M., Allali, G. & Annweiler, C. (2013). Blood pressure levels and brain volume reduction: A systematic review and meta-analysis. *Journal of Hypertension*, 31(8), 1502–1516.
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B (Statistical Methodology)*, 57(1), 289–300.
- Bohannon, A. D., Fillenbaum, G. G., Pieper, C. F., Hanlon, J. T., & Blazer, D. G. (2002). Relationship of race/ethnicity and blood pressure to change in cognitive function. *Journal of the American Geriatrics Society*, 50(3), 424–429.
- Braak, H., & Braak, E. (1991). Neuropathological staging of Alzheimer-related changes. *Acta Neuropathologica*, 82(4), 239–259.
- Callahan, C. M., Unverzagt, F. W., Hui, S. L., Perkins, A. J., & Hendrie, H. C. (2002). Six-item screener to identify cognitive impairment among potential subjects for clinical research. *Medical Care*, 40(9), 771–781.
- Calvin, R., Winters, K., Wyatt, S. B., Williams, D. R., Henderson, F. C., & Walker, E. R. (2003). Racism and cardiovascular disease in African Americans. *American Journal of the Medical Sciences*, 325(6), 315–331.
- Chand, G. B., Wu, J., Qiu, D., & Hajjar, I. (2017). Racial differences in insular connectivity and thickness and related cognitive impairment in hypertension. *Frontiers in Aging Neuroscience*, 9, 177.
- Coogan, P., Schon, K., Li, S., Cozier, Y., Bethea, T., & Rosenberg, L. (2020). Experiences of racism and subjective cognitive function in African American women. *Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring*, 12(1), e12067.
- Den Heijer, T., Launer, L. J., Prins, N. D., Van Dijk, E. J., Vermeer, S. E., Hofman, A., Koudstaal, P. J., & Breteler, M. (2005). Association between blood pressure, white matter lesions, and atrophy of the medial temporal lobe. *Neurology*, 64(2), 263–267.
- Du, A. T., Schuff, N., Amend, D., Laakso, M. P., Hsu, Y. Y., Yaffe, K., Kramer, J. H., Reed, B., Norman, D., Chui, H. C., & Weiner, M. W. (2001). Magnetic resonance imaging of the entorhinal cortex and hippocampus in mild cognitive impairment and Alzheimer's disease. *Journal of Neurology, Neurosurgery and Psychiatry*, 71(4), 441–447.
- Edmondson, D., & vonKänel, R. (2017). Post-traumatic stress disorder and cardiovascular disease. *Lancet Psychiatry*, 4(4), 320–329.
- Faraco, G., & Iadecola, C. (2013). Hypertension: A harbinger of stroke and dementia. *Hypertension*, 62(5), 810–817.
- Fischl, B. (2012). FreeSurfer. *NeuroImage*, 62(2), 774–781.
- Fischl, B., & Dale, A. M. (2000). Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proceedings of the National Academy of Sciences*, 97(20), 11050–11055.
- Frankenfield, D. C., Rowe, W. A., Cooney, R. N., Smith, J. S., & Becker, D. (2001). Limits of body mass index to detect obesity and predict body composition. *Nutrition*, 17(1), 26–30.
- Frieden, T. R. (2013). CDC health disparities and inequalities report—United States, 2013. Foreword. *MMWR Supplement*, 62(3), 1–2.
- Gautam, P., Cherbuin, N., Sachdev, P. S., Wen, W., & Anstey, K. J. (2013). Sex differences in cortical thickness in middle aged and early old-aged adults: Personality and total health through life study. *Neuroradiology*, 55(6), 697–707.

- Glasser, M. F., Smith, S. M., Marcus, D. S., Andersson, J. L., Auerbach, E. J., Behrens, T. E., Coalson, T. S., Harms, M. P., Jenkinson, M., Moeller, S., Robinson, E. C., Sotiropoulos, S. N., Xu, J., Yacoub, E., Ugurbil, K., & Van Essen, D. C. (2016). The human connectome project's neuroimaging approach. *Nature Neuroscience*, *19*(9), 1175.
- Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L., Xu, J., Jbabdi, S., Webster, M., Polimeni, J. R., Van Essen, D. C., Jenkinson, M., & Wu-Minn HCP Consortium. (2013). The minimal preprocessing pipelines for the Human Connectome Project. *NeuroImage*, *80*, 105–124.
- Gorelick, P. B., & Nyenhuis, D. (2012). Blood pressure and treatment of persons with hypertension as it relates to cognitive outcomes including executive function. *Journal of the American Society of Hypertension*, *6*(5), 309–315.
- Gur, R. C., Richard, J., Hughett, P., Calkins, M. E., Macy, L., Bilker, W. B., Brensinger, C., & Gur, R. E. (2010). A cognitive neuroscience-based computerized battery for efficient measurement of individual differences: standardization and initial construct validation. *Journal of Neuroscience Methods*, *187*(2), 254–262.
- Hajjar, I., Wharton, W., Mack, W. J., Levey, A. I., & Goldstein, F. C. (2016). Racial disparity in cognitive and functional disability in hypertension and all-cause mortality. *American Journal of Hypertension*, *29*(2), 185–193.
- Harding, K., Mersha, T. B., Vassalotti, J. A., Webb, F. J., & Nicholas, S. B. (2017). Current state and future trends to optimize the care of chronic kidney disease in African Americans. *American Journal of Nephrology*, *46*(2), 176–186.
- Iadecola, C., & Gottesman, R. F. (2019). Neurovascular and cognitive dysfunction in hypertension: epidemiology, pathobiology, and treatment. *Circulation Research*, *124*(7), 1025–1044.
- Korf, E. S., White, L. R., Scheltens, P., & Launer, L. J. (2004). Midlife blood pressure and the risk of hippocampal atrophy: The Honolulu Asia Aging Study. *Hypertension*, *44*(1), 29–34.
- Kunkle, B. W., Schmidt, M., Klein, H.-U., Naj, A. C., Hamilton-Nelson, K. L., Larson, E. B., Evans, D. A., De Jager, P. L., Crane, P. K., Buxbaum, J. D., Ertekin-Taner, N., Barnes, L. L., Fallin, M. D., Manly, J. J., Go, R. C. P., Obisesan, T. O., Kambh, M. I., Bennett, D. A., Hall, K. S., & Kukull, W. A. (2021). Novel Alzheimer disease risk loci and pathways in African American individuals using the African genome resources panel: a meta-analysis. *JAMA Neurology*, *78*(1), 102–113.
- Levine, D. A., Galecki, A. T., Langa, K. M., Unverzagt, F. W., Kabeto, M. U., Giordani, B., Cushman, M., McClure, L. A., Safford, M. M., & Wadley, V. G. (2019). Blood pressure and cognitive decline over 8 years in middle-aged and older black and white Americans. *Hypertension*, *73*(2), 310–318.
- Levine, D. A., Gross, A. L., Briceño, E. M., Tilton, N., Kabeto, M. U., Hingtgen, S. M., Giordani, B. J., Sussman, J. B., Hayward, R. A., Burke, J. F., Elkind, M. S. V., Manly, J. J., Moran, A. E., Kulick, E. R., Gottesman, R. F., Walker, K. A., Yano, Y., Gaskin, D. J., Sidney, S., ... Galecki, A. T. (2020). Association between blood pressure and later-life cognition among black and white individuals. *JAMA Neurology* *77*(7):810–819.
- Mahinrad, S., Kurian, S., Garner, C. R., Sedaghat, S., Nemeth, A. J., Moscufo, N., Higgins, J. P., Jacobs Jr, D. R., Hausdorff, J. M., Lloyd-Jones, D. M., & Sorond, F. A. (2020). Cumulative blood pressure exposure during young adulthood and mobility and cognitive function in midlife. *Circulation*, *141*(9), 712–724.
- Menon, V., & Uddin, L. Q. (2010). Saliency, switching, attention and control: A network model of insula function. *Brain Structure & Function*, *214*(5), 655–667.
- Morillas, P., deAndrade, H., Castillo, J., Quiles, J., Bertomeu-González, V., Cordero, A., Tarazón, E., Roselló, E., Portolés, M., Rivera, M., & Bertomeu-Martínez, V. (2012). Inflammation and apoptosis in hypertension. Relevance of the extent of target organ damage. *Revista Española De Cardiología*, *65*(9), 819–825.
- Muller, M., van derGraaf, Y., Visseren, F. L., Mali, W. P. T. M., & Geerlings, M. I.; Group SS (2012). Hypertension and longitudinal changes in cerebral blood flow: The SMART-MR study. *Annals of Neurology*, *71*(6), 825–833.
- Pan, L., Galuska, D. A., Sherry, B., Hunter, A. S., Rutledge, G. E., Dietz, W. H., & Balluz, L. S. (2009). Differences in prevalence of obesity among black, white, and Hispanic adults—United States, 2006–2008. *Morbidity and Mortality Weekly Report*, *58*(27), 740–744.
- Raz, N. (2000). Aging of the brain and its impact on cognitive performance: Integration of structural and functional findings. In F.I.M.Craik & T.A.Salthouse (Eds.), *The handbook of aging and cognition* (pp. 1–90). Lawrence Erlbaum Associates Publishers.
- Redmond, N., Baer, H. J., & Hicks, L. S. (2011). Health behaviors and racial disparity in blood pressure control in the national health and nutrition examination survey. *Hypertension*, *57*(3), 383–389.
- Reuben, D. B., Magasi, S., McCreath, H. E., Bohannon, R. W., Wang, Y.-C., Bubela, D. J., Rymer, W. Z., Beaumont, J., Rine, R. M., Lai, J. S., & Gershon, R. C. (2013). Motor assessment using the NIH toolbox. *Neurology*, *80*(11 Suppl 3), S65–S75.
- Savic, I., & Arver, S. (2014). Sex differences in cortical thickness and their possible genetic and sex hormonal underpinnings. *Cerebral Cortex*, *24*(12), 3246–3257.
- Saxby, B. K., Harrington, F., McKeith, I. G., Wesnes, K., & Ford, G. A. (2003). Effects of hypertension on attention, memory, and executive function in older adults. *Health Psychology*, *22*(6), 587.
- Seo, S. W., Lee, J.-M., Im, K., Park, J.-S., Kim, S.-H., Kim, S. T., Ahn, J. H., Kim, M.-J., Kim, G. H., Kim, J. H., Roh, J. H., Cheong, H.-K., & Na, D. L. (2012). Cardiovascular risk factors cause cortical thinning in cognitively impaired patients: relationships among cardiovascular risk factors, white matter hyperintensities, and cortical atrophy. *Alzheimer Disease and Associated Disorders*, *26*(2), 106–112.
- SIVERS GHBAH. (1998). Cognitive impact of traumatic events. *Development and Psychopathology*, *10*, 625–653.
- Suhr, J. A., Stewart, J. C., & France, C. R. (2004). The relationship between blood pressure and cognitive performance in the Third National Health and Nutrition Examination Survey (NHANES III). *Psychosomatic Medicine*, *66*(3), 291–297.
- Van Essen, D. C., Smith, S. M., Barch, D. M., Behrens, T. E., Yacoub, E., Ugurbil, K., & WU-Minn HCP Consortium (2013). The WU-Minn human connectome project: an overview. *NeuroImage*, *80*, 62–79.
- Van Essen, D. C., Ugurbil, K., Auerbach, E., Barch, D., Behrens, T. E. J., Bucholz, R., Chang, A., Chen, L., Corbetta, M., Curtiss, S. W., Penna, S. D., Feinberg, D., Glasser, M. F., Harel, N., Heath, A. C., Larson-Prior, L., Marcus, D., Michalareas, G., Moeller, S., & WU-Minn HCP Consortium. (2012). The Human Connectome Project: A data acquisition perspective. *NeuroImage*, *62*(4), 2222–2231.
- Weintraub, S., Dikmen, S. S., Heaton, R. K., Tulsky, D. S., Zelazo, P. D., Bauer, P. J., Carlozzi, N. E., Slotkin, J., Blitz, D., Wallner-Allen, K., Fox, N. A., Beaumont, J. L., Mungas, D., Nowinski, C. J., Richler, J., Deocampo, J. A., Anderson, J. E., Manly, J. J., Borosh, B., Havlik, R., & Gershon, R. C. (2013). Cognition assessment using the NIH Toolbox. *Neurology*, *80*(11 Suppl 3), S54–S64.
- Won, J., Alfini, A. J., Weiss, L. R., Nyhuis, C. C., Spira, A. P., Callow, D. D., & Smith, J. C. (2019). Caudate volume mediates the interaction

between total sleep time and executive function after acute exercise in healthy older adults. *Brain Plasticity*, 5(1), 69–82.

- Young, J. H., Klag, M. J., Muntner, P., Whyte, J. L., Pahor, M., & Coresh, J. (2002). Blood pressure and decline in kidney function: Findings from the Systolic Hypertension in the Elderly Program (SHEP). *Journal of the American Society of Nephrology*, 13(11), 2776–2782.
- Zola-Morgan, S., Squire, L. R., & Amaral, D. G. (1986). Human amnesia and the medial temporal region: enduring memory impairment following a bilateral lesion limited to field CA1 of the hippocampus. *Journal of Neuroscience*, 6(10), 2950–2967.

How to cite this article: Won J, Ranadive SM, Callow DD, Chen S, Carson Smith J. Blood pressure-related differences in brain health between young African Americans and Caucasian Americans. *Physiol Rep*. 2021;9:e14819. <https://doi.org/10.14814/phy2.14819>