

The Use of an Iterative Strategy of Cognitive Interview and Expert Consultation to Revise the Quality of Life Scale for Patients with Aplastic Anemia (QLS-AA)

Min Xu^{1,*}, Ting Liu^{2,*}, Menghua Ye¹, Xiaoxue Tan¹, Qiu Hua Sun²

¹The First Affiliated Hospital of Zhejiang Chinese Medical University, Hangzhou, People's Republic of China; ²The School of Nursing, Zhejiang Chinese Medical University, Hangzhou, People's Republic of China

*These authors contributed equally to this work

Correspondence: Qiu Hua Sun, Zhejiang Chinese Medical University, The School of Nursing, 548 Binwen Road, Binjiang District, Hangzhou City, Zhejiang Province, People's Republic of China, Email sunqiu Hua@zcmu.edu.cn

Background: Aplastic anemia is characterized by anemia, hemorrhage and infection, and is accompanied by a variety of complications and psychological burden. Therefore, the quality of life of AA patients is not optimistic. Our team is committed to developing an assessment tool for the quality of life of AA patients, and adopting an iterative strategy of cognitive interview and expert consultation to solve the challenges encountered in item revision.

Purpose: We aim to use the strategy of cognitive interview and expert consultation to inform revision of the QLS-AA into a user-friendly tool with unambiguous items and improve the content validity of the scale.

Methods: We used an iterative strategy of cognitive interview and expert consultation. Two rounds of cognitive interview were conducted to identify problems with item comprehension, recall and other cognitive processes. As well as, a multi-disciplinary group of expert consultation was consulted to review the rationality of item revisions.

Results: In the first round of cognitive interview, 16 participants responded to 107 items. Among them, the most common problems were “clarification” and “item duplication”. Based on the results of the first round of interview, an expert consultation was organized. A total of 16 amendments were put forward by the expert and 14 were adopted. In the second round of cognitive interviews, A total of 5 participants were included and 64 items were evaluated. Two items were suggested to be revised, and the remaining items were accurately understood and recognized by all participants.

Conclusion: This study highlights the key issues to consider when incorporating patient perspectives into quality measurement. The revision of QLS-AA through the strategy of cognitive interview and expert consultation may provide valuable insights into the measurement of quality of life in aplastic patients.

Trial Registration Number: ChiCTR2100047575.

Keywords: cognitive interview, expert consultation, aplastic anemia, quality of life

Introduction

Aplastic anaemia (AA) is a rare, life-threatening and heterogeneous disorder. It is defined as pancytopenia with a hypocellular bone marrow in the absence of an abnormal infiltrate or marrow fibrosis.¹ The excessive apoptosis of hematopoietic stem cells caused by cytotoxic T cells and/or lymphokines is the main pathogenesis of acquired AA.² The incidence of AA in the United States and Europe is below 2.5/million, while the incidence of AA in Asia is 2–3 times higher.^{3,4} In China, the annual incidence of AA is 7.4/10⁵ population, and has a biphasic distribution, with peaks at 10–25 years and over 60 years.² Anemia, hemorrhage and infection are the initial manifestations of AA, and with the

development of the disease, the clinical symptoms are gradually worsen, which may be accompanied by refractory anemia, multi-organ hemorrhage and sepsis, which pose a great threat to their life and health.^{3,5}

Treatment plans for AA include hematopoietic stem cell transplantation (HSCT), immunosuppressive therapy (IST), androgen-based hematopoietic therapy and new attempts of new drugs for refractory AA.⁶ In recent years, with the development of new drugs, the progress of HSCT technology and the maturity of IST treatment, more and more patients can obtain the best treatment plan, and the survival time of AA patients has been significantly prolonged. Studies have shown that the 5-year survival rate of patients with severe aplastic anemia treated with HSCT or IST can reach more than 80%, while the 9-year survival rate of patients treated with HSCT is as high as 87–89%.⁷ However, aplastic therapy is often accompanied by side effects such as decreased physical function, fatigue, diarrhea, nausea and vomiting, oral ulcer, hair loss, rash and respiratory problems. In addition, patients face a variety of risks such as disease recurrence, secondary clonal disease, infection, graft-versus-host disease (GVHD) and so on.⁸ Therefore, AA has the characteristics of long treatment cycle, repeated disease changes and heavy disease burden, resulting in about 15–35% of AA patients with depression and anxiety,⁹ and even serious consequences such as giving up treatment and suicide.¹⁰ It can be seen that as the survival time of AA patients increases, the quality of life of AA patients should no longer be neglected. Reliable assessment tools play a fundamental role in the study of the quality of life level and support strategies of AA patients, but there is no mature and specific assessment tool that can be used at present.

We have been committed to develop the Quality of Life Scale for Patients with Aplastic Anemia (QLS-AA). In the preliminary work, we obtained the item pool and formed the pretest scale through literature study and qualitative interviews.¹¹ The test approaches of the pretest scale include focus groups, cognitive interviewing and expert consultation.^{12,13} Cognitive interview is a methodology based on the interdisciplinary field termed cognitive aspects of survey methodology (CASM), initiated in the 1980s, which involves theories and results from studies on the cognitive processes involved in survey responding.¹⁴ Cognitive testing is designed to identify unobservable problems in item comprehension, recall, and other cognitive processes that can be remediated through question rewording, reordering, or more extensive instrument revision. Cognitive interview techniques were developed mainly for face-to-face interviews, and the interviews rely on intensive verbal probing of volunteer participants by a specially trained interviewer.¹⁴ It has been widely used to guide the revision and cultural adjustment of scales, the development of questionnaires and other self-report measures.^{15,16} Expert consultation is defined as a method in which some experts with highly specialized technology or experience provide personalized technical assistance, which is a core component of many research projects and can reflect our attention to special issues.¹⁷ The participation of clinical experts can provide efficient information, and expert consultation is adopted when the information is uncertain.¹⁸ However, expert consultation also has shortcomings such as strong subjectivity and empiricism, so it is mainly used to the early stage of decision-making or the soliciting opinions.

We used an iterative strategy of cognitive interview and expert consultation to reduce errors that could affect measurement validity, and included preferences for layout of the measurement.¹³ Our objective was to use the cognitive testing to identify otherwise unobservable problems with item comprehension, recall, and other cognitive processes that can be remediated through question rewording, reordering or more extensive instrument revision. We also hope to use the results of cognitive interview and expert consultation to reduce factors that may affect content validity, including differences in perceptions and patient preferences.

Methods

Participants and Recruitment Procedure

The purposive sampling method and maximum variance strategy were used. The sample size was calculated according to the recommendations of the cognitive interview.¹⁹ The first round of interviews was based on the criterion that each item was evaluated by at least 5 participants with no more than 35 items per evaluation. The second round of interviews usually requires 3–5 participants. We divided the pretest scale into 3 sets in a balanced manner to facilitate completion. AA patients who were admitted to the Department of Hematology of the First Affiliated Hospital of Zhejiang Chinese Medical University from August to October 2021 were selected as interview subjects.

Inclusion Criteria

In line with the diagnostic criteria for aplastic anemia proposed in the Standards for Diagnosis and Efficacy of Hematological Diseases;²⁰ age ≥ 18 years; the patient is aware of the diagnosis and development of the disease; has basic reading and writing skills and able to independently fill in our questionnaire; interested in participating in this research.

Exclusion Criteria

People with communication disorders; those with severe changes in heart, lung, liver, and kidney functions; those with previous mental disorders. The ethical principles of the Declaration of Helsinki were followed in this study. We actively briefed the participants on the purpose and significance of the study, and they were assured that their responses would be anonymous and their personal information confidential until they signed the informed consent form. Ethics approval was obtained from the Ethics Committee of the First Affiliated Hospital of Zhejiang Chinese Medical University (Ethical approval ID: 2021-KL-055-01).

Formation of the Pretest Scale

QLS-AA is based on the concept category of quality of life proposed by WHO and the theoretical framework of quality of life proposed by Dr. Ferrell.^{21,22} Combined with cultural background and clinical environment, our research identified a pretest scale with 107 items. We used an iterative strategy of cognitive interview and expert consultation to reduce errors that could compromise a measures validity (Figure 1).

The Outline of the Interview

Cognitive interviews are based on the cognitive theory proposed by Tourangeau,²³ in which the response process is divided into four components: comprehension (attend to questions and instructions, represent logical form of question, identify question focus, link key terms to relevant concepts); retrieval (generate retrieval strategy and cues, retrieve specific/generic memories, fill in missing details); judgment (assess completeness and relevance of memories, draw inferences based on accessibility).^{14,24} Table 1 illustrates domains, example questions and corresponding cognitive interview probes.

Data Collection

Before the interview, explained the research purpose and privacy protection to the participants in detail, signed the informed consent form and completed the general information questionnaire and QLS-AA subscale. Semi-structured interviews were administered throughout the process of interview, and the whole interview would be recorded. Verbal-probing technique was used to encourage open-ended dialogue.²⁵ Participant responses were documented during the interview using a predefined data collection sheet.



Figure 1 Schematic diagram of the iterative strategy of cognitive interview and expert consultation.

Table 1 Domains and Examples of Draft Items and Cognitive Interview Probes

Domains	Draft Items		Cognitive Interviews	
	Example Question	Response Categories	Example Probe	Probe Purpose
Physical	Hirsutism/voice changes due to medication have troubled my life.	1= Not at all 2= A little bit 3= Somewhat 4= Quite a bit 5= Very much	Please tell me in your own words what this question is asking?	Test comprehension
Psychological	In the past month, I have been feeling anxious/nervous/ distraught.	1= Never 2= Rarely 3= Sometimes 4= Often 5= Always	Are there any words that you do not understand in this item?	Test question clarity
Social relationships	Aplastic anemia interfered with my daily social activities.	1= Not at all 2= A little bit 3= Somewhat 4= Quite a bit 5= Very much	Can you talk about why you came up with this answer?	Elicit participant thought process
Spirituality	Faith in my heart comforts and encourages me most of the time.	1= Not at all 2= A little bit 3= Somewhat 4= Quite a bit 5= Very much	Was it easy or difficult for you to choose this answer?	Test rationality of question-answer
Environment	I need more ways to communicate with fellow patients.	1= Never 2= Rarely 3= Sometimes 4= Often 5= Always	I noticed you hesitated before your response, what were your thoughts?	Identify source of potential confusion

Data Analysis

The content analysis method was used to review the responses of the participants and an independent summary was formed.²⁶ Qualitative data analysis is based on hermeneutic, which was conducted within cognitive interviews framework. In 1808, Friederich proposed a circular approach to hermeneutics whereby the foundations of knowledge and understanding was to find the spirit of the whole through the individual, and to grasp the individual through the whole.²⁷ The hermeneutic circle has become the most resonant idea in hermeneutic theory, which is concerned with the dynamic relationship between the part and the whole.²⁸ The questions were independently classified by 2 researchers, and the questionable items were initially revised through special discussion to ensure validity and reliability. The revised items, especially controversial or questionable ones, would be further discussed and reviewed through expert consultation, and then proceed to the next round of cognitive interviews until all items could be understood by the participants, that is, the data is at a relatively saturated state.

Results

Characteristics of Cognitive Interview Participants and Experts

2 rounds of cognitive interviews and expert consultation have been completed. The first round of CI included 16 participants (1-S1~16), and the second round of CI included 5 participants (2-S17~21). According to the research purpose and the representativeness of the experts, 6 experts (E1-6) with multidisciplinary backgrounds were identified as the axis members of the expert consultation group. The characteristics of the participants are shown in [Table 2](#). More information on the specialist background and self-assessment of these experts is provided in [Supplementary 1](#).

Table 2 Characteristics of Cognitive Interview Participants and Experts

Characteristics		Participants (n= 21)	Experts (n=6)
Male		71% (n=15)	16.7% (n=1)
Mean age in years		31.8±11.3	41.3±9.0
Education			
	High school or less	52.4% (n=11)	/
	College graduate	38.1% (n=8)	50% (n=3)
	Master degree or above	9.5% (n=3)	50% (n=3)
Work experience (Median, IQR)		/	12.5 (10 to 21)
Disease course (Median, IQR) ^a		12.5 (6 to 37)	/
Anemia			
	Yes	76.2% (n=16)	/
	No	23.8% (n=5)	
Treatment regimen received			
	Transfusion therapy	95.2% (n=20)	/
	Cyclosporin treatment	85.7% (n=18)	
	TCM treatment ^b	71.4% (n=15)	
	Androgen therapy	61.9% (n=13)	
	HSCT therapy	47.6% (n=10)	
	Eltrombopag treatment	42.9% (n=9)	
Pecuniary condition			
	Completely independent	19.1% (n=4)	/
	Partial independence	19.1% (n=4)	
	Completely dependent	61.9% (n=13)	

Notes: ^aThe course of the disease is measured in month. ^bTCM treatment means Traditional Chinese medicine treatment.

Results of the First Round of Cognitive Interviews

The first round of cognitive interviews involved 16 patients and 107 draft items were tested, and the total interview time was about (28.88±10.18) mins. Item problems identified as “clarity” and “response categories” (vague and overlapping) were the most common types of specific question. In addition, 4 items were reflected by participants as diversified response processes and even the views are contradictory, while the linear problem coding is not conducive to understanding the problems, or even misleading the modification of the items. Therefore, we temporarily categorize it as “other” and carry out a further interpretative analysis in expert consultation along with preliminary revised terms. Table 3 shows examples of the analysis of those items and their preliminary revisions.

Results of the First Round of Expert Consultation

The expert consultation meeting reviewed the preliminary revisions undertaken by the researchers after the first round of cognitive interviews. 6 experts put forward a total of 16 revision opinions, of which 11 were revised, 2 were merged, and 4 were added (Supplementary 2). Subsequently, the 4 items with diversified response process were discussed and analyzed, finally a consensus revision was reached, of which 3 items were revised and 1 item was deleted.

Results of the Second Round of Cognitive Interviews

Items verified by expert consultation enter the second round of cognitive interviews. A total of 5 participants were involved and 64 items were evaluated by cognitive interview. The results showed that 1 item was suggested to modify by one participants, but it was not adopted after discussion by the research group (Table 4). There are no problems with other items in terms of response process.

Table 3 Examples of Analysis of Item-Specific Problems and Its Revision

Problem Coding		Example		
		Original Item	Problem Analysis	Revised Item
Clarity	Wording	Aplastic anemia and treatment have reduced my appetite and taste.	I-S1, I-S2: The expressions of “loss of appetite” and “loss of taste” are similar, which is easy to confuse.	Aplastic anemia and treatment made me feel poor appetite and/or diminished taste.
	Technical term	Drug-induced facial acne/hyperpigmentation has affected my life.	I-S2, I-S5: Does not understand the concepts of “facial acne” and “hyperpigmentation”.	Drug-induce pimple/skin dull have affected my life.
Knowledge/recall		I have some defecation problems (such as: diarrhea, constipation).	I-S15: Defecation problems cannot fully recall the entire process.	During the last 2 weeks, I have some defecation problems (such as diarrhea, constipation).
Response categories	Vague	Aplastic anemia and treatment make it difficult for me to climb stairs.	I-S14: I can climb stairs, but my physical strength is poor, and I can only climb two or three floors (stairs). I-S10: The height of the stairs should be limited.	Aplastic anemia and treatment make it difficult for me to climb stairs below three floors.
	Overlapping	Mouth ulcers making life difficult for me Bleeding of the oral mucosa or gums brings trouble to my life	I-S11, I-S16: Oral ulcers, gum bleeding, and gingival hyperplasia are all oral problems caused by aplastic anemia and treatment.	Oral problems (ulcers, infections, bleeding) bring trouble to my life.
Others	Diversification	(interpretative analysis in expert consultation)		

Table 4 Opinions and Amendments

Original Items	Opinions	Adoption	Reason
I am worried about taking medicine for the rest of my life.	2-S2 believes that the item is too colloquial.	Not taken	The colloquial expression is conducive to easier understanding.

Results of Second Round of Expert Consultation

Following the second round of expert consultation, there was a consensus amongst all experts that all revisions were appropriate. Assess consensus by inviting experts to comment on all items individually, and particular attention was paid to obtaining comments from experts who were less active. We believe that the data has reached relative saturation and the Pre-QLS-AA scale is ready for further testing.

Discussion

Cognitive interview is helpful to study test response processes, improve survey construction and map cognitive models of complex thought processes.²⁹ As we expected, cognitive testing identified otherwise unobservable problems in item comprehension, recall, and other cognitive processes that could be remediated by question rewording, reordering, or more extensive instrument revision, which largely contributed to the improvement of QLS-AA.

Approaches to qualitative analysis tend to be described in a linear, step by step manner, and cognitive interview is no exception. Classical cognitive interviews provide a unique way to capture participants’ cognitive process, and their structured questioning and linear analysis approach suffice for most studies to test response processes.³⁰ However, classical cognitive interviews are challenged in increasingly diverse cognitive process, contradictory opinions and some

sensitive health topics.³¹ Hermeneutics provides important theoretical insights into cognitive interviews, and provides a dynamic, non-linear style of thinking that can take a part-to-whole perspective on complex content and diverse responses to meet the specific needs of similar research.³² This kind of technical assistance is realized in the form of expert consultation. In particular, a key tenet of expert consultation is that the process of analysis is iterative, where we may move back and forth through a range of different ways of thinking about the data, rather than completing each step one after the other. In this work, we describe the application of the integrated cognitive interviews in the Pre-QLS-AA. Under the research context of the complexity of the test content and the diversification of cognitive responses, we propose this iterative analysis method based on hermeneutics and implement it through expert consultation. An interpretive cycle from cognitive interviewing to expert consultation until the item being tested is fully understood and accepted.

Classical coding systems tend to categorize linearly, which has the advantage of being organizational and cognitive interview.³³ But for diverse viewpoints, interpretive analysis methods appear to be more conducive to gaining a deeper understanding of the response process and linking individual items to the goals of the project as a whole.³² We achieved interpretive analysis through expert consultation, not only because expert consultation is one of the inherent methods of pre-testing, but more importantly, experts with rich clinical experience and multidisciplinary backgrounds can provide understanding and analysis from different perspectives.³⁴ This ensures that we make careful changes to every draft item. The results of the second round of cognitive interviews confirmed that all the items of Pre-QLS-AA (revised) can be understood, clear and easy to fill out, laying a foundation for the evaluation of the test efficiency in the next step.

Limitations

There are limitations to this study. First, our research method is a comprehensive exploration, although this exploration is carried out within the framework of cognitive interview, this process lacks standard procedures. Although the revision of Pre-QLS-AA has produced significant positive effects, its generalization and applicable scenarios still need to be verified and discussed. Moreover, the results of our study were reported sequentially within the framework of interpretive study, rather than the general reporting paradigm of qualitative research. How to report the results in a more standardized way and present the advantage of complementary combination of methods is a challenge that has not been fully solved in this study. In addition, our study did not conduct a national survey and interview due to resource limitations, but the impact of regional differences on patients' quality of life and their cognition should not be ignored, which may make QLS-AA lose some applicability in remote areas.

Reflection

In our study, we encountered the challenge of having a diverse response process for 4 draft items. This may be due to the lack of predictability of cognition that the researchers used the cognitive interview for the first time. In response to this existing dilemma, the research team proposed expert consultation based on interpretive analysis to clarify the relationship between diverse viewpoints and the overall purpose of the project, thereby helping us revise the project. Through reflection on our research, we think that we can provide two inspirations for other researchers: (1) Newcomers to cognitive interviews need to be very well prepared for research, especially to anticipate projects as well as keep an open mind to avoid preconceptions. Timely reflection during interview is also helpful to discover the shortcomings of the researcher himself as a research tool. (2) For the existing diversified response processes or contradictory viewpoints, the methods explored in this study may provide some references.

Conclusion

In this study, we report a revision of the Pre-QLS-AA with an comprehensive cognitive interview method. Our improvement lies in incorporating expert consultation based on interpretive analysis within the framework of cognitive interviews to address the challenges of item revision. The method showed a significant effect on the revision of Pre-QLS-AA in this study, and also provides ideas for the development of cognitive interviewing multimodality. In particular, our research methodology can inform the development of cognitive interviews or newly developed drafts.

Acknowledgments

The project was supported by grants from the Medical and Health Science and Technology Program of Zhejiang Province Health Department (2021KY822).

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Disclosure

The authors have no relevant financial or non-financial interests to disclose.

References

- Killick SB, Bown N, Cavenagh J, et al. Guidelines for the diagnosis and management of adult aplastic anaemia. *Br j Haematol*. 2016;172(2):187–207. doi:10.1111/bjh.13853
- Liu C, Shao Z. Aplastic Anemia in China. *J transl int med*. 2018;6(3):134–137. doi:10.2478/jtim-2018-0028
- Wang L, Liu H. Pathogenesis of aplastic anemia. *Hematology*. 2019;24(1):559–566. doi:10.1080/16078454.2019.1642548
- Montané E, Ibáñez L, Vidal X, et al. Epidemiology of aplastic anemia: a prospective multicenter study. *Haematologica*. 2008;93(4):518–523. doi:10.3324/haematol.12020
- Iftikhar R, Ahmad P, De Latour R, et al. Special issues related to the diagnosis and management of acquired aplastic anemia in countries with restricted resources, a report on behalf of the Eastern Mediterranean blood and marrow transplantation (EMBT) group and severe aplastic anemia working party of the European Society for blood and marrow transplantation (SAAWP of EBMT). *Bone Marrow Transplant*. 2021;56(10):2518–2532. doi:10.1038/s41409-021-01332-8
- Zhu XF, He HL, Wang SQ, et al. Current treatment patterns of aplastic anemia in China: a prospective cohort registry study. *Acta Haematol*. 2019;142(3):162–170. doi:10.1159/000499065
- Xu ZL, Xu LP, Wu DP, et al. Comparable long-term outcomes between upfront haploidentical and identical sibling donor transplant in aplastic anemia: a national registry-based study. *Haematologica*. 2022;107(12):2918–2927. doi:10.3324/haematol.2022.280758
- Wang LL, Mo WJ, Zhang YP, et al. Clinical analysis of CMV infection in patients with severe aplastic anemia after allogeneic hematopoietic stem cell transplantation. *J Exp Hematol*. 2021;29(3):944–950. doi:10.19746/j.cnki.issn1009-2137.2021.03.046
- Wu XL, Qiu YL, Wang L, et al. Analysis of self-perceived burden and influencing factors in patients with aplastic anemia. *Hosp Manag Forum*. 2018;35(12):23–26. doi:10.3969/j.issn.1671-9069.2018.12.007
- Tang J, Qian Y. Analysis and nursing experience of 1 case of severe aplastic anemia patient with suicide. *Chin Commun Physician*. 2019;35(18):146+9. doi:10.3969/j.issn.1007-614x.2019.18.106
- Streiner DL, Norman GR. *Health Measurement Scales: A Practical Guide to Their Development and Use*. 4th ed. Oxford, UK: Oxford University Press; 2008.
- Bobrovitz N, Santana MJ, Kline T, et al. The use of cognitive interviews to revise the Quality of Trauma Care Patient-Reported Experience Measure (QTAC-PREM). *Qual Life Res*. 2015;724(8):1911–1919. doi:10.1007/s11136-015-0919-5
- Collins D. Pretesting survey instruments: an overview of cognitive methods. *Qual Life Res*. 2003;12(3):229–238. doi:10.1023/a:1023254226592
- Watt T, Rasmussen AK, Groenvold M, et al. Improving a newly developed patient-reported outcome for thyroid patients, using cognitive interviewing. *Qual Life Res*. 2008;17(7):1009–1017. doi:10.1007/s11136-008-9364-z
- Murtagh FE, Addington-Hall JM, Higginson IJ. The value of cognitive interviewing techniques in palliative care research. *Palliat Med*. 2007;21(2):87–93. doi:10.1177/0269216306075367
- Fortune-Greeley AK, Flynn KE, Jeffery DD, et al. Using cognitive interviews to evaluate items for measuring sexual functioning across cancer populations: improvements and remaining challenges. *Qual Life Res*. 2009;18(8):1085–1093. doi:10.1007/s11136-009-9523-x
- Pillay T, Pillay M. Contextualising clinical reasoning within the clinical swallow evaluation: a scoping review and expert consultation. *S Afr J Commun Disord*. 2021;68(1):e1–e12. doi:10.4102/sajcd.v68i1.832
- Hu XH, Long L. Reflection and reconstruction of expert demonstration system for administrative decision-making. *J Jishou Univ*. 2017;38(5):95–102. doi:10.13438/j.cnki.jdx.2017.05.013
- DeWalt DA, Rothrock N, Yount S, et al. Evaluation of item candidates: the PROMIS qualitative item review. *Med Care*. 2007;45(5 Suppl 1):S12–S21. doi:10.1097/01.mlr.0000254567.79743.e2
- Shen T, Zhao YQ. *Criteria for Diagnosis and Efficacy of Hematological Diseases*. 4th ed. Beijing: Science Press; 2018.
- Whoqol Group T; The World Health Organization Quality of Life Assessment (WHOQOL). Development and general psychometric properties. *Soc Sci Med*. 1998;46(12):1569–1585. doi:10.1016/s0277-9536(98)00009-4
- Ferrell BR, Dow KH, Grant M. Measurement of the quality of life in cancer survivors. *Qual Life Res*. 1995;4(6):523–531. doi:10.1007/BF00634747
- Tourangeau R, Rips LJ, Rasinski K. *The Psychology of Survey Response*. Cambridge, UK: Cambridge University Press; 2000.
- Jobe JB. Cognitive psychology and self-reports: models and methods. *Qual Life Res*. 2003;12(3):219–227. doi:10.1023/A:1023279029852
- Meyer R, Drewniak D, Hovorka T, et al. Questioning the questionnaire: methodological challenges in measuring subjective quality of life in nursing homes using cognitive interviewing techniques. *Qual Health Res*. 2019;29(7):972–986. doi:10.1177/1049732318812042

26. Elo S, Kyngäs H. The qualitative content analysis process. *J adv nurs*. 2008;62(1):107–115. doi:10.1111/j.1365-2648.2007.04569.x
27. Whiteley I, Gullick J. The embodied experience of pregnancy with an ileostomy. *J Clin Nurs*. 2018;27(21–22):3931–3944. doi:10.1111/jocn.14601
28. Priest H. An approach to the phenomenological analysis of data. *Nurse Res*. 2002;10(2):50–63. PMID: 12518666. doi:10.7748/nr2003.01.10.2.50.c5888
29. Sherman KJ, Eaves ER, Ritenbaugh C, et al. Cognitive interviews guide design of a new CAM patient expectations questionnaire. *BMC Complement Altern Med*. 2014;14(39):1–16. doi:10.1186/1472-6882-14-39
30. Meadows K. Cognitive interviewing methodologies. *Clin Nurs Res*. 2021;30(4):375–379. doi:10.1177/10547738211014099
31. Arthur EK, Menon U, Browning K, et al. Challenges of cognitive interviewing in sensitive health topic research. *Nurs Res*. 2021;70(5):376–382. doi:10.1097/NNR.0000000000000530
32. Nyholm L, Nystrom L, Lindstrom U-A. Toward understanding through hermeneutic letterwriting. *Nurs Sci Q*. 2018;31(2):160–165. doi:10.1177/0894318418755731
33. Buers C, Triemstra M, Bloemendal E, et al. The value of cognitive interviewing for optimizing a patient experience survey. *Int J Soc Res Method*. 2013;17(4):325–340. doi:10.1080/13645579.2012.750830
34. Wang J, Lloyd-Evans B, Giacco D, et al. Social isolation in mental health: a conceptual and methodological review. *Soc Psychiatry Psychiatr Epidemiol*. 2017;52(12):1451–1461. doi:10.1007/s00127-017-1446-1

Patient Preference and Adherence

Dovepress

Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focusing on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize clinical outcomes for existing disease states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/patient-preference-and-adherence-journal>