



Cardiac troponin I in healthy Norwegian Forest Cat, Birman and domestic shorthair cats, and in cats with hypertrophic cardiomyopathy

Sofia Hanås^{1,2} , Anders Larsson³ , Jesper Rydén⁴, Inger Lilliehöök¹, Jens Häggström¹ , Anna Tidholm^{1,5}, Katja Höglund⁶, Ingrid Ljungvall¹ and Bodil S Holst¹

Journal of Feline Medicine and Surgery
2022, Vol. 24(10) e370–e379
© The Author(s) 2022



Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1098612X221117115
journals.sagepub.com/home/jfm

This paper was handled and processed by the European Editorial Office (ISFM) for publication in *JFMS*



Abstract

Objectives The aims of this study were to assess the potential associations between the serum cardiac troponin I (cTnI) concentration in healthy cats and feline characteristics, systolic blood pressure, heart rate (HR), echocardiographic measurements and storage time; and to compare cTnI concentrations in healthy cats with concentrations in cats with hypertrophic cardiomyopathy (HCM), with or without left atrial enlargement (LAE) and in cats with HCM, to assess potential associations between cTnI concentration and echocardiographic variables.

Methods Cardiac TnI was analysed using an Abbott ARCHITECT ci16200 analyser in serum from prospectively included healthy Norwegian Forest Cat (NF; $n = 33$), Birman ($n = 33$) and domestic shorthair (DSH; $n = 30$) cats, and from 39 cats with HCM, with or without LAE.

Results In healthy cats, higher cTnI concentrations were found in Birman cats than in NF cats ($P = 0.014$) and in neutered male cats than in intact females ($P = 0.032$). Cardiac TnI was positively associated with HR ($P < 0.0001$). In cats with HCM, cTnI concentration was positively associated with left ventricular wall thickness and with left atrial-to-aortic root ratio (all $P \leq 0.010$). Cats with HCM had higher cTnI concentrations than healthy cats, and cTnI concentrations were higher in cats with HCM and LAE than in those with HCM without LAE (all $P = 0.0003$).

Conclusions and relevance Breed and sex may affect serum cTnI concentrations in healthy cats. The cTnI concentration increased with increasing severity of HCM.

Keywords: hs-cTnI; breed; biomarker; feline; heart

Accepted: 12 July 2022

Introduction

Cardiac troponin I (cTnI) is a sensitive biomarker for myocardial injury, and high-sensitivity cTnI (hs-cTnI) assays allow detection of low concentrations of cTnI in healthy cats,^{1–4} dogs^{1,5} and humans.⁶ An hs-cTnI assay enables the detection of cTnI concentrations in >50% of healthy humans, with a coefficient of variation (CV) of $\leq 10\%$ for the 99th percentile.⁶ In humans, different cut-off values are used for the available hs-cTnI assays because of varying capture and detection antibodies and lack of standardisation.^{7,8} Only one hs-cTnI assay has been validated in cats.¹ In humans, male sex is positively associated with cTnI concentration, and sex-specific reference intervals (RIs) are used.^{9–11} Breed differences have been found in healthy dogs,^{12,13} and a previous report in cats

¹Department of Clinical Sciences, Swedish University of Agricultural Sciences, Uppsala, Sweden

²Evidensia Specialist Animal Hospital Strömsholm, Strömsholm, Sweden

³Department of Medical Sciences, Clinical Chemistry, Uppsala University, Uppsala, Sweden

⁴Department of Energy and Technology, Swedish University of Agricultural Sciences, Uppsala, Sweden

⁵Anicura Albano Animal Hospital, Stockholm, Sweden

⁶Department of Anatomy, Physiology and Biochemistry, Swedish University of Agricultural Sciences, Uppsala, Sweden

Corresponding author:

Sofia Hanås DVM, PhD, Department of Clinical Sciences, Swedish University of Agricultural Sciences, Box 7054, Uppsala 750 07, Sweden
Email: sofia.hanas@slu.se

with hypertrophic cardiomyopathy (HCM) indicated a breed effect.¹⁴ In healthy cats, cTnI concentrations have neither been associated with sex nor with breed or body condition score (BCS).^{2,3,14} Results in cats regarding associations between cTnI concentrations and age and body weight (BW) are conflicting.^{2,3,15}

HCM is a common cardiac disease in cats, with a reported prevalence rate of approximately 15%.^{16,17} The disease is characterised by left ventricular hypertrophy in the absence of other explanations for wall thickening,¹⁸ and may lead to congestive heart failure (CHF), arterial thromboembolism (ATE), arrhythmia or sudden death.^{19–21} Using conventional cTnI assays, significantly higher concentrations have been detected in cats with HCM than in healthy cats,^{22,23} in which cTnI concentrations have been reported to be commonly below the detection limit.^{23,24} In cats with HCM, positive associations between cTnI concentrations and hypertrophy of the left ventricle (LV), and with left atrial enlargement (LAE), have been shown.^{2–4,23} Although cTnI is specific for the myocardium, increased cTnI concentrations are not specific for primary cardiac disease but have also been shown in cats with hyperthyroidism,²⁵ hypertension,²⁶ renal disease²⁷ and critical illness.²⁸

The aims of this study were to assess potential associations between serum concentration of cTnI in healthy cats and feline characteristics (breed, sex, age, BW and BCS), systolic blood pressure (SBP), heart rate (HR) at auscultation, echocardiographic measurements and storage time. A further aim was to compare cTnI concentrations in healthy cats with concentrations in cats with HCM, with and without LAE, and in cats with HCM, to assess potential associations between serum cTnI concentration in cats with HCM and echocardiographic variables. We hypothesised that the serum concentration of cTnI in healthy cats may be influenced by feline characteristics, and that cTnI concentrations may differ between healthy cats and cats with HCM, with or without LAE.

Materials and methods

This prospective observational study was approved by the Uppsala Animal Experiment Ethics Board, Sweden (C137/13). Client-owned cats were recruited through information distributed online, at seminars or at recruiting clinics. Cats were examined at the Evidensia Animal Clinic in Västerås, Sweden, between September 2014 and June 2017. Clinically healthy cats and cats with murmurs or previously diagnosed HCM were examined for possible inclusion. Informed written consent was obtained from the owner of each cat. The population has, in part, been described previously.^{29,30}

Inclusion criteria

Apparently healthy Norwegian Forest Cat (NF), Birman and domestic shorthair (DSH) cats aged 1–14 years, with normal echocardiograms, were included, as were cats of any breed with preclinical HCM or clinical HCM

stabilised following CHF therapy. A diagnosis of HCM was based on characteristic findings on an echocardiogram, as outlined below.

Exclusion criteria

Cats with mean SBP >160 mmHg, or increased serum concentrations of total thyroxine (TT4), creatinine or alanine aminotransferase were excluded. Cats with decompensated CHF, ATE, congenital cardiac disease, other acquired cardiovascular disorders, equivocal findings concerning the presence of LV hypertrophy, or significant organ-related or systemic diseases other than HCM were excluded. All cats receiving medical treatment other than standard CHF treatment, standard antithrombotic treatment or medroxyprogesterone acetate were excluded.

Physical examination and blood pressure measurements

All examinations were performed according to a standardised protocol in a quiet examination room by an experienced veterinarian (SH). SBP was measured indirectly using a high-definition oscillometric device (VET Memodiagnostic HDO monitor; S+B medVET),³¹ followed by a general physical examination, including assessment of BCS on a nine-point scale,³² echocardiography and blood sampling for haematology, biochemistry profiles, and TT4 and fructosamine levels, as previously described.²⁹

Echocardiography

Echocardiographic examination was performed using an ultrasound unit (IE33; Philips Ultrasound) equipped with a 4–12 MHz phased-array probe, and with continuous electrocardiogram monitoring.³³ The left atrial-to-aortic root diameter ratio (LA:Ao) was measured from the right two-dimensional short-axis view.³⁴ Left atrial enlargement was defined as LA:Ao \geq 1.5.³⁵ End-diastolic and systolic LV dimensions (interventricular septum in diastole [IVSd], LV internal diameter in diastole [LVIDd], LV free wall in diastole [LVFWd] and LV internal diameter in systole [LVIDs]) were measured from M-mode and two-dimensional images, and fractional shortening (FS) was calculated.^{33,36}

A diagnosis of HCM was made when subjective impression of hypertrophy (diffuse or regional) with a non-dilated LV chamber was supported by increased M-mode and two-dimensional diastolic LV wall dimensions of the IVSd, LVFWd or both, as previously described.²⁹ Expected BW-dependent values for the IVSd, LVIDd and LVFWd, as well as percentage deviations from these expected values, were calculated for both healthy cats and cats with HCM according to previously generated formulas for cats.³⁵ The calculated deviations and LA:Ao were used to classify cats into three groups – healthy controls, HCM without LAE and HCM with LAE – as described previously.²⁹

Analytical performance of hs-cTnI

The validation study was set up in accordance with the Clinical and Laboratory Standards Institute document EP15-A3, evaluating precision using a one-way ANOVA.³⁷ In the imprecision study, two concentrations of pooled feline serum (cTnI concentrations 22.8 ng/l and 312.7 ng/l, respectively) and two human control samples (Liquicheck Cardiac Marker LT1 and Liquicheck Cardiac Marker Cardio 2 [Bio-Rad Laboratories]) were used. Each sample was analysed three times daily for 5 days to determine within-run (CV_R), between-run (CV_B) and within-laboratory (CV_{WL}) repeatability. Limit of blank (LOB) was evaluated using 20 replicates of physiological saline solution. Linearity under dilution was performed in duplicate through the dilution of two concentrations of pooled feline serum (45 ng/l and 173 ng/l, respectively) with physiological saline solution as serial dilution and the addition of an extra sample consisting of 75% feline serum and 25% physiological saline solution. Recovery ([observed/expected concentration] \times 100) was calculated. Stability was determined by keeping fresh feline serum samples with initial cTnI concentrations of 1070, 1040, 761 and 91 ng/l, respectively, at 20°C in the dark for 0, 3, 5 and 7 days, respectively, and then stored at -80°C until thawed for batched analysis. The effect of freezing and thawing was determined by assaying aliquots of fresh feline serum samples subjected to three cycles of freezing at -80°C for 24 h and thawing to room temperature for 30 mins.

Analysis of cTnI in cats

Whole blood was centrifuged and the serum stored at -80°C within 60 mins of collection. Samples were transported from the clinic to the laboratory at -80°C using a portable freezer. Serum for cTnI was batch-analysed in duplicate at an accredited laboratory at the Department of Clinical Chemistry and Pharmacology, Uppsala University Hospital, using a two-step, double-monoclonal chemiluminescent microparticle immunoassay for the detection of cTnI concentration (Abbott ARCHITECT ci16200 analyser; Abbott Laboratories). The reported assay interval for cTnI concentration was 2–50,000 ng/l. Concentrations <2 ng/l were assigned a value of 2 ng/l in the calculations.

Statistical analysis

Statistical analyses were performed using Rstudio³⁸ and JMP (version 12.2.0; SAS Institute). A P value <0.05 was considered to be statistically significant. Descriptive statistics for continuous variables (age, BW, SBP, HR, basic echocardiographic measurements and basic laboratory variables) were presented as mean \pm SD. Within each of these continuous variables, multiple comparisons were performed between breeds for healthy cats, and between healthy cats and cats with HCM, using one-way

ANOVA followed by Tukey's honestly significant difference (HSD) test.

Data on cTnI concentrations were presented as median and interquartile range (IQR). Potential associations between cTnI concentration in healthy cats, as well as in cats with HCM and the variables breed, sex, BCS, age, BW, SBP, HR, echocardiographic measurements $IVSd_{inc\%}$, $LVIDd_{inc\%}$, $LVPWd_{inc\%}$, $FS\%$ and LA/Ao , and the duration of storage at -80°C of serum samples, were analysed using univariable linear regression for continuous variables and one-way ANOVA for categorical variables. BCS was divided into normal (BCS 4–5) and overweight (BCS 6–7), and sex into four classes (male, female, neutered male and neutered female). Variables with a P value <0.2 were included in a linear multiple regression analysis and, as cTnI is a marker of myocardial injury, a variable for the LV wall – $LVPWd_{inc\%}$ – was also included in the model for healthy cats. A model selection was performed, excluding covariate echocardiographic variables for the LV, using the echocardiographic variable with the lowest P value in the model. The variable with the highest P value was then removed until all remaining variables had a P value <0.05. The model with the lowest Akaike information criterion was chosen. All variables were assessed as main effects. The interactions breed \times BW and breed \times sex were evaluated but were not significant and thus not included in the model. All multiple regression analyses were performed after logarithmic transformation of cTnI concentrations, to have normally distributed residuals. For multiple comparisons between sex and breed categories, respectively, Tukey's HSD test was used as implemented by the routine general linear hypotheses in the R package 'multcomp'.

Comparisons of cTnI concentrations between the healthy, HCM without LAE and HCM with LAE groups were done using the Kruskal–Wallis test, followed by post-hoc Wilcoxon tests for each pair.

Results

Validation of the hs-cTnI immunoassay

The CV_R , CV_B and CV_{WL} were 4.0–8.6% for the feline samples and 5.1–8.4% for the human control samples. Assay results were adequately linear after dilution and the recovery was 64–116%. The LOB was 0.25 ± 0.12 ng/l. After 3 days of storage at room temperature, the concentrations of cTnI were still within 10% in comparison with the initial values (see Supplementary Appendix 1 and Supplementary Figure 1 in the supplementary material).

Study population

Healthy cats The 96 healthy cats comprised Birman, NF and DSH cats. The characteristics of the healthy cats are presented in Table 1 and their basic echocardiographic data in Supplementary Table 1; excluded cats are detailed in Figure 1.

Table 1 Feline characteristics, heart rate (HR) at auscultation and systolic blood pressure (SBP) in 96 healthy cats

Group	Birman (n = 33)	DSH (n = 30)	NF (n = 33)
Sex			
Female	12	0	10
Male	5	0	3
Female neutered	9	14	12
Male neutered	7	16	8
Age (years)	4.8 ± 4.0 ^a	7.0 ± 4.4 ^a	5.0 ± 3.0 ^a
Body weight (kg)	3.6 ± 0.7 ^a	4.7 ± 1.1 ^b	5.4 ± 1.6 ^b
Normal*	21	12	14
Overweight†	11‡	18	19
HR (bpm)	159 ± 20 ^a	151 ± 27 ^a	165 ± 27 ^a
SBP (mmHg)	125 ± 11 ^a	140 ± 10 ^b	134 ± 13 ^b

Data are provided mean ± SD for continuous variables. Within each row, values with different superscripts differ significantly between groups. Statistical significance was set at $P < 0.05$. Multiple comparisons within each independent variable were corrected using Tukey's method

*Body condition score (BCS) 4–5/9

†BCS 6–7/9

‡One missing value

DSH = domestic shorthair; NF = Norwegian Forest Cat; LAE = left atrial enlargement; HR = heart rate at auscultation; bpm = beats/min; SBP = systolic blood pressure

Cats with HCM The 39 cats with HCM comprised 18 DSH and cats from 10 different breeds (five Persian, four NF, two Bengal, two Maine Coon, two Ragdoll and one each of the following breeds: British Shorthair, Cornish

Rex, Devon Rex, Exotic Shorthair, Siberian and Sphynx). Of 32 cats with HCM without LAE, one female was treated with medroxyprogesterone acetate, and three were treated with enalapril. Of the seven cats with HCM and LAE, three cats with stabilised CHF were treated with furosemide and enalapril, one of which also received clopidogrel. The characteristics of the included cats with HCM are presented in Table 2 and their basic echocardiographic data in Supplementary Table 2; excluded cats are detailed in Figure 1.

Association between cTnI and feline characteristics, and clinical and echocardiographic variables in healthy cats

cTnI was $>2\text{ng/l}$ in 89/96 samples (93%). cTnI was significantly associated with breed and sex, and positively associated with HR (Tables 3 and 4) but not with age, SBP within the normal range or LFWd_{inc%}. Higher cTnI concentrations were found in Birman cats than in NF cats ($P = 0.014$), and in neutered males than in intact females ($P = 0.032$; Tables 3 and 4). The final multiple regression model included breed, sex, HR and LFWd_{inc%} (Table 4).

Association between cTnI and HCM

The median cTnI serum concentration in cats with HCM was 37 ng/l (IQR 17.5–81.5), and was higher in cats with HCM with LAE than in cats with HCM without LAE or in healthy cats ($P < 0.0001$; Table 5). All three groups differed (all $P = 0.0003$; Table 5). For the analysis of the 39 cats with HCM, the final model included the variables LFWd_{inc%} and LA/Ao, both positively associated with cTnI (Table 6).




Figure 1 Flow chart of the study, including the reasons for the exclusion of 30 cats

Table 2 Feline characteristics, heart rate (HR) at auscultation, systolic blood pressure (SBP) and selected echocardiographic data in 96 healthy cats and 39 cats with hypertrophic cardiomyopathy (HCM)

Group	Healthy (n = 96)	HCM without LAE (n = 32)	HCM with LAE (n = 7)
Sex			
Female	22	3	1
Male	8	4	0
Female neutered	35	9	0
Male neutered	31	16	6
Age (years)	5.6 ± 3.9 ^a	6.1 ± 3.4 ^a	6.3 ± 2.7 ^a
Weight (kg)	4.6 ± 1.4 ^a	5.1 ± 1.2 ^a	4.8 ± 1.1 ^a
Normal*	47	6	5
Overweight†	48‡	22§	2
HR (bpm)	158 ± 25 ^a	167 ± 26 ^a	169 ± 31 ^a
SBP (mmHg)	133 ± 13 ^a	138 ± 12 ^a	128 ± 11 ^a
LA:Ao	1.1 ± 0.1 ^{a,¶}	1.1 ± 0.1 ^{a,¶}	1.6 ± 0.1 ^{b,¶}
LVFWd (mm)	3.8 ± 0.5 ^{a,¶}	5.4 ± 1.3 ^{b,¶}	7.6 ± 0.5 ^{c,¶}
LVFWd _{inc%}	-0.9 ± 9.5 ^{a,¶}	39.6 ± 33.2 ^{b,¶}	99.8 ± 16.7 ^{c,¶}

Mean ± SD is provided for continuous variables. Within each row, values with different superscripts differ significantly between groups. Multiple comparisons within each independent variable were corrected using Tukey's method. The level of statistical significance was set at $P < 0.05$

*Body condition score (BCS) 4–5/9

†Overweight = BCS 6–7/9

‡One missing value

§Four missing values

¶Expected differences due to echocardiographic classification of cats into three groups: healthy controls, HCM without LAE, and HCM with LAE. HR = heart rate at auscultation; LAE = left atrial enlargement; bpm = beats/min; SBP = systolic blood pressure; LA:Ao = left atrial-to-aortic root diameter ratio; LVFWd = left ventricular free wall in diastole; LVFWd_{inc%} = percentage increase left ventricular free wall in diastole

Table 3 Serum concentration of cardiac troponin I (cTnI) in 96 healthy cats

Group	n	Median (IQR) cTnI (ng/l)	Range (ng/l)
Breed			
NF	33	4.0 (2.6–7.7) ^a	<2.0–42.0
Birman	33	7.6 (3.3–14.4) ^b	<2.0–49.0
DSH	30	7.7 (3.0–16.1) ^{a,b}	<2.0–156.0
Sex			
Intact female	22	3.7 (2.1–7.5) ^a	<2.0–20.0
Intact male	8	7.8 (5.5–10.5) ^{a,b}	<2.0–24.0
Female neutered	35	6.3 (3.3–11.5) ^{a,b}	<2.0–156.0
Male neutered	31	7.5 (3.0–16.0) ^b	<2.0–61.5
All cats	96	5.7 (2.8–11.0)	<2.0–156.0

Different superscripts within a column denote differences in cTnI concentrations between breed and sex, respectively. The level of statistical significance level was set at $P < 0.05$. Multiple comparisons within breed and sex were adjusted using Tukey's method. IQR = interquartile range; NF = Norwegian Forest Cat; DSH = domestic shorthair

Discussion

In the present study, the hs-cTnI assay was precise and accurate for use in cats. The main findings were a significant association between serum cTnI concentrations and breed, sex and HR in healthy cats. There was no significant association between cTnI concentration and age, or

with SBP within the normal range. There were significant differences in serum cTnI concentration between healthy cats, cats with HCM without LAE and cats with HCM with LAE, which is in accordance with previous studies in cats.^{2,3}

In healthy cats, higher cTnI concentrations were found in Birman cats than in NF cats. Interbreed variations have previously been reported in dogs, with higher cTnI concentrations in Greyhound and Boxer dogs than in other breeds,^{12,13} but not in healthy cats.^{2,3,14} In cats with HCM, a previous study indicated a breed effect with higher cTnI concentrations in British Shorthair cats than in Maine Coons.¹⁴ However, echocardiographic variables indicating the severity of HCM were not included in the statistical analysis in that study,¹⁴ thus the British Shorthair cats may simply have had more severe HCM at inclusion than the Maine Coons. An effect of sex was also found in healthy cats, with higher cTnI concentrations in male neutered cats than in intact female cats. In humans, women have been described as having lower cTnI concentrations than men.^{9,39} In the present study, intact female cats had lower concentrations of cTnI than intact males. The observed difference was not statistically significant, but this may have been a type II statistical error due to the small number of intact cats included in the study. Healthy cats had comparatively lower serum cTnI concentrations, and the breed and sex associations were probably detected owing to the more sensitive cTnI assay used in the present study,

Table 4 Final model in the multiple regression analysis of association between the effects of feline characteristics and echocardiographic variables and heart rate (HR) at auscultation on the serum concentration of cardiac troponin I (cTnI) in 96 healthy cats

Variable	Estimate	<i>P</i> value	Adjusted <i>P</i> value	95% CI
NF	Baseline breed			
Birman	0.589	0.005	0.014	0.133–1.045
DSH	0.521	0.023	0.058	0.080–0.961
Female sex	Baseline sex			
Male sex	0.417	0.22	0.599	–0.245 to 1.079
Female neutered	0.483	0.045	0.178	0.017–0.948
Male neutered	0.708	0.0067	0.032	0.209–1.208
HR (bpm)	0.017	<0.0001		0.010–0.024
LVFWd _{inc%}	0.012	0.19		–0.006 to 0.030

The estimate column gives the estimated regression coefficients in the fitted regression model. cTnI is a marker of myocardial injury, and thus a variable for the left ventricular wall (LVFWd_{inc%}) was included in the model. The level of statistical significance was set at *P* < 0.05. Multiple comparisons within breed and sex, respectively, were adjusted using Tukey's method. The final model had an *r*² of 0.23 and an overall *P* value of < 0.0001

CI = confidence interval; NF = Norwegian Forest Cat; DSH = domestic shorthair; HR = heart rate at auscultation; bpm = beats/min; LVFWd_{inc%} = percentage increase left ventricular free wall in diastole (percentage)

Table 5 Serum concentration of cardiac troponin I (cTnI) vs echocardiographic classification in the study population of 135 cats

Group	n	Median (IQR) cTnI (ng/l)	Range (ng/l)
Healthy controls	96	5.7 (2.8–11.0) ^a	<2.0–156.0
HCM without LAE	32	29.3 (13.3–46.5) ^b	3.1–318.0
HCM with LAE	7	296 (92.0–642.0) ^c	56.0–1880.0

Different superscripts within a column denote differences in cTnI concentrations between the groups (healthy controls, HCM without LAE and HCM with LAE). The level of statistical significance was set at *P* < 0.05. For multiple comparisons, Kruskal–Wallis test was used, followed by post-hoc Wilcoxon tests for each pair. HCM = hypertrophic cardiomyopathy; LAE = left atrial enlargement; IQR = interquartile range

as compared with previously used assays in cats.^{2,3,14} However, despite being statistically significant, differences in cTnI concentrations among breed and sex were much smaller than between healthy cats and cats with HCM, and likely not of relevance in a clinical situation. Further studies would be needed to investigate if breed- and sex-specific RIs are beneficial.

The lack of association between age and cTnI concentration is in contrast to reports on healthy people,^{40,41} healthy dogs⁴² and dogs with mild myxomatous mitral valve disease,⁴³ where cTnI has been positively associated with age. In cats, both a positive association¹⁵ and no association^{2,3} with age have been previously described. In the present study, the mean age of the healthy cats was 5.6 years, with few older cats, which possibly contributed to the lack of association.

In the present study, a positive association between cTnI concentration and HR was found in healthy cats. Prolonged exercise or stress increases HR and blood

Table 6 Final model in the multiple regression analysis of association between the effects of echocardiographic variables on the serum concentration of cardiac troponin I in 39 cats with hypertrophic cardiomyopathy

Variable	Estimate	<i>P</i> value	95% CI
LVFWd _{inc%}	0.016	0.004	0.006–0.026
LA:Ao	2.41	0.010	0.66–4.15

The estimate column gives the estimated regression coefficients in the fitted regression model. The level of statistical significance was set at *P* < 0.05. The final model had an adjusted *r*² of 0.46 and an overall *P* value of < 0.0001

CI = confidence interval; LA:Ao = left atrial-to-aortic root diameter ratio; LVFWd_{inc%} = percentage increase of the left ventricular free wall in diastole

pressure.^{44–46} In humans⁴⁷ and in horses,^{48–50} post-exercise-induced cTnI elevations have been reported after normal physical activity, with a peak at 3–6 h post-exercise and normalisation within 24 h.^{49,51} The increase of circulating cTnI after exercise has been suggested to arise from a proportion of cTnI that is free in the cytosol,⁵² and thus not a sign of cardiac damage. In the present study, blood sampling was performed a couple of hours after leaving home, and situational stress might be an explanation for higher HR.

cTnI concentration differed significantly between healthy cats, cats with HCM without LAE and cats with HCM with LAE. Previous investigations in humans,⁵³ cats^{4,54} and dogs⁴³ suggest that cTnI correlates with the severity of HCM and myxomatous mitral valve disease. Echocardiography is used for the diagnosis of LV hypertrophy and the HCM phenotype in cats.^{18,55–57} Diagnosis is based on a subjective impression of LV hypertrophy supported by measurement of maximal end-diastolic wall thicknesses via two-dimensional or

M-mode echocardiography.^{58–60} Fixed diagnostic cutoffs defining normal diastolic LV wall thickness have been used.^{55,60,61} Several studies have recommended allometric scaling and BW-based 95% prediction intervals, in particular in small and large cats.^{35,62,63} In the present study, BW-dependent echocardiographic values, as well as percentage deviations from these expected values, were calculated and used. Evaluations of the cTnI concentration for assessing HCM in cats categorised using echocardiographic measurements adjusted for BW have, to our knowledge, not been published previously. Cats with HCM and LAE have a more severe disease than cats with HCM without LAE.^{20,64–66} In a study with a median follow-up of 3.1 years, cats with HCM and LAE were four times more likely to experience a cardiac event than cats with a normal atrial size.⁶⁷ In accordance with previous studies in cats, concentrations of cTnI in cats with HCM were positively associated with LVFWd_{inc%} and LA:AO,^{2,3} possibly caused by myocardial ischaemia due to myocyte death or by an imbalance between hypertrophy of the myocardium and insufficient coronary arterial supply.^{68,69} Although cTnI is heart-specific, increases are not specific for conditions affecting the heart, and increased cTnI concentrations have been described in cats with hyperthyroidism,²⁵ hypertension,²⁶ renal disease²⁷ and critical illness.²⁸

Concentrations of cTnI differ substantially between different hs-cTnI assays,⁷ and concentrations obtained with different hs-cTnI assays are therefore difficult to compare. In the present study, cTnI was detectable in 93% of healthy cats, which is higher than previously reported in cats evaluated using another hs-cTnI assay.² The CVs in our study were comparable to those previously reported for cTnI assays in humans,^{70,71} dogs^{1,72} and cats.¹ The concentration of cTnI in samples stored at 20°C for 3 days changed by less than 10%, possibly enabling the transport of fresh feline serum samples to a central laboratory. However, stability was only studied for cTnI concentrations above 90 ng/l, and studies of concentrations closer to the clinical cutoff would be valuable. In the present study, cTnI analyses for the healthy cats and the cats with HCM were performed after up to 5 years of serum storage at –80°C, but storage time was not associated with cTnI concentration in the statistical analysis.⁷¹

The design of our study, with different criteria for selection of the healthy controls and the cats with HCM (ie, three specific breeds were included as healthy controls and any breed was permitted in the HCM cat group), is a limitation. The different breed distributions may have affected the results because breed influences the cTnI concentration. Only healthy cats and cats with HCM, but no other known diseases, were included in the study. Cats with equivocal echocardiographic findings and measurements, or other comorbidities were excluded, which introduces the spectrum effect for cTnI concentration.⁷³ In a mixed cat population, including cats with comorbidities,

the rate of elevated cTnI results would increase and the accuracy for the test for discriminating healthy cats from cats with HCM is likely to decrease. The present study was intended as an exploratory study, and further research in a mixed cat population is warranted for a discriminatory analysis. Body-weight dependent echocardiographic values were used for both healthy cats and cats with HCM, which affected the ability to compare the results to those of previous studies that have used a fixed diagnostic cutoff for HCM. A fixed cutoff may have increased the inclusion of mild HCM in the healthy control group, especially if the cats were small, and extremely large cats may have been falsely included as mild HCM cats.

Conclusions

Analytical performance of the hs-cTnI analysis allows its clinical use in cats. There was an effect of both breed and sex on the serum concentration of cTnI. In cats with HCM, cTnI concentrations increased with increasing LV wall thickness, and cats with LAE had higher cTnI concentrations than cats with HCM without LAE and than healthy cats.

Acknowledgements The authors would like to thank the dedicated veterinary technicians at Evidensia Animal Clinic in Västerås for their important role in completing this study. Special thanks to all participating cats and their owners.

Author note This work was a part of Sofia Hanås's PhD thesis. This paper was presented, in part, at the 31st ECVIM Online Congress as a short abstract and a poster in 2021.

Supplementary material The following files are available online:

Supplementary Appendix 1: Stability study.

Supplementary Figure 1: The stability of cardiac troponin I concentration in serum samples from three cats with hypertrophic cardiomyopathy after storage in the dark at 20°C.

Supplementary Table 1: Auscultation, basic echocardiographic and laboratory variables in 96 healthy Birman, domestic short-hair and Norwegian Forest Cats.

Supplementary Table 2: Auscultation, basic echocardiographic and laboratory variables in 96 healthy cats and 39 cats with hypertrophic cardiomyopathy.


Conflict of interest The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.


Funding This study was supported by the Foundation Strömsholms Djursjukvård, Agria and Swedish Kennel Club's Research Fund, SLU Companion Animals Research Fund, Svealand Research Fund, and Michael Forsgren Foundation.

Ethical approval The work described in this manuscript involved the use of non-experimental (owned or unowned) animals. Established internationally recognised high standards ('best practice') of veterinary clinical care for the individual patient were always followed and/or this work involved the


use of cadavers. Ethical approval from a committee was therefore not specifically required for publication in *JFMS*. Although not required, where ethical approval was still obtained, it is stated in the manuscript.

Informed consent Informed consent (either verbal or written) was obtained from the owner or legal custodian of all animal(s) described in this work (non-experimental animals, including cadavers) for all procedure(s) undertaken (prospective studies). No animals or people are identifiable within this publication, and therefore additional informed consent for publication was not required.

ORCID iD Sofia Hanås  <https://orcid.org/0000-0002-6758-8190>

Anders Larsson  <https://orcid.org/0000-0003-3161-0402>

Jens Häggström  <https://orcid.org/0000-0003-3402-023X>

Bodil S Holst  <https://orcid.org/0000-0001-7623-7141>

References

- Langhorn R, Willesen JL, Tarnow I, et al. **Evaluation of a high-sensitivity assay for measurement of canine and feline serum cardiac troponin I.** *Vet Clin Path* 2013; 42: 490–498.
- Hertzsch S, Roos A and Wess G. **Evaluation of a sensitive cardiac troponin I assay as a screening test for the diagnosis of hypertrophic cardiomyopathy in cats.** *J Vet Intern Med* 2019; 33: 1242–1250.
- Hori Y, Iguchi M, Heishima Y, et al. **Diagnostic utility of cardiac troponin I in cats with hypertrophic cardiomyopathy.** *J Vet Intern Med* 2018; 32: 922–929.
- Langhorn R, Tarnow I, Willesen JL, et al. **Cardiac troponin I and T as prognostic markers in cats with hypertrophic cardiomyopathy.** *J Vet Intern Med* 2014; 28: 1485–1491.
- Winter RL, Saunders AB, Gordon SG, et al. **Analytical validation and clinical evaluation of a commercially available high-sensitivity immunoassay for the measurement of troponin I in humans for use in dogs.** *J Vet Cardiol* 2014; 16: 81–89.
- Apple FS and Collinson PO. **Analytical characteristics of high-sensitivity cardiac troponin assays.** *Clin Chem* 2012; 58: 54–61.
- Apple FS, Wu AHB, Sandoval Y, et al. **Sex-specific 99th percentile upper reference limits for high sensitivity cardiac troponin assays derived using a universal sample bank.** *Clin Chem* 2020; 66: 434–444.
- Clerico A, Zaninotto M, Ripoli A, et al. **The 99th percentile of reference population for cTnI and cTnT assay: methodology, pathophysiology and clinical implications.** *Clin Chem Lab Med* 2017; 55: 1634–1651.
- Kimenai DM, Janssen E, Eggers KM, et al. **Sex-specific versus overall clinical decision limits for cardiac troponin I and T for the diagnosis of acute myocardial infarction: a systematic review.** *Clin Chem* 2018; 64: 1034–1043.
- Mueller T, Egger M, Peer E, et al. **Evaluation of sex-specific cut-off values of high-sensitivity cardiac troponin I and T assays in an emergency department setting – results from the Linz Troponin (LITROP) study.** *Clin Chim Acta* 2018; 487: 66–74.
- Love SA, Sandoval Y, Smith SW, et al. **Incidence of undetectable, measurable, and increased cardiac troponin I concentrations above the 99th percentile using a high-sensitivity vs a contemporary assay in patients presenting to the emergency department.** *Clin Chem* 2016; 62: 1115–1119.
- LaVecchio D, Marin LM, Baumwart R, et al. **Serum cardiac troponin I concentration in retired racing greyhounds.** *J Vet Intern Med* 2009; 23: 87–90.
- Baumwart RD, Orvalho J and Meurs KM. **Evaluation of serum cardiac troponin I concentration in Boxers with arrhythmogenic right ventricular cardiomyopathy.** *Am J Vet Res* 2007; 68: 524–528.
- Langhorn R, Willesen JL, Tarnow I, et al. **Cardiac troponin I in three cat breeds with hypertrophic cardiomyopathy.** *Vet Rec* 2016; 178: 532. DOI: 10.1136/vr.103549.
- Serra M, Papakonstantinou S, Adamcova M, et al. **Veterinary and toxicological applications for the detection of cardiac injury using cardiac troponin.** *Vet J* 2010; 185: 50–57.
- Payne JR, Brodbelt DC and Luis Fuentes V. **Cardiomyopathy prevalence in 780 apparently healthy cats in rehoming centres (the CatScan study).** *J Vet Cardiol* 2015; 17 Suppl 1: S244–S257.
- Paige CF, Abbott JA, Elvinger F, et al. **Prevalence of cardiomyopathy in apparently healthy cats.** *J Am Vet Med Assoc* 2009; 234: 1398–1403.
- Luis Fuentes V, Abbott J, Chetboul V, et al. **ACVIM consensus statement guidelines for the classification, diagnosis, and management of cardiomyopathies in cats.** *J Vet Intern Med* 2020; 34: 1062–1077.
- Atkins CE, Gallo AM, Kurzman ID, et al. **Risk factors, clinical signs, and survival in cats with a clinical diagnosis of idiopathic hypertrophic cardiomyopathy: 74 cases (1985–1989).** *J Am Vet Med Assoc* 1992; 201: 613–618.
- Rush JE, Freeman LM, Fenollosa NK, et al. **Population and survival characteristics of cats with hypertrophic cardiomyopathy: 260 cases (1990–1999).** *J Am Vet Med Assoc* 2002; 220: 202–207.
- Payne JR, Borgeat K, Connolly DJ, et al. **Prognostic indicators in cats with hypertrophic cardiomyopathy.** *J Vet Intern Med* 2013; 27: 1427.
- Herndon WE, Kittleson MD, Sanderson K, et al. **Cardiac troponin I in feline hypertrophic cardiomyopathy.** *J Vet Intern Med* 2002; 16: 558–564.
- Connolly DJ, Cannata J, Boswood A, et al. **Cardiac troponin I in cats with hypertrophic cardiomyopathy.** *J Feline Med Surg* 2003; 5: 209–216.
- Langhorn R, Yrfelt JD, Stjernegaard CS, et al. **Analytical validation of a conventional cardiac troponin I assay for dogs and cats.** *Vet Clin Path* 2019; 48: 36–41.
- Sangster JK, Panciera DL, Abbott JA, et al. **Cardiac biomarkers in hyperthyroid cats.** *J Vet Intern Med* 2014; 28: 465–472.
- Bijsmans ES, Jepson RE, Wheeler C, et al. **Plasma N-terminal probrain natriuretic peptide, vascular endothelial growth factor, and cardiac troponin I as novel biomarkers of hypertensive disease and target organ damage in cats.** *J Vet Intern Med* 2017; 31: 650–660.
- Langhorn R, Jessen LR, Kloster AS, et al. **Cardiac troponin I in cats with compromised renal function.** *J Feline Med Surg* 2019; 21: 985–991.
- Sharpe AN, Gunther-Harrington CT, Epstein SE, et al. **Cats with thermal burn injuries from California wildfires**

- show echocardiographic evidence of myocardial thickening and intracardiac thrombi. *Sci Rep* 2020; 10: 2648–2648.
- 29 Hanås S, Holst BS, Höglund K, et al. Effect of feline characteristics on plasma N-terminal-prohormone B-type natriuretic peptide concentration and comparison of a point-of-care test and an ELISA test. *J Vet Intern Med* 2020; 34: 1187–1197.
- 30 Hanås S, Holst BS, Ljungvall I, et al. Influence of clinical setting and cat characteristics on indirectly measured blood pressure and pulse rate in healthy Birman, Norwegian Forest, and domestic shorthair cats. *J Vet Intern Med* 2021; 35: 801–811.
- 31 Martel E, Egner B, Brown SA, et al. Comparison of high-definition oscillometry – a non-invasive technology for arterial blood pressure measurement – with a direct invasive method using radio-telemetry in awake healthy cats. *J Feline Med Surg* 2013; 15: 1104–1113.
- 32 Laflamme D. Development and validation of a body condition score system for cats: a clinical tool. *Feline Pract* 1997; 25: 13–18.
- 33 Thomas WP, Gaber CE, Jacobs GJ, et al. Recommendations for standards in transthoracic two-dimensional echocardiography in the dog and cat. Echocardiography Committee of the Specialty of Cardiology, American College of Veterinary Internal Medicine. *J Vet Intern Med* 1993; 7: 247–252.
- 34 Hansson K, Haggstrom J, Kvart C, et al. Left atrial to aortic root indices using two-dimensional and M-mode echocardiography in cavalier King Charles spaniels with and without left atrial enlargement. *Vet Radiol Ultrasound* 2002; 43: 568–575.
- 35 Häggström J, Andersson AO, Falk T, et al. Effect of body weight on echocardiographic measurements in 19,866 pure-bred cats with or without heart disease. *J Vet Intern Med* 2016; 30: 1601–1611.
- 36 Sahn DJ, DeMaria A, Kisslo J, et al. Recommendations regarding quantitation in M-mode echocardiography: results of a survey of echocardiographic measurements. *Circulation* 1978; 58: 1072–1083.
- 37 Wayne P. User verification of precision and estimation of bias; approved guideline. CLSI document EP15-A3. 3rd ed. Wayne, PA: CLSI, 2014.
- 38 R Core Team. **R: a language and environment for statistical computing.** Vienna: R Foundation for Statistical Computing, 2020.
- 39 Giannitsis E, Mueller-Hennessen M, Zeller T, et al. Gender-specific reference values for high-sensitivity cardiac troponin T and I in well-phenotyped healthy individuals and validity of high-sensitivity assay designation. *Clin Biochem* 2020; 78: 18–24.
- 40 Clerico A, Ripoli A, Masotti S, et al. Evaluation of 99th percentile and reference change values of a high-sensitivity cTnI method: a multicenter study. *Clin Chim Acta* 2019; 493: 156–161.
- 41 Venge P, Johnston N, Lagerqvist B, et al. Clinical and analytical performance of the liaison cardiac troponin I assay in unstable coronary artery disease, and the impact of age on the definition of reference limits. A FRISC-II sub-study. *Clin Chem* 2003; 49: 880–886.
- 42 Oyama MA and Sisson DD. Cardiac troponin-I concentration in dogs with cardiac disease. *J Vet Intern Med* 2004; 18: 831–839.
- 43 Ljungvall I, Höglund K, Tidholm A, et al. Cardiac troponin I is associated with severity of myxomatous mitral valve disease, age, and C-reactive protein in dogs. *J Vet Intern Med* 2010; 24: 153–159.
- 44 Quimby JM, Smith ML and Lunn KF. Evaluation of the effects of hospital visit stress on physiologic parameters in the cat. *J Feline Med Surg* 2011; 13: 733–737.
- 45 Belew AM, Barlett T and Brown SA. Evaluation of the white-coat effect in cats. *J Vet Intern Med* 1999; 13: 134–142.
- 46 Heinonen I, Kalliokoski KK, Hannukainen JC, et al. Organ-specific physiological responses to acute physical exercise and long-term training in humans. *Physiol J* 2014; 29: 421–436.
- 47 Nie J, Tong TK, Shi Q, et al. Serum cardiac troponin response in adolescents playing basketball. *Int J Sports Med* 2008; 29: 449–452.
- 48 Pourmohammad R, Mohri M, Seifi HA, et al. Evaluation of cardiac troponin I, atrial natriuretic peptide and some oxidative/antioxidative biomarkers in the serum and hemolysate of trained Arabian horses after exercise. *Iran J Vet Res* 2020; 21: 211–215.
- 49 Rossi TM, Kavsak PA, Maxie MG, et al. Post-exercise cardiac troponin I release and clearance in normal Standardbred racehorses. *Equine Vet J* 2019; 51: 97–101.
- 50 Nostell K and Häggström J. Resting concentrations of cardiac troponin I in fit horses and effect of racing. *J Vet Cardiol* 2008; 10: 105–109.
- 51 Gresslien T and Agewall S. Troponin and exercise. *Int J Cardiol* 2016; 221: 609–621.
- 52 Stavroulakis GA and George KP. Exercise-induced release of troponin. *Clin Cardiol* 2020; 43: 872–881.
- 53 Kubo T, Kitaoka H, Okawa M, et al. Combined measurements of cardiac troponin I and brain natriuretic peptide are useful for predicting adverse outcomes in hypertrophic cardiomyopathy. *Circ J* 2011; 75: 919–926.
- 54 Borgeat K, Sherwood K, Payne JR, et al. Plasma cardiac troponin I concentration and cardiac death in cats with hypertrophic cardiomyopathy. *J Vet Intern Med* 2014; 28: 1731–1737.
- 55 Fox PR, Liu SK and Maron BJ. Echocardiographic assessment of spontaneously occurring feline hypertrophic cardiomyopathy. An animal model of human disease. *Circulation* 1995; 92: 2645–2651.
- 56 Klues HG, Schiffrers A and Maron BJ. Phenotypic spectrum and patterns of left ventricular hypertrophy in hypertrophic cardiomyopathy: morphologic observations and significance as assessed by two-dimensional echocardiography in 600 patients. *J Am Coll Cardiol* 1995; 26: 1699–1708.
- 57 Maron BJ, McKenna WJ, Danielson GK, et al. American College of Cardiology/European Society of Cardiology clinical expert consensus document on hypertrophic cardiomyopathy. A report of the American College of Cardiology Foundation Task Force on Clinical Expert Consensus Documents and the European Society of Cardiology Committee for Practice Guidelines. *J Am Coll Cardiol* 2003; 42: 1687–1713.

- 58 Häggstrom J, Luis Fuentes V and Wess G. **Screening for hypertrophic cardiomyopathy in cats.** *J Vet Cardiol* 2015; 17 Suppl 1: S134–S149.
- 59 Wagner T, Fuentes VL, Payne JR, et al. **Comparison of auscultatory and echocardiographic findings in healthy adult cats.** *J Vet Cardiol* 2010; 12: 171–182.
- 60 Marz I, Wilkie LJ, Harrington N, et al. **Familial cardiomyopathy in Norwegian Forest Cats.** *J Feline Med Surg* 2015; 17: 681–691.
- 61 Mottet E, Amberger C, Doherr MG, et al. **Echocardiographic parameters in healthy young adult Sphynx cats.** *Schweiz Arch Tierheilkd* 2012; 154: 75–80.
- 62 Scansen BA and Morgan KL. **Reference intervals and allometric scaling of echocardiographic measurements in Bengal cats.** *J Vet Cardiol* 2015; 17 Suppl 1: S282–S295.
- 63 Schober K, Savino S and Yildiz V. **Reference intervals and allometric scaling of two-dimensional echocardiographic measurements in 150 healthy cats.** *J Vet Med Sci* 2017; 79: 1764–1771.
- 64 Schober KE, Zientek J, Li X, et al. **Effect of treatment with atenolol on 5-year survival in cats with preclinical (asymptomatic) hypertrophic cardiomyopathy.** *J Vet Cardiol* 2013; 15: 93–104.
- 65 Payne JR, Borgeat K, Brodbelt DC, et al. **Risk factors associated with sudden death vs. congestive heart failure or arterial thromboembolism in cats with hypertrophic cardiomyopathy.** *J Vet Cardiol* 2015; 17 Suppl 1: S318–S328.
- 66 Payne J, Fuentes VL, Boswood A, et al. **Population characteristics and survival in 127 referred cats with hypertrophic cardiomyopathy (1997 to 2005).** *J Small Anim Pract* 2010; 51: 540–547.
- 67 Ironside VA, Tricklebank PR and Boswood A. **Risk indicators in cats with preclinical hypertrophic cardiomyopathy: a prospective cohort study.** *J Feline Med Surg* 2021; 23: 149–159.
- 68 Maron MS, Olivotto I, Maron BJ, et al. **The case for myocardial ischemia in hypertrophic cardiomyopathy.** *J Am Coll Cardiol* 2009; 54: 866–875.
- 69 Falk T, Ljungvall I, Zois NE, et al. **Cardiac troponin-I concentration, myocardial arteriosclerosis, and fibrosis in dogs with congestive heart failure because of myxomatous mitral valve disease.** *J Vet Intern Med* 2013; 27: 500–506.
- 70 Krintus M, Kozinski M, Boudry P, et al. **European multi-center analytical evaluation of the Abbott ARCHITECT STAT high sensitive troponin I immunoassay.** *Clin Chem Lab Med* 2014; 52: 1657–1665.
- 71 Egger M, Dieplinger B and Mueller T. **One-year in vitro stability of cardiac troponins and galectin-3 in different sample types.** *Clin Chim Acta* 2018; 476: 117–122.
- 72 Klüser L, Maier ET and Wess G. **Evaluation of a high-sensitivity cardiac troponin I assay compared to a first-generation cardiac troponin I assay in Doberman Pinschers with and without dilated cardiomyopathy.** *J Vet Intern Med* 2019; 33: 54–63.
- 73 Usher-Smith JA, Sharp SJ and Griffin SJ. **The spectrum effect in tests for risk prediction, screening, and diagnosis.** *BMJ* 2016; 353: i3139. DOI: 10.1136/bmj.i3139.