

# Comparing Patient Satisfaction with Automated Drug Dispensing System and Traditional Drug Dispensing System: A Cross-Sectional Study

Palanisamy Amirthalingam<sup>1</sup>, Abdulrahman Sulaiman Alruwaili<sup>2</sup>, Omar Ahmed Albalawi<sup>2</sup>, Fayez Mohammed Alatawi<sup>2</sup>, Saleh F Alqifari<sup>1</sup>, Ahmed D Alatawi<sup>3</sup>, Ahmed Aljabri<sup>4</sup>

<sup>1</sup>Department of Pharmacy Practice, Faculty of Pharmacy, University of Tabuk, Tabuk, Saudi Arabia; <sup>2</sup>Pharm.D Program, Faculty of Pharmacy, University of Tabuk, Tabuk, Saudi Arabia; <sup>3</sup>Department of Clinical Pharmacy, College of Pharmacy, Jouf University, Sakaka, Saudi Arabia; <sup>4</sup>Department of Pharmacy Practice, Faculty of Pharmacy, King Abdulaziz University, Jeddah, Saudi Arabia

Correspondence: Palanisamy Amirthalingam, Department of Pharmacy Practice, Faculty of Pharmacy, University of Tabuk, P.O. Box 741, Tabuk, 71491, Saudi Arabia, Email pchettiar@ut.edu.sa

**Introduction:** The adoption of automated drug dispensing systems (ADDS) in hospital pharmacies is a global trend, driven by its potential to reduce dispensing errors, minimize prescription filling time, and ultimately, improve patient care services. However, a significant research gap exists in the field, as a comprehensive assessment of patient satisfaction with ADDS is currently lacking. This study, with its comprehensive approach, aims to fill this gap by comparing patient satisfaction between hospital pharmacies implementing ADDS and traditional drug dispensing systems (TDDS).

**Patients and Methods:** The cross-sectional study was conducted in hospitals adopting ADDS and TDDS. All the outpatients aged 18 or above who visited the pharmacy were included, and severely ill patients were excluded from the study. A 17-item, 5-point Likert scale questionnaire assessed the participant's satisfaction. The questionnaire has four domains: pharmacy administration, dispensing practice, patient education, and dispensing system.

**Results:** The demographics of the study participants were normally distributed between ADDS and TDDS according to chi-square analysis. The mean participant satisfaction was significantly ( $P < 0.05$ ) higher in ADDS than in TDDS regarding all the items of dispensing practice and dispensing system domains. Three items related to the pharmacy administration domain showed significant participant satisfaction with ADDS. However, the participants' satisfaction showed no significant difference ( $p = 0.176$ ) between ADDS and TDDS in terms of the cleanliness of the pharmacy. Also, the participant's satisfaction between ADDS and TDDS was not statistically significant regarding the pharmacist's explanation of the side effects ( $p = 0.850$ ) and provision of all necessary information to the patient ( $p = 0.061$ ) in the patient education domain.

**Conclusion:** Patient satisfaction was higher in the ADDS participants than in TDDS regarding pharmacy administration, patient education, dispensing practice, and systems. However, pharmacists in ADDS need to be motivated to transfer the advantages of ADDS to patient care, including comprehensive patient education, particularly on side effects.

**Keywords:** automated drug dispensing system, dispensing practice, dispensing system, patient satisfaction, pharmacy administration, traditional drug dispensing system

## Introduction

The Pharmacist's role in dispensing medication is crucial and integral to patient care. It involves a complex process that combines technologies and human interaction. Dispensing errors are one component of medication errors that disrupt the achievement of therapeutic outcomes. It is a "discrepancy between a prescription and the medicine that the pharmacy delivers to the patient or distributes to the ward based on this prescription, including dispensing a medicine with inferior pharmaceutical or informational quality."<sup>1,2</sup> The most common dispensing errors reported by the previous studies include wrong medication dispensed, wrong medication strength, and labeling errors.<sup>3</sup> A recent meta-analysis, which includes 62 studies published between 2010 and 2023, reported that the pooled prevalence of dispensing error is 1.6% (95%

confidence interval 1.2% to 2.1%). They added that the prevalence of dispensing errors ranged from country to country between 0 to 33%.<sup>4</sup> Previous reports stated that 5.8% of medication errors were hazardous, and 0.8% were fatal.<sup>5</sup>

According to a systematic review, the introduction of automated drug dispensing systems (ADDS) globally has shown significant potential in reducing errors and associated risks, thereby enhancing medication safety.<sup>6</sup> Pharmacists and nurses identify 30 to 70% of medication errors in traditional drug dispensing systems (TDDS), which are almost nullified by ADDS and improve patient safety.<sup>7,8</sup>

Pharmacists have expressed a remarkably positive perception of its effectiveness in medication dispensing. They also believe that the time saved through ADDS can be redirected towards patient care, such as patient education and addressing patient queries. This potential to free up more time for direct patient care is a promising aspect of ADDS that can enhance the overall patient experience.<sup>9</sup> In ADDs, the pharmacist had much more time to collaborate with the physician and nurses to check their orders related to patients' drug profiles, medication reconciliation, and ward round participation. Moreover, the ADDS was cost-effective compared to the TDDS.<sup>10</sup> Another advantage of ADDS is that it requires smaller warehouses with improved inventory control and reduced storage errors.<sup>11,12</sup>

Although ADDS was effective in dispensing, it had several barriers and pitfalls before and during implementation.<sup>5</sup> First, as per the recent study reports, 0.018% of dispensing errors, including wrong drugs and quantities, were still detected even after ADDS was implemented.<sup>13,14</sup> Second, it increases the workload for pharmacy technicians in the cart-fill process since collaborating with them was reported as critical.<sup>13,15</sup>

The successful implementation of ADDS is independent of the hospital according to its policies and procedures. Therefore, the outcomes of ADDS implementation vary from one hospital to another.<sup>6-8</sup> Hence, the careful implementation of ADDs will have higher benefits. Obtaining patient satisfaction will play an important role in successfully implementing the drug dispensing system by identifying pitfalls at the hospital level.<sup>16</sup> To our knowledge, patient satisfaction with ADDS is not widely established in this context. Hence, the present study aimed to establish patient satisfaction with ADDS by comparing it with TDDS.

## Materials and Methods

### Study Design and Site

A cross-sectional study was carried out to compare the patient perceptions regarding outpatient pharmacy services between ADDS and TDDS for six months from February to July 2023. Two Governmental Hospitals belonging to the Ministry of Health in Tabuk city, one that adopted an ADDS and the other that embraced a TDDS, were included. Both hospitals had similarities regarding governance, bed capacity, medical departments, laboratory facilities, number of in-patients, out-patients, and healthcare practitioners.<sup>9</sup>

### Participants

This study included all participants aged 18 years or above who visited the outpatient pharmacy during the study period. It excluded the severely ill patients (eg, prolonged cough, severe pain, etc).

### Questionnaire

To obtain patient satisfaction with the drug dispensing system, a previously validated 5-point Likert scale questionnaire with 17 items was used.<sup>17</sup> The necessary permission was obtained from the original authors of the questionnaire. The questionnaire has 17 items in four domains: pharmacy administration (4 items), dispensing practice (3 items), patient education (5 items), and dispensing system (5 items). The response was recorded as strongly agree, agree, neutral, disagree, and strongly disagree, with scores of 5,4,3,2 and 1, respectively. The questionnaire also has demographic details of the participants, including gender, age, nationality, education, residence, marital status, employment, type of care, and number of visits to the pharmacy per year.

### Sampling and Data Collection Method

Following the inclusion and exclusion criteria in the participant section mentioned above, a convenient sampling method was used to recruit the study participants. After obtaining their medications, patients who visited the outpatient pharmacy

were requested to participate in the study. The reasons for refusals were also documented. The participants were thoroughly explained about the study objectives and expected outcomes. The consented participants were included in the study.


## Statistics

Chi-square statistics were used to analyze the distribution demographics of the study population. The Mean ( $\pm$  Standard Deviation) value of the participant's perception was compared between the ADDS and TDDS using the Mann–Whitney *U*-test. A confidence interval of 95% was used, and  $p < 0.05$  was considered statistically significant. A statistical package of social sciences (SPSS Version 25.0) was used for the statistical analysis.

## Results

### Recruitment Process of the Study Population

Figure 1 shows the details of the recruitment process for the study participants. A total of 503 and 496 participants visited the pharmacies of ADDS and TDDS, respectively. According to the inclusion and exclusion criteria, 79 patients in ADDS and 75 patients in TDDS were deemed not eligible for the study. Predominantly aged  $< 18$  years, participants were excluded from both ADDS ( $n=79$ ) and TDDS ( $n=75$ ). Eighty-seven participants in ADDS and 78 participants in TDDS



**Figure 1** Details of inclusion of participants for a survey.

refused to participate in the study. Uninterested was the main reason for the refusal in both ADDS (n=39) and TDDS (n=35). Finally, 337 participants in the ADDS and 343 in the TDDS consented to participate in the study.

## Demographics of the Study Population

Table 1 shows the demographics of the study population. According to chi-square statistics, there is no significant statistical difference in the distribution of the participants' demographics between ADDS and TDDS. ADDS and TDDS had a predominant distribution of males over 65, Saudi nationals who had school education, were married, had private employment, were in acute care, and visited the pharmacy twice a year. The predominant population in ADDS (53.41%) resides in the city, and TDDS (53.93%) resides outside the city. However, this result has no significant statistical difference ( $\chi^2=3.672$ ;  $df=1$ ;  $p=0.055$ ).

## Level of Participant's Satisfaction Between ADDS and TDDS

The mean (standard deviation) comparison of participant's satisfaction levels between ADDS and TDDS was analyzed using the Mann-Whitney *U*-test, represented in Table 2. The ADDS participants were significantly more satisfied

**Table 1** Demographics of the Study Population

Demographics	Total (680) n (%)	ADDS (337) n (%)	TDDS (343) n (%)	$\chi^2$	df	p
Gender						
Male	376 (55.29)	184 (54.60)	192 (55.98)	0.130	1	0.718
Female	304 (44.70)	153 (45.40)	151 (44.02)			
Age (in years)						
Less than 30	85 (12.5)	36 (10.68)	49 (14.28)	2.019	2	0.364
31–65	225 (33.08)	114 (33.82)	111 (32.36)			
More than 65	370 (54.41)	187 (55.48)	183 (53.35)			
Nationality						
Saudi	648 (95.29)	318 (94.36)	330 (96.21)	1.294	1	0.255
Non-Saudi	32 (4.70)	19 (5.64)	13 (3.79)			
Education						
Illiterate	6 (0.88)	2 (0.59)	4 (1.16)	3.922	2	0.141
School	385 (56.61)	203 (60.23)	182 (53.06)			
Graduates/Diploma	289 (42.5)	132 (39.16)	157 (45.77)			
Residence						
City	338 (49.70)	180 (53.41)	158 (46.06)	3.672	1	0.055
Outside City	342 (50.29)	157 (46.59)	185 (53.93)			
Marital Status						
Single	204 (30)	114 (33.82)	90 (26.23)	7.255	3	0.064
Married	459 (67.5)	213 (63.20)	246 (71.72)			
Divorcee	9 (1.32)	4 (1.19)	5 (1.46)			
Widow/Widower	8 (1.17)	6 (1.78)	2 (0.58)			
Employment						
Government	114 (16.76)	54 (16.02)	60 (17.49)	6.416	5	0.268
Military	42 (6.17)	21 (6.23)	21 (6.12)			
Private	369 (54.26)	182 (54)	187 (54.51)			
Business	45 (6.61)	29 (8.60)	16 (4.66)			
Student	84 (12.35)	36 (10.68)	48 (13.99)			
Retired	26 (3.82)	15 (4.45)	11 (3.20)			

(Continued)

**Table 1** (Continued).

Demographics	Total (680) n (%)	ADDS (337) n (%)	TDDS (343) n (%)	$\chi^2$	df	p
Patient care						
Acute	501 (73.67)	238 (70.62)	263 (76.68)	3.212	1	0.073
Chronic	179 (26.32)	99 (29.37)	80 (23.32)			
Number of visits/years						
1	111 (16.32)	59 (17.50)	52 (15.16)	3.251	4	0.517
2	296 (43.52)	151 (44.80)	145 (42.27)			
3	132 (19.41)	60 (17.80)	72 (20.99)			
4	91 (13.38)	40 (11.87)	51 (14.86)			
Five or more than 5	50 (7.35)	27 (8.01)	23 (6.71)			

**Note:**  $p < 0.05$  considered as statistically significant (mentioned in bold letters).

**Abbreviations:** ADDS, Automated Drug Dispensing System; TDDS, Traditional Drug Dispensing System; df, Degrees of freedom.

**Table 2** Level of Participant's Satisfaction Between ADDs and TDDs

Domain	Item	ADDS Mean (SD)	TDDS Mean (SD)	Z	p
Pharmacy Administration	Maintenance of the pharmacy is good	4.04 (0.68)	3.91 (0.71)	-2.318	<b>0.020</b>
	The pharmacy area is neat and hygienic	3.59 (0.96)	3.50 (0.80)	-1.353	0.176
	The Pharmacy has sufficient space for the drug dispensing process	3.41 (1.51)	3.01 (0.98)	-4.901	<b>0.000</b>
	A sufficient number of pharmacists working in the pharmacy	4.00 (0.87)	3.26 (0.98)	-9.963	<b>0.000</b>
Dispensing Practice	The medication label is clear and understandable	4.07 (0.67)	3.84 (0.71)	-4.298	<b>0.000</b>
	The Pharmacist dispensed all the medications in the correct quantity	4.09 (0.71)	3.88 (0.66)	-4.283	<b>0.000</b>
	The Pharmacist verifies all the medications before dispensing	3.59 (1.01)	3.35 (0.39)	-3.227	<b>0.001</b>
Patient Education	The Pharmacist explains how to take medications	4.07 (0.65)	3.87 (0.67)	-4.011	<b>0.000</b>
	The Pharmacist explains the side effects of medications	3.40 (1.15)	3.41 (1.04)	-0.189	0.850
	The Pharmacist provides all the necessary information for the patient	3.85 (0.82)	3.74 (0.72)	-1.876	0.061
	The Pharmacist always listened to the patients and cleared doubts	3.78 (0.81)	3.63 (0.70)	-2.494	<b>0.013</b>
	The Pharmacist ensures that the patient fully understands the explanation given	3.67 (0.94)	3.42 (0.86)	-3.693	<b>0.000</b>
Dispensing System	This drug dispensing system is efficient	3.83 (0.80)	3.59 (0.73)	-4.289	<b>0.000</b>
	The drug dispensing system is safe for the patient	3.82 (0.79)	3.60 (0.69)	-3.574	<b>0.000</b>
	This dispensing system reduces patient time	3.64 (0.96)	3.41 (0.87)	-3.326	<b>0.001</b>
	This dispensing system reduces pharmacists' time	3.74 (0.79)	3.46 (0.74)	-4.150	<b>0.000</b>
	This dispensing system is helpful to the hospital	3.91 (0.82)	3.67 (0.70)	-4.678	<b>0.000</b>

**Note:**  $p < 0.05$  considered as statistically significant (mentioned in bold letters).

**Abbreviations:** ADDS, Automated drug dispensing system; TDDS, Traditional Drug Dispensing System.

( $p < 0.05$ ) than the participants in TDDS with all the items regarding the dispensing practice and dispensing system domains. The participants in ADDS believed that their dispensing systems were efficient, safe for the patient, helpful to the hospital, and reduced patient and pharmacist time. Also, they added that regarding the dispensing practice domain, the medication label was clear and understandable, the pharmacist dispensed all the medications in the correct quantity, and the drugs were readily available.

In the pharmacy administration domain, the participants in ADDS were significantly more satisfied with three out of four items than those in TDDS. They felt the pharmacy's maintenance was good; sufficient seating was available in the waiting area near the pharmacy, and a sufficient number of pharmacists were working there. Meanwhile, the satisfaction level regarding the neatness and hygiene of the pharmacy was higher in ADDS participants than in TDDS participants; it did not reach a statistically significant difference ( $p = 0.176$ ).

Meanwhile, two items in the patient education domain also have participant satisfaction levels between ADDS and TDDS, which have no statistical significance. Those were where the pharmacists explained the side effects of the medications ( $p = 0.850$ ), and the pharmacist provided all the necessary information for the patient ( $p = 0.061$ ). On the other hand, the participants in ADDS had statistically more satisfaction than TDDS in the items, including the pharmacist explaining how to take the medication, the pharmacists always listening to the patients and clearing doubts, and the pharmacist ensuring that the patient fully understood the explanation.

## Discussion

This survey addresses patient satisfaction using a validated 17-item questionnaire for the first time. The survey was implemented simultaneously in pharmacies that adopted ADDs and TDDs, and the level of patient satisfaction was compared between them. According to the demographics, the study participants were normally distributed between ADDS and TDDS. The participants were significantly more satisfied with ADDs than TDDs in all items related to dispensing practice and system domain. The participants in ADDS had significantly higher satisfaction than those in TDDS regarding the dispensing system's efficiency, safety, and helpfulness to the hospital. These results authenticate the previous findings regarding the efficiency of automated drug dispensing system.<sup>7,8,11,15,18–21</sup>

The participants in ADDS believed that automation could reduce the patient and pharmacist time since it minimizes the workload of pharmacists by reducing the prescription filling time, and this finding substantiates the previous reports.<sup>7,8,13,20,22–24</sup> This saved time could be translated into patient care by providing appropriate patient education to improve the therapeutic outcomes, as already emphasized by the previous researchers.<sup>9,13</sup> This study authenticates those findings since participants in ADDS showed higher satisfaction regarding the pharmacist's involvement in educating the patients on taking medications, listening to them, clearing their doubts, and ensuring their understanding of their medications. However, the participants in ADDS had no significantly higher satisfaction than TDDS participants regarding the pharmacist's involvement in explaining side effects and providing all necessary information for the patient. Hence, pharmacists in ADDS need more motivation to translate their freed-up time, as has already been addressed by previous research.<sup>9</sup> Only a very low proportion of pharmacists (30.4%) explain possible side effects<sup>25</sup> that can be improved by effectively implementing ADDS. Provision of all necessary information to the patient regarding the drug inarguably improves medication safety and patient compliance.<sup>26</sup> In this regard, the institutions adopting ADDS should ensure improved patient care by providing good quality patient education from the pharmacists.

The participants perceived the dispensing practice in ADDS more than in the TDDS regarding pharmacist verification of medications before dispensing, clear medication labeling, and dispensing of medications in the correct quantity. These results are consistent with the previous reports, which addressed the efficient dispensing process with reduced errors.<sup>10,20,21,27,28</sup>

In this study, both hospitals belong to the Ministry of Health, and healthcare facilities are almost identical. However, one hospital adopted ADDS, and another one used TDDS. According to participants' satisfaction levels, the hospital adopting ADDS had better pharmacy administration regarding maintenance, sufficient space for dispensing, and a sufficient number of pharmacists. Adopting ADDS helps improve pharmacy administration by minimizing the shortage of pharmacists' workload, and prescription filling time.<sup>22–24,29–31</sup>

Patient satisfaction with pharmacy services is multifactorial and varies from hospital to hospital. It includes pharmacy administration, patient education services, dispensing practice, and systems.<sup>32–35</sup> Therefore, the institutions adopting ADDS should assess the level of patient satisfaction for effective implementation.

## Strength and Limitations

The following are considered strengths of this study: 1. The study adopted a validated questionnaire for the first time to assess patient satisfaction between ADDS and TDDS. 2. Both hospitals that adopted ADDS and TDDS belong to similar governance and are almost equal in facilities. 3. The demographic findings revealed that the study participants were normally distributed between ADDS and TDDS.

The study also has some limitations, such as it does not include ethnicity, details of comorbidities, and the number of drugs in a prescription, which could be potential factors in patient satisfaction with pharmacy services. The efficiency of the dispensing process can be influenced directly by the number of drugs in a prescription and indirectly by individuals' comorbidities (which determine the number of drugs in each prescription). Future researchers in ADDS can address these characteristics.

## Conclusion

Patient satisfaction was higher in the ADDS participants than in TDDS regarding pharmacy administration, patient education, dispensing practice, and systems. However, pharmacists in ADDS need to be motivated to transfer the advantages of ADDS to patient care, including comprehensive patient education, particularly on side effects. The hospital adopting ADDS can periodically assess the acquired benefits of ADDS with patient satisfaction.

## Data Sharing Statement

The datasets used and analyzed during the current study are available from the corresponding author upon reasonable request.

## Ethical Approval and Patient Consent

The Institutional Review Board, Ministry of Health, approved the study (TU-077/023/182). This study complies with the Declaration of Helsinki. Before enrolment, written consent was obtained from all participants. The study was conducted in two Government hospitals (study sites) belonging to the Ministry of Health (MOH); hence, obtaining IRB approval from the MOH is mandatory in Saudi Arabia. Meanwhile, healthcare professionals at the study sites did not contribute to or co-author the paper. However, we acknowledged the hospital administrators for permitting us to conduct the study. I assure you there is no conflict of interest regarding authorship and affiliation.

## Acknowledgments

The authors thank the hospital administrators and the participants for conducting this study.

## Disclosure

The authors declare no conflict of interest.

## References

1. Aldhwaihi K, Schifano F, Pezzolesi C, Umaru N. A systematic review of the nature of dispensing errors in hospital pharmacies. *Integr Pharm Res Pract.* 2016;5:1–10. doi:10.2147/IPRP.S95733
2. Cheung KC, Bouvy ML, De Smet PA. Medication errors: the importance of safe dispensing. *Br J Clin Pharmacol.* 2009;67(6):676–680. doi:10.1111/j.1365-2125.2009.03428.x
3. Maharaj S, Brahim A, Brown H, et al. Identifying dispensing errors in pharmacies in a medical science school in Trinidad and Tobago. *J Pharm Policy Pract.* 2020;13(1):67. doi:10.1186/s40545-020-00263-x
4. Um IS, Clough A, Tan ECK. Dispensing error rates in pharmacy: a systematic review and meta-analysis. *Res Social Adm Pharm.* 2024;20(1):1–9. doi:10.1016/j.sapharm.2023.10.003
5. Mulac A, Taxis K, Hagesaether E, Gerd Granas A. Severe and fatal medication errors in hospitals: findings from the Norwegian incident reporting system. *Eur J Hosp Pharm.* 2021;28(Suppl 2):e56–e61. doi:10.1136/ejhpharm-2020-002298



6. Ahtiainen HK, Kallio MM, Airaksinen M, Holmström AR. Safety, time and cost evaluation of automated and semi-automated drug distribution systems in hospitals: a systematic review. *Eur J Hosp Pharm.* 2020;27(5):253–262. doi:10.1136/ejpharm-2018-001791
7. Tariq RA, Vashisht R, Sinha A, et al. Medication dispensing errors and prevention. In: *StatPearls*. Treasure Island (FL): StatPearls Publishing; 2024.
8. Almalki A, Jambi A, Elbehiry B, Albuti H. Improving inpatient medication dispensing with an automated system. *Glob J Qual Saf Healthc.* 2023;6(4):117–125. doi:10.36401/JQSH-23-15
9. Alanazi MF, Shahein MI, Alsharif HM, et al. Impact of automated drug dispensing system on patient safety. *Pharm Pract.* 2022;20(4):2744. doi:10.18549/PharmPract.2022.4.2744
10. Chapuis C, Bedouch P, Detavernier M, et al. Automated drug dispensing systems in the intensive care unit: a financial analysis. *Crit Care.* 2015;19(1):318. doi:10.1186/s13054-015-1041-3
11. Tsao NW, Lo C, Babich M, Shah K, Bansback NJ. Decentralized automated dispensing devices: systematic review of clinical and economic impacts in hospitals. *Can J Hosp Pharm.* 2014;67(2):138–148. doi:10.4212/cjhp.v67i2.1343
12. Oldland AR, Golightly LK, May SK, Barber GR, Stolpmann NM. Electronic inventory systems and barcode technology: impact on pharmacy technical accuracy and error liability. *Hosp Pharm.* 2015;50(1):34–41. doi:10.1310/hpj5001-34
13. Takase T, Masumoto N, Shibatani N, et al. Evaluating the safety and efficiency of robotic dispensing systems. *J Pharm Health Care Sci.* 2022;8(1):24. doi:10.1186/s40780-022-00255-w
14. Bagattini AM, Borges JLA, Riera R, de Carvalho DCMF. Automation of a tertiary hospital pharmacy drug dispensing system in a lower-middle-income country: a case study and preliminary results. *Explor Res Clin Soc Pharm.* 2022;6:100151. doi:10.1016/j.rcsop.2022.100151
15. Temple J, Ludwig B. Implementation and evaluation of carousel dispensing technology in a university medical center pharmacy. *Am J Health Syst Pharm.* 2010;67(10):821–829. doi:10.2146/ajhp090307
16. Ali MAS, Amirthalingam P, Alatawi Y, Aljabri A. Patient satisfaction of ambulatory care pharmacy services. *J Patient Exp.* 2022;9:23743735221112631. doi:10.1177/23743735221112631
17. Amirthalingam P, Alharbe UA, Almfalh HSS, et al. Validation of a questionnaire to assess patient satisfaction with an automated drug dispensing system. *Healthcare.* 2024;12(16):1598. doi:10.3390/healthcare12161598
18. Kobayashi K, Uenoyama K, Ito T, et al. Efforts in facilitating work through the introduction of a powdered medicine dispensing robot. *J Jpn Soc Hosp Pharm.* 2019;55:402–408.
19. Tabata H, Kaya N, Inagaki Y, Shibasaki A. Survey on use of a dispensing medicine robot (DimeRo) and considerations for efficient operation. *J Jpn Soc Hosp Pharm.* 2018;54:175–179.
20. Rodriguez-Gonzalez CG, Herranz-Alonso A, Escudero-Vilaplana V, Ais-Larigoitia MA, Iglesias-Peinado I, Sanjurjo-Saez M. Robotic dispensing improves patient safety, inventory management, and staff satisfaction in an outpatient hospital pharmacy. *J Eval Clin Pract.* 2019;25(1):28–35. doi:10.1111/jep.13014
21. Berdot S, Korb-Savoldelli V, Jaccoulet E, et al. A centralized automated-dispensing system in a French teaching hospital: return on investment and quality improvement. *Int J Qual Health Care.* 2019;31(3):219–224. doi:10.1093/intqhc/mzy152
22. Roman C, Poole S, Walker C, Smit de V, Dooley MJ. A ‘time and motion’ evaluation of automated dispensing machines in the emergency department. *Australas Emerg Nurs J.* 2016;19(2):112–117. doi:10.1016/j.aenj.2016.01.004
23. Momattin H, Arafa S, Momattin S, Rahal R, Waterson J. Robotic pharmacy implementation and outcomes in Saudi Arabia: a 21-Month Usability Study. *JMIR Hum Factors.* 2021;8(3):e28381. doi:10.2196/28381
24. Lin AC, Huang YC, Panches G, Chen Y. Effect of a robotic prescription-filling system on pharmacy staff activities and prescription-filling time. *Am J Health Syst Pharm.* 2007;64(17):1832–1839. doi:10.2146/ajhp060561
25. Tadesse YB, Sendekie AK, Mekonnen BA, Denberu FG, Kassaw AT. Pharmacists’ medication counseling practices and knowledge and satisfaction of patients with an outpatient hospital pharmacy service. *Inquiry.* 2023;60:469580231219457. doi:10.1177/00469580231219457
26. Ejeta F, Feyisa D, Kebede O, et al. Medication counseling practices in medicine retail outlets found in bench sheko zone, southern nations, nationalities, and peoples’ region, South West Ethiopia. *Pragmat Obs Res.* 2021;12:105–117. doi:10.2147/POR.S322407
27. Oswald S, Caldwell R. Dispensing error rate after implementation of an automated pharmacy carousel system. *Am J Health Syst Pharm.* 2007;64:1427–1431. doi:10.2146/ajhp060313
28. van Doormaal JE, Rommers MK, Kosterink JG, Teepe-Twiss IM, Haaijer-Ruskamp FM, Mol PG. Comparison of methods for identifying patients at risk of medication-related harm. *Qual Saf Health Care.* 2010;19(6):e26. doi:10.1136/qshc.2009.033324
29. Ramachandram DS, Tan CH, Selvaraj A, Appalasamy JR, Kow CS. The workload and satisfaction of pharmacists and pharmacy assistants after installation of an automated tablet dispensing and packaging system in a tertiary university hospital in Malaysia. *Int J Pharm Pract.* 2024;32(1):100–103. doi:10.1093/ijpp/riad085
30. Shao SC, Chan YY, Lin SJ, et al. Workload of pharmacists and the performance of pharmacy services. *PLoS One.* 2020;15(4):e0231482. doi:10.1371/journal.pone.0231482
31. Kilonzi M, Mutagonda RF, Mlyuka HJ, et al. Barriers and facilitators of integration of pharmacists in the provision of clinical pharmacy services in Tanzania. *BMC Prim Care.* 2023;24(1):72. doi:10.1186/s12875-023-02026-4
32. Altarifi D, Harb T, Abualhasan M. Patient satisfaction with pharmaceutical services at primary healthcare centers under the Palestinian Ministry of Health. *BMC Health Serv Res.* 2024;24(1):514. doi:10.1186/s12913-024-10983-4
33. Yehualaw A, Tafere C, Demsie DG, et al. Determinants of patient satisfaction with pharmacy services at Felege Hiwot comprehensive specialized hospital, Bahir Dar, Ethiopia. *Ann Med Surg.* 2023;85(12):5885–5891. doi:10.1097/MS9.0000000000001161
34. Alanazi AS, Shah S, Abbas G, et al. Assessing patient satisfaction with community pharmacy services: a large regional study at Punjab, Pakistan. *Patient Prefer Adherence.* 2023;17:13–22. doi:10.2147/PPA.S389053
35. Molla M, Sisay W, Andargie Y, Kefale B, Singh P. Patients’ satisfaction with outpatient pharmacy services and associated factors in Debre Tabor comprehensive specialized hospital, Northwest Ethiopia: a cross-sectional study. *PLoS One.* 2022;17(1):e0262300. doi:10.1371/journal.pone.0262300



## Patient Preference and Adherence

Dovepress

**Publish your work in this journal**

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focusing on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize clinical outcomes for existing disease states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/patient-preference-and-adherence-journal>