

POSTER PRESENTATION

Open Access

Abnormalities in aortic arch geometry do not lead to reduced exercise performance: a comparison study between patients with transposition of the great arteries repaired by arterial switch operation and normal controls

Hopewell Ntsinjana^{1,2*}, Giovanni Biglino^{1,2}, Claudio Capelli^{1,2}, Alessandro Giardini², Graham Derrick², Silvia Schievano^{1,2}, Andrew M Taylor^{1,2}

From 16th Annual SCMR Scientific Sessions
San Francisco, CA, USA. 31 January - 3 February 2013

Background

Abnormal aortic arch geometry, in particular the 'gothic arch', has been associated with reduced exercise performance in patients with repaired aortic coarctation. Following the arterial switch operation (ASO) for transposition of the great arteries (TGA), the morphology of the aorta is also known to be abnormal, presenting with dilated aortic root and 'gothic' deformity of the aortic arch. In this study, our aim was to compare aortic geometry between normal and ASO patients to establish if aortic arch shape influenced exercise performance.

Methods

We studied 17 TGA patients repaired by ASO (age=15.3 ±1.8 years, BSA=1.8±0.2 m²) and 17 matched controls (age=14.9±1.9 years, BSA=1.6±0.2 m²). All subjects signed informed consent. 3D b-SSFP whole heart CMR images were acquired for 3D volume reconstruction. 3D anatomical models of the left heart including LV, LVOT, aortic root, ascending aorta, aortic arch and descending aorta to the level of diaphragm were created using commercial software (Mimics). Geometrical analysis (Figure 1) was performed measuring: (i) the angle between the line connecting the centre of the valve with the apex of the LV and the aortic valve plane; (ii) the angle between the aortic valve plane and the sinotubular junction plane; (iii) the

indexed length of the vessel centreline starting from the centre of the aortic valve to the level of the diaphragm; (iv) the indexed curvature of the aortic arch, from the inverse of the radius (r) of the maximum circumference fitted at the highest point of the centreline ($=1/r$). On the same day as the CMR scan, cardiopulmonary exercise test was performed in all subjects using an electronically braked ergometer cycle. Exercise capacity, heart rate and systolic blood pressure response to exercise were measured.

Results

There were significant differences in aortic geometry between patients and controls (Table 1). However, there are no notable differences in physiological response to exercise between the two groups.

Conclusions

Using ASO as a model for alteration in aortic arch geometry without direct operation on the arch, we show that marked changes in aortic shape are not associated with changes in exercise capacity, raising the possibility that the 'gothic arch' appearances seen in aortic coarctation do not play a role in the exercise hypertensive response seen in such coarctation patients.

Funding

Commonwealth Scholarships; National Institute of Health Research UK (NIHR); Fondation Leducq; Royal Academy of Engineering and EPSRC.

¹Centre for Cardiovascular Imaging, Institute of Cardiovascular Science, University College London, London, UK
Full list of author information is available at the end of the article




Figure 1 Identification of geometrical markers and measures on 3D models of a TGA subject (A) and of a control case (B), as described in the Methods section.

Table 1 Differences in geometric variables and hemodynamic response to exercise between TGA subjects and controls. $p < 0.05$ is considered statistically significant.

Variable	TGA Subjects	Controls	p-value
LV-AV angle [°]	62 ± 9	63 ± 8	0.65
AV-STJ angle [°]	151 ± 10	172 ± 3	<0.001
Aortic centreline length [mm/m ²]	140 ± 22	123 ± 12	0.008
Aortic arch curvature [1/mm/m ²]	0.053 ± 0.016	0.020 ± 0.004	<0.001
VO ₂ max [mL/kg/min]	39.4 ± 5.0	36.6 ± 7.8	0.23
Peak blood pressure [mmHg]	151 ± 20	152 ± 21	0.96
Peak heart rate [bpm]	175 ± 10	175 ± 13	0.99

Author details

¹Centre for Cardiovascular Imaging, Institute of Cardiovascular Science, University College London, London, UK. ²Cardiorespiratory Unit, Great Ormond Street Hospital for Children NHS Foundation Trust, London, UK.

Published: 30 January 2013

doi:10.1186/1532-429X-15-S1-P291

Cite this article as: Ntsinjana *et al.*: Abnormalities in aortic arch geometry do not lead to reduced exercise performance: a comparison study between patients with transposition of the great arteries repaired by arterial switch operation and normal controls. *Journal of Cardiovascular Magnetic Resonance* 2013 **15**(Suppl 1):P291.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

