


See Article page 263.



## Commentary: There are chords in the human heart that had better not be vibrated

Giorgia Bonalumi, MD,<sup>a</sup> Ilaria Giambuzzi, MD,<sup>a,b</sup>  
Alessandro Parolari, MD, PhD,<sup>c</sup> and  
Michele Di Mauro, MD, PhD, MSc<sup>d</sup>

Posterior leaflet prolapse is the most common cause of mitral valve regurgitation in the Western world, and, notably, the easiest lesion to be repaired. The gold standard technique for its repair is resection, first introduced by Alain Carpentier,<sup>1</sup> but recently newer techniques have been proposed and have become commonly used. Perier and colleagues<sup>2</sup> introduced the “respect rather than resect” approach and, since then, many authors have published their results with the use of artificial chordae.<sup>3,4</sup> Still, a biomechanical study on the actual force needed to disrupt the artificial chordae is lacking. Mateo Marin-Cuartas and colleagues<sup>5</sup> elegantly describe the differences among 3 of the most-used techniques to implant artificial chordae: running,<sup>6</sup> standard interrupted,<sup>7</sup> and loop.<sup>8,9</sup> The neochords were attached to 3-dimensional plastic models representing papillary muscles and mitral valve leaflets and, by applying a tensile force on both fixtures, authors measured the needed force to disrupt the artificial chordae. They demonstrated that running chordae needed the greatest force to be ruptured even if that resulted in failure of the technique if there was just one running chordae. Moreover, as expected, the site of rupture was near the knot, more commonly in patients with either loop or interrupted chordae. Given these



### CENTRAL MESSAGE

Tensile forces on artificial mitral chordae have different effects depending on the employed surgical technique, showing that running chordae are more resistant than loop and interrupted ones.

findings, running chordae should be implanted in multiple sets, because, even if they are more resistant to disruptive forces, a single rupture would lead to the failure of the chordae. Nevertheless, the forces used in the test are greater than those physiologically experienced in patients.

Surgical manipulation, degeneration of artificial chordae, and chronically worsening tensile forces over time might explain late failure of artificial chordae, which is, at any rate, rare.<sup>10</sup> Recently, Mutsuga and colleagues<sup>4</sup> showed that also in vivo a less-resistant chordae (CV4) breaks easier than a more resistant one (CV5); of course, a thicker chordae is harder to manipulate in surgery. Marin-Cuartas and colleagues<sup>5</sup> demonstrated through bioengineering that not only are thicker chordae more resistant but are also more resistant when they are implanted in a running fashion. We believe Marin-Cuaratas and colleagues<sup>5</sup> open up the way to more biomechanical studies on the behavior under physiological conditions (and, therefore, to lower forces) of the different techniques of chordae implantation under physiological conditions, leading to a scientifically guided choice (not only based on the surgeon preferences) of the best technique to implant neochordae.

Paraphrasing what Charles Dickens wrote, we can say that “there are chords in the human heart that had better not be vibrated...,”<sup>11</sup> and maybe they are artificial chordae implanted not in a running fashion.

From the <sup>a</sup>Department of Cardiac Surgery, IRCCS Monzino Cardiology Center; <sup>b</sup>DISCO (Dipartimento di Scienze Cliniche e di Comunità), and <sup>c</sup>Unit of Cardiac Surgery and Translational Research, IRCCS Policlinico S. Donato, University of Milan, Milan, Italy; and <sup>d</sup>Cardio-Thoracic Surgery Unit, Heart and Vascular Centre, Maastricht University Medical Centre, Cardiovascular Research Institute Maastricht, Maastricht, The Netherlands.

Disclosures: The authors reported no conflicts of interest.

The *Journal* policy requires editors and reviewers to disclose conflicts of interest and to decline handling or reviewing manuscripts for which they may have a conflict of interest. The editors and reviewers of this article have no conflicts of interest.

Received for publication Aug 11, 2021; revisions received Aug 11, 2021; accepted for publication Aug 12, 2021; available ahead of print Sept 6, 2021.

Address for reprints: Michele Di Mauro, MD, PhD, MSc, Cardio-Thoracic Surgery Unit, Heart and Vascular Centre, Maastricht University Medical Centre (MUMC), P. Debyelaan 25, 6202 AZ, Maastricht, The Netherlands (E-mail: [mdimauro1973@gmail.com](mailto:mdimauro1973@gmail.com)).

JTCVS Open 2021;8:278-9  
2666-2736

Copyright © 2021 The Author(s). Published by Elsevier Inc. on behalf of The American Association for Thoracic Surgery. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).  
<https://doi.org/10.1016/j.jxjon.2021.08.017>

## References

1. Carpentier A. Cardiac valve surgery: the "French correction." *J Thorac Cardiovasc Surg.* 1983;86:323-7.
2. Perier P, Hohenberger W, Lakew F, Batz G, Urbanski P, Zacher M, et al. Toward a new paradigm for the reconstruction of posterior leaflet prolapse: midterm results of the "respect rather than resect" approach. *Ann Thorac Surg.* 2008;86:718-25.
3. Sakaguchi T, Hiraoka A, Totsugawa T, Hayashida A, Ryomoto M, Sekiya N, et al. Clinical impact of the repair technique for posterior mitral leaflet prolapse: resect or respect? *J Card Surg.* 2021;36:971-7.
4. Mutsuga M, Narita Y, Tokuda Y, Uchida W, Ito H, Terazawa S, et al. Predictors of failure of mitral valve repair using artificial chordae. *Ann Thorac Surg.* May 19, 2021 [Epub ahead of print].
5. Marin-Cuartas M, Imbrie-Moore AM, Zhu Y, Park MH, Wilkerson R, Leipzig M, et al. Biomechanical engineering analysis of commonly utilized mitral neochordae. *J Thorac Cardiovasc Surg Open.* 2021;8:263-75.
6. David TE. Replacement of chordae tendineae with expanded polytetrafluoroethylene sutures. *J Card Surg.* 1989;4:286-90.
7. Perier P. A new paradigm for the repair of posterior leaflet prolapse: respect rather than 347 resect. *Oper Tech Thorac Cardiovasc Surg.* 2005;10:180-93.
8. Von Oppell UO, Mohr FW. Chordal replacement for both minimally invasive and conventional mitral valve surgery using premeasured Gore-Tex loops. *Ann Thorac Surg.* 2000;70:2166-8.
9. Fortunato G, Battellini R, Raffaelli P, Kotowicz V. How to create patient-specific loops for correcting mitral valve prolapse through a minimally invasive approach. *Multimed Man Cardiothorac Surg.* December 19, 2019;2019.
10. Pfannmueller B, Misfeld M, Verevkin A, Garbade J, Holzhey DM, Davierwala P, et al. Loop neochord versus leaflet resection techniques for minimally invasive mitral valve repair: long term results. *Eur J Cardiothorac Surg.* 2021;59:180-6.
11. Dickens C. Barnaby Rudge: a tale of the riots of eighty. In: *Master Humphrey's Clock.* London: Chapman and Hall, London; 1840.