



Endovascular treatment of penetrating thoracic aorta injury – case report

Abordagem endovascular de lesão de aorta torácica penetrante – relato de caso

Lucas Mansano Sarquis¹ , Wilson Michaelis², Antonio Lacerda Santos Filho², Rogerio Akira Yokoyama^{1,2}, Mariana Vieira Delazeri^{1,2}, Antonio Luiz da Costa Martins², Rodrigo Krieger Martins¹, Bruno Berardi Gazola³

Abstract

In the current scenario, traumas with violent causes are responsible for large numbers of cases. Among these, thoracic aorta injury caused by penetrating trauma is a cause of elevated morbidity and mortality, demanding adequate diagnosis, and can now often be repaired using endovascular procedures. This treatment method has proven to be safer, with a lower rate of complications than open surgical procedures. After endovascular repair, it is necessary to conduct continuous monitoring of the patient's health and correct any complications related to the procedure that may emerge. The objective of this article is to describe a case of penetrating trauma of the thoracic aorta that was treated endovascularly, since the literature predominantly covers blunt trauma injuries.

Keywords: thoracic aorta; wounds and injuries; endovascular procedures.

Resumo

O cenário atual do trauma por causas violentas tem apresentado um elevado número de casos. Entre eles, a lesão de aorta torácica por trauma penetrante é uma condição de elevada morbimortalidade que exige diagnóstico adequado e que, nos últimos anos, tem sido corrigida com procedimentos endovasculares. Essa modalidade de tratamento se mostra mais segura, com menor número de complicações em relação ao procedimento cirúrgico aberto. Após a realização da abordagem endovascular, há necessidade de acompanhamento contínuo, que visa monitorar as condições de saúde do paciente e corrigir possíveis complicações relacionadas ao procedimento. O objetivo deste artigo é relatar um caso de traumatismo penetrante de aorta torácica, tratado por via endovascular, visto que a literatura aborda predominantemente as lesões por trauma contuso.

Palavras-chave: aorta torácica; ferimentos e lesões; procedimentos endovasculares.

How to cite: Sarquis LM, Michaelis W, Santos Filho AL, et al. Endovascular treatment of penetrating thoracic aorta injury – case report. *J Vasc Bras.* 2020;19:e20200132. <https://doi.org/10.1590/1677-5449.200132>

¹Hospital do Trabalhador, Curitiba, PR, Brasil.

²Hospital Universitário Evangélico Mackenzie, Curitiba, PR, Brasil.

³Faculdade Pequeno Príncipe, Curitiba, PR, Brasil.

Financial support: None.

Conflicts of interest: No conflicts of interest declared concerning the publication of this article.

Submitted: May 27, 2020. Accepted: September 04, 2020.

The study was carried out at Hospital do Trabalhador and at Hospital Universitário Evangélico Mackenzie, Curitiba, PR, Brazil.

■ INTRODUCTION

External causes are now the third greatest cause of mortality in Brazil, with predominance of interpersonal violence and traffic accidents, especially in the younger population.¹ Thoracic traumas can be classified as blunt or penetrating. Penetrating traumas are generally provoked by application of a mechanical force to a small area on the surface of the chest, causing a break in the continuity of the skin and chest wall.¹ The majority of thoracic aorta ruptures are caused by blunt thoracic trauma,² and, among the most common penetrating injuries, those that involve the aorta are predominantly caused by firearms or cold weapons, in which the quality of pre-hospital care and diagnostic tools is of fundamental importance.³ Penetrating injuries involving the thoracic aorta are rare and the majority need to be treated with open surgery, with few applications for endovascular methods, since patients are generally admitted hemodynamically unstable.⁴ Eighty percent of patients with thoracic aorta injuries die at the scene of trauma and seven in every 10 are male.⁵

After stabilization of the patient, diagnosis of aortic injury is a challenge for the surgeon, since in addition to identifying them it is also necessary to treat other associated traumatism. Concomitant injuries seen in cases of high energy trauma include severe traumatic brain injury (TBI), pulmonary and/or cardiac contusion, abdominal hemorrhage, fractures to long and/or pelvic bones, and ruptured diaphragms.⁶ Symptoms of aortic rupture can include pain in the chest, back, or abdomen, in addition to external signs of thoracic injury.⁷

Diagnostic suspicion begins with physical examination and chest X-ray; but both are of low sensitivity. Computed tomography angiography (angio-CT) is considered the gold standard, since in addition to diagnosis, it can be used to plan endovascular treatment. However, in order to conduct an examination, the patient must be stable from a hemodynamic point of view.⁴ Aortic injuries are classified as grade 1 – intimal lesion; grade 2 – pseudoaneurysm; grade 3 – intramural hematoma; or grade 4 – total rupture.⁸

The first report of thoracic endovascular aortic repair (TEVAR) was published in the 1990s and this has now become the technique that is most used, consisting of deployment of an endograft from a peripheral access. As lower profile devices have been developed with better navigation properties and treating teams have gained experience, complications related to the procedure have reduced.⁹ This method of treatment stands out, especially for traumatic aortic injuries and complicated acute thoracic aorta dissection, because

of the lower morbidity and mortality compared with traditional surgical repair.¹⁰

Given that the majority of studies cover injuries to the aorta caused by blunt traumas, the objective of this article is to highlight the importance of endovascular repair in patients who have suffered penetrating traumas with injuries to the aorta.

■ CASE DESCRIPTION

The patient was a 36-year-old male, admitted to a specialist trauma center on 24 December, 2019, at 21:00, having suffered wounds caused by a cold weapon. He had exhibited hemodynamic instability during transport, but after fluid resuscitation he was stable when he arrived at the hospital. The most notable findings of the physical examination conducted in the emergency room were subdued vesicular murmur in the left hemithorax, heart rate of 125 bpm, mean blood pressure (MBP) of 70-75 mmHg, and multiple wounds to the anterior and posterior chest (Figure 1).

After administration of packed red blood cells and prophylactic antibiotics, angio-CT of the thorax was conducted (Figures 2 and 3), showing contrast leakage from the thoracic aorta and left sided hemothorax. Left pleural drainage and arteriography of the thoracic aorta were performed.

The decision was taken to proceed directly to definitive endovascular treatment. Intraoperative aortography (Figure 4) was performed, confirming the damage to the thoracic aorta (at the level of T8) with adjacent tamponade, and a 20/20/82 mm straight thoracic endograft was deployed in the descending thoracic aorta. The endovascular intervention was conducted at 01:00 on December 25, 2019. Control aortography performed after implantation (Figure 5) showed no evidence of contrast leakage.

The patient remained in the intensive care unit during the immediate postoperative period. On the first postoperative (PO) day, the patient suffered acute myocardial infarction of the lateral wall, but without compromise to cardiac or hemodynamic function. He was discharged to the wards on PO day 3. The patient had a collection in the left hemithorax due to retained hemothorax, and underwent videothoracoscopy and decortication, conducted by the thoracic surgery team. His recovery was otherwise uneventful and he was discharged from hospital on the 10th day after admission. He is currently in outpatients follow-up with the specialties and is asymptomatic. He had a control chest angio-CT 30 days after the trauma, which found no evidence of complications related to the surgical procedure.




Figure 1. Diagram illustrating cold weapon wound sites.



Figure 2. Coronal angio-CT image showing contrast leakage from the thoracic aorta.




Figure 3. Axial angio-CT image showing contrast leakage from the thoracic aorta.



Figure 4. Aortography showing thoracic aorta injury.



Figure 5. Control aortography after implantation of an endograft.

■ DISCUSSION

Penetrating injuries to the thoracic aorta are potentially fatal and need medical care that is correctly coordinated, in order to define the order of priority of the care to be provided. Control of blood pressure is one of the first measures, aiming to achieve an MBP in the range of 60-70 mmHg, with systolic pressure below 130 mmHg, which is one of the fundamental factors in ensuring that there is enough time to attempt repair, whether conventional or endovascular.¹¹ After initial stabilization of the patient, as in this case, with transfusion of blood products, angio-CT is essential for identification of the aortic injury and assessment of the anatomy to decide on the possibility of endovascular treatment.

Treatment of the aortic injury can be postponed, generally by 1-2 weeks, but sometimes up to 45 days, according to an evaluation made in conjunction with other specialties. In the case of multiple trauma patients, treatment of life-threatening injuries should always be prioritized.¹² However, early intervention (less than 24 h from hospital admission) should be encouraged for patients with expanding pseudoaneurysms, incapacity to maintain target blood pressure, or leakage of contrast on angio-CT.¹²

Since the patient in the current report did not have other injuries that needed specific treatment, the aortic injury was treated immediately. Concurrent injuries, the grade of aortic injury, hemodynamic stability, and associated comorbidities are the determinant factors of the choice of time and type of repair.⁶ Although the injury was classified as grade 4, total rupture of the aorta combined with hemothorax, the favorable anatomy for endovascular treatment and the patient's hemodynamic stability made TEVAR a possible option. This aortic repair causes less blood loss, does not require cross-clamping of the aorta, and is associated with faster recovery than open procedures.¹³

The endograft is introduced using guides, via a common femoral artery access, and advanced until it reaches the thoracic aorta, where it is positioned. Essential characteristics of a successful procedure include choosing a prosthesis with appropriate size and shape and appropriate patient vascular anatomy. In healthy patients without prior vascular diseases, which is the situation in the majority of cases, it is possible to reduce the neck of the healthy aorta to 0.5-0.7 cm and seal the endograft, whereas in patients with preexisting vascular disease, the ideal would be to maintain a neck of 1.5-2.5 cm.¹¹ The type of device chosen is primarily dependent on the patient's anatomy, the angles involved, and the stock available for use at the time of trauma.⁴

After implantation, the patient will need lifelong monitoring with imaging exams. Guidelines suggest intervals of 30 days, 6 months, and 12 months after repair and, if there are no complications, annually thereafter.¹³ The open procedure, which requires thoracotomy or sternotomy, is associated with a 28% mortality rate and around 16% paraplegia,³ whereas overall 30-day mortality rates linked to TEVAR are around 8%.¹⁴ Common complications seen after endovascular treatment include endoleaks, graft migration, graft stenosis, and graft infection, in addition to systemic complications such as ischemia, cerebrovascular events, and inflammatory reactions to the material implanted.¹³ Endoleaks are one of the most common indications for reintervention after this type of repair, with incidence of around 12.1%.¹⁰ Acute myocardial infarction after endovascular treatment for aortic diseases is a rare cause of complications, but can be present.¹⁰ With regard to the acute myocardial infarction seen in the case described here, we cannot conclude that it was an event that occurred as a result of the endovascular procedure or even because of the penetrating trauma, but not identified during examinations conducted on admission.

It is thus concluded that endovascular treatment of the thoracic aorta is a viable option after penetrating injuries. Prospective and multicenter studies are needed to evaluate long-term results of placement of aortic endografts in patients who have been victims of penetrating wounds.

■ REFERENCES

1. Melo ASA, Moreira LBM, Pessoa FMC, et al. Aspectos tomográficos do trauma torácico aberto: Lesões por projéteis de arma de fogo e armas brancas. *Radiol Bras.* 2017;50(6):372-7. PMID:29307927.
2. Szostek M, Jakuczun W, Oseka M, et al. Endovascular treatment of traumatic rupture of thoracic aorta: long term results. *Int Angiol.* 2016;35(2):129-39. PMID:26221975.
3. Brand S, Breitenbach I, Bolzen P, Petri M, Krettek C, Teebken O. Open repair versus thoracic endovascular aortic repair in multiple-injured patients: observations from a level-1 trauma center. *Arch Trauma Res.* 2015;4(4):e27183. <http://dx.doi.org/10.5812/atr.27183>. PMID:26848470.
4. Chou EL, Twerdahl EH, Eagleton MJ. Endovascular management of penetrating and non-penetrating aortic injury. *Vasa.* 2019;48(1):23-33. <http://dx.doi.org/10.1024/0301-1526/a000740>. PMID:30227809.
5. Vargas TB, Bergo R, Ferreira R, Melo Q, Almeida L, Starling SV. Lesão da aorta torácica por trauma fechado. *Rev Médica Minas Gerais.* 2016;26(Supl 4):20-3.
6. Cavalcante EF Jr, Avezum VAPAF, Silva PJLS, Vasconcellos VRB, Falavigna GG, Espada PC. Lesão aórtica e pélvica em vítima de politrauma. *Relatos Casos Cir.* 2017;3.
7. Kumar R, Raja J, Munirathinam GK, Mishra AK, Singh RS, Thingnam SKS. A case of traumatic thoracic aorta rupture: a life threatening emergency. *J Cardiovasc Thorac Res.* 2019;11(3):248-50. <http://dx.doi.org/10.15171/jcvtr.2019.41>. PMID:31579467.

8. Lee WA, Matsumura JS, Mitchell RS, et al. Endovascular repair of traumatic thoracic aortic injury: clinical practice guidelines of the Society for Vascular Surgery. *J Vasc Surg.* 2011;53(1):187-92. <http://dx.doi.org/10.1016/j.jvs.2010.08.027>. PMID:20974523.
9. Brenner M, Teeter W, Hadud M, et al. Long-term outcomes of thoracic endovascular aortic repair: a single institution's 11-year experience. *J Trauma Acute Care Surg.* 2017;82(4):687-93. <http://dx.doi.org/10.1097/TA.0000000000001365>. PMID:28129260.
10. Quintas A, Bastos Gonçalves F, Rodrigues H, et al. Tratamento endovascular de patologia da aorta torácica: experiência institucional. *Angiol Cir Vasc.* 2016;12(1):3-11. <http://dx.doi.org/10.1016/j.ancv.2015.12.002>.
11. Kolbeck KJ, Kaufman JA. Endovascular stent grafts in urgent blunt and penetrating thoracic aortic trauma. *Semin Intervent Radiol.* 2011;28(1):98-106. <http://dx.doi.org/10.1055/s-0031-1273944>. PMID:22379280.
12. Botta L, Russo V, Savini C, et al. Endovascular treatment for acute traumatic transection of descending aorta: focus on operative timing and left subclavian artery management. *J Thorac Cardiovasc Surg.* 2008;136(6):1558-63. <http://dx.doi.org/10.1016/j.jtcvs.2008.07.040>. PMID:19114206.
13. Daye D, Walker TG. Complications of endovascular aneurysm repair of the thoracic and abdominal aorta: Evaluation and management. *Cardiovasc Diagn Ther.* 2018;8(Supl 1):S138-56. <http://dx.doi.org/10.21037/cdt.2017.09.17>. PMID:29850426.
14. Newman J, Mattia A, Manetta F. Indications for Thoracic Endovascular Aortic Repair (TEVAR): A Brief Review. *Int J Angiol.* 2018;27(4):177-84. <http://dx.doi.org/10.1055/s-0038-1666972>. PMID:30410287.

Correspondence

Lucas Mansano Sarquis
Hospital do Trabalhador
Rua Augusto Stelfeld, 1908 - Bigorrrilho
CEP 80730-150 - Curitiba (PR), Brasil
Tel.: +55 (41) 3240-5227
E-mail: lucas_sarquis@hotmail.com

Author information

LMS – Vascular surgery resident, Hospital do Trabalhador (HT).
WM – Chief and Preceptor of Vascular surgery service, Hospital Universitário Evangélico Mackenzie (HUEM).
ALSF – Preceptor of Vascular surgery service, Hospital Universitário Evangélico Mackenzie (HUEM).
RAY – Preceptor of Vascular surgery service, Hospital Universitário Evangélico Mackenzie (HUEM); Preceptor of Vascular surgery service, Hospital do Trabalhador (HT).
MVD – Preceptor of Vascular surgery service, Hospital Universitário Evangélico Mackenzie (HUEM); Preceptor of Vascular surgery service, Hospital do Trabalhador (HT).
ALCM – Endovascular surgery resident, Vascular surgery service, Hospital Universitário Evangélico Mackenzie (HUEM).
RKM – General surgery residente, Hospital do Trabalhador (HT).
BBG - Medical student, Faculdade Pequeno Príncipe (FPP).

Author contributions

Conception and design: LMS, WM, ALSF, RAY, MVD
Analysis and interpretation: LMS, WM, ALSF, RAY, MVD
Data collection: LMS, ALCM, MVD, RKM, BBG
Writing the article: LMS, WM, ALSF, RAY, MVD, ALCM, RKM, BBG
Critical revision of the article: LMS, WM, ALSF, RAY, MVD
Final approval of the article*: LMS, WM, ALSF, RAY, MVD, ALCM, RKM, BBG
Statistical analysis: LMS, ALSF, MVD, ALCM, BBG
Overall responsibility: LMS

*All authors have read and approved of the final version of the article submitted to *J Vasc Bras*.



Abordagem endovascular de lesão de aorta torácica penetrante – relato de caso

Endovascular treatment of penetrating thoracic aorta injury – case report

Lucas Mansano Sarquis¹ , Wilson Michaelis², Antonio Lacerda Santos Filho², Rogerio Akira Yokoyama^{1,2}, Mariana Vieira Delazeri^{1,2}, Antonio Luiz da Costa Martins², Rodrigo Krieger Martins¹, Bruno Berardi Gazola³

Resumo

O cenário atual do trauma por causas violentas tem apresentado um elevado número de casos. Entre eles, a lesão de aorta torácica por trauma penetrante é uma condição de elevada morbimortalidade que exige diagnóstico adequado e que, nos últimos anos, tem sido corrigida com procedimentos endovasculares. Essa modalidade de tratamento se mostra mais segura, com menor número de complicações em relação ao procedimento cirúrgico aberto. Após a realização da abordagem endovascular, há necessidade de acompanhamento contínuo, que visa monitorar as condições de saúde do paciente e corrigir possíveis complicações relacionadas ao procedimento. O objetivo deste artigo é relatar um caso de traumatismo penetrante de aorta torácica, tratado por via endovascular, visto que a literatura aborda predominantemente as lesões por trauma contuso.

Palavras-chave: aorta torácica; ferimentos e lesões; procedimentos endovasculares.

Abstract

In the current scenario, traumas with violent causes are responsible for large numbers of cases. Among these, thoracic aorta injury caused by penetrating trauma is a cause of elevated morbidity and mortality, demanding adequate diagnosis, and can now often be repaired using endovascular procedures. This treatment method has proven to be safer, with a lower rate of complications than open surgical procedures. After endovascular repair, it is necessary to conduct continuous monitoring of the patient's health and correct any complications related to the procedure that may emerge. The objective of this article is to describe a case of penetrating trauma of the thoracic aorta that was treated endovascularly, since the literature predominantly covers blunt trauma injuries.

Keywords: thoracic aorta; wounds and injuries; endovascular procedures.

Como citar: Sarquis LM, Michaelis W, Santos Filho AL, et al. Abordagem endovascular de lesão de aorta torácica penetrante – relato de caso. J Vasc Bras. 2020;19:e20200132. <https://doi.org/10.1590/1677-5449.200132>

¹Hospital do Trabalhador, Curitiba, PR, Brasil.

²Hospital Universitário Evangélico Mackenzie, Curitiba, PR, Brasil.

³Faculdade Pequeno Príncipe, Curitiba, PR, Brasil.

Fonte de financiamento: Nenhuma.

Conflito de interesse: Os autores declararam não haver conflitos de interesse que precisam ser informados.

Submetido em: Maio 27, 2020. Aceito em: Setembro 04, 2020.

O estudo foi realizado no Hospital do Trabalhador e no Hospital Universitário Evangélico Mackenzie, Curitiba, PR, Brasil.

■ INTRODUÇÃO

Atualmente, as causas externas, em que se destacam a violência interpessoal e os acidentes de trânsito, representam a terceira causa de mortalidade no Brasil, especialmente entre a população jovem¹. O trauma torácico pode ser classificado em contuso ou penetrante. O penetrante geralmente é provocado por aplicação de força mecânica sobre uma área pequena na superfície torácica, que gera uma solução de continuidade da pele e da parede torácica¹. A ruptura da aorta torácica é majoritariamente causada por trauma contuso de tórax², e, entre as principais lesões penetrantes, o acometimento da aorta é predominantemente causado por armas de fogo ou por armas brancas, sendo de fundamental importância a qualidade da assistência pré-hospitalar e das ferramentas diagnósticas³. As lesões penetrantes de aorta torácica são raras; além disso, na sua maioria, necessitam de abordagem cirúrgica, com pouco espaço para abordagem endovascular, visto que o paciente geralmente é admitido instável hemodinamicamente⁴. Oitenta por cento dos pacientes com lesão da aorta torácica morrem no local do trauma e sete em cada 10 são do sexo masculino⁵.

Após a estabilização do paciente, o diagnóstico das lesões de aorta passa a ser um desafio para o cirurgião, visto que, além de identificá-las, é preciso tratar outros traumatismos associados. Entre as lesões concomitantes nos casos de trauma de alta energia, destacam-se o traumatismo cranioencefálico (TCE) grave, contusão pulmonar e/ou cardíaca, hemorragia abdominal, fraturas de ossos longos e/ou pélvica e ruptura diafragmática⁶. Os sintomas de ruptura aórtica podem se apresentar como dor no peito, no dorso ou no abdome, além de sinais externos de lesão torácica⁷.

A suspeição do diagnóstico se inicia com exame físico e radiografia de tórax; porém, ambos têm baixa sensibilidade. A angiotomografia computadorizada (angio-TC) é considerado o padrão-ouro, visto que, além de fazer o diagnóstico, pode servir para planejar o tratamento endovascular. Entretanto, para a realização do exame é necessário que o paciente esteja estável do ponto de vista hemodinâmico⁴. A classificação das lesões de aorta se divide em: grau 1 - lesão da íntima; grau 2 - pseudoaneurisma; grau 3 - hematoma intramural; e grau 4 - ruptura total⁸.

O primeiro relato de reparo endovascular de aorta torácica (TEVAR) foi registrado nos anos 90, sendo atualmente a técnica mais utilizada, que consiste na colocação de endoprótese a partir de uma localização periférica. Com a evolução do material para perfis mais baixos e com melhor navegação, além da experiência das equipes, houve uma redução de complicações relacionadas ao procedimento⁹. Essa modalidade de tratamento vem se destacando, especialmente para

lesões aórticas traumáticas e dissecação aórtica torácica aguda complicada, devido à redução da morbidade e mortalidade em comparação ao reparo operatório tradicional¹⁰.

Dado que a maioria dos estudos aborda as lesões de aorta por traumas contusos, este trabalho tem o objetivo de ressaltar a importância do reparo endovascular aos pacientes vítimas de trauma torácico penetrante com lesão da aorta.

■ DESCRIÇÃO DO CASO

Trata-se de um paciente masculino, 36 anos, admitido em um serviço de referência em trauma no dia 24 de dezembro de 2019 às 21:00 h, vítima de múltiplos ferimentos por arma branca. Ele apresentou instabilidade hemodinâmica durante o transporte, mas, após ressuscitação volêmica, chegou estável ao hospital. Durante o exame físico em sala de emergência, se destacaram murmúrio vesicular reduzido em hemitórax esquerdo, frequência cardíaca (FC) 125 bpm, pressão arterial média (PAM) 70-75 mmHg e múltiplos ferimentos em região anterior e posterior de tórax (Figura 1).

Após uso de concentrado de hemácias e antibioticoprofilaxia, optou-se pela realização de angio-TC de tórax (Figuras 2 e 3), a qual identificou extravasamento de contraste em aorta torácica e hemotórax a esquerda. Foi optado por drenagem pleural a esquerda e arteriografia de aorta torácica.

Optou-se por proceder ao tratamento definitivo via endovascular. Durante o procedimento, a aortografia (Figura 4) confirmou a lesão em aorta torácica (nível de T8) com tamponamento adjacente, sendo implantada endoprótese torácica reta 20/20/82 mm em aorta torácica descendente. A abordagem endovascular foi realizada à 01:00 h do dia 25 de dezembro de 2019. O controle pós-implante (Figura 5) não evidenciou extravasamento de contraste.

O paciente permaneceu em Unidade de Terapia Intensiva no pós-operatório (PO) imediato. No 1º PO, o paciente apresentou infarto agudo do miocárdio (IAM) de parede lateral, porém, sem comprometimento da função cardíaca ou hemodinâmica. No 3º PO, recebeu alta para a enfermaria. O paciente encontrou-se com uma coleção em hemitórax esquerdo por hemotórax retido, sendo submetido a videotoroscopia e decorticação pela equipe de cirurgia torácica, evoluindo sem demais intercorrências, e recebeu alta hospitalar no 10º dia de internação. O paciente atualmente se encontra em seguimento ambulatorial com as especialidades e assintomático. Já realizou angio-TC de tórax de controle com 30 dias pós-trauma, sem evidências de complicações relacionadas ao procedimento cirúrgico.




Figura 1. Ilustração dos locais com ferimento de arma branca.




Figura 2. Corte coronal de angio-TC identificando extravasamento de contraste em aorta torácica.




Figura 3. Corte axial de angio-TC identificando extravasamento de contraste em aorta torácica.




Figura 4. Aortografia com identificação de lesão em aorta torácica.



Figura 5. Aortografia de controle após colocação de endoprótese.

■ DISCUSSÃO

As lesões penetrantes da aorta torácica são potencialmente fatais e necessitam de assistência médica devidamente coordenada, a fim de elencar a ordem de prioridade dos cuidados a serem prestados. Uma das medidas a serem realizadas no atendimento inicial é o controle da pressão arterial, buscando uma PAM entre 60-70 mmHg e sistólica abaixo de 130 mmHg, sendo um dos fatores fundamentais para permitir que haja tempo hábil até que o reparo, convencional ou endovascular, seja realizado¹¹. Após a estabilização inicial do paciente, como neste caso, com a transfusão de hemoderivados, a angio-TC é essencial para identificação da lesão de aorta e avaliação da anatomia para uma possível abordagem endovascular.

O tratamento da lesão de aorta pode ser postergado, geralmente entre 1-2 semanas, podendo chegar até 45 dias, conforme avaliação conjunta com as demais especialidades. Nos casos de pacientes politraumatizados, sempre priorizar o tratamento das lesões que colocam o paciente em risco de vida¹². Entretanto, a abordagem precoce (em menos de 24 h da admissão hospitalar) deve ser estimulada para pacientes que apresentem aumento do pseudoaneurisma, incapacidade de manter pressão arterial (PA) alvo ou extravasamento de contraste na angio-TC¹².

Visto que o paciente do presente relato não apresentava demais lesões que demandassem tratamento específico, partiu-se para o tratamento da lesão de aorta. As lesões associadas, o grau da lesão de aorta, a estabilidade hemodinâmica e as comorbidades associadas são fatores determinantes para definir o momento e o tipo de reparo escolhidos⁶. Apesar de a lesão ser classificada como grau 4 – ruptura total da aorta associada com hemotórax, a anatomia favorável para tratamento endovascular e a estabilidade hemodinâmica do paciente possibilitaram a TEVAR. Esse reparo da aorta gera menor perda sanguínea, dispensa o pinçamento cruzado da aorta e tem recuperação mais rápida quando comparado ao procedimento aberto¹³.

A colocação da endoprótese é feita através de guias, com acesso pela artéria femoral comum, até que chegue à aorta torácica, onde é posicionada. Entre as características essenciais para o sucesso do procedimento, está a escolha de uma prótese com tamanho e configuração adequados, bem como a adequação anatômica vascular do próprio paciente. Tratando-se de pacientes saudáveis e sem doenças vasculares prévias, como ocorre na maioria das vezes, é possível diminuir o colo de aorta sadia para selar a endoprótese entre 0,5-0,7 cm, enquanto, para pacientes com doença vascular prévia, o ideal seria manter um

colo entre 1,5-2,5 cm¹¹. A escolha do tipo de material está relacionada principalmente com a anatomia do paciente, as angulações e a disponibilidade para uso no momento do trauma⁴.

Após o implante, há a necessidade de acompanhamento ao longo da vida por meio de exames de imagem. As diretrizes sugerem periodicidade de 30 dias, 6 e 12 meses após o reparo e, em seguida, anualmente, caso não haja complicações¹³. O procedimento aberto, que necessita de toracotomia ou esternotomia, está associado a uma taxa de 28% de mortalidade e de cerca de 16% de paraplegia³, enquanto as taxas de mortalidade geral em 30 dias, relacionadas ao TEVAR, giram em torno de 8%¹⁴. Entre as complicações comumente encontradas após abordagem endovascular, estão vazamentos (*endoleaks*), migração, estenose e infecção do enxerto, além de complicações sistêmicas como isquemias, eventos cerebrovasculares e reações inflamatórias contra o material implantado¹³. Os *endoleaks* representam uma das indicações mais comuns de reintervenções nesse tipo de reparo, com uma incidência em torno de 12,1%¹⁰. A presença de IAM pós-tratamento endovascular de doenças aórticas é uma causa rara de complicações, mas que pode estar presente¹⁰. Sobre o IAM apresentado no caso em questão, não podemos concluir se foi um evento que ocorreu em decorrência do procedimento endovascular ou ainda decorrente do trauma penetrante sem identificação nos exames de admissão.

Conclui-se, portanto, que o tratamento endovascular da aorta torácica para lesões penetrantes é uma alternativa viável. Estudos prospectivos e multicêntricos são necessários para avaliar os resultados em longo prazo da colocação de endoprótese aórtica em casos de pacientes vítimas de ferimentos penetrantes.

■ REFERÊNCIAS

1. Melo ASA, Moreira LBM, Pessoa FMC, et al. Aspectos tomográficos do trauma torácico aberto: Lesões por projéteis de arma de fogo e armas brancas. *Radiol Bras.* 2017;50(6):372-7. PMID:29307927.
2. Szostek M, Jakuczun W, Oseka M, et al. Endovascular treatment of traumatic rupture of thoracic aorta: long term results. *Int Angiol.* 2016;35(2):129-39. PMID:26221975.
3. Brand S, Breitenbach I, Bolzen P, Petri M, Krettek C, Teebken O. Open repair versus thoracic endovascular aortic repair in multiple-injured patients: observations from a level-1 trauma center. *Arch Trauma Res.* 2015;4(4):e27183. <http://dx.doi.org/10.5812/atr.27183>. PMID:26848470.
4. Chou EL, Twerdahl EH, Eagleton MJ. Endovascular management of penetrating and non-penetrating aortic injury. *Vasa.* 2019;48(1):23-33. <http://dx.doi.org/10.1024/0301-1526/a000740>. PMID:30227809.
5. Vargas TB, Bergamo R, Ferreira R, Melo Q, Almeida L, Starling SV. Lesão da aorta torácica por trauma fechado. *Rev Médica Minas Gerais.* 2016;26(Supl 4):20-3.
6. Cavalcante EF Jr, Avezum VAPAF, Silva PJLS, Vasconcellos VRB, Falavigna GG, Espada PC. Lesão aórtica e pélvica em vítima de politrauma. *Relatos Casos Cir.* 2017;3.

7. Kumar R, Raja J, Munirathinam GK, Mishra AK, Singh RS, Thingnam SKS. A case of traumatic thoracic aorta rupture: a life threatening emergency. *J Cardiovasc Thorac Res.* 2019;11(3):248-50. <http://dx.doi.org/10.15171/jcvtr.2019.41>. PMID:31579467.
8. Lee WA, Matsumura JS, Mitchell RS, et al. Endovascular repair of traumatic thoracic aortic injury: clinical practice guidelines of the Society for Vascular Surgery. *J Vasc Surg.* 2011;53(1):187-92. <http://dx.doi.org/10.1016/j.jvs.2010.08.027>. PMID:20974523.
9. Brenner M, Teeter W, Hadud M, et al. Long-term outcomes of thoracic endovascular aortic repair: a single institution's 11-year experience. *J Trauma Acute Care Surg.* 2017;82(4):687-93. <http://dx.doi.org/10.1097/TA.0000000000001365>. PMID:28129260.
10. Quintas A, Bastos Gonçalves F, Rodrigues H, et al. Tratamento endovascular de patologia da aorta torácica: experiência institucional. *Angiol Cir Vasc.* 2016;12(1):3-11. <http://dx.doi.org/10.1016/j.ancv.2015.12.002>.
11. Kolbeck KJ, Kaufman JA. Endovascular stent grafts in urgent blunt and penetrating thoracic aortic trauma. *Semin Intervent Radiol.* 2011;28(1):98-106. <http://dx.doi.org/10.1055/s-0031-1273944>. PMID:22379280.
12. Botta L, Russo V, Savini C, et al. Endovascular treatment for acute traumatic transection of descending aorta: focus on operative timing and left subclavian artery management. *J Thorac Cardiovasc Surg.* 2008;136(6):1558-63. <http://dx.doi.org/10.1016/j.jtcvs.2008.07.040>. PMID:19114206.
13. Daye D, Walker TG. Complications of endovascular aneurysm repair of the thoracic and abdominal aorta: Evaluation and management. *Cardiovasc Diagn Ther.* 2018;8(Supl 1):S138-56. <http://dx.doi.org/10.21037/cdt.2017.09.17>. PMID:29850426.
14. Newman J, Mattia A, Manetta F. Indications for Thoracic Endovascular Aortic Repair (TEVAR): A Brief Review. *Int J Angiol.* 2018;27(4):177-84. <http://dx.doi.org/10.1055/s-0038-1666972>. PMID:30410287.

Correspondência

Lucas Mansano Sarquis
Hospital do Trabalhador
Rua Augusto Stelfeld, 1908 - Bigorrrilho
CEP 80730-150 - Curitiba (PR), Brasil
Tel.: (41) 3240-5227
E-mail: lucas_sarquis@hotmail.com

Informações sobre os autores

LMS - Residente em Cirurgia Vascular, Hospital do Trabalhador (HT).
WM – Chefe e Preceptor do Serviço de Cirurgia Vascular, Hospital Universitário Evangélico Mackenzie (HUEM).
ALSF – Preceptor do Serviço de Cirurgia Vascular, Hospital Universitário Evangélico Mackenzie (HUEM).
RAY – Preceptor do Serviço de Cirurgia Vascular, Hospital Universitário Evangélico Mackenzie (HUEM); Preceptor do Serviço de Cirurgia Vascular, Hospital do Trabalhador (HT).
MVD - Preceptor do Serviço de Cirurgia Vascular, Hospital Universitário Evangélico Mackenzie; Preceptor do Serviço de Cirurgia Vascular, Hospital do Trabalhador (HT).
ALCM – Residente em Cirurgia Endovascular, Serviço de Cirurgia Vascular, Hospital Universitário Evangélico Mackenzie (HUEM).
RKM – Residente em Cirurgia Geral, Hospital do Trabalhador (HT).
BBG - Acadêmico de Medicina, Faculdade Pequeno Príncipe (FPP).

Contribuições dos autores

Concepção e desenho do estudo: LMS, WM, ALSF, RAY, MVD
Análise e interpretação dos dados: LMS, WM, ALSF, RAY, MVD
Coleta de dados: LMS, ALCM, MVD, RKM, BBG
Redação do artigo: LMS, WM, ALSF, RAY, MVD, ALCM, RKM, BBG
Revisão crítica do texto: LMS, WM, ALSF, RAY, MVD
Aprovação final do artigo*: LMS, WM, ALSF, RAY, MVD, ALCM, RKM, BBG
Análise estatística: LMS, ALSF, MVD, ALCM, BBG
Responsabilidade geral pelo estudo: LMS

*Todos os autores leram e aprovaram a versão final submetida ao *J Vasc Bras*.