

# Positive Fecal Immunochemical Test Strongly Predicts Adenomas in Younger Adults With Fatty Liver and Metabolic Syndrome

Jen-Hao Yeh, MD<sup>1,2,3</sup>, Chih-Wen Lin, MD, PhD<sup>1,3,4</sup>, Wen-Lun Wang, MD, PhD<sup>3,4</sup>, Ching-Tai Lee, MD<sup>3,4</sup>, Jen-Chieh Chen, MD<sup>1,5</sup>, Chia-Chang Hsu, MD<sup>3,5</sup> and Jaw-Yuan Wang, MD, PhD<sup>6,7,8,9</sup>

**INTRODUCTION:** The incidence of early-onset colorectal cancer is increasing. This study explored the feasibility of fecal immunochemical test (FIT) and risk factors for predicting colorectal neoplasm in younger adults.

**METHODS:** This single-center study included 6,457 participants who underwent health examination from 2013 to 2016 including index colonoscopy (3,307 individuals aged 30–49 years as the younger adult group and 3,150 aged ≥50 years as the average-risk group). Primary outcomes were adenoma detection rate (ADR) and advanced ADR (AADR). Findings of younger participants were stratified by the results of FIT and clinical risk factors and were compared with those of the average-risk group.

**RESULTS:** Among participants aged 30–49 years, a positive FIT was associated with significantly higher ADR (28.5% vs 15.5,  $P < 0.001$ ) and AADR (14.5% vs 3.7%,  $P < 0.001$ ) than a negative FIT. Moreover, a positive FIT was associated with higher AADR in younger participants than in average-risk counterparts (14.5% vs 9.8%,  $P = 0.028$ ). Although no single risk factor predicted FIT positivity in younger participants, nonalcoholic fatty liver disease was independently associated with higher ADR (odds ratio = 2.60, 95% confidence interval = 1.27–5.34,  $P = 0.001$ ), and metabolic syndrome was independently predictive of higher AADR in younger participants than in average-risk participants (odds ratio = 3.46, 95% confidence interval = 1.66–7.21,  $P = 0.001$ ).

**DISCUSSION:** A positive FIT in people aged 30–49 years implies a higher risk of colorectal neoplasm, particularly among patients with nonalcoholic fatty liver disease and metabolic syndrome.

**SUPPLEMENTARY MATERIAL** accompanies this paper at <http://links.lww.com/CTG/A496>

*Clinical and Translational Gastroenterology* 2021;12:e00305. <https://doi.org/10.14309/ctg.0000000000000305>

## INTRODUCTION

Colorectal cancer (CRC) is one of the most common malignancies worldwide (1). In general, the target population of CRC screening is people aged younger than or equal to 50 years because the risk in this population is higher than that in younger people. Although effective screening policies in some countries have led to an overall stability or even decline of CRC incidence, the incidence of early-onset CRC, that is CRC at younger than 50 years of age, seems to be increasing (2,3). By 2030, the incidence of early-onset CRC is expected to double among people aged 20–34 years and increase by approximately 30% among people aged 35–49 years (4).

Few recommendations have been made regarding a screening strategy for early-onset CRC. Although some experts have advocated lowering the age of screening to 45 years (5), this is not endorsed by other experts (6). Moreover, the best screening strategy for the younger population is yet to be determined. The fecal immunochemical test (FIT) is an effective and noninvasive screening method for average-risk people (7,8), but few studies have investigated its efficacy in younger adults. Jung et al. (9) found the FIT useful to detect advanced adenoma in asymptomatic people aged 35–49 years. However, the prevalence of CRC was relatively low (0.027%). Hence, for the younger

<sup>1</sup>Division of Gastroenterology and Hepatology, Department of Internal Medicine, E-DA Dachang Hospital, Kaohsiung, Taiwan; <sup>2</sup>Department of Medical Technology, College of Medicine, I-Shou University, Kaohsiung, Taiwan; <sup>3</sup>Division of Gastroenterology and Hepatology, Department of Internal Medicine, E-DA Hospital, Kaohsiung, Taiwan; <sup>4</sup>School of Medicine, College of Medicine, I-Shou University, Kaohsiung, Taiwan; <sup>5</sup>Department of Health Examination, E-DA Hospital, Kaohsiung, Taiwan; <sup>6</sup>Division of Colorectal Surgery, Department of Surgery, Kaohsiung Medical University Hospital, Kaohsiung Medical University, Kaohsiung, Taiwan; <sup>7</sup>Department of Surgery, Faculty of Medicine, College of Medicine, Kaohsiung Medical University Hospital, Kaohsiung Medical University, Kaohsiung, Taiwan; <sup>8</sup>Graduate Institute of Clinical Medicine, College of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan; <sup>9</sup>Center for Cancer Research, Kaohsiung Medical University, Kaohsiung, Taiwan. **Correspondence:** Jaw-Yuan Wang, MD, PhD. E-mail: jawyuanwang@gmail.com.

**Received June 13, 2020; accepted December 18, 2020; published online February 3, 2021**

© 2021 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of The American College of Gastroenterology

population, a risk-based approach, if available, seems reasonable over universal screening.

Several risk factors, including age 40 years or younger, obesity, diabetes mellitus, hypertension, smoking, nonalcoholic fatty liver disease (NAFLD), and metabolic syndrome, are associated with colorectal adenoma in the younger population (10–12). However, little is known about their impact in the presence of a positive FIT. From the perspective of cost-effectiveness, younger adults who present a higher-risk counterpart than average-risk counterparts will benefit most from screening. The study investigated the efficacy of the FIT in healthy younger adults and explored the additional risk factors for colorectal neoplasm in individuals with a positive FIT.

## MATERIAL AND METHODS

### Study design

This retrospective, single-center study included participants visiting the health examination department of E-Da Hospital, Kaohsiung, Taiwan, between January 2013 and December 2016. It was approved by the Institutional Review Board of E-Da Hospital (No. EMRP40108N). Similar to the previous studies (9,13), participants were asymptomatic people who underwent health checkups at their own expense or following national labor laws requirements.

The inclusion criteria were individuals aged 20–49 years (younger adult group) who received both FIT and index colonoscopy during the checkup and participants aged younger than or equal to 50 years (average-risk group) who received index colonoscopy, regardless of whether FIT was performed. In either group, colonoscopy within 6 months after FIT was considered in the analysis. The exclusion criteria were as follows: (i) duplicated cases owing to previous checkups, (ii) a history of CRC or colorectal resection, (iii) a history of colonoscopy within 10 years, and (iv) poor colon preparation.

### FIT and colonoscopy procedures

The specimen for FIT (OC-Sensor kit; Eiken Chemical, Tokyo, Japan) was obtained and sent to the hospital within 1 week before the examination. The results are expressed in nanograms of hemoglobin per milliliter buffer (ng Hb/mL), and the cutoff value of positivity was set at 100 ng Hb/mL.

All colonoscopies were performed by well-experienced endoscopists who had performed >1,000 colonoscopy examinations, with  $\geq 300$  procedures conducted annually, using the EvisLucera CV-260 colonoscope (Olympus Medical Systems, Tokyo, Japan). Bowel preparation regimens were split-dose sodium phosphate or 2 L of polyethylene glycol solution. During the examination, the decision of biopsy, snare polypectomy, or endoscopic mucosal resection for the colorectal neoplasm was at the discretion of the endoscopist. However, the presence of adenoma, advanced adenoma, or CRC was diagnosed through histologic evaluation. In the case of difficult-to-treat polyps, subsequent endoscopic or surgical resection was performed within 6 months after index colonoscopy. Accordingly, polyps and lesions that were not removed at our institute within 6 months of index colonoscopy were not included in the analysis, regardless of endoscopic diagnosis.

### Outcome assessment

Primary outcomes were adenoma detection rate (ADR) and advanced ADR (AADR) of younger participants with positive FIT as

compared to those with negative FIT and average-risk group. ADR and AADR were calculated as the proportion of cases with detected adenomas and advanced adenomas, respectively, divided by the total number of participants in each group. Advanced adenoma refers to polyps (i) with high-grade dysplasia, carcinoma *in situ*, or intramucosal carcinoma; (ii)  $\geq 1$  cm in size; or (iii) with  $>25\%$  villous component.

Participants' baseline characteristics, including age, sex, underlying disease, laboratory, and endoscopic findings, were thoroughly recorded. Obesity was defined as body mass index  $>27$  kg/m<sup>2</sup>. NAFLD was defined as ultrasonographically detected fatty liver disease in the absence of substantial alcohol intake. Preexisting diagnosis of any metabolic diseases, such as hypertension, diabetes mellitus, dyslipidemia, and metabolic syndrome, was noted. In addition, participants with systolic blood pressure  $\geq 180$  mm Hg or diastolic blood pressure  $\geq 120$  mm Hg with sustained high-abnormal readings at 2 consecutive examination were considered to have hypertension. Diabetes mellitus was defined as fasting glucose  $\geq 126$  mg/dL or glycosylated hemoglobin  $\geq 6.5\%$ . Dyslipidemia was defined as either triglyceride  $\geq 150$  mg/dL, total cholesterol  $\geq 200$  mg/dL, high-density lipoprotein cholesterol  $<40$  mg/dL in men or  $<50$  mg/dL in women, or low-density lipoprotein cholesterol  $\geq 130$  mg/dL.

Metabolic syndrome was defined as the presence of at least 3 of the following findings: central obesity (waist circumference  $\geq 90$  cm in men or  $\geq 80$  cm in women), elevated blood pressure (systolic blood pressure  $\geq 130$  mm Hg or diastolic blood pressure  $\geq 80$  mm Hg in 2 consecutive readings), fasting glucose impairment ( $>100$  mg/dL), elevated triglyceride ( $\geq 150$  mg/dL), and reduced high-density lipoprotein cholesterol ( $<40$  mg/dL in men or  $<50$  mg/dL in women). A family history of CRC was not confined to first-degree relatives.

### Statistical analysis

In this study, continuous variables are presented as mean  $\pm$  SD and were compared using Student *t* tests. Categorical variables are presented as numbers (percentages) and were compared using the  $\chi^2$  test.  $P < 0.05$  was considered statistically significant. For the comparison of primary outcomes, odds ratio (OR) with 95% confidence interval (CI) was calculated. All statistical analyses were conducted using SPSS version 26.0 (IBM Corp, Armonk, NY).

To evaluate the impact of FIT results for ADR and AADR, younger adult participants were stratified by age (20–29, 30–39, and 40–49 years) and the FIT results (positive or negative). Because the primary outcomes were not statistically significant for participants aged 20–29 years in the preliminary analysis, further analysis included only participants aged 30–49 years versus the average-risk counterparts.

Clinical risk factors, including age, sex, obesity, smoking habits, hypertension, diabetes mellitus, dyslipidemia, metabolic syndrome, and family history of CRC, were analyzed using the  $\chi^2$  test to identify factors associated with increased ADR and AADR. Factors that were statistically significant were included in the multivariable binary logistic regression model with backward selection. The diagnostic power of FIT was evaluated using the receiver operating characteristic curve and was displayed as the area under the curve (AUC), and sensitivity analysis was performed to define to best cutoff value, with the highest accuracy (minimal false-negative and false-positive results) determined according to the AUC.

## RESULTS

### Study participants and baseline characteristics

Overall, 59,413 records were retrieved from the database. The flowchart of participant selection is presented in Figure 1. After applying the exclusion criteria, 6,906 participants (men, 57.2%) were included in the final analysis. Among these participants, 3,756 were younger adults (of whom 3,307 were 30–49 year old), and 3,150 were included in the average-risk group.

The baseline characteristics of each group are summarized in Table 1. Younger adults had a mean age of 40.2 years, and the average-risk group had a mean age of 58.0 years. Slight male predominance was found in the younger adult group (59.5% vs 55.7%,  $P < 0.001$ ). Compared with younger adults, a significantly higher proportion of participants in the average-risk group had adenoma (16.2% vs 28.5%,  $P = 0.001$ ), advanced adenoma (4.3% vs 9.6%,  $P < 0.001$ ), and CRC (0.2% vs 0.8%,  $P < 0.001$ ). Notably, all younger adults with CRC were aged 30–49 years.

In addition, compared with those in the younger adult group, participants in the average-risk group were more likely to have a positive FIT (5.6% vs 8.2%,  $P < 0.001$ ) and metabolic diseases including hypertension (6.3% vs 24.7%,  $P < 0.001$ ), diabetes mellitus (4.2% vs 17.2%,  $P < 0.001$ ), dyslipidemia (64.0% vs 76.1%,  $P < 0.001$ ), metabolic syndrome (24.6% vs 37.5%,  $P < 0.001$ ), and NAFLD (40.0% vs 45.6%,  $P < 0.001$ ). However, a higher proportion of younger adults were current smokers (21.7% vs 15.6%,  $P < 0.001$ ).


### Association of ADR and AADR with FIT among different age groups

ADR was compared between younger adult participants with positive and negative FIT results and average-risk group counterparts (Figure 2a). ADR of average-risk participants was 28.5%. ADRs among participants aged 20–29, 30–39, 40–49, and 30–49 years with a positive FIT were 5.6%, 20.8%, 33.9%, and 28.5%, respectively, and with a negative FIT was 3.2% ( $P = 0.594$ ), 9.6% ( $P = 0.002$ ), 20.1% ( $P = 0.001$ ), and 15.5% ( $P < 0.001$ ), respectively.

AADR was 9.6% among average-risk participants. AADR among younger adult participants aged 20–29, 30–39, 40–49, and 30–49 years with positive FITs were 0%, 11.7%, 16.5%, and 14.5%, respectively, and with negative FITs were 0.9%, 1.9%, 5.2%, and 3.7%, respectively (Figure 2b). Similar to the comparative results of ADR, AADR was significantly associated with FIT among participants aged 30–39, 40–49, and 30–49 years (all  $P < 0.001$ ), but not among those aged 20–29 years ( $P = 0.681$ ).

### Association of ADR and AADR with any clinical factor among younger adults

We explored the association of ADR and AADR (Table 2) with the clinical risk factors according to different age groups among younger adults. For participants aged 20–29 years, no factor significantly affected ADR and AADR. However, for participants aged 30–49 years, ADR was significantly increased in the presence of male sex (20.5% vs 10.0%,  $P < 0.001$ ), obesity (21.8% vs 16.2%,  $P < 0.001$ ), diabetes mellitus (25.0% vs 15.9%,  $P = 0.004$ ), hypertension (21.5% vs 15.9%,  $P = 0.032$ ), and current smoking (28.0% vs 13.0%,  $P < 0.001$ ). AADR was significantly increased in the presence of male sex (5.7% vs 2.2%,  $P < 0.001$ ), obesity (6.3% vs 3.9%,  $P = 0.007$ ), metabolic syndrome (6.1% vs 3.7%,  $P = 0.003$ ), diabetes mellitus (9.3% vs 4.1%,  $P = 0.003$ ), NAFLD (5.4%



**Figure 1.** Flowchart of participant selection. FIT, fecal immunochemical test.

vs 3.6%,  $P = 0.01$ ), and current smoking (7.7% vs 3.4%,  $P < 0.001$ ). No single risk factor, including positive FIT, caused the ADR of younger adults to be significantly higher than that of average-risk groups. However, a positive FIT was associated with the significantly higher AADR of the younger adults than average-risk counterparts (14.5% vs 9.8%,  $P = 0.028$ ).

Furthermore, we compared the ADR (Table 3) and AADR (Table 4) of younger, FIT-positive participants with specific clinical factors to that of all average-risk participants. Univariable analysis revealed that male sex (OR = 1.56, 95% CI = 1.04–2.32,  $P = 0.028$ ), obesity (OR = 2.51, 95% CI = 1.22–5.15,  $P = 0.012$ ), metabolic syndrome (OR = 2.11, 95% CI = 1.11–4.09,  $P = 0.022$ ), and NAFLD (OR = 1.91, 95% CI = 1.12–3.05,  $P = 0.006$ ) were associated with a higher ADR in younger adults than in average-risk adults. However, multivariable analysis indicated NAFLD (OR = 2.69, 95% CI = 1.36–5.33,  $P = 0.005$ ) as the only independent risk factor for a higher ADR, and female sex (OR 0.32, 95% CI = 0.16–0.64,  $P = 0.001$ ) was associated with significantly lower ADR.

Univariable analysis revealed the following factors to be associated with higher AADR: male sex (OR = 2.30, 95% CI = 1.40–3.76,  $P = 0.001$ ), obesity (OR = 2.87, 95% CI = 1.22–6.74,  $P = 0.016$ ), metabolic syndrome (OR = 3.49, 95% CI = 1.67–7.28,  $P = 0.001$ ), dyslipidemia (OR = 1.73, 95% CI = 1.00–2.99,  $P = 0.048$ ), NAFLD (OR = 2.52, 95% CI = 1.46–4.33,  $P = 0.001$ ), and current smoking (OR = 2.43, 95% CI = 1.34–5.10,  $P = 0.027$ ). However, multivariable analysis revealed that metabolic syndrome (OR = 3.46, 95% CI = 1.66–7.21,  $P = 0.001$ ) was the only independent risk factor associated with increased AADR among younger adults.

### Diagnostic yield of FIT and its association with clinical risk factors in younger adults

In the exploration of the association between FIT positivity and clinical risk factors in the younger adults, no single factor predicted significantly higher FIT positivity (see Supplementary Table 1, Supplementary Digital Content 1, <http://links.lww.com/CTG/A496>). The presence of dyslipidemia was associated with a lower proportion of FIT positivity (4.9% vs 6.8%,  $P = 0.02$ ).

For the diagnostic ability of FIT, the sensitivities of FIT for adenoma detection among participants aged 30–39, 40–49, and  $\geq 50$  years were 10.8%, 9.5%, and 14.0%, respectively, and the specificities were 95.3%, 95.1%, and 94.1%, respectively; for advanced adenoma

**Table 1. Baseline characteristics of the included patients**

Group (years-old)	Young-age group				Average-risk ≥50	P value
	20–29	30–39	40–49	30–49		
No. of patients	449	1,450	1857	3,307	3,150	
Age (mean ± SD)	26.2 ± 2.5	35.3 ± 2.7	44.3 ± 2.8	40.2 ± 5.3	58.0 ± 6.1	<0.001 <sup>a</sup>
Male sex, n (%)	228 (50.8)	789 (54.4)	1,179 (63.5)	1968 (59.5)	1755 (55.7)	0.002 <sup>a</sup>
Body mass index (kg/m <sup>2</sup> , mean ± SD)	22.7 ± 4.6	23.5 ± 4.1	24.2 ± 3.6	23.9 ± 3.8	24.4 ± 3.4	<0.001 <sup>a</sup>
Obesity, n (%)	67 (14.9)	248 (17.1)	370 (19.9)	618 (18.7)	614 (19.5)	0.400
Smoking habits						<0.001 <sup>a</sup>
Current smoker, n (%)	75 (16.7)	284 (19.7)	430 (23.2)	714 (21.7)	488 (15.6)	<0.001 <sup>a</sup>
Ex-smoker, n (%)	9 (2.0)	47 (3.3)	103 (5.6)	150 (4.6)	198 (6.3)	0.015 <sup>a</sup>
Hypertension, n (%)	1 (0.2)	39 (2.7)	170 (9.2)	209 (6.3)	778 (24.7)	<0.001 <sup>a</sup>
Diabetes mellitus, n (%)	4 (0.9)	20 (1.4)	120 (6.5)	140 (4.2)	541 (17.2)	<0.001 <sup>a</sup>
Fasting sugar (mg/dL, mean ± SD)	90.8 ± 9.4	95.1 ± 14.4	101.5 ± 22.2	98.7 ± 19.4	108.7 ± 28.7	<0.001 <sup>a</sup>
HbA1C (%), mean ± SD)	5.2 ± 0.4	5.3 ± 0.5	5.5 ± 0.7	5.4 ± 0.7	5.9 ± 1.0	<0.001 <sup>a</sup>
Dyslipidemia, n (%)	178 (39.6)	818 (56.6)	1,289 (69.7)	2,107 (64.0)	2,333 (76.1)	<0.001 <sup>a</sup>
Metabolic syndrome (%)	42 (9.4)	271 (18.7)	544 (29.4)	815 (24.6)	1,181 (37.5)	<0.001 <sup>a</sup>
AST (IU/ml, mean ± SD)	23.4 ± 12.2	26.3 ± 15.7	28.1 ± 15.6	27.3 ± 15.7	30.4 ± 27.9	<0.001 <sup>a</sup>
ALT (IU/ml, mean ± SD)	23.4 ± 20.9	31.4 ± 29.7	32.9 ± 25.6	32.3 ± 27.5	31.6 ± 39.9	0.447
NAFLD	77 (17.1)	466 (32.1)	857 (46.2)	1,323 (40.0)	1,435 (45.6)	<0.001 <sup>a</sup>
Positive FIT	18 (4.0%)	77 (5.3%)	109 (5.9%)	186 (5.6)	207 (8.2%) <sup>b</sup>	<0.001 <sup>a</sup>
CRC family history	36 (8.0)	117 (8.1)	118 (6.4)	235 (7.1)	209 (6.6)	0.454
Adenoma (%)	15 (3.3)	148 (10.2)	389 (20.9)	537 (16.2)	897 (28.5)	<0.001 <sup>a</sup>
Advanced adenoma (%)	4 (0.9)	35 (2.4)	108 (5.8)	143 (4.3)	302 (9.6)	<0.001 <sup>a</sup>
Serrated adenoma (%)	4 (0.9)	32 (2.2)	54 (2.9)	86 (2.6)	83 (2.6)	0.929
CRC (%)	0	3 (0.2)	5 (0.3)	8 (0.1)	24 (0.8)	0.003 <sup>a</sup>

Data were compared between the younger adults and the average-risk group.  
 AST, aspartate aminotransferase; ALT, alanine aminotransferase; CRC, colorectal cancer; FIT, fecal immunochemical study; NAFLD, nonalcoholic fatty liver disease.  
<sup>a</sup>*P* < 0.05.  
<sup>b</sup>Only 2,518 participants in the average-risk group received FIT.

detection, the sensitivities were 11.7%, 16.5%, and 25.9%, respectively, and the specificities were 95.2%, 94.8%, and 93.7%, respectively. For participants aged 30–49 years, the sensitivity, specificity, and positive and negative predictive values were 9.9%, 95.2%, 28.5%, and 84.5%, respectively, for adenoma and 18.9%, 95.0%, 14.5%, and 96.3%, respectively, for advanced adenoma.

Sensitivity analysis was performed by comparing the AUC for advanced adenoma for participants aged 30–49 years with different cutoff values of FIT. These cutoff values selected were 50, 100, and 200 ng Hb/mL. When the cutoff was set at 100 ng Hb/mL, the AUC for advanced adenoma of FIT was 0.569. However, the AUC was not significantly different when the cutoff value was set at either 50 ng Hb/mL (0.570, *P* = 0.957) or 150 ng Hb/mL (0.557, *P* = 0.204). In addition, the presence of metabolic syndrome did not improve the AUC (0.573 vs 0.517, *P* = 0.330).


## DISCUSSION

Nearly 10%–20% of CRC cases are diagnosed before the age of 50 years (14,15), and early-onset CRC has become a leading cause of cancer death in that population (16). Although the effectiveness of screening tests is largely unknown in the younger population, it

is reasonable to screen at-risk younger adults. The incidence of CRC is low in this population, and studies have suggested that younger adults with advanced adenoma have a similar risk of metachronous advanced neoplasia as the average-risk counterparts (17,18); therefore, the current study used both ADR and AADR as the primary outcomes of evaluation.

Our findings revealed that a positive FIT in people aged 30–49 was predictive of significantly higher ADR and AADR with the index colonoscopy and that the risk of advanced adenoma was even higher in people aged 40–49 years. Moreover, we found that NAFLD and metabolic syndrome were independent risk factors for higher ADR and AADR, respectively. The first part of the findings is similar to the results of Jung et al. (9) although they used the age of 35–49 years as the cutoff point. However, they did not explore parameters other than male sex and smoking, such as metabolic diseases, which were considered in the present study.

Similar to CRC in average-risk people, diabetes mellitus, obesity, and metabolic syndrome are risk factors for early-onset CRC, and insulin resistance is probably the mutual pathophysiology. NAFLD, however, is a precursor of metabolic syndrome and diabetes mellitus (19), and there has been some evidence



**Figure 2.** (a) Comparison of ADR based on the results of FIT among different age groups. \*\**P* < 0.01. (b) Comparison of AADR based on the results of the FIT among different age groups. \*\**P* < 0.01. ADR, adenoma detection rate; AADR, advanced ADR; FIT, fecal immunochemical test.

linking its presence to colorectal neoplasms (20–22). Although diabetes mellitus and family history of CRC are established risk factors (5,6), we did not find the association between them and the risk of adenoma in younger FIT-positive adults. This may be explained by the low prevalence of diabetes mellitus in this

subgroup (8/182, 4.3%). Similarly, family history of CRC was also not significant in this cohort likely because the family history in the database was not confined to first-degree relatives.

In this study, our findings suggest that positive FIT is correlated with higher colorectal neoplasms especially for younger

**Table 2.** Adenoma and advanced adenoma detection rate in the presence and absence of the clinical risk factors

Risk factor	Adenoma detection rate in each age group, %									
	20–29		30–39		40–49		30–49		50+	
	+	–	+	–	+	–	+	–	+	–
Male	4.4	2.3	12.0 <sup>a</sup>	8.0 <sup>a</sup>	26.1 <sup>a</sup>	11.9 <sup>a</sup>	20.5 <sup>a</sup>	10.0 <sup>a</sup>	35.5 <sup>a</sup>	19.6 <sup>a</sup>
FIT	5.6	3.2	20.8 <sup>a</sup>	9.6 <sup>a</sup>	33.9 <sup>a</sup>	20.1 <sup>a</sup>	28.5 <sup>a</sup>	15.5 <sup>a</sup>	48.8 <sup>a</sup>	26.8 <sup>a</sup>
Obesity	6.0	2.9	8.5	10.6	30.8 <sup>a</sup>	18.5 <sup>a</sup>	21.8 <sup>a</sup>	16.2 <sup>a</sup>	33.6 <sup>a</sup>	27.3 <sup>a</sup>
MS	4.8	3.2	12.2	9.8	27.2 <sup>a</sup>	18.4 <sup>a</sup>	22.2	14.3	33.3 <sup>a</sup>	25.6 <sup>a</sup>
Dyslipidemia	4.5	2.6	11.1	8.6	22.5 <sup>a</sup>	17.1 <sup>a</sup>	18.1	12.6	28.6	27.9
Diabetes mellitus	0	3.4	20	10.1	25.8	20.6	25.0 <sup>a</sup>	15.9 <sup>a</sup>	30.2	28.0
Hypertension	0	3.3	7.7	10.3	24.7	20.6	21.5 <sup>a</sup>	15.9 <sup>a</sup>	33.3 <sup>a</sup>	26.9 <sup>a</sup>
NAFLD	6.5	2.7	10.9	9.9	25.0 <sup>a</sup>	17.4 <sup>a</sup>	20.0	13.7	31.7 <sup>a</sup>	25.7 <sup>a</sup>
Family history	0	3.6	6.8	10.5	27.1	20.5	17.0	16.2	31.1	28.3
Current smoker	4.0	3.2	19.7 <sup>a</sup>	7.8 <sup>a</sup>	33.5 <sup>a</sup>	17.2 <sup>a</sup>	28.0 <sup>a</sup>	13.0 <sup>a</sup>	39.8 <sup>a</sup>	26.4 <sup>a</sup>

Risk factor	Advanced adenoma detection rate in each age group, %									
	20–29		30–39		40–49		30–49		50+	
	+	–	+	–	+	–	+	–	+	–
Male	1.8*	0*	2.9	1.8	9.6 <sup>a</sup>	2.7 <sup>a</sup>	5.7 <sup>a</sup>	2.2 <sup>a</sup>	12.5 <sup>a</sup>	5.9 <sup>a</sup>
FIT	0	0.9	11.7 <sup>a</sup>	1.9 <sup>a</sup>	16.5 <sup>a</sup>	5.2 <sup>a</sup>	14.5 <sup>a</sup>	3.7 <sup>a</sup>	30.9 <sup>a</sup>	7.9 <sup>a</sup>
Obesity	1.5	0.8	2.0	2.5	9.2 <sup>a</sup>	5.0 <sup>a</sup>	6.3 <sup>a</sup>	3.9 <sup>a</sup>	12.5 <sup>a</sup>	8.9 <sup>a</sup>
MS	0	1.0	2.6	2.4	7.9 <sup>a</sup>	5.0 <sup>a</sup>	6.1 <sup>a</sup>	3.7 <sup>a</sup>	12.4 <sup>a</sup>	7.9 <sup>a</sup>
Dyslipidemia	1.1	0.7	2.2	2.6	6.3	4.6	4.7	3.5	9.5	9.8
Diabetes mellitus	0	0.9	5.0	2.4	10 <sup>a</sup>	5.5 <sup>a</sup>	9.3 <sup>a</sup>	4.1 <sup>a</sup>	10.9	9.3
Hypertension	0	0.9	0	2.5	7.6	5.6	6.2	4.2	12.9 <sup>a</sup>	8.5 <sup>a</sup>
NAFLD	2.6	0.5	3.2	2.0	6.7	5.1	5.4 <sup>a</sup>	3.6 <sup>a</sup>	11.4 <sup>a</sup>	8.1 <sup>a</sup>
Family history	0	0.9	2.6	2.4	9.3	5.6	6.0	4.2	7.7	9.7
Current smoker	0	1.1	3.9	2.2	10.2	4.5	7.7	3.4	16.4 <sup>a</sup>	8.4 <sup>a</sup>

FIT, fecal immunochemical study; MS, metabolic syndrome; NAFLD, nonalcoholic fatty liver disease.

<sup>a</sup>Statistically significant (*P* value < 0.05) for the difference of rate between the specific positive and negative clinical risk factors in the same-age group.



**Table 3.** Adenoma detection rate in younger participants (30–49 years of age) with a positive FIT with any specific risk factor versus average-risk counterparts

Risk factor	ADR, %	OR (95% CI): 30–49 year-old FIT (+) vs average risk			
		Univariable analysis	P value	Multivariable analysis <sup>b</sup>	P value
Male	38.3	1.56 (1.04–2.32)	0.028 <sup>a</sup>	0.91 (0.52–1.61)	0.768
Obesity	50	2.51 (1.22–5.15)	0.012 <sup>a</sup>	1.84 (0.76–4.44)	0.175
Metabolic syndrome	45.9	2.13 (1.11–4.09)	0.022 <sup>a</sup>	1.30 (0.51–3.31)	0.577
Dyslipidemia	31.1	1.13 (0.74–1.73)	0.567		
Diabetes mellitus	50	2.51 (0.62–10.06)	0.193		
Hypertension	50	2.51 (0.87–7.18)	0.086		
NAFLD	43.2	1.91 (1.20–3.05)	0.006 <sup>a</sup>	2.69 (1.36–5.33)	0.005 <sup>a</sup>
Family history	30.8	1.11 (0.34–3.63)	0.855		
Current smoker	35.9	1.40 (0.72–2.71)	0.310		
Average-risk	28.5	1 (reference)		1 (reference)	

FIT, fecal immunochemical study; NAFLD, nonalcoholic fatty liver disease.  
<sup>a</sup>P < 0.05.  
<sup>b</sup>Binary logistic regression.

adults with NAFLD and metabolic syndrome. As demonstrated in the results, for most younger adults who had a negative FIT, the risk of colorectal adenoma was generally low. Although individuals with multiple clinical risk factors may remain at a high risk of developing adenoma (11), most of them do not require early colonoscopy. Interestingly, a cost-effective model by Wong et al. (23) also suggested FIT rather than colonoscopy as the screening tool for younger adults with NAFLD. In addition to the lower cost, FIT is also more convenient and less invasive than colonoscopy. Despite the several advantages of FIT, more studies are necessary to evaluate the utility of FIT as the screening tool for the younger patients, and the comparison versus colonoscopy in the cost and efficacy also has to be performed to determine the optimal screening strategy.

The study has some limitations. First, this was a retrospective study. Second, we did not consider the effectiveness of previous FIT, and the surveillance interval for a negative FIT remains unclear. Third, we did not include the location and numbers of lesions in the analysis. Although a fecal occult blood test is traditionally believed to be more sensitive for distal colon lesions traditionally, one study found FIT to be equally sensitive for proximal and distal colon neoplasia (24). Fourth, we did not analyze sessile serrated adenomas owing to the generally low prevalence in this cohort. Finally, the clinical risk factors evaluated in this study might have confounding effects on each other even in multivariable analysis. For example, patients with obesity are more likely to have fatty liver and metabolic syndrome, and

**Table 4.** Advanced adenoma detection rate in younger participants (30–49 years of age) with positive FIT with any specific risk factor versus average-risk counterparts

Risk factor	AADR, %	OR (95% CI): 30–49 year-old FIT (+) vs average risk			
		Univariable analysis	P value	Multivariable analysis <sup>b</sup>	P value
Male	13.6	2.30 (1.40–3.76)	0.001 <sup>a</sup>	1.74 (0.95–3.17)	0.068
Obesity	23.3	2.87 (1.22–6.74)	0.016 <sup>a</sup>	0.85 (0.25–2.85)	0.802
Metabolic syndrome	27.0	3.49 (1.67–7.28)	0.001 <sup>a</sup>	3.46 (1.66–7.21)	0.001 <sup>a</sup>
Dyslipidemia	15.5	1.73 (1.00–2.99)	0.048 <sup>a</sup>	0.66 (0.26–1.69)	0.389
Diabetes mellitus	12.5	1.34 (0.16–10.98)	0.781		
Hypertension	21.4	2.57 (0.71–9.27)	0.149		
NAFLD	21.6	2.60 (1.47–4.58)	0.001 <sup>a</sup>	1.69 (0.82–4.23)	0.255
Family history	23.1	2.82 (0.77–10.33)	0.116		
Current smoker	20.5	2.43 (1.34–5.10)	0.027 <sup>a</sup>	0.98 (0.35–2.74)	0.972
Average-risk	9.6	1 (reference)		1 (reference)	

FIT, fecal immunochemical study; NAFLD, nonalcoholic fatty liver disease.  
<sup>a</sup>P < 0.05.  
<sup>b</sup>Binary logistic regression.

metabolic syndrome is associated with diabetes mellitus occurrence owing to underlying insulin resistance. Further research is necessary to validate our findings and develop a prediction model to more accurately identify younger adults who would be candidates for FIT.

In conclusion, a positive FIT in individuals aged 30–49 years may imply a higher risk of colorectal neoplasm, especially for patients with NAFLD and metabolic syndrome. Future studies are warranted to determine the utility of FIT and optimal time interval of surveillance with an initially negative test.

## CONFLICTS OF INTEREST

**Guarantor of the article:** Jaw-Yuan Wang, MD, PhD.

**Specific author contributions:** Chia-Chang Hsu, MD and Jaw-Yuan Wang, MD, PhD, contributed equally to this work. J.H.Y. performed data retrieval, statistical analysis, and wrote the study. C.W.L. and W.L.W. reviewed the data and made suggestions for manuscript writing. C.T.L., J.C.C., and C.C.H. performed colonoscopy examination for the participants in the study. C.C.H. is the chief of health examination department in E-Da Hospital who supervised the study, and he also contributed to the study design and analysis. J.Y.W. designed the study and revised the study critically. All authors approved the final version of the article, including the authorship list.

**Financial support:** This study was supported by the EDHP109009 project of E-Da Hospital granted to Dr. Jen-Hao Yeh. This work was also supported by grants through funding from the Ministry of Science and Technology (MOST 109-2314-B-037-035, MOST 109-2314-B-037-040, MOST 109-2314-B-037-046-MY3) and the Ministry of Health and Welfare (MOHW107-TDU-B-212-123006, MOHW107-TDU-B-212-114026B, MOHW108-TDU-B-212-133006, MOHW109-TDU-B-212-134026, MOHW109-TDU-B-212-114006) and funded by the health and welfare surcharge of on tobacco products, and the Kaohsiung Medical University Hospital (KMUH109-9R32, KMUH109-9R33, KMUH109-9R34, KMUH109-9M30, KMUH109-9M31, KMUH109-9M32, KMUH109-9M33, KMUH-DK109003, KMUH-DK109005~3, KMUHS10903, KMUHSA10903) and KMU Center for Cancer Research (KMU-TC108A04-2) as well as and a KMU Center for Liquid Biopsy and Cohort Research Center Grant (KMU-TC109B05), Kaohsiung Medical University. In addition, this study was supported by the Grant of Taiwan Precision Medicine Initiative, Academia Sinica, Taiwan, R.O.C.

**Potential competing interests:** None to report.

## Study Highlights

### WHAT IS KNOWN

- ✓ Early onset CRC has become more frequent.
- ✓ FIT predicts colorectal adenoma in younger adults.
- ✓ The performance of FIT in different risk groups are largely unknown.

### WHAT IS NEW HERE

- ✓ FIT is the strongest factor to predict advanced adenomas in younger adults.
- ✓ Positive FIT is associated with higher risk in adults aged 30 and more.
- ✓ Metabolic syndrome and fatty liver with positive FIT are among highest risk.

## ACKNOWLEDGMENTS

We appreciate the assistance of Ms Pao-Shu Hsu in data collection and Ms Tzu-Shan Chen for advice on statistical analysis.

## REFERENCES

1. Bray F, Ferlay J, Soerjomataram I, et al. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018;68:394–424.
2. Araghi M, Soerjomataram I, Bardot A, et al. Changes in colorectal cancer incidence in seven high-income countries: A population-based study. *Lancet Gastroenterol Hepatol* 2019;4:511–8.
3. Siegel RL, Fedewa SA, Anderson WF, et al. Colorectal cancer incidence patterns in the United States, 1974–2013. *J Natl Cancer Inst* 2017;109:djw322.
4. Bailey CE, Hu CY, You YN, et al. Increasing disparities in the age-related incidences of colon and rectal cancers in the United States, 1975–2010. *JAMA Surg* 2015;150:17–22.
5. Wolf AMD, Fontham ETH, Church TR, et al. Colorectal cancer screening for average-risk adults: 2018 guideline update from the American Cancer Society. *CA Cancer J Clin* 2018;68:250–81.
6. Force UPST. Screening for colorectal cancer: US Preventive Services Task Force Recommendation Statement. *JAMA* 2016;315:2564–75.
7. Katsoula A, Paschos P, Haidich AB, et al. Diagnostic accuracy of fecal immunochemical test in patients at increased risk for colorectal cancer: A meta-analysis. *JAMA Intern Med* 2017;177:1110–8.
8. Lee YC, Fann JCY, Chiang TH, et al. Time to colonoscopy and risk of colorectal cancer in patients with positive results from fecal immunochemical tests. *Clin Gastroenterol Hepatol* 2019;17:1332–40.e3.
9. Jung YS, Park CH, Kim NH, et al. Colorectal cancer screening with the fecal immunochemical test in persons aged 30 to 49 years: Focusing on the age for commencing screening. *Gastrointest Endosc* 2017;86:892–9.
10. Chen KC, Chung CS, Hsu WF, et al. Identification of risk factors for neoplastic colonic polyps in young adults with bloody stool in comparison with those without symptom. *J Gastroenterol Hepatol* 2018;33:1335–40.
11. Jung YS, Park CH, Kim NH, et al. Clinical risk stratification model for advanced colorectal neoplasia in persons with negative fecal immunochemical test results. *PLoS One* 2018;13:e0191125.
12. Jung YS, Ryu S, Chang Y, et al. Risk factors for colorectal neoplasia in persons aged 30 to 39 years and 40 to 49 years. *Gastrointest Endosc* 2015;81:637–45.e7.
13. Chang LC, Shun CT, Hsu WF, et al. Fecal immunochemical test detects sessile serrated adenomas and polyps with a low level of sensitivity. *Clin Gastroenterol Hepatol* 2017;15:872–9.e1.
14. Ahnen DJ, Wade SW, Jones WF, et al. The increasing incidence of young-onset colorectal cancer: A call to action. *Mayo Clin Proc* 2014;89:216–24.
15. Siegel RL, Miller KD, Fedewa SA, et al. Colorectal cancer statistics, 2017. *CA Cancer J Clin* 2017;67:177–93.
16. Bhandari A, Woodhouse M, Gupta S. Colorectal cancer is a leading cause of cancer incidence and mortality among adults younger than 50 years in the USA: A SEER-based analysis with comparison to other young-onset cancers. *J Invest Med* 2017;65:311–5.
17. Kim HG, Cho Y-S, Cha JM, et al. Risk of metachronous neoplasia on surveillance colonoscopy in young patients with colorectal neoplasia. *Gastrointest Endosc* 2018;87:666–73.
18. Nagpal SJS, Mukhija D, Sanaka M, et al. Metachronous colon polyps in younger versus older adults: A case-control study. *Gastrointest Endosc* 2018;87:657–65.
19. Adams LA, Waters OR, Knudman MW, et al. NAFLD as a risk factor for the development of diabetes and the metabolic syndrome: An eleven-year Follow-up study. *Am J Gastroenterol* 2009;104:861–7.
20. Huang KW, Leu HB, Wang YJ, et al. Patients with nonalcoholic fatty liver disease have higher risk of colorectal adenoma after negative baseline colonoscopy. *Colorectal Dis* 2013;15:830–5.
21. Cho Y, Lim SK, Joo SK, et al. Nonalcoholic steatohepatitis is associated with a higher risk of advanced colorectal neoplasm. *Liver Int* 2019;39:1722–31.
22. Mantovani A, Dauriz M, Byrne CD, et al. Association between nonalcoholic fatty liver disease and colorectal tumours in asymptomatic adults undergoing screening colonoscopy: A systematic review and meta-analysis. *Metabolism* 2018;87:1–12.

23. Wong MC, Ching JY, Chan VC, et al. Screening strategies for colorectal cancer among patients with nonalcoholic fatty liver disease and family history. *Int J Cancer* 2016;138:576–83.
24. de Wijkerslooth TR, Stoop EM, Bossuyt PM, et al. Immunochemical fecal occult blood testing is equally sensitive for proximal and distal advanced neoplasia. *Am J Gastroenterol* 2012;107:1570–8.

---

**Open Access** This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.