


# Mutations in two ERAD E3 ubiquitin ligase enzymes reduce spontaneous reversal frequency in *Caenorhabditis elegans*

Mackenzi Oswald<sup>1</sup>, Heino Hulsey-Vincent<sup>1</sup> and Caroline (Lina) Dahlberg<sup>1§</sup>

<sup>1</sup>Department of Biology, Western Washington University, Bellingham, WA, 98225, USA

§To whom correspondence should be addressed: lina.dahlberg@wwu.edu



**Figure 1. Mutations in E3 ubiquitin ligase genes and *glr-1* affect spontaneous reversal frequencies:** **A.** Spontaneous reversal frequency assays were performed using *C. elegans* strains harboring mutations in E3 ubiquitin ligases. Box and whisker plots show the average reversals per minute, bounded by quartiles; the line in each box represents the median of the average reversals per minute for each genotype. N=23 individual animals measured for WT, *hrd-1*, and *hrdl-1*; N=24 individual animals for *marc-6*. \* =  $p < 0.05$  ( $p=0.00069$  for *marc-6*;  $p=0.032$  for *hrd-1*); for *hrdl-1*,  $p=0.14$ . Significance was calculated using the Tukey-Kramer test following a one-way ANOVA. The data is normally distributed (Shapiro-Wilk test,  $p=0.37$ ) and groups show equal variance (residuals vs. fitted plot). **B.** Spontaneous reversal frequency for wild-type and *glr-1* (n2461) animals. N=20 individual animals for both genotypes; significance was calculated using Student's t-test ( $p=0.0063$ ). The data is normally distributed (Shapiro-Wilk test,  $p=0.52$ ) and groups show equal variance (residuals vs. fitted plot). **C.** Schematics of E3 ubiquitin ligase genes and *glr-1* (Oswald *et al.*, 2020). Schematics were made and annotated using <http://wormweb.org/exonintron>. The mutant E3 ubiquitin ligase strain *hrd-1* contains an indel, *hrdl-1* contains a deletion and *marc-6* contains a nonsense mutation resulting in a premature stop codon.

## Description

Secretory and membrane-bound proteins must be properly folded and matured at the endoplasmic reticulum (ER). Despite the molecular machinery dedicated to these processes, up to 1/3 of proteins are destroyed within minutes of their synthesis (Hirsch *et al.* 2009; Schubert *et al.* 2000). Misfolded proteins in the endoplasmic reticulum can accumulate and disrupt

proteostasis, which can contribute to neurodegenerative diseases. The Endoplasmic Reticulum Associated Degradation (ERAD) pathway relies on E2 ubiquitin-conjugating enzymes and E3 ubiquitin ligases to ubiquitylate misfolded proteins, signaling for degradation of these misfolded proteins by the proteasome (Vembar and Brodsky 2008). Three putative E3 ligases that are expected to be involved in ERAD in *C. elegans* are HRDL-1, HRD-1, and MARC-6 (Sasagawa *et al.* 2007). We used strains harboring mutations in *hrdl-1*, *hrd-1* and *marc-6* genes to determine if these proteins are required for regulating spontaneous reversal behavior in *C. elegans*.

Spontaneous reversals are a *C. elegans* behavior whose frequency is regulated by well-defined circuitry and neurotransmitter receptors, including the glutamate receptor, *glr-1* (Brockie *et al.* 2001; Burbea *et al.* 2002; Dahlberg and Juo 2014, Hart *et al.* 1995; Kowalski *et al.* 2011; Zheng *et al.* 1999). We hypothesized that the spontaneous reversal frequency behavior of *C. elegans* would be affected by E3 ligase mutations if they are important for normal spontaneous reversal behavior. *glr-1* animals reverse significantly less than wild-type animals and were used as a positive control (Figure 1B) (Hart *et al.* 1995; Kowalski *et al.* 2011). Animals harboring mutations in *marc-6* and *hrd-1* also reverse significantly less than wild-type animals (Figure 1A). Animals lacking full-length *hrdl-1* reversed less than wild-type animals, but this was not statistically significant (Figure 1A).

The primary motivation for our work was to ask if the glutamate receptor, GLR-1, might be regulated by these E3s, but further research must be done in order to determine molecular mechanisms that cause the differences in behavior that we report. Because ERAD E2 and E3 proteins can compensate for each other's absence, we hypothesize that in the absence of any one E3 ligase, others are upregulated either through protein activity or gene expression (Bays *et al.* 2001; Weber *et al.* 2016). For example, if HRD-1 is upregulated to compensate for the putative loss-of-function of HRDL-1 in *hrdl-1* mutant animals, this could explain why *hrdl-1* animals did not show a statistically significant reduction in reversals/minute compared to wild-type animals. However, our results from the *hrd-1* and *marc-6* mutants suggest that there is not complete redundancy between the E3 ligases. Future experiments will focus on testing this hypothesis using double and triple mutants in the E3 ligase genes.

## Methods

[Request a detailed protocol](#)

**Reversal assay protocol:** Each *C. elegans* strain was grown at 21.4°C in the same bin on separate NGM agar plates seeded with OP50. Before performing the reversal assays, young adult hermaphroditic nematodes from each strain were picked onto separate seeded NGM agar plates and coded. This allows the later analysis of videotaped trials to be blind in order to reduce potential bias when scoring reversal assays. One animal at a time was picked from its coded seeded plate onto an unseeded NGM agar plate using halocarbon oil (a non-food substance) to induce food-seeking behavior. The animal was allowed to move around on the plate for two minutes before videotaping began. If the animal did not move away from its initial position on the plate after two minutes it was discarded and not counted in the data set. Each animal was recorded for five minutes and then discarded. At least one of each strain was observed during each experimental session in order to allow for any potential variations in temperature and humidity to be accounted for equally across all strains. The total N values listed represent measurements from multiple experimental sessions.

**Scoring reversals:** After recording all reversal assay trials for an experiment, the recorded coded trials were viewed and scored blindly by one researcher. A reversal was only counted if the animal moved backwards at least 1/6th of its body-length. The posterior pharyngeal bulb position was used as a marker for that distance.

**Recording setup:** Videos were recorded using an Olympus SZ61 microscope attached to a TLB 4000 Series Substage Illuminator base. The microscope was connected to The Imaging Source DFK 31AF03 color camera, which was connected to a computer running Windows OS. The software used to record the videos was Debut Professional by NCH Software.

**Genotyping:** The genotype of KP4 was confirmed using DNA sequencing. The genotypes of FJ861 and FJ863 were confirmed by PCR. The genotype of CLD33 was confirmed using PCR and restriction digest mapping using EarI (method noted on Million Mutation Project website, <http://genome.sfu.ca/mmp/about.html>).

Note on background transgene in CLD33: although the *marc-6* mutation is in a GLR-1::GFP background, GLR-1::GFP animals show similar responsiveness to nose-touch assays (another GLR-1 dependent behavior) compared to wild-type animals (Rongo *et al.* 1998), and we do not expect that it affected the response of the animals.

## Reagents

**Table 1.** *C. elegans* strains used.

Strain Name	Genotype	Description	Reference
N2		Wild-type	

KP4	<i>glr-1(n2461)</i> III	<i>glr-1</i> putative knockout, nonsense mutation	Hart AC <i>et al.</i> 1995
FJ861	<i>hrd-1(tm1743)</i> V*6	E3 ubiquitin ligase gene <i>hrd-1</i> putative knockout, indel, backcrossed 6 times	<i>C. elegans</i> Deletion Mutant Consortium 2012
FJ863	<i>hrdl-1(gk28)</i> I*6	E3 ubiquitin ligase gene <i>hrdl-1</i> putative knockout, deletion. Original strain VC35, backcrossed 6 times	<i>C. elegans</i> Deletion Mutant Consortium 2012
CLD33	<i>nuIs24(IV);marc-6(gk121407)</i> I*6	Transgene <i>Pglr-1::GLR-1::GFP</i> crossed with E3 ubiquitin ligase gene <i>marc-6</i> putative knockout, nonsense mutation, backcrossed 6 times. Original <i>marc-6</i> strain is VC20284. Please also see the note in Methods.	WormBase ID: WBTransgene00001321  WormBase ID: WBVar00344650.

**Acknowledgments:** We thank the students of Western Washington University's Biology 487 course (Spring 2020) and Dr. Jacqueline Rose for critical readings of this manuscript.

## References

- Bays NW, Gardner RG, Seelig LP, Joazeiro CA, Hampton RY. 2001. Hrd1p/Der3p is a membrane-anchored ubiquitin ligase required for ER-associated degradation. *Nat Cell Biol* 3: 24-9. PMID: 11146622.
- Brockie PJ, Mellem JE, Hills T, Madsen DM, Maricq AV. 2001. The *C. elegans* glutamate receptor subunit NMR-1 is required for slow NMDA-activated currents that regulate reversal frequency during locomotion. *Neuron* 31: 617-30. PMID: 11545720.
- Burbea M, Dreier L, Dittman JS, Grunwald ME, Kaplan JM. 2002. Ubiquitin and AP180 regulate the abundance of GLR-1 glutamate receptors at postsynaptic elements in *C. elegans*. *Neuron* 35: 107-20. PMID: 12123612.
- C. elegans* Deletion Mutant Consortium. 2012. large-scale screening for targeted knockouts in the *Caenorhabditis elegans* genome. *G3 (Bethesda)* 2: 1415-25. PMID: 23173093.
- Dahlberg CL, Juo P. 2014. The WD40-repeat proteins WDR-20 and WDR-48 bind and activate the deubiquitinating enzyme USP-46 to promote the abundance of the glutamate receptor GLR-1 in the ventral nerve cord of *Caenorhabditis elegans*. *J Biol Chem* 289: 3444-56. PMID: 24356955.
- Hart AC, Sims S, Kaplan JM. 1995. Synaptic code for sensory modalities revealed by *C. elegans* GLR-1 glutamate receptor. *Nature* 378: 82-5. PMID: 7477294.
- Hirsch C, Gauss R, Horn SC, Neuber O, Sommer T. 2009. The ubiquitylation machinery of the endoplasmic reticulum. *Nature* 458: 453-60. PMID: 19325625.
- Kowalski JR, Dahlberg CL, Juo P. 2011. The deubiquitinating enzyme USP-46 negatively regulates the degradation of glutamate receptors to control their abundance in the ventral nerve cord of *Caenorhabditis elegans*. *J Neurosci* 31: 1341-54. PMID: 21273419.
- Oswald, M; Hulsey-Vincent, H; Dahlberg, C (2020). Individual point mutations in two ERAD E2 ubiquitin-conjugating enzymes do not affect *Caenorhabditis elegans* spontaneous reversal frequency. *microPublication Biology*. PMID: 10.17912/micropub.biology.000328. .
- Rongo C, Whitfield CW, Rodal A, Kim SK, Kaplan JM. 1998. LIN-10 is a shared component of the polarized protein localization pathways in neurons and epithelia. *Cell* 94: 751-9. PMID: 9753322.
- Sasagawa Y, Yamanaka K, Ogura T. 2007. ER E3 ubiquitin ligase HRD-1 and its specific partner chaperone BiP play important roles in ERAD and developmental growth in *Caenorhabditis elegans*. *Genes Cells* 12: 1063-73. PMID: 17825049.
- Schubert U, Antón LC, Gibbs J, Norbury CC, Yewdell JW, Bennink JR. 2000. Rapid degradation of a large fraction of newly synthesized proteins by proteasomes. *Nature* 404: 770-4. PMID: 10783891.
- Vembar SS, Brodsky JL. 2008. One step at a time: endoplasmic reticulum-associated degradation. *Nat Rev Mol Cell Biol* 9: 944-57. PMID: 19002207.
- Weber A, Cohen I, Popp O, Dittmar G, Reiss Y, Sommer T, Ravid T, Jarosch E. 2016. Sequential Poly-ubiquitylation by Specialized Conjugating Enzymes Expands the Versatility of a Quality Control Ubiquitin Ligase. *Mol Cell* 63: 827-39. PMID: 27570077.
- WormBase web site, <http://www.wormbase.org>. WormBase ID: WBTransgene00001321.

WormBase web site, <http://www.wormbase.org>. WormBase ID: WBVar00344650.

Zheng Y, Brockie PJ, Mellem JE, Madsen DM, Maricq AV. 1999. Neuronal control of locomotion in *C. elegans* is modified by a dominant mutation in the GLR-1 ionotropic glutamate receptor. *Neuron* 24: 347-61. PMID: 10571229.

**Funding:** Partial funding was provided by a Research and Creative Opportunities Grant for Undergraduate Students from the Office of Research & Sponsored Programs at Western Washington University.

**Author Contributions:** Mackenzi Oswald: Formal analysis, Funding acquisition, Investigation, Methodology, Writing - original draft, Writing - review and editing. Heino Hulsev-Vincent: Formal analysis, Visualization, Writing - review and editing. Caroline (Lina) Dahlberg : Conceptualization, Formal analysis, Funding acquisition, Methodology, Project administration, Resources, Supervision, Writing - review and editing.

**Reviewed By:** Eric Luth and Anonymous

**History:** Received June 2, 2020 Revision received November 9, 2020 Accepted November 10, 2020 Published November 19, 2020

**Copyright:** © 2020 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Citation:** Oswald, M; Hulsev-Vincent, H; Dahlberg, C (2020). Mutations in two ERAD E3 ubiquitin ligase enzymes reduce spontaneous reversal frequency in *Caenorhabditis elegans*. *microPublication Biology*. <https://doi.org/10.17912/micropub.biology.000329>