



Cochrane
Library

Cochrane Database of Systematic Reviews

Interventions to improve water quality for preventing diarrhoea (Review)

Clasen TF, Alexander KT, Sinclair D, Boisson S, Peletz R, Chang HH, Majorin F, Cairncross S

Clasen TF, Alexander KT, Sinclair D, Boisson S, Peletz R, Chang HH, Majorin F, Cairncross S.

Interventions to improve water quality for preventing diarrhoea.

Cochrane Database of Systematic Reviews 2015, Issue 10. Art. No.: CD004794.

DOI: [10.1002/14651858.CD004794.pub3](https://doi.org/10.1002/14651858.CD004794.pub3).

www.cochranelibrary.com

Interventions to improve water quality for preventing diarrhoea (Review)

Copyright © 2015 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

WILEY

TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	6
OBJECTIVES	6
METHODS	7
RESULTS	9
Figure 1.	9
Figure 2.	12
Figure 3.	13
Figure 4.	14
Figure 5.	15
Figure 6.	16
DISCUSSION	17
AUTHORS' CONCLUSIONS	18
ACKNOWLEDGEMENTS	18
REFERENCES	20
CHARACTERISTICS OF STUDIES	32
DATA AND ANALYSES	98
Analysis 1.1. Comparison 1 Water quality intervention versus control, Outcome 1 Diarrhoea: all ages.	98
Analysis 1.2. Comparison 1 Water quality intervention versus control, Outcome 2 Diarrhoea: children < 5 years.	100
Analysis 2.1. Comparison 2 Source: water supply improvement versus control, Outcome 1 Diarrhoea: CBA studies subgrouped by age.	102
Analysis 2.2. Comparison 2 Source: water supply improvement versus control, Outcome 2 Diarrhoea: CBA studies subgrouped by age.	102
Analysis 3.1. Comparison 3 POU: water chlorination versus control, Outcome 1 Diarrhoea: subgrouped by study design.	104
Analysis 3.2. Comparison 3 POU: water chlorination versus control, Outcome 2 Diarrhoea: cluster-RCTs: subgrouped by age. ..	105
Analysis 3.3. Comparison 3 POU: water chlorination versus control, Outcome 3 Diarrhoea: cluster-RCTs; subgrouped by adherence.	106
Analysis 3.4. Comparison 3 POU: water chlorination versus control, Outcome 4 Diarrhoea: cluster-RCTs by risk of bias by blinding of participants.	107
Analysis 3.5. Comparison 3 POU: water chlorination versus control, Outcome 5 Diarrhoea: cluster-RCTs; subgrouped by additional water storage intervention.	108
Analysis 3.6. Comparison 3 POU: water chlorination versus control, Outcome 6 Diarrhoea: cluster-RCTs: subgrouped by sufficiency of water quantity.	109
Analysis 3.7. Comparison 3 POU: water chlorination versus control, Outcome 7 Diarrhoea: cluster-RCTs: subgrouped by water source.	109
Analysis 3.8. Comparison 3 POU: water chlorination versus control, Outcome 8 Diarrhoea: cluster-RCTs: subgrouped by sanitation level.	110
Analysis 3.9. Comparison 3 POU: water chlorination versus control, Outcome 9 Diarrhoea: cluster-RCTs; subgrouped by length of follow-up.	111
Analysis 4.1. Comparison 4 POU: flocculation and disinfection versus control, Outcome 1 Diarrhoea: cluster-RCTs.	113
Analysis 4.2. Comparison 4 POU: flocculation and disinfection versus control, Outcome 2 Diarrhoea: cluster-RCTs: subgrouped by age; excluding Doocy 2006 LBR.	114
Analysis 4.3. Comparison 4 POU: flocculation and disinfection versus control, Outcome 3 Diarrhoea: cluster-RCTs: subgrouped by adherence.	114
Analysis 4.4. Comparison 4 POU: flocculation and disinfection versus control, Outcome 4 Diarrhoea: cluster-RCTs: subgrouped by additional storage container.	115
Analysis 4.5. Comparison 4 POU: flocculation and disinfection versus control, Outcome 5 Diarrhoea: cluster-RCTs: subgrouped by sufficiency of water quantity.	115
Analysis 4.6. Comparison 4 POU: flocculation and disinfection versus control, Outcome 6 Diarrhoea: cluster-RCTs: subgrouped by water source.	116

Analysis 4.7. Comparison 4 POU: flocculation and disinfection versus control, Outcome 7 Diarrhoea: cluster-RCTs: subgrouped by sanitation level.	117
Analysis 4.8. Comparison 4 POU: flocculation and disinfection versus control, Outcome 8 Diarrhoea: cluster-RCTs: subgrouped by length of follow-up.	117
Analysis 5.1. Comparison 5 POU: filtration versus control, Outcome 1 Diarrhoea: cluster-RCTs: subgrouped by age.	120
Analysis 5.2. Comparison 5 POU: filtration versus control, Outcome 2 Diarrhoea: cluster-RCTs: subgrouped by type of filtration.	121
Analysis 5.3. Comparison 5 POU: filtration versus control, Outcome 3 Diarrhoea: cluster-RCTs: subgrouped by blinding of participants.	122
Analysis 5.4. Comparison 5 POU: filtration versus control, Outcome 4 Diarrhoea: ceramic filter studies subgrouped by water source.	123
Analysis 5.5. Comparison 5 POU: filtration versus control, Outcome 5 Diarrhoea: ceramic filter studies subgrouped by sanitation level.	123
Analysis 5.6. Comparison 5 POU: filtration versus control, Outcome 6 Diarrhoea: sand filter studies: subgrouped by water source.	124
Analysis 5.7. Comparison 5 POU: filtration versus control, Outcome 7 Diarrhoea: sand filter studies: subgrouped by sanitation level.	125
Analysis 5.8. Comparison 5 POU: filtration versus control, Outcome 8 Diarrhoea: cluster-RCTs: subgrouped by adherence.	125
Analysis 5.9. Comparison 5 POU: filtration versus control, Outcome 9 Diarrhoea: cluster-RCTs: subgrouped by additional water storage intervention.	126
Analysis 5.10. Comparison 5 POU: filtration versus control, Outcome 10 Diarrhoea: cluster-RCTs; subgrouped by length of follow-up.	127
Analysis 6.1. Comparison 6 POU: solar disinfection versus control, Outcome 1 Diarrhoea: subgrouped by study design.	129
Analysis 6.2. Comparison 6 POU: solar disinfection versus control, Outcome 2 Diarrhoea: cluster-RCTs; subgrouped by age. ...	130
Analysis 6.3. Comparison 6 POU: solar disinfection versus control, Outcome 3 Diarrhoea: cluster-RCTs; subgrouped by adherence.	131
Analysis 6.4. Comparison 6 POU: solar disinfection versus control, Outcome 4 Diarrhoea: cluster-RCTs; subgrouped by sufficiency of water supply level.	131
Analysis 6.5. Comparison 6 POU: solar disinfection versus control, Outcome 5 Diarrhoea: cluster-RCTs; subgrouped by water source.	132
Analysis 6.6. Comparison 6 POU: solar disinfection versus control, Outcome 6 Diarrhoea: cluster-RCTs; subgrouped by sanitation level.	132
Analysis 6.7. Comparison 6 POU: solar disinfection versus control, Outcome 7 Diarrhoea: cluster-RCTs; subgrouped by length of follow-up.	133
Analysis 7.1. Comparison 7 POU: UV disinfection versus control, Outcome 1 Diarrhoea: cluster-RCT.	134
Analysis 8.1. Comparison 8 POU: improved storage versus control, Outcome 1 Diarrhoea: cluster-RCTs: subgrouped by age. ...	134
ADDITIONAL TABLES	134
APPENDICES	170
WHAT'S NEW	172
HISTORY	172
CONTRIBUTIONS OF AUTHORS	173
DECLARATIONS OF INTEREST	173
SOURCES OF SUPPORT	173
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	173
INDEX TERMS	173

[Intervention Review]

Interventions to improve water quality for preventing diarrhoea

Thomas F Clasen¹, Kelly T Alexander¹, David Sinclair², Sophie Boisson³, Rachel Peletz⁴, Howard H Chang⁵, Fiona Majorin³, Sandy Cairncross⁶

¹Department of Environmental Health, Rollins School of Public Health, Emory University, Atlanta, GA, USA. ²Department of Clinical Sciences, Liverpool School of Tropical Medicine, Liverpool, UK. ³Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London, UK. ⁴Aquaya Institute, Nairobi, Kenya. ⁵Department of Biostatistics and Bioinformatics, Rollins School of Public Health, Emory University, Atlanta, GA, USA. ⁶Department of Disease Control, Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, London, UK

Contact: Thomas F Clasen, Department of Environmental Health, Rollins School of Public Health, Emory University, 1518 Clifton Road NE, Atlanta, GA, 30322, USA. thomas.clasen@lshtm.ac.uk.

Editorial group: Cochrane Infectious Diseases Group.

Publication status and date: Unchanged, published in Issue 10, 2015.

Citation: Clasen TF, Alexander KT, Sinclair D, Boisson S, Peletz R, Chang HH, Majorin F, Cairncross S. Interventions to improve water quality for preventing diarrhoea. *Cochrane Database of Systematic Reviews* 2015, Issue 10. Art. No.: CD004794. DOI: [10.1002/14651858.CD004794.pub3](https://doi.org/10.1002/14651858.CD004794.pub3).

Copyright © 2015 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration. This is an open access article under the terms of the [Creative Commons Attribution-Non-Commercial](https://creativecommons.org/licenses/by-nc/4.0/) Licence, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

ABSTRACT

Background

Diarrhoea is a major cause of death and disease, especially among young children in low-income countries. In these settings, many infectious agents associated with diarrhoea are spread through water contaminated with faeces.

In remote and low-income settings, source-based water quality improvement includes providing protected groundwater (springs, wells, and bore holes), or harvested rainwater as an alternative to surface sources (rivers and lakes). Point-of-use water quality improvement interventions include boiling, chlorination, flocculation, filtration, or solar disinfection, mainly conducted at home.

Objectives

To assess the effectiveness of interventions to improve water quality for preventing diarrhoea.

Search methods

We searched the Cochrane Infectious Diseases Group Specialized Register (11 November 2014), CENTRAL (the Cochrane Library, 7 November 2014), MEDLINE (1966 to 10 November 2014), EMBASE (1974 to 10 November 2014), and LILACS (1982 to 7 November 2014). We also handsearched relevant conference proceedings, contacted researchers and organizations working in the field, and checked references from identified studies through 11 November 2014.

Selection criteria

Randomized controlled trials (RCTs), quasi-RCTs, and controlled before-and-after studies (CBA) comparing interventions aimed at improving the microbiological quality of drinking water with no intervention in children and adults.

Data collection and analysis

Two review authors independently assessed trial quality and extracted data. We used meta-analyses to estimate pooled measures of effect, where appropriate, and investigated potential sources of heterogeneity using subgroup analyses. We assessed the quality of evidence using the GRADE approach.

Main results

Forty-five cluster-RCTs, two quasi-RCTs, and eight CBA studies, including over 84,000 participants, met the inclusion criteria. Most included studies were conducted in low- or middle-income countries (LMICs) (50 studies) with unimproved water sources (30 studies) and unimproved or unclear sanitation (34 studies). The primary outcome in most studies was self-reported diarrhoea, which is at high risk of bias due to the lack of blinding in over 80% of the included studies.

Source-based water quality improvements

There is currently insufficient evidence to know if source-based improvements such as protected wells, communal tap stands, or chlorination/filtration of community sources consistently reduce diarrhoea (one cluster-RCT, five CBA studies, *very low quality evidence*). We found no studies evaluating reliable piped-in water supplies delivered to households.

Point-of-use water quality interventions

On average, distributing water disinfection products for use at the household level may reduce diarrhoea by around one quarter (Home chlorination products: RR 0.77, 95% CI 0.65 to 0.91; 14 trials, 30,746 participants, *low quality evidence*; flocculation and disinfection sachets: RR 0.69, 95% CI 0.58 to 0.82, four trials, 11,788 participants, *moderate quality evidence*). However, there was substantial heterogeneity in the size of the effect estimates between individual studies.

Point-of-use filtration systems probably reduce diarrhoea by around a half (RR 0.48, 95% CI 0.38 to 0.59, 18 trials, 15,582 participants, *moderate quality evidence*). Important reductions in diarrhoea episodes were shown with ceramic filters, biosand systems and LifeStraw® filters; (Ceramic: RR 0.39, 95% CI 0.28 to 0.53; eight trials, 5763 participants, *moderate quality evidence*; Biosand: RR 0.47, 95% CI 0.39 to 0.57; four trials, 5504 participants, *moderate quality evidence*; LifeStraw®: RR 0.69, 95% CI 0.51 to 0.93; three trials, 3259 participants, *low quality evidence*). Plumbed in filters have only been evaluated in high-income settings (RR 0.81, 95% CI 0.71 to 0.94, three trials, 1056 participants, fixed effects model).

In low-income settings, solar water disinfection (SODIS) by distribution of plastic bottles with instructions to leave filled bottles in direct sunlight for at least six hours before drinking probably reduces diarrhoea by around a third (RR 0.62, 95% CI 0.42 to 0.94; four trials, 3460 participants, *moderate quality evidence*).

In subgroup analyses, larger effects were seen in trials with higher adherence, and trials that provided a safe storage container. In most cases, the reduction in diarrhoea shown in the studies was evident in settings with improved and unimproved water sources and sanitation.

Authors' conclusions

Interventions that address the microbial contamination of water at the point-of-use may be important interim measures to improve drinking water quality until homes can be reached with safe, reliable, piped-in water connections. The average estimates of effect for each individual point-of-use intervention generally show important effects. Comparisons between these estimates do not provide evidence of superiority of one intervention over another, as such comparisons are confounded by the study setting, design, and population.

Further studies assessing the effects of household connections and chlorination at the point of delivery will help improve our knowledge base. As evidence suggests effectiveness improves with adherence, studies assessing programmatic approaches to optimising coverage and long-term utilization of these interventions among vulnerable populations could also help strategies to improve health outcomes.

16 April 2019

Update pending

Studies awaiting assessment

The CIDG is currently examining a new search conducted in April 2019 for potentially relevant studies. These studies have not yet been incorporated into this Cochrane Review.

PLAIN LANGUAGE SUMMARY

Interventions to improve water quality and prevent diarrhoea

This Cochrane Review summarizes trials evaluating different interventions to improve water quality and prevent diarrhoea. After searching for relevant trials up to 11 November 2014, we included 55 studies enrolling over 84,000 participants. Most included studies were conducted

Interventions to improve water quality for preventing diarrhoea (Review)

Copyright © 2015 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

in low- or middle-income countries (LMICs) (50 studies), with unimproved water sources (30 studies), and unimproved or unclear sanitation (34 studies).

What causes diarrhoea and what water quality interventions might prevent diarrhoea?

Diarrhoea is a major cause of death and disease, especially among young children in low-income countries where the most common causes are faecally contaminated water and food, or poor hygiene practices.

In remote and low-income settings, source-based water quality improvement may include providing protected groundwater (springs, wells, and bore holes) or harvested rainwater as an alternative to surface sources (rivers and lakes). Alternatively water may be treated at the point-of-use in people's homes by boiling, chlorination, flocculation, filtration, or solar disinfection. These point-of-use interventions have the potential to overcome both contaminated sources and recontamination of safe water in the home.

What the research says

There is currently insufficient evidence to know if source-based improvements in water supplies, such as protected wells and communal tap stands or treatment of communal supplies, consistently reduce diarrhoea in low-income settings (*very low quality evidence*). We found no trials evaluating reliable piped-in water supplies to people's homes.

On average, distributing disinfection products for use in the home may reduce diarrhoea by around one quarter in the case of chlorine products (*low quality evidence*), and around a third in the case of flocculation and disinfection sachets (*moderate quality evidence*).

Water filtration at home probably reduces diarrhoea by around a half (*moderate quality evidence*), and effects were consistently seen with ceramic filters (*moderate quality evidence*), biosand systems (*moderate quality evidence*) and LifeStraw® filters (*low quality evidence*). Plumbed-in filtration has only been evaluated in high-income settings (*low quality evidence*).

In low-income settings, distributing plastic bottles with instructions to leave filled bottles in direct sunlight for at least six hours before drinking probably reduces diarrhoea by around a third (*moderate quality evidence*).

Research assessing the effects of household connections and chlorination at the point of delivery will help improve our knowledge base. Evidence indicates the more people use the various interventions for improving water quality, the larger the effects, so research into practical approaches to increase coverage and help assure long term use of them in poor groups will help improve impact.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. Summary of findings table 1

Point-of-use water quality interventions for preventing diarrhoea in rural settings in low- and middle-income countries

Patient or population: adults and children

Settings: low- and middle-income countries in rural areas

Intervention: point of use water quality interventions

Comparison: no intervention

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	Number of participants (trials)	Quality of the evidence (GRADE)
	Assumed risk	Corresponding risk			
Diarrhoea episodes	No intervention	Chlorination	RR 0.77 (0.65 to 0.91)	30,746 (14 trials)	⊕⊕⊕⊕ low 1,2,3,4
	3 episodes per person per year	2.3 episodes (2.0 to 2.7)			
	No intervention	Flocculation/disinfection	RR 0.69 (0.58 to 0.82)	11,788 (4 trials)	⊕⊕⊕⊕ moderate 1,3,4,5,6
	3 episodes per person per year	2.1 episodes (1.7 to 2.5)			
	No intervention	Filtration	RR 0.48 (0.38 to 0.59)	15,582 (18 trials)	⊕⊕⊕⊕ moderate 1,3,4,5
	3 episodes per person per year	1.4 episodes (1.1 to 1.8)			
	No intervention	Solar disinfection (SODIS)	RR 0.62 (0.42 to 0.94)	3460 (4 trials)	⊕⊕⊕⊕ moderate 1,3,4,5
	3 episodes per person per year	1.9 episodes (1.3 to 2.8)			

The **assumed risk** is taken from [Fischer Walker 2012](#) and represents an estimated average for the incidence of diarrhoea in low- and middle-income countries. The **corresponding risk** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **RR:** risk ratio.

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: we are very uncertain about the estimate.

¹Downgraded by 1 for serious risk of bias: the outcome was measured as self-reported episodes of diarrhoea, and is susceptible to bias as most studies were unblinded.

²Downgraded by 1 for serious inconsistency: statistical heterogeneity was very high with six out of fourteen trials having point estimates close to no effect. A subgroup analysis by adherence with the intervention (assessed by measurements of residual chlorine in drinking water) found larger effects in the studies with better adherence but the results remained inconsistent.

³No serious indirectness: these studies are mainly from low- and middle-income countries, in settings with both improved and unimproved water sources and sanitation.

⁴No serious imprecision: The analysis is adequately powered to detect this effect.

⁵No serious inconsistency: The evidence of benefit is consistent across trials, but there is substantial statistical heterogeneity in the size of the effect.

⁶ This analysis excludes one additional study which found a much larger effect than seen in the other four trials and was considered an outlier ([Doocy 2006 LBR](#)).

BACKGROUND

Description of the condition

Diarrhoeal disease is the third leading cause of mortality in low-income countries, causing an estimated 1.4 million deaths in 2012 (WHO 2014;GBD 2015). Young children are especially vulnerable, with diarrhoea accounting for more than a quarter of all deaths in children aged under five years in Africa and Southeast Asia (Murray 2012; Lanata 2013; Walker 2013).

The bacterial, viral, and protozoan pathogens causing diarrhoeal disease are primarily transmitted via the faecal-oral route, through the consumption of faecally contaminated food and water (Byers 2001). Among the most important of these are rotavirus, *Cryptosporidium* sp., *Escherichia coli*, *Salmonella* sp., *Shigella* sp., *Campylobacter jejuni*, *Vibrio cholerae*, norovirus, *Giardia lamblia*, and *Entamoeba histolytica* (Leclerc 2002; Kotloff 2013), though the relative importance of these varies among settings, seasons, and population groups.

An estimated 1.1 billion people worldwide rely on water supplies that are at high risk of faecal contamination (Bain 2014). Moreover, nearly half the world's population lack household water connections (WHO/UNICEF 2015), and are at increased risk of unsafe water due to contamination during collection, storage, and use in the home (Wright 2004).

Description of the intervention

Interventions to improve the microbiological quality of water can be grouped into four main categories:

- Physical removal of pathogens (for example, filtration, adsorption, or sedimentation).
- Chemical treatment to kill or deactivate pathogens (most commonly with chlorine).
- Disinfection by heat (for example, boiling or pasturization) or ultraviolet (UV) radiation (for example, solar disinfection, or artificial UV lamps).
- Combination of these approaches (for example, filtration or flocculation combined with disinfection).

In higher-income countries, and in many urban settings worldwide, drinking water is treated centrally at the source of supply and distributed to consumers through a network of pipes and household taps. Alternatively, water may be treated at any point in the distribution network, or at the 'point-of-use' (POU) in people's homes, schools, or workplaces.

In remote and low-income settings, source-based water quality improvement may include providing protected groundwater (springs, wells, and bore holes) or harvested rainwater as an alternative to surface sources (rivers and lakes). These improvements frequently also improve both the quantity and access to water by increasing the volume or frequency of water delivery or reducing the time spent in collecting water. This may result in significant benefits not only in health but also in economic and social welfare (Hutton 2013; Stelmach 2015).

Potential and widely used POU interventions for remote or low-income settings include boiling, filtration, chlorination, flocculation, and solar disinfection. These interventions have

the potential to overcome both contaminated sources and recontamination of safe water in the home (Wright 2004). A review commissioned by the World Health Organization (WHO) identified a wide variety of options for household-based water treatment and assessed the available evidence on their microbiological effectiveness, health impact, acceptability, affordability, sustainability, and scalability (Sobsey 2002).

How the intervention might work

Health authorities generally accept that microbiologically safe water plays an important role in preventing outbreaks of waterborne diseases (Reynolds 2008). Moreover, there is evidence that chlorination and filtration of municipal water supplies contributed to substantial health gains in the late 19th and early 20th century (Cutler 2005).

However, much of the epidemiological evidence for increased health benefits following improvements in the quality of drinking water has been equivocal, particularly in low-income settings (Clasen 2006; Waddington 2009; Cairncross 2010).

This may be due to the variety of alternative transmission pathways, such as ingestion of contaminated food, person-to-person contact, or direct contact with infected faeces. In addition, interventions which only target the home may fail if unsafe water is consumed at work or school. Consequently, effective programmes may require combined interventions to address not only water quality, but also water quantity and access, the proper disposal of human faeces (sanitation), and the promotion of hand washing and hygiene practices within communities.

The effectiveness of individual water quality interventions may also vary between settings due to the varied prevalence of the organisms causing diarrhoea. For instance, ceramic filters are only marginally protective against viral illness, while chlorination may provide little protection against *Cryptosporidium*.

Why it is important to do this review

This is an update of a Cochrane Review that was first completed in 2006 (Clasen 2006). The review concluded that, in general, interventions to improve microbiological quality of drinking water are effective in preventing diarrhoea, and that interventions at the household level were more effective than those at the source.

New studies have been recently published, and other unpublished studies have been made available to us. In this Cochrane Review update, we have reapplied the inclusion criteria, repeated data extraction, added new studies, and used the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach to assess the quality of the evidence. We were also able to apply statistical methods to unify the measures of effect and to apply additional criteria for subgrouping based on study design, setting, and length of follow-up.

OBJECTIVES

To assess the effectiveness of interventions to improve water quality for preventing diarrhoea.

Interventions to improve water quality for preventing diarrhoea (Review)

METHODS

Criteria for considering studies for this review

Types of studies

Cluster-randomized controlled trials (cluster-RCTs), quasi-randomized controlled trials (quasi-RCTs) and controlled before-and-after studies (CBAs).

Types of participants

Children and adults.

Types of interventions

Intervention

Any intervention aimed at improving the microbiological quality of drinking water.

We included interventions that combined improvements in water quality with hygiene or health promotion, but excluded studies that combined water quality interventions with other water, sanitation, and hygiene (WASH) interventions, such as improvements in water quantity or sanitation. We also excluded studies where the water quality intervention was implemented away from the home, such as at schools, clinics, markets, or workplaces.

Control

No intervention, or a dummy intervention.

Types of outcome measures

Primary

- Diarrhoea episodes among individuals, whether or not confirmed by microbiological examination.

The WHO's definition of diarrhoea is three or more loose or fluid stools (that take the shape of the container) in a 24-hour period ([WHO 1993](#)). We defined diarrhoea and an episode in accordance with the case definitions used in each trial. In the 'Summary of findings' tables, we have converted the results to episodes per year from a baseline of three episodes/child year in 2010 ([Fischer Walker 2012](#)).

Secondary

- Death.
- Adverse events.

We excluded studies that had no clinical outcomes; for example, studies that only report on microbiological pathogens in the stool.

Search methods for identification of studies

We attempted to identify all relevant studies regardless of language or publication status (published, unpublished, in press, and in progress).

Electronic searches

We searched the following databases using the search terms and strategy described in [Appendix 1](#): Cochrane Infectious Diseases Group Specialized Register (11 November 2014); Cochrane Central Register of Controlled Trials (CENTRAL), published in the Cochrane Library (7 November, 2014); MEDLINE (1966 to 10 November 2014);

EMBASE (1974 to 10 November 2014); and LILACS (1982 to 7 November 2014).

Searching other resources

Conference proceedings

We searched the conference proceedings of the following organizations for relevant abstracts: International Water Association (IWA) (1990 to 11 November 2014); and Water, Engineering and Development Centre, Loughborough University, UK (WEDC) (1973 to 11 November 2014).

Researchers and organizations

We contacted individual researchers working in the field and the following organizations for unpublished and ongoing studies: Water, Sanitation and Health Programme of the WHO; World Bank Water and Sanitation Program; UNICEF Water, Sanitation and Hygiene; and IRC International Water and Sanitation Centre; Foodborne and Diarrhoeal Diseases Branch, Division of Bacterial and Mycotic Diseases, Centers for Disease Control and Prevention (CDC); US Agency for International Development (USAID), including its Environmental Health Project (EHP); and the UK Department for International Development (DFID).

Reference lists

We checked the reference lists of all studies identified by the above methods.

Data collection and analysis

Selection of studies

Two review authors (RP and SB) independently reviewed the titles and abstracts located in the searches and selected all potentially relevant studies. After obtaining the full-text articles, they independently determined whether they met the inclusion criteria. Where they were unable to agree, they consulted a third review author (TFC) and arrived at a consensus. We have listed the potentially relevant studies that were ultimately excluded together with the reasons for exclusion in the '[Characteristics of excluded studies](#)' section.

Data extraction and management

Two review authors (RP and SB) used a pre-piloted form to extract and record the data described in [Appendix 2](#). One review author entered the extracted data into [Review Manager \(RevMan\)](#) (KA).

Assessment of risk of bias in included studies

Two review authors (KA and FM) independently assessed the risk of bias of the included studies and resolved differences of opinion through discussion.

For cluster-RCTs we used the Cochrane 'Risk of bias' assessment tool ([Higgins 2011](#)). We followed the guidance to assess whether adequate steps were taken to reduce the risk of bias across five domains: sequence generation; allocation concealment; blinding of participants and personnel; blinding of outcome assessors; and incomplete outcome data.

For sequence generation and allocation concealment, we reported the methods used. For blinding, we described who was blinded and the blinding method. For incomplete outcome data, we reported

Interventions to improve water quality for preventing diarrhoea (Review)

the percentage and proportion of participants lost to follow-up. For selective outcome reporting, any discrepancies between the methods used and the results were stated in terms of the outcomes measured or the outcomes reported. For other biases, we described any other trial features that could have affected the trial result (for example, if the trial was stopped early).

We categorized our 'Risk of bias' judgements as 'low', 'high', or 'unclear'. Where risk of bias was unclear, we attempted to contact the study authors for clarification and we resolved any differences of opinion through discussion. We classified the inclusion of randomized participants in the analysis as 'low risk' if 90% or more of all participants randomized to the study were included in the analysis.

For quasi-RCTs and CBA studies, we used two additional criteria:

1. Comparability of baseline characteristics: we classified studies as 'low risk' if there were no substantial differences between groups with respect to water quality, diarrhoeal morbidity, age, socioeconomic status, access to water, hygiene practices, and sanitation facilities.
2. Contemporaneous data collection: we classified studies as 'low risk' if data were collected at similar points in time, 'unclear' if the relative timing was not reported or not clear from trial, or 'high risk' if data were not collected at similar points in time.

Measures of treatment effect

Two review authors independently extracted and, where necessary, calculated the measure of effect of the intervention on diarrhoea. We extracted the measure of effect as reported by the authors of each study, whether it be risk ratios (RRs), rate ratios, odds ratios (ORs), longitudinal prevalence ratios, or means ratios. In using these various measures of effect, we noted the design effect in treating all such measures of effect as equivalent for common outcomes such as diarrhoea and the debate about methodologies for converting such measures of effect into a single measure (Zhang 1998; McNutt 2003).

For purposes of analysis, we transformed ORs into RRs using the assumed control risk and the formula prescribed in Higgins 2011 (Section 12.5.4.4).

Unit of analysis issues

A number of the included studies had multiple intervention arms (for example, treating water with bleach or with a flocculant and disinfectant) and compared two or more intervention groups against a single control group. In some analyses, we included multiple comparisons from the same study, which double counts the control group participants and yields results in the meta-analysis that are artificially precise. Unfortunately, because of the way data was presented in included studies, it was not possible to correct for this error by dividing the control group participants between multiple groups.

Dealing with missing data

When data was missing or incomplete we attempted to contact the study authors.

Assessment of heterogeneity

We assessed the statistical heterogeneity between trials by visually examining the forest plots for overlapping confidence intervals (CIs), applying the Chi² test with a 10% level of statistical significance, and using the I² statistic with a value of 50% to denote moderate levels of heterogeneity.

Assessment of reporting biases

When there were sufficient studies, we assessed the possibility of publication bias by constructing funnel plots and looking for asymmetry.

Data synthesis

We entered the estimates of effect using the generic inverse variance method on the log scale (Higgins 2006), and analysed the data using Review Manager (RevMan).

We stratified our primary analysis by intervention type, and study design (cluster-RCT, quasi-RCT, or CBA). When appropriate we used meta-analyses to derive pooled estimates of effect using a random-effects model because of the substantial heterogeneity in study settings, interventions, and outcome measures.

We summarized the evidence using 'Summary of findings' tables that we created using the GRADE Guideline Development Tool (GRADEpro GDT). The quality of evidence was rated using the GRADE approach, which consists of five factors that are used to assess the quality of the evidence: study limitations (risk of bias), inconsistency, indirectness, imprecision, and publication bias (Guyatt 2008).

Subgroup analysis and investigation of heterogeneity

We investigated the potential causes of heterogeneity by conducting the following subgroup analyses: age (all ages versus children under five years old); adherence with intervention (< 50%, 50% to 85%, > 85%); water source; water access; water quantity; sanitation conditions; country income level; and length of follow-up.

In the subgroup analyses based on water source, we followed terminology used by the WHO/UNICEF Joint Monitoring Programme (JMP) on Water and Sanitation (WHO/UNICEF 2015), using 'unimproved' to extend to unprotected wells or springs, vendor- or tanker-provided water or bottled water, and 'improved' to extend to household connections, public standpipes, boreholes, protected dug wells or springs, or rainwater collection; we categorized studies as 'unclear' with respect to water supply if they contained insufficient information.

We used the same definitions from the WHO/UNICEF JMP criteria to classify sanitation conditions as 'improved' (connection to a public sewer or septic system, pour-flush latrine, simple pit latrine, ventilated improved pit latrine) or 'unimproved' (service or bucket latrines, public latrines, open latrines); where the necessary information was unclear or unreported, we categorized the sanitation facilities as 'unclear'.

To subgroup studies based on access to water source, we used the classifications defined by the Sphere Project 2011, classifying access as 'sufficient' if a consistently available source was located within 500 m, with queuing no more than 15 minutes and filling

time for a 20 L container no more than three minutes, 'insufficient' if any access failed any such criteria, and 'unclear' if such criteria was unreported or unclear.

The quantity of water available to study participants was considered 'sufficient' if consisting of a minimum of 15 L per person per day. For country income level, we used the World Bank classification of country income levels (high, upper middle, lower middle, low) ([World Bank Country and Lending Groups](#)).

Sensitivity analysis

We conducted a sensitivity analysis to investigate the robustness of the results to each of the 'Risk of bias' components by including only studies that were at low risk of bias. We used this information to guide our judgements on the quality of the evidence.

In addition, we explored the impact of non-blinding of POU interventions using a Bayesian meta-analysis with bias correction. For this purpose, we assumed the true log relative risks from non-blinding studies are subject to a multiplicative bias that results

in the observed relative risks being inflated in magnitude. We assumed the bias is normally distributed with a mean 1.48 or 1.65 and a corresponding standard deviation (SD) of 0.17 or 0.13. These values were derived from the additive bias correction employed in [Wood 2008](#) and [Savović 2012](#). While we believe an attempt to adjust for non-blinding is appropriate, we urge caution in relying on these adjusted estimates since the basis for the adjustment is from clinical (mainly drug) studies that may not be transferable to field studies of environmental interventions and because methodology for the adjustment has not been validated.


RESULTS

Description of studies

Results of the search

The search strategy identified 1088 titles and abstracts, 1076 from the databases and 12 from the other sources ([Figure 1](#)). We screened these titles and abstracts, and obtained the full-text articles of 161 studies for further assessment.

Figure 1. Study flow diagram.



Included studies

Fifty-five studies, including 84,023 participants, met the inclusion criteria (see [Characteristics of included studies](#)). Of these, six studies had two relevant intervention arms (Austin 1993; URL 1995; Luby 2004; Crump 2005; Brown 2008; Lindquist 2014), two had three arms (Luby 2006; Opryszko 2010), and one had four arms (Reller 2003), making a total of 65 discrete comparisons. Three included studies had inadequate information on disease morbidity to include in the quantitative analysis (Torun 1982 GTM; Kremer 2011 KEN; Patel 2012 KEN). We contacted the study authors for further information, but no data could be provided. Therefore we have only described these three studies and their results, but have not integrated these studies into the analysis.

Study design and length

Forty-five studies were cluster-RCTs, two were quasi-RCTs, and eight were CBA studies. Most included cluster-RCTs used households as the unit of randomization, though some used neighbourhoods, villages, or communities. Most CBA studies used villages or communities as the unit of allocation. The intervention period ranged from eight weeks to four years. The duration of the cluster-RCTs (median seven months, range 9.5 weeks to 18 months) tended to be shorter than in the CBA studies (median 12 months, range two to 60 months). Studies of source-based interventions were also longer (median 24 months, range eight weeks to two years) than those of POU interventions (median six months, range 9.5 weeks to 17 months).

Participants and settings

Nine studies included data only for children under five years of age, and three studies included data only on adults. The other studies enrolled and presented results for all ages of participants.

Most studies were undertaken in lower middle or low-income countries based on World Bank criteria, but three studies were conducted in the USA (Colford 2002 USA; Colford 2005 USA; Colford 2009 USA), one in Australia (Rodrigo 2011 AUS), and one in Saudi Arabia (Mahfouz 1995 KSA). Five studies were conducted in urban settings (Semenza 1998 UZB; Colford 2002 USA; Colford 2005 USA; Colford 2009 USA; Rodrigo 2011 AUS), five in peri-urban settings (Quick 1999 BOL; Quick 2002 ZMB; du Preez 2010 ZAF; Jain 2010 GHA; Peletz 2012 ZMB), two in urban informal or squatter settlements (Handzel 1998 BGD; Luby 2004), two in camps for refugees or displaced persons (Roberts 2001 MWI; Doocy 2006 LBR), five in multiple settings (URL 1995; Clasen 2005 COL; Stauber 2009 DOM; du Preez 2011 KEN; Boisson 2013 IND), and the others in villages or other rural settings.

Primary drinking water supply and sanitation facilities

The primary drinking water supply before the intervention was 'unimproved' in 30 studies, 'improved' in 15 studies, and 'unclear' or unreported in five studies. Sanitation facilities in trial settings were 'improved' in 12 studies, 'unimproved' in 15 studies, and 'unclear' or unreported in 19 studies. Access to a water source was deemed 'sufficient' in 14 studies, 'insufficient' in four studies, and 'unclear' or unreported in the remaining studies. The quantity of water available to study participants was considered 'sufficient' in eight studies, 'insufficient' in four studies, and 'unclear' in 43 studies.

Seventeen studies measured water quality before the introduction of the intervention as an indication of the ambient risk and the microbiological quality of the water consumed by the control group. Details on the indicators used varied among the studies (see [Table 1](#)). Thirty-five studies measured colony-forming units (CFUs) of thermotolerant coliforms, faecal coliforms, or *E. coli*, reporting geometric means, arithmetic means, number of CFUs/100 mL, mean faecal coliforms/100 mL, *E. coli* most probable number, median, or log₁₀CFUs/100 mL. Other studies measured the frequency of samples containing such bacteria, or the CFU of total coliforms or other indicators of microbial contamination. None continually measured the microbiological performance of their interventions against the full range of bacterial, viral, and protozoan pathogens known to cause diarrhoea.

Eight studies did not report actually having measured microbiological water quality at all (Alam 1989 BGD; Xiao 1997 CHN; Luby 2006; Mäusezhal 2009 BOL; Opryszko 2010; Majuru 2011 ZAF; Rodrigo 2011 AUS; Lindquist 2014). Thus, it cannot be concluded definitively that the interventions investigated in these studies actually resulted in an improvement in drinking water quality.

Among the eight studies investigating interventions to improve water quality at the point of distribution, only four tested microbiological water quality (Torun 1982 GTM; Gasana 2002 RWA; Jensen 2003 PAK; Kremer 2011 KEN). As these tests were at the source or point of distribution and not the POU, their results do not reflect possible post-collection contamination.

Interventions

Eight studies evaluated source-based interventions: improved wells or boreholes (Alam 1989 BGD; Xiao 1997 CHN; Opryszko 2010b AFG; Opryszko 2010c AFG) or improved community sources and distribution to public tap stands (Torun 1982 GTM; Gasana 2002 RWA; Jensen 2003 PAK; Kremer 2011 KEN; Majuru 2011 ZAF); none evaluated reliable piped-in household connections.

Forty-seven studies evaluated POU interventions: chlorination (17 studies), filtration (20 studies), combined flocculation and disinfection (five studies), SODIS solar disinfection (six studies), combination UV disinfection and filtration (one study), and improved storage (two studies). Significantly, there were no eligible studies that investigated the impact of boiling, even though that is by far the most common type of POU water treatment (Rosa 2010).

Many studies provided a supplementary hygiene education or instruction beyond the use of the intervention itself, and among POU interventions the primary intervention was often combined with some form of improved storage. In only three multiple-intervention arm studies did study authors establish different intervention groups with and without hygiene or other non-water improvement steps in order to isolate the impact of water quality (URL 1995; Opryszko 2010; Lindquist 2014).

Except in blinded trials involving placebos, control arms generally continued to use their pre-trial water supply and treatment practices. In one trial of POU chlorination plus a safe storage container, however, control households also received the container (Jain 2010 GHA). In two of the solar disinfection studies (Conroy 1996 KEN; Conroy 1999 KEN) both intervention and control households received plastic bottles for storing their drinking water.

Interventions to improve water quality for preventing diarrhoea (Review)

The intervention group was instructed to place the bottles on roofs to expose them to the sun, while the control group was told to keep the filled bottles indoors. It is important to note that since improved storage even in the absence of treatment has been shown to improve microbial water quality (Wright 2004), the comparison between the intervention and control in these studies may understate the effectiveness of the intervention when compared to the controls following customary water handling practices.

Adherence with the intervention

Studies of source water interventions tended to assume adherence based on the fact that the primary water supply had been improved. Some studies of POU water treatment undertook indirect assessments of adherence by measuring residual chlorine levels in stored water, comparing microbiological water quality of intervention and control groups, conducting periodic or post-study surveys, or counting the amount of intervention product used. Most other studies measured adherence only by occasional observation, while eight cluster-RCTs did not report on adherence.

The studies of chlorine residuals reported adherence ranging from a high of 95% (Doocy 2006 LBR) to a low of 11% (Opryszko 2010a AFG). Even among these studies, however, investigators acknowledged that it was not possible to know to what extent intervention group participants actually consumed treated water or avoided consuming untreated water. For those studies that reported on adherence, three took the additional step of investigating and reporting on continued consumption of untreated water (Boisson 2010 DRC; Peletz 2012 ZMB; Boisson 2013 IND) a source of exposure that could be masked by less direct metrics of adherence.

Outcome measures

The studies' main outcome measure was diarrhoeal disease, but different methods were used to define, assess, and report this. Thirty-six studies used the WHO's definition of diarrhoea, while other studies used the following definitions: the mother's or respondent's definition (Austin 1993; Gasana 2002 RWA; Reller 2003; Crump 2005; Chiller 2006 GTM); 'watery diarrhoea as a component of gastroenteritis' (Colford 2002 USA; Colford 2005 USA; Colford 2009 USA; Rodrigo 2011 AUS); the local term (Conroy 1996 KEN; Conroy 1999 KEN; Boisson 2009 ETH); "significant change in bowel habits towards decreased consistency or increased frequency" (Kirchhoff 1985 BRA); or dysentery (du Preez 2010 ZAF; du Preez 2011 KEN). Four studies did not report the case definition used for diarrhoea (Torun 1982 GTM; Xiao 1997 CHN; Günther 2013 BEN; Lindquist 2014).

The method of diarrhoea surveillance and assessment also varied. In most cases, participants were visited on a periodic basis, either weekly (19 studies), fortnightly (16 studies), or more infrequently (14 studies). Participants were asked to recall and report on cases of diarrhoea during a previous period, usually seven days (30 studies) or 14 days (six studies), with four studies having recall periods of one to four days and one trial having a recall period of four weeks

(Günther 2013 BEN). Twelve studies asked each participant or a designated household to keep a log or record to indicate days with or without diarrhoea, one procured data on diarrhoea from family records and disease registries (Mahfouz 1995 KSA), or used paediatricians to assess the participants during regular medical checkups (Gasana 2002 RWA). Only one trial did not report the method (Xiao 1997 CHN).

Using these data, study authors reported diarrhoeal disease using one or more of the following epidemiological measures of disease frequency: incidence (34 studies); period prevalence (12 studies); and longitudinal prevalence (nine studies). The studies also reported other measures of disease, including incidence of persistent diarrhoea, gastrointestinal illness, including specific symptoms thereof, incidence or prevalence of bloody diarrhoea, and days of work or school lost due to diarrhoea (Lule 2005 UGA). Seven studies also reported on mortality (Crump 2005; Colford 2009 USA; Boisson 2010 DRC; du Preez 2011 KEN; Kremer 2011 KEN; Peletz 2012 ZMB; Boisson 2013 IND). None reported adverse events.

None of these studies were primarily designed to investigate the impact of the intervention on death, and as such most were underpowered to evaluate this outcome.

Data presentation

Forty-three studies presented results both for children aged under five years (or a subgroup thereof) and for all ages or older age groups, three presented results only for adults, and nine presented results only for children under five years (or a subgroup thereof). Most of the studies adjusted raw data to account for possible covariates, including age, sex, sanitation or hygiene practices, area of residence, household income or proxies thereof, education or maternal literacy, age and occupation of the head of household, number of participants in the household or absent there from, baseline diarrhoea or conditions at baseline, or other variables associated with the household environment and participant behaviour.

Most studies of interventions at the POU also used statistical methods to adjust their results, either for repeated episodes of diarrhoea by the same participant or for clustering within the household, village or both. The studies that did not adjust for clustering may receive excess weight in meta-analysis due to artificial precision (Kirchhoff 1985 BRA; Austin 1993; Mahfouz 1995 KSA; URL 1995).


Excluded studies

We excluded 108 studies for the reasons given in the [Characteristics of excluded studies](#) table. Two studies that appear to meet this review's inclusion criteria are currently ongoing (see [Characteristics of ongoing studies](#)).

Risk of bias in included studies

We have summarized our judgements about the risk of bias of included studies in [Figure 2](#).

Figure 2. Risk of bias graph: summary of authors' judgements about each 'Risk of bias' item presented as percentages across all included studies.



Allocation

The allocation sequence was generated using an adequate method and classified as 'low risk' in 36 of the 45 cluster-RCTs, 'high risk' in two, and 'unclear' in seven [Figure 2](#). The method of allocation concealment was 'low risk' in 34 trials and 'high risk' in two and 'unclear' in nine.

Comparability of baseline characteristics (confounding bias)

All the quasi-RCTs and CBA studies were judged to be at low risk of bias for this criteria except [Gasana 2002 RWA](#), which was at 'unclear' risk.

Contemporaneous data collection

We judged all the quasi-RCTs and CBA studies to be at low risk of bias for this criteria except [Gasana 2002 RWA](#), which was at 'unclear' risk.

Blinding

Nine trials were blinded at the participant level ([Kirchhoff 1985 BRA](#); [Austin 1993](#); [Colford 2002 USA](#); [Colford 2005 USA](#); [Colford 2009 USA](#); [Boisson 2010 DRC](#); [Jain 2010 GHA](#); [Rodrigo 2011 AUS](#); [Boisson 2013 IND](#)); all but two of these were blinded at the assessor level as well ([Kirchhoff 1985 BRA](#); [Austin 1993](#)). The others followed an open design, classified as 'high risk' of bias. One of the principal objectives of [Colford 2002 USA](#) was to assess the effectiveness of its blinding methodology; it therefore provides the most comprehensive analysis of these issues. [Colford 2002 USA](#), [Colford 2005 USA](#), [Boisson 2010 DRC](#) and [Rodrigo 2011 AUS](#) all used household sham water filters. [Austin 1993](#), [Kirchhoff 1985 BRA](#), [Jain 2010 GHA](#) and [Boisson 2013 IND](#), which were assessing the effectiveness of home-based chlorination, provided placebos to control households.

Incomplete outcome data

Twenty four studies were at 'low risk' of bias, 18 at 'high risk', and three studies were unclear.

Effects of interventions

See: [Summary of findings for the main comparison Summary of findings table 1](#)

Analysis 1: Any water quality intervention versus no intervention

Diarrhoea episodes

An overall pooled analysis, across different trial designs, interventions and settings, finds the risk of diarrhoea to be lower with any water quality intervention compared to no intervention, both among all ages (RR 0.59, 95% CI 0.51 to 0.69, 81215 participants; 52 studies [Analysis 1.1](#)), and under fives (RR 0.61, 95% CI 0.49 to 0.75 [Analysis 1.2](#)). However, as would be expected given the diverse nature of the trials, statistical heterogeneity between trials is very high (I^2 statistic = 98% and 97%, respectively). Our primary analysis is therefore stratified by the specific intervention type (for example, interventions at water source, POU chlorination, POU filtration), and by study design (for example, cluster-RCT, quasi-RCT, CBAs).

Mortality

Only nine studies reported any deaths among study participants. Five reported the number of deaths in each study arm without differences evident (see [Table 2](#)). Two studies reported the total number of deaths without stating how many occurred in each group ([du Preez 2010 ZAF](#); [Boisson 2013 IND](#)), and two reported recording deaths but the numbers were not presented in the papers ([Boisson 2009 ETH](#); [Kremer 2011 KEN](#)).

None of these studies were primarily designed to investigate the impact of the intervention on mortality, and all were underpowered to investigate these effects.

Adverse events

No trial reported adverse events from the interventions.


Analysis 2: Interventions at the water source

One cluster-RCT and five CBA studies evaluated interventions at the water source (Table 3). All but one study were from settings with 'unimproved' water sources (unprotected wells or surface water), and all had unclear levels of sanitation. Three studies evaluated improved wells or boreholes, two evaluated chlorination or filtration of community water sources, and one evaluated an improved community piped supply. No studies evaluated reliable household connections to a clean water source (see Table 4 and Table 5 for a description of study settings and interventions).

The single cluster-RCT from Afghanistan reported no statistically significant difference in diarrhoea with improved wells compared to no intervention (one trial, 3266 participants; Analysis 2.1; very low quality evidence).

The CBA studies evaluated different interventions, had variable findings, and were all at unclear risk of multiple sources of bias (see Figure 3). Three of the five studies reported statistically significant effects on diarrhoea (Analysis 2.1; Analysis 2.2): in Bangladesh, provision of one hand pump per four to six households (three times as many as control areas) was associated with a small reduction in diarrhoea over three-years follow-up (RR 0.83, 95% CI 0.71 to 0.97); in remote areas in South Africa a new community piped water supply was associated with around a 50% reduction in diarrhoea compared to untreated river water (RR 0.43, 95% CI 0.24 to 0.77); and in China structural well improvements were also associated with around a 50% reduction in diarrhoea (RR 0.45, 95% CI 0.43 to 0.47). In contrast, chlorination and filtration of community water supplies were not associated with positive benefits in Rwanda and Pakistan respectively. Overall, the body of evidence is judged to be of very low quality (Table 3). Given the variability in interventions, further subgroup analyses to try to understand the heterogeneity were not useful.

Figure 3. Forest plot of comparison: 2 Source: water supply improvement versus control, outcome: 2.1 Diarrhoea: CBA studies subgrouped by age.



Footnotes

(1) Opryszko 2010-ii AFG: Provided one well per 25 households providing 25 litres/person/day


Analysis 3. POU chlorination

Fourteen cluster-RCTs, with 16 comparisons, evaluated POU chlorination versus control. Chlorine was delivered to households free of charge every one to four weeks, with instructions on how to use it, and in eight trials a water storage container was also provided (see Table 6 and Table 7 for a description of study settings and interventions).

On average, POU chlorination in cluster RCTs reduced the risk of diarrhoea episodes by around a quarter, both for all ages (RR 0.77, 95% CI 0.65 to 0.91; 14 trials, 30,746 participants; Analysis 3.2) and for children under five years of age (RR 0.77, 95% CI 0.64 to 0.92; Analysis 3.2). However, there was substantial heterogeneity in the size of the effect which was not well explained by a series of subgroup analyses (Analysis 3.2 to Analysis 3.9).

As might be expected from an effective intervention, the trials finding larger effects from chlorination tended to be those where adherence with the intervention was higher (as measured by residual chlorine) (Analysis 3.3; Figure 4), but in the four trials which had adequate blinding no effects of water chlorination were seen (Analysis 3.4). A subgroup analysis looking at interventions with and without the provision of water storage containers did not find statistical evidence of subgroup differences (Analysis 3.5). Effects were seen in trials with 3, 6, and 12 months of follow-up, but no effect was demonstrated in the two trials with follow-up longer than 12 months (Analysis 3.9). The funnel plot for this comparison has some asymmetry which may be the result of publication bias (see Figure 5). The overall quality of the evidence was therefore judged to be low (Table 8).


Figure 4. Forest plot of comparison: 3 POU: water chlorination versus control, outcome: 3.3 Diarrhoea: cluster-RCTs; subgrouped by adherence.



Footnotes

- (1) Handzel 1998 BGD: Free chlorine was measureable in 77% of samples - Unclear whether testing was during unannounced visits
- (2) Crump 2005-i KEN: Free chlorine residuals > 0.1 mg/L in 85% of samples during scheduled visits and 61% of samples during unannounced visits
- (3) Jain 2010 GHA: Free chlorine residuals > 0.2 mg/L in 74-89% of samples - Unclear whether testing was during unannounced visits
- (4) Mengistie 2013 ETH: Free chlorine residuals > 0.2 mg/L in 76-77% of samples - Testing was during unannounced visits
- (5) Quick 1999 BOL: The proportion of stored water samples with detectable levels of total chlorine increased from 71 % at the time of the first observation to...
- (6) Semenza 1998 UZB: Chlorine was detected in 73% of household samples at the end of the study.
- (7) Boisson 2013 IND: Free chlorine was measureable in 32% of samples - Unclear whether testing was during unannounced visits
- (8) Reller 2003-ii GUA: Participants had free chlorine >0.1 mg/mL in 36% of samples - Testing during unannounced visits
- (9) Reller 2003-iii GUA: Participants had free chlorine >0.1 mg/mL in 44% of samples - Testing during unannounced visit
- (10) Kirchhoff 1985 BRA: The chlorination was performed daily by blinded health staff.
- (11) Lule 2005 UGA: Compliance not reported
- (12) Mahfouz 1995 KSA: The average free residual chlorine is reported as 0.13 ppm
- (13) Opryszko 2010-iii AFG: Self reported use of Chlorine in the previous two weeks was 82%

Figure 5. Funnel plot of comparison: 3 POU: water chlorination versus control, outcome: 3.1 Diarrhoea: subgrouped by study design.



An additional two CBA studies evaluated POU chlorination but only provide very low quality evidence of any effect (Analysis 3.1, Table 8).

Analysis 4. POU combined flocculation and disinfection

Five cluster-RCTs from low-income settings evaluated interventions where sachets of flocculant and disinfectant were distributed to households to treat water from unimproved sources (three trials), improved sources (one trial), and unclear sources (one trial). Four trials also provided water containers and mixing equipment (see Table 9 and Table 10 for a description of study settings and interventions). None of the trials blinded the outcome assessment.

Four of the five trials found statistically significant reductions in diarrhoea with the intervention (Table 11), but statistical heterogeneity in the size of this effect made pooling the data difficult (I^2 statistic = 99%; Analysis 4.1). This heterogeneity relates to one trial from Liberia IDP camps, Doocy 2006 LBR, where the flocculation and disinfection kits reduced diarrhoea by 88% (RR 0.12, 95% CI 0.11 to 0.13; one trial, 2191 participants). Exclusion of this potential outlier finds a more modest effect with the other four trials both for all ages (RR 0.69, 95% CI 0.58 to 0.82; four trials, 11788 participants; Analysis 4.2) and for children under five years of age (RR 0.71, 95% CI 0.61 to 0.84; Analysis 4.2).


Adherence with the intervention, as measured by residual chlorine, was generally low (< 50%), but higher in the trial from Liberia showing large effects (Analysis 4.3). Larger effects tended to also be seen in the trials also providing water storage containers (Analysis 4.4). The effects were present in trials with both improved and unimproved water source and sanitation (Analysis 4.5; Analysis 4.6; Analysis 4.7). None of the trials had follow-up longer than 12 months (Analysis 4.8).

Analysis 5. POU filtration

Overall 20 cluster-RCTs evaluated POU filtration: ceramic filtration (nine trials), biosand filtration (five trials), LifeStraw® filters (three trials), and plumbed-in filtration (three trials) (see Table 12 and Table 13 for a description of study settings and interventions).

On average, POU filtration technologies reduced diarrhoea by around a half, both for all ages (RR 0.48, 95% CI 0.38 to 0.59; 18 trials, 15,582 participants; Analysis 5.1) and for children under five years of age (RR 0.49, 95% CI 0.38 to 0.62; Analysis 5.1). However, the number of trials and the quality of evidence was different for each specific intervention (Analysis 5.2; Figure 6). The lack of blinding in these studies is a major concern: of the five trials with adequate blinding only one found a statistically significant effect (Analysis 5.3). The quality of evidence was therefore downgraded for all types of filters due to risk of bias (Table 14).

Figure 6. Forest plot of comparison: 4 POU: filtration versus control, outcome: 4.2 Diarrhoea: cluster-RCTs: subgrouped by type of filtration.



POU ceramic filters reduced diarrhoea by around 60% in nine trials mainly from low- or middle-income countries, regardless of whether the water source or sanitation was classified as improved or unimproved (RR 0.39, 95% CI 0.29 to 0.53, eight trials, 5763 participants; [Analysis 5.3](#); [Analysis 5.4](#); *moderate quality evidence*).

Similarly, biosand filtration reduced diarrhoea by around a half consistently across five trials from low- or middle-income settings, again regardless of whether the water source or sanitation was improved or unimproved (RR 0.47, 95% CI 0.39 to 0.57, four trials, 5504 participants; [Analysis 5.6](#); [Analysis 5.7](#); *moderate quality evidence*).

On average, the use of LifeStraw® filters reduced diarrhoea by around a third in three trials from low-income settings with unimproved water sources (RR 0.69, 95% CI 0.51 to 0.93; three trials, 3259 participants; [Analysis 5.2](#); *low quality evidence*).

Plumbed-in filtration has only been evaluated in high-income settings (USA). There is a modest effect in all three trials, although

only one reaches standard levels of statistical significance. The overall meta-analysis has similar effect sizes with both fixed effects and random effects models, but wider confidence intervals with random effects (Fixed-effects: RR 0.81, 95% CI 0.70 to 0.94; Random-effects: RR 0.73, 95% CI 0.52 to 1.03; three trials, 1056 participants; [Analysis 5.2](#); *moderate quality evidence*).

Adherence with the filtration systems was reported by 14 trials, of which eight assessed this by self-reported use which is at high risk of bias due to the lack of blinding. Adherence was generally reported as high, and larger effects were apparent in trials with higher adherence ([Analysis 5.8](#)). A subgroup analysis looking at filtration interventions with and without the provision of water storage containers (excluding the trials evaluating plumbed in filtration), found larger effects in the nine trials providing containers ([Analysis 5.9](#)). Effects were seen in trials with 3, 6, and 12 months of follow-up, but no effect was demonstrated in the one trial with follow-up longer than 12 months ([Analysis 5.10](#)).

Interventions to improve water quality for preventing diarrhoea (Review)

Analysis 6. POU solar disinfection (SODIS)

Four cluster-RCTs and two quasi-RCTs evaluated solar disinfection of water from improved sources (one study) and unimproved sources (five studies) in low-income settings. Plastic bottles were distributed to households with instructions to leave filled bottles in direct sunlight for at least six hours before drinking (see [Table 15](#) and [Table 16](#) for a description of study settings and interventions).

Overall in the cluster-RCTs, solar disinfection reduced diarrhoea by around a third for all ages (RR 0.62, 95% CI 0.42 to 0.94; four trials, 3460 participants; [Analysis 6.1](#)), and almost a half in children under five years of age (RR 0.55, 95% CI 0.34 to 0.91; [Analysis 6.2](#)). The largest effect was seen in the trial with the highest adherence ([Analysis 6.3](#)). The quality of evidence was downgraded to moderate due to the lack of blinding and the inherent risk of bias ([Table 17](#)).

In the quasi-RCTs the observed effect was lower (RR 0.82, 95% CI 0.69 to 0.97; two trials, 555 participants; [Analysis 6.1](#)).

Analysis 7. POU UV disinfection

One cluster-RCT from Mexico evaluated an UV tube disinfection technology ([Gruber 2013 MEX](#); see [Table 18](#) and [Table 19](#) for a description of the study setting and intervention).

The effect on diarrhoea among all age populations did not reach standard levels of statistical significance (RR 0.79, 95% CI 0.49 to 1.27; one trial, 1913 participants; [Analysis 7.1](#)), and did not report separately for children under five years of age.

Analysis 8. POU improved storage

Two trials from Malawi and Benin evaluated the distribution of improved water storage containers in settings with improved water sources (see [Table 20](#) and [Table 21](#) for a description of the study setting and intervention).

Overall, there was no statistically significant effect on diarrhoea for all ages (RR 0.91, 95% CI 0.74 to 1.11; two trials, 1871 participants; [Analysis 8.1](#)), or children under five years of age (RR 0.69, 95% CI 0.47 to 1.01; [Analysis 8.1](#)). Both studies were at high risk of bias due to being non-blinded, and the overall quality of the evidence was judged to be low ([Table 22](#)).

Analyses adjusted for non-blinding

In [Table 23](#) we have presented meta-analysis results adjusted for non-blinding using an approach described in the [Methods](#) section and based in part on those employed by other researchers ([Hunter 2009](#); [Wolf 2014](#)). In these analyses, the effects of POU chlorination and filtration are smaller but remain statistically significant; the effect of POU solar disinfection becomes borderline non-significant.

DISCUSSION

Summary of main results

There is currently insufficient evidence to know if source-based improvements such as protected wells, communal tap stands, or chlorination/filtration of community sources consistently reduce diarrhoea (*very low quality evidence*).

The distribution and promotion of point-of-use water chlorination products may reduce diarrhoea by around one quarter (*low quality evidence*). Similarly, distribution and promotion of flocculation and disinfection sachets probably reduces diarrhoea but had highly variable effects (*moderate quality evidence*).

Point-of-use filtration systems probably reduce diarrhoea by around a half (*moderate quality evidence*). This reduction was apparent for ceramic filters, biosand systems and LifeStraw® filters, but plumbed in filtration has only been evaluated in high-income settings and a statistically significant effect has not been demonstrated.

In low-income settings, distribution of plastic bottles with instructions to leave filled bottles in direct sunlight for at least six hours before drinking (SODIS) probably reduces diarrhoea by around a third (*moderate quality evidence*).

In subgroup analyses, larger effects were seen in trials with higher adherence, and trials that provided a safe storage container.

Overall completeness and applicability of evidence

Fifty-five studies met the inclusion criteria, of which most studies were conducted in low- or middle-income countries (50 studies), with unimproved water sources (30 studies), and unimproved or unclear sanitation (34 studies).

For water source interventions, there are simply too few studies to make conclusions about what may or may not be effective in different settings. While protective effects were seen in some individual trials, it is unclear whether these effects could be expected to be reproducible in other settings, and all of the trials had multiple potential sources of bias. Significantly, we found no studies evaluating reliable, piped-in water supplies.

In contrast, some POU interventions do appear to be broadly protective against diarrhoea across many settings regardless of whether water sources and sanitation are 'improved' or 'unimproved'. This finding affirms the current strategy of the WHO and UNICEF to promote POU water treatment and safe storage, even though this will not increase the number of households with access to improved water supplies and therefore will not contribute towards achieving current international water targets ([WHO 2011](#)). The effectiveness of POU interventions in settings without improved sanitation contradicts earlier findings that interventions to improve water quality are effective only where sanitation has already been addressed ([Esrey 1986](#); [VanDerSluis 1995](#)), or that environmental interventions to prevent diarrhoea are effective only by employing an integrated approach ([Eisenberg 2007](#)).

Although we provide average estimates of effect for each individual POU intervention, we recommend caution in using these estimates to conclude the superiority of one intervention over another. Such an observational analysis would be highly susceptible to confounding by study setting and population, and may not represent true differences in the size of the effects. Head-to-head trials would be necessary to reliably conclude superiority and these were not the focus of this review.

As few studies continued follow-up beyond 12 months, we are unable to comment reliably on the long-term sustainability of these effects. While pooled estimates of studies with follow-up periods

under 12 months were generally protective, those with follow-up periods in excess of 12 months were not.

Quality of the evidence

The quality of evidence for the effects of the individual interventions on diarrhoea ranged from moderate (for ceramic filters and biosand filtration), to low (for distribution of chlorination kits, flocculation and disinfection sachets, and LifeStraw® filters), to very low (for water source improvements).

The primary reason for downgrading the quality of evidence was the risk of bias inherent in unblinded studies evaluating the efficacy of an intervention on a self-reported outcome. Notably, only one of the nine blinded trials reported a statistically significant protective effect, but this observation may be explained by other confounding factors present in these nine trials (see [Table 24](#)):

1. Four studies were conducted in high-income countries where the water was of good microbiological quality even in the control groups ([Colford 2002 USA](#); [Colford 2005 USA](#); [Colford 2009 USA](#); [Rodrigo 2011 AUS](#)).
2. One further trial from Ghana found very low levels of faecal contamination of water supplies in the control group which were likely to present only minimal risk ([Jain 2010 GHA](#)).
3. Three studies had either low adherence with the intervention ([Austin 1993](#); [Boisson 2013 IND](#)), or very high reported use of drinking untreated water from other sources ([Boisson 2010 DRC](#)).
4. Two studies employed control interventions which may have improved water quality: [Boisson 2010 DRC](#) employed a "placebo" that actually removed one log (90%) of faecal indicator bacteria and [Jain 2010 GHA](#) provided control households with safe storage.

The second common reason for downgrading the quality of evidence was unexplained heterogeneity. For some of the POU interventions, the protective effect varied considerably across studies. Some of this variability could be explained by adherence with the intervention, with larger effects in studies with higher adherence, but some variability remained which we were unable to explain despite multiple subgroup analyses. This is likely to reflect important underlying clinical heterogeneity: the aetiology and epidemiology of diarrhoea is complex and variable, transmission pathways are multiple, and even the portion of diarrhoea that is waterborne is not well understood ([Eisenberg 2012](#)).

There was also some evidence of possible publication bias in the trials evaluating home chlorination but this was not strong enough to further downgrade the quality of evidence.

Potential biases in the review process

A number of the included studies had multiple intervention arms comparing two or more intervention groups against a single control group. In some analyses, we included multiple comparisons from the same trial which double counts the control group participants and yields results in the meta-analysis that are artificially precise. However, this bias is unlikely to have significantly impacted the overall quality of evidence or conclusions.

Agreements and disagreements with other studies or reviews

Our results are generally consistent with the prior version of this Cochrane Review ([Clasen 2006](#)) and with other reviews of water quality interventions ([Fewtrell 2005](#); [Arnold 2007](#); [Waddington 2009](#); [Cairncross 2010](#); [Wolf 2014](#)).

One additional review of water quality interventions reports no effect with POU interventions once blinding is taken into account ([Engell 2013](#)). While we share the concerns about the lack of blinding in many of these trials (and have downgraded the quality of evidence accordingly), and also found no effect in any of the trials with adequate blinding, we have identified several possible confounders in this observation (discussed above), and retain low to moderate confidence that these interventions are effective.

Although we found no controlled trials evaluating piped-in water supplies, a recent review that also included some observational studies reported some evidence of a protective effect with this intervention ([Wolf 2014](#)).

The finding of larger effects with increased adherence is consistent with modelling data based on quantitative microbial risk assessment which suggest a dose-response association between water quality and diarrhoea ([Brown 2012](#); [Enger 2013](#)).

AUTHORS' CONCLUSIONS

Implications for practice

Interventions that address the microbial contamination of water at the POU are important interim measures to improve drinking water quality until homes can be reached with safe, reliable, household piped-water connections.

Implications for research

Rigorously conducted RCTs that compare various approaches to improving drinking water quality will help clarify the potential for water quality interventions to prevent endemic diarrhoea. It is particularly important that such trials be designed to minimize reporting bias, such as through the use of objective outcomes.

Among source-based interventions, there is a need for studies to assess household connections and other approaches (such as chlorination at the point of delivery) that are more likely to ensure safe drinking water from source through to the POU.

There is also a need for longer-term studies in programmatic settings on approaches to optimise the coverage and long-term utilization of these interventions among vulnerable populations.

ACKNOWLEDGEMENTS

We gratefully acknowledge co-authors of the previous version of this Cochrane Review: Ian Roberts, Tamer Rabie, and Wolf-Peter Schmidt ([Clasen 2006](#)). We are also grateful to the following people for their research, advice, assistance, and other valuable contributions: Greg Allgood, Jamie Bartram, Joseph Brown, Jack Colford, John Crump, Tom Chiller, Val Curtis, Shannon Doocy, Lorna Fewtrell, Carrol Gamble, Bruce Gordon, Stephen Gundry, Paul Hunter, Bruce Keswick, Steve Luby, Rob Quick, Mark Sobsey, Sara Thomas, and James Wright. We also appreciate the assistance

and help provided by CIDG members and by the referees of both this review and the protocol.

The editorial base for the Cochrane Infectious Diseases Group is funded by the UK Department for International Development (DFID) for the benefit of developing countries.

REFERENCES

References to studies included in this review

Abebe 2014 ZAF {published data only}

Abebe LS, Smith JA, Narkiewicz S, Oyanedel-Craver V, Conaway M, Singo A, et al. Ceramic water filters impregnated with silver nanoparticles as a point-of-use water-treatment intervention for HIV-positive individuals in Limpopo Province, South Africa: a pilot study of technological performance and human health benefits. *Journal of Water and Health* 2014;**12**(2):288-300.

Alam 1989 BGD {published data only}

Alam N, Wojtyniak B, Henry FJ, Rahaman MM. Mothers' personal and domestic hygiene and diarrhoea incidence in young children in rural Bangladesh. *International Journal of Epidemiology* 1989;**18**(1):242-7.

Austin 1993a GMB {unpublished data only}

Austin CJ. Investigation of in-house water chlorination and its effectiveness for rural areas of the Gambia [dissertation]. New Orleans: Tulane University School of Public Health and Tropical Medicine, 1993.

Austin 1993b GMB {unpublished data only}

Austin CJ. Investigation of in-house water chlorination and its effectiveness for rural areas of the Gambia [dissertation]. New Orleans: Tulane University School of Public Health and Tropical Medicine, 1993.

Boisson 2009 ETH {published data only}

Boisson S, Schmidt WP, Berhanu T, Gezahegn H, Clasen T. Randomized controlled trial in rural Ethiopia to assess a portable water treatment device. *Environmental Science & Technology* 2009;**43**(15):5934-9.

Boisson 2010 DRC {published data only}

Boisson S, Kiyombo M, Sthreshley L, Tumba S, Makambo J, Clasen T. Field assessment of a novel household-based water filtration device: a randomised, placebo-controlled trial in the Democratic Republic of Congo. *PLoS One* 2010;**5**(9):e12613.

Boisson 2013 IND {published data only}

Boisson S, Stevenson M, Shapiro L, Kumar V, Singh LP, Ward D, et al. Effect of household-based drinking water chlorination on diarrhoea among children under five in Orissa, India: a double-blind randomised placebo-controlled trial. *PLoS Medicine* 2013;**10**(8):e1001497.

Brown 2008a KHM {published data only}

Brown J, Sobsey MD, Loomis D. Local drinking water filters reduce diarrheal disease in Cambodia: a randomized, controlled trial of the ceramic water purifier. *American Journal of Tropical Medicine and Hygiene* 2008;**79**(3):394-400.

Brown 2008b KHM {published data only}

Brown J, Sobsey MD, Loomis D. Local drinking water filters reduce diarrheal disease in Cambodia: a randomized, controlled trial of the ceramic water purifier. *American Journal of Tropical Medicine and Hygiene* 2008;**79**(3):394-400.

Chiller 2006 GTM {published data only}

Chiller TM, Mendoza CE, Lopez MB, Alvarez M, Hoekstra RM, Keswick BH, et al. Reducing diarrhoea in Guatemalan children: randomized controlled trial of flocculant-disinfectant for drinking-water. *Bulletin of the World Health Organization* 2006;**84**(1):28-35.

Clasen 2004b BOL {published and unpublished data}

Clasen TF, Brown J, Collin SM. Preventing diarrhoea with household ceramic water filters: assessment of a pilot project in Bolivia. *International Journal of Environmental Health Research* 2006;**16**(3):231-9.

Clasen 2004c BOL {published data only}

Clasen TF, Brown J, Collin S, Suntura O, Cairncross S. Reducing diarrhea through the use of household-based ceramic water filters: a randomized, controlled trial in rural Bolivia. *American Journal of Tropical Medicine and Hygiene* 2004;**70**(6):651-7.

Clasen 2005 COL {published data only}

Clasen T, Parra GG, Boisson S, Collin S. Household-based ceramic water filters for the prevention of diarrhea: a randomized, controlled trial of a pilot program in Colombia. *American Society of Tropical Medicine and Hygiene* 2005;**73**(4):790-5.

Colford 2002 USA {published data only}

Colford JM Jr, Rees JR, Wade TJ, Khalakdina A, Hilton JF, Ergas IJ, et al. Participant blinding and gastrointestinal illness in a randomized, controlled trial of an in-home drinking water intervention. *Emerging Infectious Diseases* 2002;**8**(1):29-36.

Colford 2005 USA {published data only}

Colford JM Jr, Saha SR, Wade TJ, Wright CC, Vu M, Charles S, et al. A pilot randomized, controlled trial of an in-home drinking water intervention among HIV+ persons. *Journal of Water and Health* 2005;**3**(2):173-84.

Colford 2009 USA {published data only}

Colford JM Jr, Hilton JF, Wright CC, Arnold BF, Saha S, Wade TJ, et al. The Sonoma water evaluation trial: a randomized drinking water intervention trial to reduce gastrointestinal illness in older adults. *American Journal of Public Health* 2009;**99**(11):1988-95.

Conroy 1996 KEN {published data only}

Conroy RM, Elmore-Meegan M, Joyce T, McGuigan KG, Barnes J. Solar disinfection of drinking water and diarrhoea in Maasai children: a controlled field trial. *Lancet* 1996;**348**(9043):1695-7.

Conroy 1999 KEN {published data only}

Conroy RM, Meegan ME, Joyce T, McGuigan K, Barnes J. Solar disinfection of water reduces diarrhoeal disease: an update. *Archives of Disease in Childhood* 1999;**81**(4):337-8.

Crump 2005a KEN {published data only}

Crump JA, Otieno PO, Slutsker L, Keswick BH, Rosen DH, Hoekstra RM, et al. Household based treatment of drinking water with flocculant-disinfectant for preventing diarrhoea in

areas with turbid source water in rural western Kenya: cluster randomised controlled trial. *BMJ* 2005;**331**(7515):478.

Crump 2005b KEN {unpublished data only}

Crump JA, Otieno PO, Slutsker L, Keswick BH, Rosen DH, Hoekstra RM, et al. Otieno PO, Slutsker L, Keswick BH, Rosen DH, Hoekstra RM, et al. Household based treatment of drinking water with flocculant-disinfectant for preventing diarrhoea in areas with turbid source water in rural western Kenya: cluster randomised controlled trial. *BMJ* 2005;**331**(7515):478.

Doocy 2006 LBR {published data only}

Doocy S, Burnham G. Point-of-use water treatment and diarrhoea reduction in the emergency context: an effectiveness trial in Liberia. *Tropical Medicine & International Health* 2006;**11**(10):1542-52.

du Preez 2008 ZAF/ZWE {published and unpublished data}

du Preez M, Conroy RM, Wright JA, Moyo S, Potgieter N, Gundry SW. Use of ceramic water filtration in the prevention of diarrheal disease: a randomized controlled trial in rural South Africa and Zimbabwe. *American Journal of Tropical Medicine and Hygiene* 2008;**79**(5):696-701.

du Preez 2010 ZAF {published data only}

du Preez M, Mcguigan KG, Conroy RM. Solar disinfection of drinking water In the prevention of dysentery in South African children aged under 5 years: the role of participant motivation. *Environmental Science & Technology* 2010;**44**(22):8744-9.

du Preez 2011 KEN {published data only}

du Preez M, Conroy RM, Ligondo S, Hennessy J, Elmore-Meegan M, Soita A, et al. Randomized intervention study of solar disinfection of drinking water in the prevention of dysentery in Kenyan children aged under 5 years. *Environmental Science & Technology* 2011;**45**(21):9315-23.

Fabiszewski 2012 HND {published data only}

Fabiszewski de Aceituno AM, Stauber CE, Walters AR, Meza Sanchez RE, Sobsey MD. A randomized controlled trial of the plastic-housing BioSand filter and its impact on diarrheal disease in Copan, Honduras. *American Journal of Tropical Medicine and Hygiene* 2012;**86**(6):913-21.

Gasana 2002 RWA {published data only}

Gasana J, Morin J, Ndikuyeze A, Kamoso P. Impact of water supply and sanitation on diarrheal morbidity among young children in the socioeconomic and cultural context of Rwanda (Africa). *Environmental Research* 2002;**90**(2):76-88.

Gruber 2013 MEX {published data only}

Gruber JS, Reygadas F, Arnold BF, Ray I, Nelson K, Colford JM Jr. A stepped wedge, cluster-randomized trial of a household UV-disinfection and safe storage drinking water intervention in rural Baja California Sur, Mexico. *American Journal of Tropical Medicine and Hygiene* 2013;**89**(2):238-45.

Günther 2013 BEN {published data only}

Günther I, Schipper Y. Pumps, germs and storage: the impact of improved water containers on water quality and health. *Health Economics* 2013;**22**(7):757-74.

Handzel 1998 BGD {published and unpublished data}

* Handzel T. The effect of improved drinking water quality on the risk of diarrhoeal disease in an urban slum of Dhakar, Bangladesh: a home chlorination intervention trial [dissertation]. Chapel Hill (NC): University of North Carolina at Chapel Hill, 1998.

Sobsey MD, Handzel T, Venczel L. Chlorination and safe storage of household drinking water in developing countries to reduce waterborne disease. *Water Science and Technology* 2003;**47**(3):221-8.

Jain 2010 GHA {published data only}

Jain S, Sahanoon OK, Blanton E, Schmitz A, Wannemuehler KA, Hoekstra RM, et al. Sodium dichloroisocyanurate tablets for routine treatment of household drinking water in periurban Ghana: a randomized controlled trial. *American Journal of Tropical Medicine and Hygiene* 2010;**82**(1):16-22.

Jensen 2003 PAK {published data only}

Jensen PK, Ensink JH, Jayasinghe G, van der Hoek W, Cairncross S, Dalsgaard A. Effect of chlorination of drinking-water on water quality and childhood diarrhoea in a village in Pakistan. *Journal of Health, Population, and Nutrition* 2003;**21**(1):26-31.

Kirchhoff 1985 BRA {published data only}

Kirchhoff LV, McClelland KE, Do Carmo Pinho M, Araujo JG, De Sousa MA, Guerrant RL. Feasibility and efficacy of in-home water chlorination in rural North-eastern Brazil. *Journal of Hygiene* 1985;**94**(2):173-80.

Kremer 2011 KEN {published and unpublished data}

Kremer M, Leino J, Miguel E, Zwane AP. Spring cleaning: rural water impacts, valuation, and property rights institutions. *Quarterly Journal of Economics* 2011;**126**(1):145-205.

Lindquist 2014a BOL {published data only}

Lindquist ED, George CM, Perin J, Neiswender de Calani KJ, Norman WR, Davis TP Jr, et al. A cluster randomized controlled trial to reduce childhood diarrhea using hollow fiber water filter and/or hygiene-sanitation educational interventions. *American Journal of Tropical Medicine and Hygiene* 2014;**91**(1):190-7.

Lindquist 2014b BOL {published data only}

Lindquist ED, George CM, Perin J, Neiswender de Calani KJ, Norman WR, Davis TP Jr, et al. A cluster randomized controlled trial to reduce childhood diarrhea using hollow fiber water filter and/or hygiene-sanitation educational interventions. *American Journal of Tropical Medicine and Hygiene* 2014;**91**(1):190-7.

Luby 2004a PAK {published data only}

Luby SP, Agboatwalla M, Hoekstra RM, Rahbar MH, Billhimer W, Keswick BH. Delayed effectiveness of home-based interventions in reducing childhood diarrhea, Karachi, Pakistan. *American Journal of Tropical Medicine and Hygiene* 2004;**71**(4):420-7.

Luby 2004b PAK {published data only}

Luby SP, Agboatwalla M, Hoekstra RM, Rahbar MH, Billhimer W, Keswick BH. Delayed effectiveness of home-based interventions

in reducing childhood diarrhea, Karachi, Pakistan. *American Journal of Tropical Medicine and Hygiene* 2004;**71**(4):420-7.

Luby 2006a PAK {published and unpublished data}

Luby SP, Agboatwalla M, Painter J, Altaf A, Billhimer W, Keswick B, et al. Combining drinking water treatment and hand washing for diarrhoea prevention, a cluster randomised controlled trial. *Tropical Medicine & International Health* 2006;**11**(4):479-89.

Luby 2006b PAK {unpublished data only}

Luby SP, Agboatwalla M, Painter J, Keswick B, Billhimer W, Altaf A, et al. Effect of a novel water treatment and handwashing promotion on childhood diarrhea in Pakistan: a randomized controlled trial. *Tropical Medicine & International Health* 2006;**11**(4):479-89.

Luby 2006c PAK {unpublished data only}

Luby SP, Agboatwalla M, Painter J, Keswick B, Billhimer W, Altaf A, et al. Effect of a novel water treatment and handwashing promotion on childhood diarrhea in Pakistan: a randomized controlled trial. *Tropical Medicine & International Health* 2006;**11**(4):479-89.

Lule 2005 UGA {published data only}

Lule JR, Mermin J, Ekwaru JP, Malamba S, Downing R, Ransom R, et al. Effect of home-based water chlorination and safe storage on diarrhea among persons with human immunodeficiency virus in Uganda. *American Journal of Tropical Medicine and Hygiene* 2005;**73**(5):926-33.

Mahfouz 1995 KSA {published data only}

Mahfouz AA, Abdel-Moneim M, al-Erian RA, al-Amari OM. Impact of chlorination of water in domestic storage tanks on childhood diarrhoea: a community trial in the rural areas of Saudi Arabia. *American Journal of Tropical Medicine and Hygiene* 1995;**98**(2):126-30.

Majuru 2011 ZAF {published data only}

Majuru B, Michael Mokoena M, Jagals P, Hunter PR. Health impact of small-community water supply reliability. *International Journal of Hygiene and Environmental Health* 2011;**214**(2):162-6.

Mäusezhal 2009 BOL {published data only}

Mäusezahl D, Christen A, Pacheco GD, Tellez FA, Iriarte M, Zapata ME, et al. Solar Drinking Water Disinfection (SODIS) to reduce childhood diarrhoea in rural Bolivia: a cluster-randomized, controlled trial. *PLoS Medicine* 2009;**6**(8):e1000125.

McGuigan 2011 KHM {published data only}

McGuigan KG, Samaiyar P, du Preez M, Conroy RM. High compliance randomized controlled field trial of solar disinfection of drinking water and its impact on childhood diarrhea in rural Cambodia. *Environmental Science & Technology* 2011;**45**(18):7862-7.

Mengistie 2013 ETH {published data only}

Mengistie B, Berhane Y, Worku A. Household water chlorination reduces incidence of diarrhea among under-five children in

rural Ethiopia: a cluster randomized controlled trial. *PLoS One* 2013;**8**(10):e77887.

Opryszko 2010a AFG {published data only}

Opryszko MC, Majeed SW, Hansen PM, Myers JA, Baba D, Thompson RE, et al. Water and hygiene interventions to reduce diarrhoea in rural Afghanistan: a randomized controlled study. *Journal of Water and Health* 2010;**8**(4):687-702.

Opryszko 2010b AFG {published data only}

Opryszko MC, Majeed SW, Hansen PM, Myers JA, Baba D, Thompson RE, et al. Water and hygiene interventions to reduce diarrhoea in rural Afghanistan: a randomized controlled study. *Journal of Water and Health* 2010;**8**(4):687-702.

Opryszko 2010c AFG {published data only}

Opryszko MC, Majeed SW, Hansen PM, Myers JA, Baba D, Thompson RE, et al. Water and hygiene interventions to reduce diarrhoea in rural Afghanistan: a randomized controlled study. *Journal of Water and Health* 2010;**8**(4):687-702.

Patel 2012 KEN {published and unpublished data}

Patel MK, Harris JR, Juliao P, Nygren B, Were V, Kola S, et al. Impact of a hygiene curriculum and the installation of simple handwashing and drinking water stations in rural Kenyan primary schools on student health and hygiene practices. *American Journal of Tropical Medicine and Hygiene* 2012;**87**(4):594-601.

Peletz 2012 ZMB {published data only}

Peletz R, Simunyama M, Sarenje K, Baisley K, Filteau S, Kelly P, et al. Assessing water filtration and safe storage in households with young children of HIV-positive mothers: a randomized, controlled trial in Zambia. *PLoS One* 2012;**7**(10):e46548.

Quick 1999 BOL {published and unpublished data}

* Quick RE, Venczel LV, Mintz ED, Soletto L, Aparicio J, Gironaz M, et al. Diarrhoea prevention in Bolivia through point-of-use water treatment and safe storage: a promising new strategy. *Epidemiology and Infection* 1999;**122**(1):83-90.

Venczel L. Evaluation and application of a mixed oxidant disinfection system for waterborne disease prevention [dissertation]. Chapel Hill (NC): University of North Carolina at Chapel Hill, 1997.

Quick 2002 ZMB {published data only}

Quick RE, Kimura A, Thevos A, Tembo M, Shamputa I, Hutwagner L, et al. Diarrhea prevention through household-level water disinfection and safe storage in Zambia. *American Journal of Tropical Medicine and Hygiene* 2002;**66**(5):584-9.

Reller 2003a GTM {published data only}

Reller ME, Mendoza CE, Lopez MB, Alvarez M, Hoekstra AR, Olson CA, et al. A randomized controlled trial of household-based flocculant-disinfectant drinking water treatment for diarrhoea prevention in rural Guatemala. *American Journal of Tropical Medicine and Hygiene* 2003;**69**(4):411-9.

Reller 2003b GTM {published data only}

Reller ME, Mendoza CE, Lopez MB, Alvarez M, Hoekstra AR, Olson CA, et al. A randomized controlled trial of household-based flocculant-disinfectant drinking water treatment for diarrhoea prevention in rural Guatemala. *American Journal of Tropical Medicine and Hygiene* 2003;**69**(4):411-9.

Reller 2003c GTM {published data only}

Reller ME, Mendoza CE, Lopez MB, Alvarez M, Hoekstra AR, Olson CA, et al. A randomized controlled trial of household-based flocculant-disinfectant drinking water treatment for diarrhoea prevention in rural Guatemala. *American Journal of Tropical Medicine and Hygiene* 2003;**69**(4):411-9.

Reller 2003d GTM {published data only}

Reller ME, Mendoza CE, Lopez MB, Alvarez M, Hoekstra AR, Olson CA, et al. A randomized controlled trial of household-based flocculant-disinfectant drinking water treatment for diarrhoea prevention in rural Guatemala. *American Journal of Tropical Medicine and Hygiene* 2003;**69**(4):411-9.

Roberts 2001 MWI {published data only}

Roberts L, Chartier Y, Chartier O, Malenga G, Toole M, Rodka H. Keeping clean water clean in a Malawi refugee camp: a randomized intervention trial. *Bulletin of the World Health Organization* 2001;**79**(4):280-7.

Rodrigo 2011 AUS {published data only}

Rodrigo S, Sinclair M, Forbes A, Cunliffe D, Leder K. Drinking rainwater: a double-blinded, randomized controlled study of water treatment filters and gastroenteritis incidence. *American Journal of Public Health* 2011;**101**(5):842-7.

Semenza 1998 UZB {published data only}

Semenza JC, Roberts L, Henderson A, Bogan J, Rubin CH. Water distribution system and diarrheal disease transmission: a case study in Uzbekistan. *American Journal of Tropical Medicine and Hygiene* 1998;**59**(6):941-6.

Stauber 2009 DOM {published and unpublished data}

Stauber CE, Ortiz GM, Loomis DP, Sobsey MD. A randomized controlled trial of the concrete Biosand filter and its impact on diarrheal disease in Bonao, Dominican Republic. *American Journal of Tropical Medicine and Hygiene* 2009;**80**(2):286-93.

Stauber 2012a KHM {published data only}

Stauber CE, Printy ER, McCarty FA, Liang KR, Sobsey MD. Cluster randomized controlled trial of the plastic BioSand water filter in Cambodia. *Environmental Science & Technology* 2012;**46**(2):722-8.

Stauber 2012b GHA {published and unpublished data}

Stauber C, Kominek B, Liang K, Osman M, Sobsey M. Evaluation of the impact of the plastic BioSand filter on health and drinking water quality in rural Tamale, Ghana. *International Journal of Environmental Research and Public Health* 2012;**9**(11):3806-23.

Tiwari 2009 KEN {published data only}

Tiwari SS, Schmidt WP, Darby J, Kariuki ZG, Jenkins MW. Intermittent slow sand filtration for preventing diarrhoea among children in Kenyan households using unimproved

water sources: randomized controlled trial. *Tropical Medicine & International Health* 2009;**14**(11):1374-82.

Torun 1982 GTM {published data only}

Torun B. Environmental and educational interventions against diarrhea in Guatemala. In: Chen LC, Scrimshaw NS editor(s). *Diarrhea and Malnutrition: Interactions, Mechanisms and Interventions*. New York: Plenum Press, 1982:235-66.

URL 1995a GTM {published data only}

Universidad Rafael Landivar. Preventing infant morbidity: artisanal filters and education [Contra la morbilidad infantil: filtros artesanales y education]. *Revista de Estudios Sociales* 1995;**IV**(53):1-66.

URL 1995b GTM {published data only}

Universidad Rafael Landivar. Preventing infant morbidity: artisanal filters and education [Contra la morbilidad infantil: filtros artesanales y education]. *Revista de Estudios Sociales* 1995;**IV**(53):1-66.

Xiao 1997 CHN {published data only}

Xiao S, Lin C, Chen K. Evaluation of effectiveness of comprehensive control for diarrhea diseases in rural areas of east Fujian and analysis of its cost-benefit [Chinese]. *Zhonghua Yu Fang Yi Xue Za Zhi [Chinese Journal of Preventive Medicine]* 1997;**31**(1):40-1.

References to studies excluded from this review
Ahoyo 2011 {published data only}

Ahoyo TA, Fatombi KJ, Boco M, Aminou T, Dramane KL. Impact of water quality and environmental sanitation on the health of schoolchildren in a suburban area of Benin: findings in the Savalou-Banté and Dassa-Glazoué sanitary districts [Impact de la qualité de l'eau et de l'assainissement sur la santé des enfants en milieu périurbain au Bénin: cas des zones sanitaires Savalou-Banté et Dassa-Glazoué]. *Médecine tropicale* 2011;**71**(3):281-5.

Aiken 2011 {published data only}

Aiken BA, Stauber CE, Ortiz GM, Sobsey MD. An assessment of continued use and health impact of the concrete biosand filter in Bonao, Dominican Republic. *American Journal of Tropical Medicine & Hygiene* 2011;**85**(2):309-17.

Alexander 2013 {published data only}

Alexander KT, Dreibelbis R, Freeman MC, Ojony B, Rheingans R. Improving service delivery of water, sanitation, and hygiene in primary schools: a cluster-randomized trial in western Kenya. *Journal of Water and Health* 2013;**11**(3):507-19.

Arnold 2009 {published data only}

Arnold B, Arana B, Mäusezahl D, Hubbard A, Colford JM Jr. Evaluation of a pre-existing, 3-year household water treatment and handwashing intervention in rural Guatemala. *International Journal of Epidemiology* 2009;**38**(6):1651-61.

Arnold 2012a {published data only}

Arnold BF, Mäusezahl D, Schmidt WP, Christen A, Colford, JM. Comment on randomized intervention study of solar

disinfection of drinking water in the prevention of dysentery in Kenyan children aged under 5 years. *Environmental Science & Technology* 2012;**46**(5):3031-2.

Arnold 2013 {published data only}

Arnold BF, Null C, Luby SP, Unicomb L, Stewart CP, Dewey KG, et al. Cluster-randomised controlled trials of individual and combined water, sanitation, hygiene and nutritional interventions in rural Bangladesh and Kenya: the WASH Benefits study design and rationale. *BMJ Open* 2013;**3**(8):e003476.

Asaolu 2002 {published data only}

Asaolu SO, Ofoezie IE, Odumuyiwa PA, Sowemimo OA, Ogunniyi TA. Effect of water supply and sanitation on the prevalence and intensity of *Ascaris lumbricoides* among pre-school-age children in Ajebandele and Ifewara, Osun State, Nigeria. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 2002;**96**(6):600-4.

Aziz 1990 BGD {published data only}

* Aziz KM, Hoque BA, Hasan KZ, Patwary MY, Huttly SR, Rahaman MM, et al. Reduction in diarrhoeal diseases in children in rural Bangladesh by environmental and behavioural modifications. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1990;**84**(3):433-8.

Hasan KZ, Briend A, Aziz KM, Hoque BA, Patwary MY, Huttly SR. Lack of impact of a water and sanitation intervention on the nutritional status of children in rural Bangladesh. *European Journal of Clinical Nutrition* 1989;**43**(12):837-43.

Henry FJ, Huttly SR, Patwary Y, Aziz KM. Environmental sanitation, food and water contamination and diarrhoea in rural Bangladesh. *Epidemiology and Infection* 1990;**104**(2):253-9.

Azurin 1974 {published data only}

Azurin JC, Alvero M. Field evaluation of environmental sanitation measures against cholera. *Bulletin of the World Health Organization* 1974;**51**(1):19-26.

Bahl 1976 {published data only}

Bahl MR. Impact of piped water supply on the incidence of typhoid fever and diarrhoeal diseases in Lusaka. *Medical Journal of Zambia* 1976;**10**(4):98-9.

Bajer 2012 {published data only}

Bajer A, Toczylowska B, Bednarska M, Sinski, E. Effectiveness of water treatment for the removal of *Cryptosporidium* and *Giardia* spp. *Epidemiology and Infection* 2012;**140**(11):2014-22.

Barreto 2007 {published data only}

Barreto ML, Genser B, Strina A, Teixeira MG, Assis AMO, Rego RF, et al. Effect of city-wide sanitation programme on reduction in rate of childhood diarrhoea in northeast Brazil: assessment by two cohort studies. *Lancet* 2007;**370**(9599):1622-8.

Barzilay 2011 {published data only}

Barzilay EJ, Aghoghovbia TS, Blanton EM, Akinpelumi AA, Coldiron ME, Akinfolayan O, et al. Diarrhea prevention in people living with HIV: an evaluation of a point-of-use water quality intervention in Lagos, Nigeria. *AIDS Care* 2011;**23**(3):330-9.

Bersh 1985 {published data only}

Bersh D, Osorio MM. Studies of diarrhoea in Quindio (Colombia): problems related to water treatment. *Social Science & Medicine* 1985;**21**(1):31-9.

Boubacar 2014 {published data only}

Boubacar Mainassara H, Tohon Z. Assessing the health impact of the following measures in schools in Maradi (Niger): construction of latrines, clean water supply, establishment of hand washing stations, and health education. *Journal of Parasitology Research* 2014;**2014**:190451.

Brown 2012a {published data only}

Brown J, Clasen T. High adherence is necessary to realize health gains from water quality interventions. *PLoS One* 2012;**7**(5):e36735.

Capuno 2011 {published data only}

Capuno JJ, Tan CAR Jr, Fabella VM. Do piped water and flush toilets prevent child diarrhea in rural Philippines?. *Asia-Pacific Journal of Public Health* 2011;**December**:1-11.

Cavallaro 2011 {published data only}

Cavallaro EC, Harris JR, da Goia MS, dos Santos Barrado JC, Alves da Nóbrega A, Carvalho de Alvarenga Júnior I, et al. Evaluation of pot-chlorination of wells during a cholera outbreak, Bissau, Guinea-Bissau, 2008. *Journal of Water and Health* 2011;**9**(2):394-402.

Chang 2012 {published data only}

Chang WK, Ryu J, Yi Y, Lee W-C, Lee C-W, Kang D, et al. Improved water quality in response to pollution control measures at Masan Bay, Korea. *Marine Pollution Bulletin* 2012;**64**(2):427-35.

Chongsuvivatwong 1994 {published data only}

Chongsuvivatwong V, Mo-suwan L, Chompikul J, Vitsupakorn K, McNeil D. Effects of piped water supply on the incidence of diarrhoeal diseases in children in southern Thailand. *Southeast Asian Journal of Tropical Medicine and Public Health* 1994;**25**(4):628-32.

Christen 2011 {published data only}

Christen A, Duran Pancheco G, Hattendorf J, Arnold BF, Cevallos M, Indergand S, et al. Factors associated with compliance among users of solar water disinfection in rural Bolivia. *BMC Public Health* 2011;**11**:210.

Clasen 2012 {published data only}

Clasen T, Boisson S, Routray P, Cumming O, Jenkins M, Ensink JHJ, et al. The effect of improved rural sanitation on diarrhoea and helminth infection: design of a cluster-randomized trial in Orissa, India. *Emerging Themes in Epidemiology* 2012;**9**:7.

Colford 2005 {published data only}

Colford JM Jr, Wade TJ, Sandhu SK, Wright CC, Lee S, Shaw S, et al. A randomized, controlled trial of in-home drinking water intervention to reduce gastrointestinal illness. *American Journal of Epidemiology* 2005;**161**(5):472-82.

Colwell 2003 {published data only}

Colwell RR, Huq A, Islam MS, Aziz KM, Yunus M, Khan NH, et al. Reduction of cholera in Bangladeshi villages by simple filtration. *Proceedings of the National Academy of Sciences of the United States of America* 2003;**100**(3):1051-5.

Conroy 2001 {published data only}

Conroy RM, Meegan ME, Joyce T, McGuigan K, Barnes J. Solar disinfection of drinking water protects against cholera in children under 6 years of age. *Archives of Disease in Childhood* 2001;**85**(4):293-5.

Coulliette 2013 {published data only}

Coulliette AD, Enger KS, Weir MH, Rose JB. Risk reduction assessment of waterborne Salmonella and Vibrio by a chlorine contact disinfectant point-of-use device. *International Journal of Hygiene and Environmental Health* 2013;**216**(3):355-61.

Crump 2007 {published data only}

Crump JA, Mendoza CE, Priest JW, Glass RI, Monroe SS, Dauphin LA, et al. Comparing serologic response against enteric pathogens with reported diarrhea to assess the impact of improved household drinking water quality. *American Journal of Tropical Medicine and Hygiene* 2007;**77**(1):136-41.

Davis 2011 {published data only}

Davis J, Pickering AJ, Rogers K, Mamuya S, Boehm AB. The effects of informational interventions on household water management, hygiene behaviors, stored drinking water quality, and hand contamination in peri-urban Tanzania. *American Journal of Tropical Medicine and Hygiene* 2011;**84**(2):184-91.

Deb 1986 {published data only}

Deb BC, Sircar BK, Sengupta PG, De SP, Mondal SK, Gupta DN, et al. Studies on interventions to prevent eltor cholera transmission in urban slums. *Bulletin of the World Health Organization* 1986;**64**(1):127-31.

Denslow 2010 {published data only}

Denslow SA, Edwards J, Horney J, Peña R, Wurzelmann D, Morgan D. Improvements to water purification and sanitation infrastructure may reduce the diarrheal burden in a marginalized and flood prone population in remote Nicaragua. *BMC International Health and Human Rights* 2010;**10**:30.

Devoto 2011 {published data only}

Devoto F, Duflo E, Dupas P, Pariente W, Pons V. Happiness on Tap: Piped Water Adoption in Urban Morocco. *National Bureau of Economic Research* 2011;**Working paper: 16933**:1-39.

Dorevitch 2011 {published data only}

Dorevitch S, Doi M, Hsu FC, Lin KT, Roberts JD, Liu LC, et al. A comparison of rapid and conventional measures of indicator bacteria as predictors of waterborne protozoan pathogen presence and density. *Journal of Environmental Monitoring* 2011;**13**(9):2427-35.

Dreibelbis 2014a KEN {published data only}

Dreibelbis R, Freeman MC, Greene LE, Saboori S, Rheingans R. The impact of school water, sanitation, and hygiene interventions on the health of younger siblings of pupils: a

cluster-randomized trial in Kenya. *American Journal of Public Health* 2014;**104**(1):e91-7.

Dreibelbis 2014b KEN {published data only}

Dreibelbis R, Freeman MC, Greene LE, Saboori S, Rheingans R. The impact of school water, sanitation, and hygiene interventions on the health of younger siblings of pupils: a cluster-randomized trial in Kenya. *American Journal of Public Health* 2014;**104**(1):e91-7.

Dreibelbis 2014c KEN {published data only}

Dreibelbis R, Freeman MC, Greene LE, Saboori S, Rheingans R. The impact of school water, sanitation, and hygiene interventions on the health of younger siblings of pupils: a cluster-randomized trial in Kenya. *American Journal of Public Health* 2014;**104**(1):e91-7.

du Preez 2012 {published data only}

du Preez M, Conroy RM, McGuigan KG. Response to comment on "A randomized intervention study of solar disinfection of drinking water (SODIS) in the prevention of dysentery in Kenyan children aged under 5 years". *Environmental Science and Technology* 2012;**46**(5):3036-7.

Eisenberg 2006 {published data only}

Eisenberg JNS, Hubbard A, Wade TJ, Sylvester MD, LeChevallier MW, Levy DA, et al. Inferences drawn from a risk assessment compared directly with a randomized trial of a home drinking water intervention. *Environmental Health Perspectives* 2006;**114**(8):1199-204.

Enger 2012 {published data only}

Enger KS, Nelson KL, Clasen T, Rose JB, Eisenberg JNS. Linking quantitative microbial risk assessment and epidemiological data: informing safe drinking water trials in developing countries. *Environmental Science and Technology* 2012;**46**(9):5160-7.

Esrey 1988 {published data only}

Esrey SA, Habicht JJP, Latham MC, Sisler DG, Casella G. Drinking water source, diarrhoeal morbidity, and child growth in villages with both traditional and improved water supplies in rural Lesotho, southern Africa. *American Journal of Public Health* 1988;**78**(11):1451-5.

Fewtrell 1994 {published data only}

Fewtrell L, Kay D, Dunlop J, O'Neill G, Wyer M. Infectious diseases and water-supply disconnections. *Lancet* 1994;**343**(8909):1370.

Fewtrell 1997 {published data only}

Fewtrell L, Kay D, Wyer M, O'Neill G. An investigation into the possible links between shigellosis and hepatitis A and public water supply disconnections. *Public Health* 1997;**111**(3):179-81.

Firth 2010 {published data only}

Firth J, Balraj V, Muliyl J, Roy S, Rani LM, Chandrasekhar R, et al. Point-of-use interventions to decrease contamination of drinking water: a randomized, controlled pilot study on efficacy, effectiveness, and acceptability of closed containers, Moringa oleifera, and In-home chlorination in rural South

Interventions to improve water quality for preventing diarrhoea (Review)

India. *American Journal of Tropical Medicine and Hygiene* 2010;**82**(5):759-65.

Fisher 2011 {published data only}

Fisher S, Kabir B, Lahiff E, MacLachlan M. Knowledge, attitudes, practices and implications of safe water management and good hygiene in rural Bangladesh: assessing the impact and scope of the BRAC WASH programme. *Journal of Water and Health* 2011;**9**(1):80-93.

Freeman 2012 {published data only}

Freeman MC, Greene LE, Dreibelbis R, Saboori S, Muga R, Brumback B, et al. Assessing the impact of a school-based water treatment, hygiene and sanitation programme on pupil absence in Nyanza Province, Kenya: a cluster-randomized trial. *Tropical Medicine & International Health* 2012;**17**(3):380-91.

Freeman 2014a KEN {published data only}

* Freeman MC, Clasen T, Dreibelbis R, Saboori S, Greene L, Brumback B, et al. The impact of a school-based water supply and treatment, hygiene, and sanitation programme on pupil diarrhoea: a cluster-randomized trial. *Epidemiology and Infection* 2014;**142**(2):340-51.

Freeman 2014b KEN {published data only}

Freeman MC, Clasen T, Dreibelbis R, Saboori S, Greene L, Brumback B, et al. The impact of a school-based water supply and treatment, hygiene, and sanitation programme on pupil diarrhoea: a cluster-randomized trial. *Epidemiology and Infection* 2014;**142**(2):340-51.

Freeman 2014c KEN {published data only}

Freeman MC, Clasen T, Dreibelbis R, Saboori S, Greene L, Brumback B, et al. The impact of a school-based water supply and treatment, hygiene, and sanitation programme on pupil diarrhoea: a cluster-randomized trial. *Epidemiology and Infection* 2014;**142**(2):340-51.

Fry 2010 {published data only}

Fry LM, Cowden JR, Watkins DW Jr, Clasen T, Mihelcic JR. Quantifying health improvements from water quantity enhancement: an engineering perspective applied to rainwater harvesting in West Africa. *Environmental Science and Technology* 2010;**44**(24):9535-41.

Galiani 2009 {published data only}

Galiani S, Gonzalez-Rozada M, Schargrodsky E. Water expansions in shantytowns: health and savings. *Economica* 2009;**76**(104):607-22.

Garrett 2008 KEN {published data only}

Garrett V, Ogutu P, Mabonga P, Ombeki S, Mwaki A, Aluoch G, et al. Diarrhoea prevention in a high-risk rural Kenyan population through point-of-use chlorination, safe water storage, sanitation, and rainwater harvesting. *Epidemiology and Infection* 2008;**136**(11):1463-71.

Ghannoum 1981 {published data only}

Ghannoum MA, Moore KE, Al-Dulaimi M, Nasr M. The incidence of water-related diseases in the Brak area, Libya from 1977 to 1979, before and after the installation of water treatment

plants. *Zentralblatt für Bakteriologie, Mikrobiologie und Hygiene. 1. Abt. Originale B, Hygiene* 1981;**173**(6):501-8.

Gómez-Couso 2012 {published data only}

Gómez-Couso H, Fontán-Sainza M, Navntoft C, Fernández-Ibáñez P, Ares-Mazás E. Comparison of different solar reactors for household disinfection of drinking water in developing countries: evaluation of their efficacy in relation to the waterborne enteropathogen *Cryptosporidium parvum*. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 2012;**106**(11):645-52.

Gorelick 2011 {published data only}

Gorelick MH, McLellan SL, Wagner D, Klein J. Water use and acute diarrhoeal illness in children in a United States metropolitan area. *Epidemiology & Infection* 2011;**139**(2):295-301.

Greene 2012 {published data only}

Greene LE, Freeman MC, Akoko D, Saboori S, Moe C, Rheingans R. Impact of a school-based hygiene promotion and sanitation intervention on pupil hand contamination in Western Kenya: a cluster randomized trial. *American Journal of Tropical Medicine and Hygiene* 2012;**87**(3):385-93.

Habib 2013 {published data only}

Habib MA, Soofi S, Sadiq K, Samejo T, Hussain M, Mirani M, et al. A study to evaluate the acceptability, feasibility and impact of packaged interventions ("Diarrhea Pack") for prevention and treatment of childhood diarrhea in rural Pakistan. *BMC Public Health* 2013;**13**:922.

Harris 2009 {published data only}

Harris JR, Greene SK, Thomas TK, Ndivo R, Okanda J, Masaba R, et al. Effect of a point-of-use water treatment and safe water storage intervention on diarrhea in infants of HIV-infected mothers. *Journal of Infectious Diseases* 2009;**200**(8):1186-93.

Harshfield 2012 {published data only}

Harshfield E, Lantagne D, Turbes A, Null C. Evaluating the sustained health impact of household chlorination of drinking water in rural Haiti. *American Journal of Tropical Medicine and Hygiene* 2012;**87**(5):786-95.

Hartinger 2011 {published data only}

Hartinger SM, Lanata CF, Hattendorf J, Gil AI, Verastegui H, Ochoa T, et al. A community randomised controlled trial evaluating a home-based environmental intervention package of improved stoves, solar water disinfection and kitchen sinks in rural Peru: rationale, trial design and baseline findings. *Contemporary Clinical Trials* 2011;**32**(6):864-73.

Hartinger 2012 {published data only}

Hartinger SM, Lanata CF, Gil AI, Hattendorf J, Verastegui H, Mäusezahl D. Combining interventions: improved chimney stoves, kitchen sinks and solar disinfection of drinking water and kitchen clothes to improve home hygiene in rural Peru. *Journal of Field Actions* 2012;**6**:1-10.

Hellard 2001 {published data only}

Hellard ME, Sinclair MI, Forbes AB, Fairley CK. A randomized, blinded, controlled trial investigating the gastrointestinal health effects of drinking water quality. *Environmental Health Perspectives* 2001;**109**(8):773-8.

Hoque 1996 {published data only}

Hoque BA, Juncker T, Sack RB, Ali M, Aziz KM. Sustainability of a water, sanitation and hygiene education project in rural Bangladesh: a 5-year follow-up. *Bulletin of the World Health Organization* 1996;**74**(4):431-7.

Huda 2012 {published data only}

Huda TMN, Unicomb L, Johnston RB, Halder AK, Yushuf Sharker MA, Luby SP. Interim evaluation of a large scale sanitation and water improvement programme on childhood diarrhea and respiratory disease in rural Bangladesh. *Social Science and Medicine* 2012;**75**(4):604-11.

Hunter 2010 {published data only}

Hunter PR, Ramírez Toro GI, Minnigh HA. Impact on diarrhoeal illness of a community educational intervention to improve drinking water quality in rural communities in Puerto Rico. *BMC Public Health* 2010;**10**:219.

Iijima 2001 {published data only}

Iijima Y, Karama M, Oundo JO, Honda T. Prevention of bacterial diarrhea by pasteurization of drinking water in Kenya. *Microbiology and Immunology* 2001;**45**(6):413-6.

Islam 2011 {published data only}

Islam MS, Mahmud ZH, Uddin MH, Islam K, Yunus M, Islam MS, et al. Purification of household water using a novel mixture reduces diarrhoeal disease in Matlab, Bangladesh. *Transactions of the Royal Society of Tropical Medicine & Hygiene* 2011;**105**(6):341-5.

Jensen 2002 {published data only}

Jensen PK, Ensink JH, Jayasinghe G, van der Hoek W, Cairncross S, Dalsgaard A. Domestic transmission routes of pathogens: the problem of in-house contamination of drinking water during storage in developing countries. *Tropical Medicine & International Health* 2002;**7**(7):604-9.

Kariuki 2012 {published data only}

Kariuki JG, Magambo KJ, Njeruh MF, Muchiri EM, Nzioka SM, Kariuki S. Effects of hygiene and sanitation interventions on reducing diarrhoea prevalence among children in resource constrained communities: case study of Turkana District, Kenya. *Journal of Community Health* 2012;**37**(6):1178-84.

Karon 2011 {published data only}

Karon AE, Hanni KD, Mohle-Boetani JC, Beretti RA, Hill VR, Arrowood M, et al. Giardiasis outbreak at a camp after installation of a slow-sand filtration water-treatment system. *Epidemiology and Infection* 2011;**139**(5):713-7.

Keraita 2007 {published data only}

Keraita B, Konradsen F, Drechsel P, Abaidoo RC. Reducing microbial contamination on wastewater-irrigated lettuce by

cessation of irrigation before harvesting. *Tropical Medicine & International Health* 2007;**12**(Suppl 2):8-14.

Khan 1984 {published data only}

Khan MU, Khan MR, Hossan B, Ahmed QS. Alum potash in water to prevent cholera. *Lancet* 1984;**2**(8410):1032.

Luby 2008 {published data only}

Luby SP, Gupta SK, Sheikh MA, Johnston RB, Ram PK, Islam MS. Tubewell water quality and predictors of contamination in three flood-prone areas in Bangladesh. *Journal of Applied Microbiology* 2008;**105**(4):1002-8.

Luoto 2011 {published data only}

Luoto J, Najnin N, Mahmud M, Albert J, Islam MS, Luby SP, et al. What point-of-use water treatment products do consumers use? Evidence from a randomized controlled trial among the urban poor in Bangladesh. *PLoS One* 2011;**6**(10):e26132.

Luoto 2012 {published data only}

Luoto J, Mahmud M, Albert J, Luby SP, Najnin N, Unicomb L, et al. Learning to dislike safe water products: results from a randomized controlled trial of the effects of direct and peer experience on willingness to pay. *Environmental Science & Technology* 2012;**46**(11):6244-51.

Macy 1998 {published data only}

Macy JT, Quick RE. Evaluation of a novel drinking water treatment and storage intervention in Nicaragua. *Revista Panamericana de Salud Pública (Pan American Journal of Public Health)* 1998;**3**(2):135-6.

Mäusezahl 2003 {unpublished data only}

Mäusezahl D, Tanner M, Hobbins M. The SODIS health impact study (as supplied 15 March 2004). Data on file.

McCabe 1957 {published data only}

McCabe LJ, Haines TW. Diarrheal disease control by improved human excreta disposal. *Public Health Reports* 1957;**72**(10):921-8.

Mertens 1990 {published data only}

Mertens TE, Cousens SN, Fernando MA, Kirkwood BR, Merkle F, Korte R, et al. Health impact evaluation of improved water supplies and hygiene practices in Sri Lanka: background and methodology. *Tropical Medicine and Parasitology* 1990;**41**(1):79-88.

Messou 1997 {published data only}

Messou E, Sangaré SV, Jossieran R, Le Corre C, Guélain J. Effect of hygiene measures, water, sanitation and oral rehydration therapy on diarrhoea in children less than five years old in the south of Ivory Coast [Effet de l'observance des mesures d'hygiène, d'approvisionnement en eau et de la thérapie par voie orale sur les diarrhées chez les enfants de moins de 5 ans dans le sud de la Côte D'Ivoire]. *Bulletin de la Société de Pathologie Exotique (1990)* 1997;**90**(1):44-7.

Nanan 2003 {published data only}

Nanan D, White F, Azam I, Afsar H, Hozhari S. Evaluation of a water, sanitation, and hygiene education intervention on

diarrhoea in northern Pakistan. *Bulletin of the World Health Organization* 2003;**81**(3):160-5.

Nerkar 2014 {published data only}

Nerkar SS, Tamhankar AJ, Khedar SU, Lundborg CS. Quality of water and antibiotic resistance of *Escherichia coli* from water sources of hilly tribal villages with and without integrated watershed management—a one year prospective study. *International Journal of Environmental Research and Public Health* 2014;**11**(6):6156-70.

Nnane 2011 {published data only}

Nnane DE, Ebdon JE, Taylor HD. Integrated analysis of water quality parameters for cost-effective faecal pollution management in river catchments. *Water Research* 2011;**45**(6):2235-46.

Oluyeye 2011 {published data only}

Oluyeye JO, Koko AE, Aregbesola OA. Bacteriological and physico-chemical quality assessment of household drinking water in Ado-Ekiti, Nigeria. *Water Science & Technology: Water Supply* 2011;**11**(1):79-84.

Palit 2012 {published data only}

Palit A, Batabyal P, Kanungo S, Sur D. In-house contamination of potable water in urban slum of Kolkata, India: a possible transmission route of diarrhea. *Water Science and Technology* 2012;**66**(2):299-303.

Pavlinac 2014 {published data only}

Pavlinac PB, Naulikha JM, Chaba L, Kimani N, Sangaré LR, Yuhas K, et al. Water filter provision and home-based filter reinforcement reduce diarrhea in Kenyan HIV-infected adults and their household members. *American Journal of Tropical Medicine and Hygiene* 2014;**91**(2):273-80.

Payment 1991a {published data only}

Payment P, Franco E, Richardson L, Siemiatycki J. Gastrointestinal health effects associated with the consumption of drinking water produced by point-of-use domestic reverse-osmosis filtration units. *Applied and Environmental Microbiology* 1991;**57**(4):945-8.

Payment 1991b {published data only}

Payment P, Richardson L, Siemiatycki J, Dewar R, Edwardes M, Franco E. A randomized trial to evaluate the risk of gastrointestinal disease due to consumption of drinking water meeting current microbiological standards. *American Journal of Public Health* 1991;**81**(6):703-8.

Peletz 2013 {published data only}

Peletz R, Simuyandi M, Simunyama M, Sarenje K, Kelly P, Clasen T. Follow-up study to assess the use and performance of household filters in Zambia. *American Journal of Tropical Medicine and Hygiene* 2013;**89**(6):1190-4.

Pinfold 1990 {published data only}

Pinfold JV. Faecal contamination of water and fingertip-rinses as a method for evaluating the effect of low-cost water supply and sanitation activities on faeco-oral disease transmission.

II. A hygiene intervention study in rural north-east Thailand. *Epidemiology and Infection* 1990;**105**(2):377-89.

Psutka 2012 {published data only}

Psutka R, Peletz R, Michelo S, Kelly P, Clasen T. Assessing the microbiological performance and potential cost of boiling drinking water in urban Zambia. *Environmental Science & Technology* 2011;**45**(14):6095-101.

Rosa 2014 {published data only}

Rosa G, Majorin F, Boisson S, Barstow C, Johnson M, Kirby M, et al. Assessing the impact of water filters and improved cook stoves on drinking water quality and household air pollution: a randomised controlled trial in Rwanda. *PLoS One* 2014;**9**(3):e91011.

Rose 2006 {published data only}

Rose A, Roy S, Abraham V, Homgren G, George K, Balraj V, et al. Solar disinfection of water for diarrhoeal prevention in Southern India. *Archives of Disease in Childhood* 2006;**91**(2):139-41.

Rubenstein 1969 {published data only}

Rubenstein A, Boyle J, Odoroff CL, Kunitz SJ. Effect of improved sanitary facilities on infant diarrhea in a Hopi village. *Public Health Reports* 1969;**84**(12):1093-7.

Russo 2012 {published data only}

Russo ET, Sheth A, Menon M, Wannemuehler K, Weinger M, Kudzala AC, et al. Water treatment and handwashing behaviors among non-pregnant friends and relatives of participants in an antenatal hygiene promotion program in Malawi. *American Journal of Tropical Medicine & Hygiene* 2012;**86**(5):860-5.

Sathe 1996 {published data only}

Sathe AA, Hinge DV, Watve MG. Water treatment and diarrhoea. *Lancet* 1996;**348**(9023):335-6.

Shah 2012 {published data only}

Shah D, Choudhury P, Gupta P, Mathew JL, Gera T, Gogia S, et al. Promoting appropriate management of diarrhea: a systematic review of literature for advocacy and action: UNICEF-PHFI series on newborn and child health, India. *Indian Pediatrics* 2012;**49**(8):627-49.

Sharan 2011 {published data only}

Sharan R, Chhibber S, Heed RH. Inactivation and sub-lethal injury of salmonella typhi, salmonella typhimurium and vibrio cholerae in copper water storage vessels. *BMC Infectious Diseases* 2011;**11**:204.

Sheth 2010 {published data only}

Sheth AN, Russo ET, Menon M, Wannemuehler K, Weinger M, Kudzala AC, et al. Impact of the integration of water treatment and handwashing incentives with antenatal services on hygiene practices of pregnant women in Malawi. *American Journal of Tropical Medicine and Hygiene* 2010;**83**(6):1315-21.

Shiffman 1978 {published data only}

Shiffman MA, Schneider R, Faigenblum JM, Helms R, Turner A. Field studies on water, sanitation and health education in

relation to health status in Central America. *Progress in Water Technology* 1978;**11**:143-50.

Shrestha 2006 {published data only}

Shrestha RK, Marseill E, Kahn JG, Lule JG, Pitter C, Blandford JM, et al. Cost-effectiveness of home-based chlorination and safe water storage in reducing diarrhea among HIV-affected households in Uganda. *American Journal of Tropical Medicine and Hygiene* 2006;**74**(5):884-90.

Shum 1971 {published data only}

Shum H, Sum CY, Chan-Teo CH. Water-borne dysentery due to *Shigella sonnei* in Hong Kong. *Southeast Asian Journal of Tropical Medicine and Public Health* 1971;**2**(2):180-5.

Sima 2012 {published data only}

Sima LC, Desai MM, McCarty KM, Elimelech M. Relationship between use of water from community-scale water treatment refill kiosks and childhood diarrhea in Jakarta. *American Journal of Tropical Medicine and Hygiene* 2012;**87**(6):979-84.

Sorvillo 1994 {published data only}

Sorvillo FJ, Lieb LE, Nahlen B, Miller J, Mascola L, Ash LR. Municipal drinking water and cryptosporidiosis among persons with AIDS in Los Angeles County. *Epidemiology and Infection* 1994;**113**(2):313-20.

Stauber 2013 {published data only}

Stauber CE, Walters A, Fabiszewski de Aceituno AM, Sobsey MD. Bacterial contamination on household toys and association with water, sanitation and hygiene conditions in Honduras. *International Journal of Environmental Research and Public Health* 2013;**10**(4):1586-97.

Sutha 2011 {published data only}

Sutha S. Contaminated drinking water and rural health perspectives in Rajasthan, India: an overview of recent case studies. *Environmental Monitoring and Assessment* 2011;**173**(1-4):837-49.

Tonglet 1992 {published data only}

Tonglet R, Isu K, Mpese M, Dramaix M, Hennart P. Can improvements in water supply reduce childhood diarrhoea?. *Health Policy and Planning* 1992;**7**(3):260-8.

Trivedi 1971 {published data only}

Trivedi BK, Gandhi HS, Shukla NK. Bacteriological water quality and incidence of water borne diseases in a rural population. *Indian Journal of Medical Sciences* 1971;**25**(11):795-801.

VanDerslice 1995 {published data only}

VanDerslice J, Briscoe J. Environmental interventions in developing countries: interactions and their implications. *American Journal of Epidemiology* 1995;**141**(2):135-44.

Varghese 2002 {unpublished data only}

Varghese A. Point-of-use water treatment systems in rural Haiti: human health and water quality impact assessment [dissertation]. Cambridge (MA): Massachusetts Institute of Technology, 2002.

Wiedenmann 2006 {published data only}

Wiedenmann A, Krüger P, Dietz K, López-Pila JM, Szewzyk R, Botzenhart K. A randomized controlled trial assessing infectious disease risks from bathing in fresh recreational waters in relation to the concentration of *Escherichia coli*, intestinal Enterococci, *Clostridium perfringens*, and somatic coliphages. *Environmental Health Perspectives* 2006;**114**(2):228-36.

Wolf 2014 {published data only}

Wolf J, Prüss-Ustün A, Cumming O, Bartram J, Bonjour S, Cairncross S, et al. Assessing the impact of drinking water and sanitation on diarrhoeal disease in low- and middle-income settings: systematic review and meta-regression. *Tropical Medicine & International Health* 2014;**19**(8):928-42.

Wood 2012 {published data only}

Wood S, Foster J, Kols A. Understanding why women adopt and sustain home water treatment: Insights from the Malawi antenatal care program. *Social Science Medicine* 2012;**75**(4):634-42.

Wu 2011 {published data only}

Wu J, van Geen A, Ahmed KM, Alam YAJ, Culligan PJ, Escamilla V, et al. Increase in diarrheal disease associated with arsenic mitigation in Bangladesh. *PLoS One* 2011;**6**(12):e29593.

References to ongoing studies

Chlorination, Dhaka {published data only}

Impact of Low-Cost In-Line Chlorination Systems in Urban Dhaka on Water Quality and Child Health. Ongoing study Early 2015.

WASH-B, Bangladesh {published data only}

WASH Benefits Bangladesh: A Cluster Randomized Controlled Trial of the Benefits of Water, Sanitation, Hygiene Plus Nutrition Interventions on Child Growth. Ongoing study May 2012.

WASH-B, Kenya {published data only}

WASH-Benefits study, Kenya. Ongoing study September 2012.

Additional references

Arnold 2007

Arnold BF, Colford JM Jr. Treating water with chlorine at point-of-use to improve water quality and reduce child diarrhea in developing countries: a systematic review and meta-analysis. *American Journal of Tropical Medicine and Hygiene* 2007;**76**(2):354-64.

Austin 1993

Austin CJ. Investigation of in-house water chlorination and its effectiveness for rural areas of the Gambia [dissertation]. New Orleans: Tulane University School of Public Health and Tropical Medicine, 1993.

Bain 2014

Bain R, Cronk R, Wright J, Yang H, Slaymaker T, Bartram J. Fecal contamination of drinking-water in low- and middle-

income countries: a systematic review and meta-analysis. *PLoS Medicine* 2014;**11**(5):e1001644.

Brown 2008

Brown J, Sobsey MD, Loomis D. Local drinking water filters reduce diarrheal disease in Cambodia: a randomized, controlled trial of the ceramic water purifier. *American Journal of Tropical Medicine and Hygiene* 2008;**79**(3):394-400.

Brown 2012

Brown J, Clasen T. High adherence is necessary to realize health gains from water quality interventions. *PLoS One* 2012;**7**(5):e36735.

Byers 2001

Byers KE, Guerrant RL, Farr BM. Fecal-oral transmission. In: Thomas JC, Webber DJ editor(s). *Epidemiologic Methods for the Study of Infectious Diseases*. Oxford: Oxford University Press, 2001:228-48.

Cairncross 2010

Cairncross S, Hunt C, Boisson S, Bostoen K, Curtis V, Fung ICH, et al. Water, sanitation and hygiene for the prevention of diarrhoea. *International Journal of Epidemiology* 2010;**39**(Suppl 1):i193-205.

Crump 2005

Crump JA, Otieno PO, Slutsker L, Keswick BH, Rosen DH, Hoekstra RM, et al. Household based treatment of drinking water with flocculant-disinfectant for preventing diarrhoea in areas with turbid source water in rural western Kenya: cluster randomised controlled trial. *BMJ* 2005;**331**(7515):478.

Cutler 2005

Cutler D, Miller G. The role of public health improvements in health advances: the twentieth-century United States [2005]. *Demography* 2005;**42**(1):1-22.

Eisenberg 2007

Eisenberg JN, Scott JC, Porco T. Integrating disease control strategies: balancing water sanitation and hygiene interventions to reduce diarrheal disease burden. *American Journal of Public Health* 2007;**97**(5):846-52.

Eisenberg 2012

Eisenberg JN, Trostle J, Sorensen RJ, Shields KF. Toward a systems approach to enteric pathogen transmission: from individual independence to community interdependence. *Annual Review of Public Health* 2012;**33**:239-57.

Engell 2013

Engell RE, Lim SS. Does clean water matter? An updated meta-analysis of water supply and sanitation interventions and diarrhoeal diseases. *Lancet* 2013;**381**(Supplement 2):S44.

Enger 2013

Enger KS, Nelson KL, Rose JB, Eisenberg JNS. The joint effects of efficacy and compliance: a study of household water treatment effectiveness against childhood diarrhea. *Water Research* 2013;**47**(3):1181-90.

Esrey 1986

Esrey SA, Habicht JP. Epidemiologic evidence for health benefits from improved water and sanitation in developing countries. *Epidemiologic Reviews* 1986;**8**:117-28.

Fewtrell 2005

Fewtrell L, Kaufmann RB, Kay D, Enanoria W, Haller L, Colford JM Jr. Water, sanitation, and hygiene interventions to reduce diarrhoea in less developed countries: a systematic review and meta-analysis. *Lancet Infectious Diseases* 2005;**5**(1):42-52.

Fischer Walker 2012

Fischer Walker CL, Perin J, Aryee MJ, Boschi-Pinto C, Black RE. Diarrhea incidence in low- and middle-income countries in 1990 and 2010: a systematic review. *BMC Public Health* 2012;**12**:220.

GBD 2015

GBD 2013 Risk Factors Collaborators. Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015;**S0140-6736**(15):128.

GRADEpro GDT [Computer program]

McMaster University (developed by Evidence Prime, Inc.). GRADEpro GDT: GRADEpro Guideline Development Tool. McMaster University (developed by Evidence Prime, Inc.), 2015.

Guyatt 2008

Guyatt GH, Oxman AD, Kunz R, Vist GE, Falck-Ytter Y, Schünemann HJ, GRADE Working Group. Rating quality of evidence and strength of recommendations: What is "quality of evidence" and why is it important to clinicians?. *BMJ* 2008;**336**(7651):995-8.

Higgins 2005

Higgins J, Green S, editors. Highly sensitive search strategies for identifying reports of randomized controlled trials in MEDLINE. *Cochrane Handbook for Systematic Reviews of Interventions* 4.2.5 [updated May 2005]; Appendix 5b (accessed 1 January 2006).

Higgins 2006

Higgins JPT, Green S, editors. Section 8: Analysing and presenting results. *Cochrane Handbook for Systematic Reviews of Interventions* 4.2.5 [updated September 2006]. <http://community.cochrane.org/sites/default/files/uploads/Handbook4.2.6Sep2006.pdf> (accessed 15 November 2014).

Higgins 2011

Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available at <http://handbook.cochrane.org/>.

Hunter 2009

Hunter PR. Household water treatment in developing countries: comparing different intervention types using meta-regression. *Environmental Science & Technology* 2009;**43**(23):8991-7.

Interventions to improve water quality for preventing diarrhoea (Review)

Hutton 2013

Hutton G. Global costs and benefits of reaching universal coverage of sanitation and drinking-water supply. *Journal of Water and Health* 2013;**11**(1):1-12.

Kotloff 2013

Kotloff KL, Nataro JP, Blackwelder WC, Nasrin D, Farag TH, Panchalingam S, et al. Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *Lancet* 2013;**382**(9888):209-22.

Lanata 2013

Lanata CF, Fischer-Walker CL, Olascoaga AC, Torres CX, Aryee MJ, Black RE, Child Health Epidemiology Reference Group of the World Health Organization and UNICEF. Global causes of diarrheal disease mortality in children <5 years of age: a systematic review. *PLoS One* 2013;**8**(9):e72788.

Leclerc 2002

Leclerc H, Schwartzbrod L, Dei-Cas E. Microbial agents associated with waterborne diseases. *Critical Reviews in Microbiology* 2002;**28**(4):371-409.

Lindquist 2014

Lindquist ED, George CM, Perin J, Neiswender de Calani KJ, Norman WR, Davis TP, et al. A cluster randomized controlled trial to reduce childhood diarrhea using hollow fiber water filter and/or hygiene-sanitation educational interventions. *American Journal of Tropical Medicine and Hygiene* 2014;**91**(1):190-7.

Luby 2004

Luby SP, Agboatwalla M, Hoekstra RM, Rahbar MH, Billhimer W, Keswick BH. Delayed effectiveness of home-based interventions in reducing childhood diarrhea, Karachi, Pakistan. *American Journal of Tropical Medicine and Hygiene* 2004;**71**(4):420-7.

Luby 2006

Luby SP, Agboatwalla M, Painter J, Altaf A, Billhimer W, Keswick B, et al. Combining drinking water treatment and hand washing for diarrhoea prevention, a cluster randomised controlled trial. *Tropical Medicine & International Health* 2006;**11**(4):479-89.

McNutt 2003

McNutt LA, Wu C, Xue X, Hafner JP. Estimating the relative risk in cohort studies and clinical trials of common outcomes. *American Journal of Epidemiology* 2003;**157**(10):940-3.

Murray 2012

Murray CJL, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012; Vol. 380, issue 9859:2197-223.

Opryszko 2010

Opryszko MC, Majeed SW, Hansen PM, Myers JA, Baba D, Thompson RE, et al. Water and hygiene interventions to reduce diarrhoea in rural Afghanistan: a randomized controlled study. *Journal of Water Health* 2010;**8**(4):687-702.

Reller 2003

Reller ME, Mendoza CE, Lopez MB, Alvarez M, Hoekstra AR, Olson CA, et al. A randomized controlled trial of household-based flocculant-disinfectant drinking water treatment for diarrhoea prevention in rural Guatemala. *American Journal of Tropical Medicine and Hygiene* 2003;**69**(4):411-9.

Review Manager (RevMan) [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.3. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014.

Reynolds 2008

Reynolds KA, Mena KD, Gerba CP. Risk of waterborne illness via drinking water in the United States. In: Whitacre DM editor(s). *Reviews of Environmental Contamination and Toxicology*. Vol. **192**, New York: Springer, 2008:117-58.

Rosa 2010

Rosa G, Clasen T. Estimating the scope of household water treatment in low- and medium-income countries. *American Journal of Tropical Medicine and Hygiene* 2010;**82**(2):289-300.

Savović 2012

Savović J, Jones HE, Altman DG, Harris RJ, Jüni P, Pildal J, et al. Influence of reported study design characteristics on intervention effect estimates from randomized, controlled trials. *Annals of Internal Medicine* 2012;**157**(6):429-38.

Sobsey 2002

Sobsey MD. Managing water in the home: accelerated health gains from improved water supply [WHO/SDE/WSH/02.07]. Geneva: World Health Organization, 2002.

Sphere Project 2011

The Sphere Project. Humanitarian Charter and Minimum Standards in Disaster Response. 3rd Edition. Rugby: Practical Action Publishing, 2011.

Stelmach 2015

Stelmach RD, Clasen T. Household water quantity and health: a systematic review. *International Journal of Environmental Research and Public Health* 2015;**12**(6):5954-74.

URL 1995

Universidad Rafael Landivar. Preventing infant morbidity: artisanal filters and education [Contra la morbilidad infantil: filtros artesanales y educación]. *Revista de Estudios Sociales* 1995;**IV**(53):1-66.

Waddington 2009

Waddington H, Snilstveit B, White H, Fewtrell L. Water, sanitation and hygiene interventions to combat childhood diarrhoea in developing countries. <http://www.3ieimpact.org/evidence/systematic-reviews/details/23/>. New Delhi, India: 3ie, (accessed 15 November 2014).

Walker 2013

Walker CL, Rudan I, Liu L, Nair H, Theodoratou E, Bhutta ZA, et al. Global burden of childhood pneumonia and diarrhoea. *Lancet* 2013;**381**(9875):1405-16.

Interventions to improve water quality for preventing diarrhoea (Review)

WHO 1993

World Health Organization. The Management and Prevention of Diarrhoea: Practical Guidelines. 3rd Edition. Geneva: World Health Organization, 1993.

WHO 2011

World Health Organization. Guidelines for drinking-water quality, 4th edition. http://apps.who.int/iris/bitstream/10665/44584/1/9789241548151_eng.pdf. online: World Health Organization, (accessed 15 November 2014).

WHO 2014

World Health Organization. The top 10 causes of death. Fact sheet N°310. <http://www.who.int/mediacentre/factsheets/fs310/en/>. online: World Health Organization, (accessed 15 January 2015).

WHO/UNICEF 2015

WHO/UNICEF. Progress on drinking water and sanitation: Joint Monitoring Programme update 2015. http://www.wssinfo.org/fileadmin/user_upload/resources/JMP-Update-report-2015_English.pdf. online: WHO/UNICEF, (accessed 2 February 2015):78.

Wood 2008

Wood L, Egger M, Gluud LL, Schulz KF, Jüni P, Altman DG, et al. Empirical evidence of bias in treatment effect estimates in

controlled trials with different interventions and outcomes: meta-epidemiological study. *BMJ* 2008;**336**(7644):601-5.

Wright 2004

Wright J, Gundry S, Conroy R. Household drinking water in developing countries: a systematic review of microbiological contamination between source and point-of-use. *Tropical Medicine & International Health* 2004;**9**(1):106-17.

Zhang 1998

Zhang J, Yu KF. What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes. *JAMA* 1998;**280**(19):1690-1.

References to other published versions of this review
Clasen 2006

Clasen T, Roberts IG, Rabie T, Schmidt WP, Cairncross S. Interventions to improve water quality for preventing diarrhoea. *Cochrane Database of Systematic Reviews* 2006, Issue 3. [DOI: [10.1002/14651858.CD004794.pub2](https://doi.org/10.1002/14651858.CD004794.pub2)]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES
Characteristics of included studies [ordered by study ID]

Abebe 2014 ZAF

Methods	RCT
Participants	Number: 74 individuals Inclusion criteria: 18 years or older, receiving anti-retroviral therapy for at least 6 months
Interventions	1. Ceramic water filter impregnated with silver nanoparticles
Outcomes	1. Incidence of diarrhoea 2. Water quality 3. Presence of <i>Cryptosporidium</i> in stool
Notes	Location: rural South Africa Length: 12 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Permuted block randomization system.

Interventions to improve water quality for preventing diarrhoea (Review)

Abebe 2014 ZAF (Continued)

Allocation concealment (selection bias)	Low risk	Permuted block randomization system.
Comparability of characteristics	Unclear risk	Not described.
Contemporaneous data collection	Unclear risk	Not described.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	> 20% loss to follow-up.

Alam 1989 BGD

Methods	Quasi-RCT
Participants	Number: 623 children Inclusion criteria: households with children aged 6 to 23 months
Interventions	1. Improved water supply and hygiene education (3 subunits) 2. Primary drinking supply (2 subunits)
Outcomes	1. Incidence of diarrhoea among children aged 6 to 23 months by water source, hygiene practices, and household socioeconomic characteristics
Notes	Location: 5 political subunits in a village in rural Bangladesh Length: 3 years Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Irrelevant to study design.
Allocation concealment (selection bias)	Low risk	Irrelevant to study design.
Comparability of characteristics	Low risk	No substantial differences at baseline.

Interventions to improve water quality for preventing diarrhoea (Review)

Alam 1989 BGD (Continued)

Contemporaneous data collection	Low risk	Data collected at similar points in time.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Irrelevant to study design.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Irrelevant to study design.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Irrelevant to study design.

Austin 1993a GMB

Methods	RCT
Participants	Number: 287 children Inclusion criteria: households with children aged 25 to 60 months (group B) from villages primarily using open, shallow wells for drinking water
Interventions	1. Sodium hypochlorite solution used at household level (11 villages) 2. Primary drinking supply (11 villages)
Outcomes	1. Longitudinal prevalence of diarrhoea 2. Change in nutritional status using weight-for-height Z-score
Notes	Location: 22 rural villages in The Gambia Length: 20 weeks Publication status: PhD dissertation

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number table.
Allocation concealment (selection bias)	Low risk	Numbers assigned to villages.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias)	Low risk	Placebo.

Interventions to improve water quality for preventing diarrhoea (Review)

Austin 1993a GMB (Continued)

All outcomes

Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	89.4% of participants included in analysis.

Austin 1993b GMB

Methods	See Austin 1993a GMB
Participants	Number: 144 children between 6 and 24 months Inclusion criteria: as above
Interventions	As above
Outcomes	As above
Notes	As above

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number table.
Allocation concealment (selection bias)	Low risk	Numbers assigned to villages.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Placebo.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	89.4% of participants included in analysis.

Interventions to improve water quality for preventing diarrhoea (Review)

Boisson 2009 ETH

Methods	RCT
Participants	<p>Number: 196 children under 5, 1516 people, 313 households</p> <p>Inclusion criteria: householders were eligible to participate in the study if (i) at least one member of the household worked away from home during the day in a setting without adequate water supply, and (ii) the household was not already practicing an effective POU water treatment method</p>
Interventions	1. LifeStraw® personal distributed to each household member over the age of six months. A special attachment was given for children under 3
Outcomes	<p>1. Incidence of diarrhoea among young children in the preceding seven days (recorded fortnightly); other health conditions also recorded</p> <p>2. Water quality, flow rate and iodine residual</p> <p>3. Acceptability and use</p>
Notes	<p>Location: rural Oromia, Ethiopia</p> <p>Length: 5 months</p> <p>Publication status: journal</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Lottery used to randomly allocate eligible households into intervention and control groups.
Allocation concealment (selection bias)	Low risk	Lottery used to randomly allocate eligible households into intervention and control groups.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	4% of person-weeks data lost to follow-up.

Boisson 2010 DRC

Methods	RCT
Participants	Number: 190 children under 5, 1144 people, 240 households

Interventions to improve water quality for preventing diarrhoea (Review)

Boisson 2010 DRC (Continued)

Inclusion criteria: unimproved water sources that tested over 1000 thermotolerant coliforms (TTC)/100 ml, reported low use of household water treatment, were easily accessible all year round and were motivated to take part in the project

Interventions	1. LifeStraw® Family filter
Outcomes	1. Incidence of diarrhoea among young children in the preceding seven days (recorded monthly); cough and fever also recorded 2. Filter and water quality monitoring 3. Compliance
Notes	Location: rural eastern province of Kasai, Democratic Republic of Congo Length: 12 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generator.
Allocation concealment (selection bias)	Low risk	"Randomisation was stratified by village and was conducted by the trial manager who played no part in the collection of the data".
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Double-blinded; however filters removed turbidity, so controls were not always successfully blinded.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Double-blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	18.2% person-weeks missing due to families moving out of study area, or not being home at time of visit.

Boisson 2013 IND

Methods	RCT
Participants	Number: 2986 children under 5, 12,454 people, 2163 households Inclusion criteria: households were eligible if there was at least one child under 5, and they lived permanently in the study area
Interventions	1. Sodium dichloroisocyanurate (NaDCC) disinfection tablets

Interventions to improve water quality for preventing diarrhoea (Review)

Boisson 2013 IND (Continued)

Outcomes	<ol style="list-style-type: none"> 1. Longitudinal prevalence of diarrhoea among children under 5 2. Diarrhoea among participants of all ages 3. Weight-for-age z-score, school absenteeism, health care expenditures; adherence; water quality
Notes	Location: informal settlements of Orissa, India Length: 12 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The randomisation list was generated using Stata 10 and was conducted by the trial manager who played no part in the collection of the data".
Allocation concealment (selection bias)	Low risk	"The randomisation list was generated using Stata 10 and was conducted by the trial manager who played no part in the collection of the data".
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"The active and placebo tablets were packaged in identical boxes of three strips containing ten tablets each".
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"The labeling of the boxes was conducted by members of staff who were neither involved in the implementation nor data collection or analysis".
Incomplete outcome data (attrition bias) All outcomes	High risk	12% days of observation lost to follow-up.

Brown 2008a KHM

Methods	RCT
Participants	Number: 239 children under 5, 1196 people, 180 households (across both interventions) Inclusion criteria: households were eligible if they stored drinking water at the household level, if they have at least one child under 5, and if the household was located in the study village
Interventions	<ol style="list-style-type: none"> 1. Iron-rich Cambodian Ceramic Water Purifier 2. Water quality
Outcomes	<ol style="list-style-type: none"> 1. Longitudinal prevalence of diarrhoea for all household members
Notes	Location: rural Kandal Province, Cambodia Length: 18 weeks

Interventions to improve water quality for preventing diarrhoea (Review)

Brown 2008a KHM (Continued)

Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random numbers table.
Allocation concealment (selection bias)	Low risk	Households were approached in group-randomized order.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	2% households lost to follow-up.

Brown 2008b KHM

Methods	See Brown 2008a KHM
Participants	As above
Interventions	1. Cambodian Ceramic Water Purifier
Outcomes	As above
Notes	As above

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random numbers table.
Allocation concealment (selection bias)	Low risk	Households were approached in group-randomized order.
Comparability of characteristics	Low risk	Irrelevant to study design.

Interventions to improve water quality for preventing diarrhoea (Review)

Brown 2008b KHM (Continued)

Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	2% households lost to follow-up.

Chiller 2006 GTM

Methods	RCT
Participants	Number: 3401 persons from 514 households Inclusion criteria: households with at least one child under 1 year
Interventions	1. Flocculant-disinfectant sachets used at household level 2. Primary drinking supply
Outcomes	1. Longitudinal prevalence of diarrhoea (portion of total days of diarrhoea out of total days of observation) among all ages 2. Incidence of persistent diarrhoea
Notes	Location: 42 neighbourhood clusters in 12 rural villages in Guatemala Length: 13 weeks Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generator used to assigned neighbourhoods to intervention or control group.
Allocation concealment (selection bias)	Low risk	Random number generator used to assigned neighbourhoods to intervention or control group.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias)	High risk	No placebo used.

Interventions to improve water quality for preventing diarrhoea (Review)

Chiller 2006 GTM (Continued)

All outcomes

Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Less than 8% of households lost to follow-up.

Clasen 2004b BOL

Methods	RCT
Participants	Number: 324 persons of all ages from 60 households Inclusion criteria: all households in the community
Interventions	1. Household gravity water filter system using imported ceramic filter elements 2. Primary drinking supply
Outcomes	1. Period prevalence of diarrhoea (7-day recall) among all ages 2. Microbial water quality
Notes	Location: rural Bolivian community Length: 9 months Publication status: unpublished

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Households were randomly allocated by names drawn from a hat in a public assembly.
Allocation concealment (selection bias)	Low risk	Households were randomly allocated by names drawn from a hat in a public assembly.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias)	Low risk	No participants lost to follow-up.

Interventions to improve water quality for preventing diarrhoea (Review)

Clasen 2004b BOL (Continued)

All outcomes

Clasen 2004c BOL

Methods	RCT
Participants	Number: 50 households with 280 persons, of which 32 (11%) were under age 5 Inclusion criteria: all households in the community
Interventions	1. Household gravity water filter system using imported ceramic filter elements 2. Primary drinking supply
Outcomes	1. Period prevalence of diarrhoea (7-day recall) among householders assessed at approximately 6-week intervals
Notes	Location: rural Bolivia Length: 6 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Households were randomly allocated by lottery, half to an intervention group and half to a control group.
Allocation concealment (selection bias)	Low risk	Households were randomly allocated by lottery, half to an intervention group and half to a control group.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Less than 1% participants lost to follow-up.

Clasen 2005 COL

Methods	RCT
---------	-----

Interventions to improve water quality for preventing diarrhoea (Review)

Copyright © 2015 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

Clasen 2005 COL (Continued)

Participants	Number: 140 children under 5, 680 people, 140 households Inclusion criteria: all households in the community
Interventions	1. Ceramic water filter
Outcomes	1. Diarrhoea prevalence during previous seven days 2. Water quality
Notes	Location: three rural villages in Colombia Length: six months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Public lottery.
Allocation concealment (selection bias)	Low risk	Lottery conducted at each study site to randomly allocate households.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	5% of households lost to follow-up.

Colford 2002 USA

Methods	RCT
Participants	Number: 236 people from 77 households Inclusion criteria: families were required to own their own homes, use municipal tap water as their main drinking water and have no seriously immunocompromised household members
Interventions	1. Household reverse osmosis filters 2. Primary drinking supply

Interventions to improve water quality for preventing diarrhoea (Review)

Colford 2002 USA (Continued)

Outcomes	<ol style="list-style-type: none"> 1. Incidence of watery diarrhoea 2. Gastrointestinal illness and various other symptoms 3. Water consumption 4. Effectiveness of blinding
Notes	Location: urban community in California, USA Length: 4 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Two random sequences generated to allocated households to intervention or control groups.
Allocation concealment (selection bias)	Low risk	Two random sequences generated to allocated households to intervention or control groups.
Comparability of characteristics	Unclear risk	Irrelevant to study design.
Contemporaneous data collection	Unclear risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	One investigator, not involved in analyses prepared coded labels for the placebo and active devices.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Triple-blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Less than 1% households lost to follow-up.

Colford 2005 USA

Methods	RCT
Participants	Number: 50 HIV+ people, all over 30 years Inclusion criteria: confirmed HIV+ status, uses tap water 75% of the time, no children residing in the home
Interventions	<ol style="list-style-type: none"> 1. Countertop water filtration device
Outcomes	<ol style="list-style-type: none"> 1. Episodes of "highly credible gastrointestinal illness" 2. Diarrhoea episodes calculated
Notes	Location: San Francisco, USA

Interventions to improve water quality for preventing diarrhoea (Review)

Colford 2005 USA *(Continued)*

Length: 12 months

Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer generated random numbers.
Allocation concealment (selection bias)	Low risk	The manufacturer provided a list of device serial numbers and their corresponding active/sham status to facilitate device assignment. All study participants, the study investigators (including clinic personnel and those performing data analysis) and the device installer were blinded throughout the trial as to device assignment.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	All study participants, the study investigators (including clinic personnel and those performing data analysis) and the device installer were blinded throughout the trial as to device assignment.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	All study participants, the study investigators (including clinic personnel and those performing data analysis) and the device installer were blinded throughout the trial as to device assignment.
Incomplete outcome data (attrition bias) All outcomes	Low risk	10% participants withdrew from study (mixed from active and sham devices).

Colford 2009 USA

Methods	Randomized controlled (crossover) trial
Participants	Number: 988 people, 714 households Inclusion criteria: households were eligible if they had one or more persons 55 or older
Interventions	1. Countertop water filtration and UV device
Outcomes	1. Episodes of "highly credible gastrointestinal illness" 2. Diarrhoea episodes calculated
Notes	Location: Sonoma County, USA Length: 13.5 months Publication status: journal

Risk of bias
Interventions to improve water quality for preventing diarrhoea (Review)

Colford 2009 USA (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Households were block-randomized in blocks of 10, with an equal probability of receiving either a sham or an active device.
Allocation concealment (selection bias)	Low risk	Households were block-randomized in blocks of 10, with an equal probability of receiving either a sham or an active device.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	All study staff involved in installation and contact with participants were blinded to device assignments throughout the trial.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Assessors blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	"Among households initially assigned to receive an active device, 89% completed cycle 1 and 83% also completed cycle 2; among households initially assigned to receive a sham device, 90% completed cycle 1 and 82% also completed cycle 2".

Conroy 1996 KEN

Methods	RCT
Participants	Number: 206 Maasai children aged 5 to 16 years in 3 adjoining areas of single province Inclusion criteria: all households in the village
Interventions	1. Solar disinfection in plastic bottles at household level 2. Primary drinking supply
Outcomes	1. Period prevalence of diarrhoea
Notes	Location: single province of rural Kenya Length: 12 weeks Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Interventions assigned by alternate household.
Allocation concealment (selection bias)	High risk	Interventions assigned by alternate household.

Interventions to improve water quality for preventing diarrhoea (Review)

Conroy 1996 KEN (Continued)

Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	No loss to follow-up.

Conroy 1999 KEN

Methods	RCT
Participants	Number: 349 Maasai children < 6 years in 140 households Inclusion criteria: all households in the village
Interventions	1. Solar disinfection in plastic bottles at household level 2. Primary drinking supply
Outcomes	1. Period prevalence of diarrhoea
Notes	Location: rural Kenya Length: 1 year Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Interventions assigned by alternate household.
Allocation concealment (selection bias)	High risk	Interventions assigned by alternative household.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.

Interventions to improve water quality for preventing diarrhoea (Review)

Conroy 1999 KEN (Continued)

Blinding of participants and personnel (performance bias) All outcomes	High risk	Not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	> 20% children lost to follow-up.

Crump 2005a KEN

Methods	RCT
Participants	Number: 6650 persons of all ages in 604 family compounds Inclusion criteria: family compounds with at least 1 child < 2 years and likely to be using highly turbid source water
Interventions	1. Sodium hypochlorite used at household level 2. Primary drinking water supply
Outcomes	1. Longitudinal prevalence (weeks with diarrhoea/weeks of observation) among all ages 2. Breastfeeding and consumption of food and water for children < 2 years 3. Deaths 4. Use of intervention 5. Mothers' knowledge of and acceptance of intervention (weeks 5 and 15) 6. Microbial water quality and turbidity 7. Mothers' knowledge of and attitudes to intervention
Notes	Location: 49 rural villages in western Kenya Length: 20 weeks Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient detail.
Allocation concealment (selection bias)	Unclear risk	Insufficient detail.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.

Interventions to improve water quality for preventing diarrhoea (Review)

Crump 2005a KEN (Continued)

Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	82% participants lost to follow-up.

Crump 2005b KEN

Methods	See Crump 2005a KEN
Participants	As above
Interventions	1. Flocculant-disinfectant sachets used at household level 2. Primary drinking water supply
Outcomes	As above
Notes	As above

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient detail.
Allocation concealment (selection bias)	Unclear risk	Insufficient detail.
Comparability of characteristics	Low risk	Irrelevant to study design,
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	82% participants lost to follow-up.

Interventions to improve water quality for preventing diarrhoea (Review)

Doocy 2006 LBR

Methods	RCT
Participants	Number: 2191 persons of all ages (1138 intervention, 1053 controls), of which 735 are children < 5 (395 intervention, 340 controls) Inclusion criteria: households in settlement area not using treated water for drinking
Interventions	1. Flocculant-disinfectant sachets used at household level, plus water storage vessel 2. Primary drinking supply; also received vessel
Outcomes	1. Longitudinal prevalence (days with diarrhoea/total days of observation) 2. Prevalence of bloody diarrhoea 3. Utilization and acceptability data from exit survey
Notes	Location: Liberian camp for displaced persons Length: 12 weeks Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random division of households by blocks and subsections.
Allocation concealment (selection bias)	Low risk	Households were systematically selected based on their assigned plot number.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	1% of households lost to follow-up.

du Preez 2008 ZAF/ZWE

Methods	RCT
Participants	Number: 115 children < 5 years

Interventions to improve water quality for preventing diarrhoea (Review)

du Preez 2008 ZAF/ZWE (Continued)

Inclusion criteria: households were randomly selected from a list of eligible households from an earlier study: if they had no in-house piped water, and if they had at least one child 12 to 24 months of age

Interventions	<ol style="list-style-type: none"> 1. Household commercial ceramic filter using imported components (60 children) 2. Primary drinking supply (55 children)
Outcomes	<ol style="list-style-type: none"> 1. Incidence of diarrhoea 2. Incidence of bloody diarrhoea and non-bloody diarrhoea 3. Microbiological water quality
Notes	Location: rural South Africa and Zimbabwe Length: 6 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Reported to be randomized, but no description of method of randomization process.
Allocation concealment (selection bias)	Unclear risk	Insufficient detail.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Insufficient detail.

du Preez 2010 ZAF

Methods	RCT
Participants	Number: 824 children, 649 households Inclusion criteria: households were eligible if they had no in-house piped water, and if they had at least one child over 6 months and under 5 years.
Interventions	<ol style="list-style-type: none"> 1. SODIS (438 children) 2. Primary drinking supply (386 children)

Interventions to improve water quality for preventing diarrhoea (Review)

du Preez 2010 ZAF (Continued)

Outcomes	<ol style="list-style-type: none"> 1. Incidence of dysentery 2. Incidence of non-dysentery diarrhoea
Notes	Location: four peri-urban districts of Gauteng Province, South Africa Length: 12 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number table.
Allocation concealment (selection bias)	Low risk	This table was not available to field workers until after the sample frame was drawn up.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	13% of children lost to follow-up.

du Preez 2011 KEN

Methods	RCT
Participants	Number: 1089 children, 765 households Inclusion criteria: eligible households stored water in containers in-house, did not have a drinking water tap in the house or yard, and had at least one child (but not more than 5) between 6 months and 5 years old residing in the house.
Interventions	<ol style="list-style-type: none"> 1. SODIS (404 households) 2. Primary drinking supply (361)
Outcomes	<ol style="list-style-type: none"> 1. Episodes of dysentery and non-dysentery diarrhoea 2. Height-for-age and weight-for-age 3. Microbial water quality
Notes	Location: three urban slums, three rural areas near Nakuru, Kenya\

Interventions to improve water quality for preventing diarrhoea (Review)

du Preez 2011 KEN (Continued)

Length: 17 months

Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random numbers between zero and one were generated and allocated to the households. If the random number allocated to a household was less than 0.5 the household was randomized to the test group. If the allocated number was above 0.5 the house was randomized to the control group.
Allocation concealment (selection bias)	Low risk	Field workers were unaware of how the numbers were allocated.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	4% children lost to follow-up.

Fabiszewski 2012 HND

Methods	RCT
Participants	Number: 230 children < 5, 1020 people, 178 households Inclusion criteria: households were eligible if they had a least one child under 5, did not have year-round access to piped water, and did not use bottled water
Interventions	1. Biosand filter (90 households) 2. Primary drinking supply (86 households)
Outcomes	1. Incidence of diarrhoea 2. Microbial water quality
Notes	Location: 11 rural communities in Copan, Honduras Length: six month follow-up Publication status: journal

Interventions to improve water quality for preventing diarrhoea (Review)

Fabiszewski 2012 HND (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generation.
Allocation concealment (selection bias)	Low risk	No one knew which group they were assigned to until the day before.
Comparability of characteristics	Low risk	Irrelevant to this study design.
Contemporaneous data collection	Low risk	Irrelevant to this study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Less than 1% lost to follow-up.

Gasana 2002 RWA

Methods	Quasi-RCT
Participants	Number: 150 children < 5 years Inclusion criteria: all households with at least one child < 5
Interventions	1. Improved source: pipes to stand post; sedimentation tank; ceramic filter; storage tank; and communal tap (95 children) 2. Primary drinking supply (55 children)
Outcomes	1. Incidence of diarrhoea
Notes	Location: rural Rwanda Length: 24 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Irrelevant to study design.

Interventions to improve water quality for preventing diarrhoea (Review)

Gasana 2002 RWA (Continued)

Allocation concealment (selection bias)	Low risk	Irrelevant to study design.
Comparability of characteristics	Unclear risk	Not described.
Contemporaneous data collection	Unclear risk	Not described.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Irrelevant to study design.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Irrelevant to study design.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Irrelevant to study design.

Gruber 2013 MEX

Methods	RCT
Participants	Number: 1916 people, 444 households Inclusion criteria: households were eligible if they did not have access to centrally treated drinking water and collected water from local sources year-round
Interventions	1. UV water treatment and storage system (Mesita Azul)
Outcomes	1. Diarrhoea prevalence 2. Microbial water quality
Notes	Location: rural Baja California Sur, Mexico Length: 15 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Eligible communities assigned a random number between zero and one by an investigator using STATA.
Allocation concealment (selection bias)	Low risk	Every 2 months another community was randomly allocated to intervention group; no one knew in advance.
Comparability of characteristics	Low risk	Irrelevant to study design.

Interventions to improve water quality for preventing diarrhoea (Review)

Gruber 2013 MEX (Continued)

Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	15% participants lost to follow-up.

Günther 2013 BEN

Methods	RCT
Participants	Number: 364 intervention households; 347 control households Inclusion criteria: all households in intervention villages
Interventions	1. Improved water vessel for fetching 2. Improved water vessel for storing
Outcomes	1. Water quality of stored water 2. Diarrhoea prevalence
Notes	Location: rural Benin Length: 3 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Insufficient detail.
Allocation concealment (selection bias)	Unclear risk	Insufficient detail.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo.

Interventions to improve water quality for preventing diarrhoea (Review)

Günther 2013 BEN (Continued)

Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	64% of sample with follow-up data (due to budgetary constraints).

Handzel 1998 BGD

Methods	RCT
Participants	Number: 447 children aged 3 to 60 months from 276 households Inclusion criteria: households with children 3 to 60 months of age using municipal water (household taps) as primary source of drinking water which had tested positive at baseline for <i>E. coli</i>
Interventions	1. Household chlorination using sodium hypochlorite solution, special storage vessel and hygiene instruction about why and how to treat water (140 households) 2. Primary drinking supply (136 households)
Outcomes	1. Incidence of diarrhoea 2. Microbial water quality
Notes	Location: informal settlement in urban Bangladesh Length: 8 months Publication status: PhD dissertation

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Lottery.
Allocation concealment (selection bias)	Low risk	Consent was obtained from participating households; none knew whether they would be placed into the intervention or comparison group.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.

Interventions to improve water quality for preventing diarrhoea (Review)

Handzel 1998 BGD *(Continued)*

Incomplete outcome data (attrition bias) All outcomes	Low risk	8% participants lost to follow-up.
---	----------	------------------------------------

Jain 2010 GHA

Methods	RCT
Participants	Number: 549 children under five, 3240 individuals, 240 households Inclusion criteria: households were eligible if there was at least one child < 5
Interventions	1. Chlorine (NaDCC) tablets (120 households) 2. Placebo-tablets without chlorine (120 households)
Outcomes	1. Diarrhoeal episodes 2. Chlorine residuals 3. Microbiological water quality
Notes	Location: peri-urban communities of Tamale, Ghana Length: 12 weeks Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number table.
Allocation concealment (selection bias)	Low risk	Only technical staff at Medentech, Ltd knew which tablets were placebo and which were NaDCC.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Triple blinded.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Triple blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Less than 1% of households lost to follow-up.

Interventions to improve water quality for preventing diarrhoea (Review)

Jensen 2003 PAK

Methods	Quasi-RCT
Participants	Number: 226 children < 5 years of age Inclusion criteria: all households that had children aged less than five years and that primarily obtained drinking-water from the water supply systems
Interventions	1. Village level chlorination of water supply using calcium hypochlorite (82 children) 2. Primary drinking supply (144 children)
Outcomes	1. Incidence of diarrhoea 2. Microbial water quality
Notes	Location: 2 villages in Pakistan Length: 6 months Publication status: journal Controlled for sanitation and water storage status of households, and for seasonality

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Irrelevant to study design.
Allocation concealment (selection bias)	Low risk	Irrelevant to study design.
Comparability of characteristics	Low risk	Water quality at baseline significantly different between intervention and control villages.
Contemporaneous data collection	Low risk	Data collected at similar points in time.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Irrelevant to study design.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Irrelevant to study design.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Irrelevant to study design.

Kirchhoff 1985 BRA

Methods	RCT
Participants	Number: 112 persons (all ages) from 20 families

Interventions to improve water quality for preventing diarrhoea (Review)

Kirchhoff 1985 BRA (Continued)

Inclusion criteria: households with at least 2 children living at home and using water from pond exclusively

Interventions	<ol style="list-style-type: none"> 1. Household level chlorination with sodium hypochlorite 2. Primary drinking supply
Outcomes	<ol style="list-style-type: none"> 1. Longitudinal prevalence of diarrhoea 2. Microbial water quality 3. Acceptability of intervention to study population
Notes	Location: rural Brazil Length: 18 weeks Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	Sequences could be related to outcomes (eligible households which agreed to participate were enrolled).
Allocation concealment (selection bias)	High risk	Sequences could be related to outcomes (eligible households which agreed to participate were enrolled).
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Study staff and participants blinded (placebo).
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	Approximately 20% participants lost to follow-up.

Kremer 2011 KEN

Methods	RCT
Participants	Number: 184 springs; 1354 households Inclusion criteria: springs that were not seasonally dry, landowner gave approval to be protected
Interventions	<ol style="list-style-type: none"> 1. Protected springs
Outcomes	<ol style="list-style-type: none"> 1. Diarrhoeal episodes

Interventions to improve water quality for preventing diarrhoea (Review)

Kremer 2011 KEN (Continued)

2. Microbiological water quality

Notes

Location: rural western Kenya

Length: 2 years

Publication status: economics quarterly journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generator assigned springs into year of treatment.
Allocation concealment (selection bias)	Low risk	Random selection of households at each intervention spring.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	95% of all households were surveyed for baseline and at least two follow-up rounds.

Lindquist 2014a BOL

Methods	RCT
Participants	Number: 330 intervention households; 279 control households Inclusion criteria: households: with children less than 60 months of age, in squatter or low-income rental housing, receive their primary drinking/household water from a non-municipal source, and no access to a direct municipal sewer line. Enrollment was limited to one child per household
Interventions	1. Filter
Outcomes	1. Diarrhoea period prevalence
Notes	Location: rural Bolivia Length: 3 months Publication status: journal

Interventions to improve water quality for preventing diarrhoea (Review)

Lindquist 2014a BOL (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generator.
Allocation concealment (selection bias)	Low risk	Randomization done at neighbourhood level.
Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	> 20% lost to follow-up.

Lindquist 2014b BOL

Methods	RCT
Participants	Number: 285 intervention households; 279 control households Inclusion criteria: as above
Interventions	1. Filter 2. WASH behaviour change education
Outcomes	As above
Notes	As above

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generator.
Allocation concealment (selection bias)	Low risk	Randomization done at neighbourhood level.

Interventions to improve water quality for preventing diarrhoea (Review)

Lindquist 2014b BOL (Continued)

Comparability of characteristics	Low risk	Irrelevant to study design.
Contemporaneous data collection	Low risk	Irrelevant to study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	> 20% lost to follow-up.

Luby 2004a PAK

Methods	Quasi-RCT
Participants	Number: 2365 persons < 15 years from 285 households Inclusion criteria: eligible households included at least one child less than five years of age and two children less than 15 years of age, had sufficient water supply for the children to bathe daily, and planned to continue to reside in their homes for at least the ensuing four months.
Interventions	1. Bleach + regular vessel (640 people) 2. Primary drinking supply (1027 people)
Outcomes	1. Longitudinal prevalence of diarrhoea 2. Use of intervention by certain household characteristics
Notes	Location: 3 neighbourhoods in squatter settlements in Karachi, Pakistan Length: 6 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Irrelevant for study design.
Allocation concealment (selection bias)	Low risk	Irrelevant for study design.
Comparability of characteristics	Low risk	Baseline characteristics did not differ significantly between groups.
Contemporaneous data collection	Low risk	Data collected at similar points in time.

Interventions to improve water quality for preventing diarrhoea (Review)

Luby 2004a PAK (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	Irrelevant for study design.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Irrelevant for study design.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Irrelevant for study design.

Luby 2004b PAK

Methods	See Luby 2004a PAK
Participants	As above
Interventions	1. Bleach + insulated vessel (697 people) 2. Primary drinking supply (1027 people)
Outcomes	As above
Notes	As above

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Irrelevant for study design.
Allocation concealment (selection bias)	Low risk	Irrelevant for study design.
Comparability of characteristics	Low risk	No substantial differences at baseline.
Contemporaneous data collection	Low risk	Data collected at similar points in time.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Irrelevant for study design.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Irrelevant for study design.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Irrelevant for study design.

Interventions to improve water quality for preventing diarrhoea (Review)

Luby 2006a PAK

Methods	RCT
Participants	Number: 5520 persons of all ages Inclusion criteria: running water at least one hour twice a week and at least one child under 5
Interventions	1. Dilute bleach + vessel (1747 people) 2. Primary drinking supply (1852 people)
Outcomes	1. Incidence and longitudinal prevalence of diarrhoea
Notes	Location: 47 squatter settlements of Karachi, Pakistan Length: 8 months Publication status: unpublished

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated random number assigned households to groups.
Allocation concealment (selection bias)	Low risk	Households consented to study before computer randomly assigned them to specific groups.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Overall less than 8% of participants lost to follow-up (averaged across all groups).

Luby 2006b PAK

Methods	See Luby 2006a PAK
Participants	As above
Interventions	1. Flocculant-disinfectant + soap (1806 in flocculant-disinfection group)

Interventions to improve water quality for preventing diarrhoea (Review)

Luby 2006b PAK (Continued)

2. Primary drinking supply (1852 people)

Outcomes	As above
Notes	As above

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated random number assigned households to groups.
Allocation concealment (selection bias)	Low risk	Households consented to study before computer randomly assigned them to specific groups.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Overall less than 8% of participants lost to follow-up (averaged across all groups).

Luby 2006c PAK

Methods	See Luby 2006a PAK
Participants	As above
Interventions	<ol style="list-style-type: none"> 1. Flocculant-disinfectant + vessel (1833 in flocculant-disinfection group) 2. Primary drinking supply (1852 people, 40.0%)
Outcomes	As above
Notes	As above

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated random number assigned households to groups.

Interventions to improve water quality for preventing diarrhoea (Review)

Luby 2006c PAK (Continued)

Allocation concealment (selection bias)	Low risk	Households consented to study before computer randomly assigned them to specific groups.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Overall less than 8% of participants lost to follow-up (averaged across all groups).

Lule 2005 UGA

Methods	RCT
Participants	Number: 2201 persons of all ages among 458 households Inclusion criteria: households without access to chlorinated municipal water; at least 1 resident of each household was HIV+
Interventions	1. Household level chlorination using sodium hypochlorite + special vessel (1097 people) 2. Primary drinking supply (1104 people) Note: hygiene education was provided to both groups
Outcomes	1. Incidence of diarrhoea 2. Days with diarrhoea (longitudinal prevalence) 3. Days lost from work or school 4. Aetiology of diarrhoea 5. Frequency of clinic visits and hospitalization 6. Mortality
Notes	Location: households in rural Uganda Length: 5 months Publication status: unpublished Succeeded by 18-month RCT that included cotrimoxazole prophylaxis

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Interventions to improve water quality for preventing diarrhoea (Review)

Lule 2005 UGA (Continued)

Random sequence generation (selection bias)	Unclear risk	Insufficient detail.
Allocation concealment (selection bias)	Unclear risk	Insufficient detail.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Less than 8% of participants lost to follow-up.

Mahfouz 1995 KSA

Methods	RCT
Participants	Number: 311 children < 5 years (among intervention households, among controls) among 171 families Inclusion criteria: households with at least one child less than 5 years of age
Interventions	1. Household level chlorination using calcium hypochlorite (159 children) 2. Primary drinking supply (152 children)
Outcomes	1. Reported cases of diarrhoea in intervention year compared with previous year
Notes	Location: 9 villages in rural Saudi Arabia Length: 6 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No description of randomization process (for villages). No description of how households were chosen.
Allocation concealment (selection bias)	Unclear risk	No description of how chosen families were selected or contacted
Comparability of characteristics	Low risk	Irrelevant for study design.

Interventions to improve water quality for preventing diarrhoea (Review)

Mahfouz 1995 KSA (Continued)

Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded
Incomplete outcome data (attrition bias) All outcomes	High risk	Large loss to follow-up in intervention and control groups

Majuru 2011 ZAF

Methods	Quasi-RCT
Participants	Number: community 1, 234 individuals; community 2, 173 individuals; reference community, 146 individuals Inclusion criteria: new community level piped water supply
Interventions	1. Community-level piped water supply (2 communities, 407 individuals) 2. Primary drinking water supply, unimproved sources (1 community, 146 individuals)
Outcomes	1. Diarrhoeal episodes
Notes	Location: rural, remote communities, Limpopo Province, South Africa Length: approximately 10 months of follow-up Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Irrelevant for study design.
Allocation concealment (selection bias)	Low risk	Irrelevant for study design.
Comparability of characteristics	Low risk	No substantial differences at baseline.
Contemporaneous data collection	Low risk	Data collected at similar points in time.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Irrelevant for study design.

Interventions to improve water quality for preventing diarrhoea (Review)

Majuru 2011 ZAF (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Irrelevant for study design.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Irrelevant for study design.

McGuigan 2011 KHM

Methods	RCT
Participants	Number: 964 children in 782 households Inclusion criteria: households were eligible if they were permanent residents in the area, had at least one child 6 months to 5 years old, and did not use other methods of household water treatment
Interventions	1. SODIS (407 households, 502 children < 5) 2. Primary drinking water supply (375 households, 426 children < 5)
Outcomes	1. Days of dysentery diarrhoea for < 5s 2. Days of non-dysentery diarrhoea for < 5s
Notes	Location: rural communities in Prey Veng and Svey Rieng provinces, Cambodia Length: 12 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomized raffle system of interested households during initial meeting.
Allocation concealment (selection bias)	Low risk	Households were randomly allocated to intervention or control groups at community meeting.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessor not blinded.
Incomplete outcome data (attrition bias)	Low risk	5% of participants had less than 10 months of follow-up.

Interventions to improve water quality for preventing diarrhoea (Review)

McGuigan 2011 KHM (Continued)

All outcomes

Mengistie 2013 ETH

Methods	RCT
Participants	Number: 36 clusters, 569 households, 845 children < 5 Inclusion criteria: households were eligible if they had at least one child < 5
Interventions	Chlorine disinfection (WaterGuard) (427 children < 5) Primary drinking supply (422 children < 5)
Outcomes	Diarrhoeal episodes for children < 5 Intervention compliance
Notes	Location: rural communities, Kersa district, Ethiopia Length: 16 weeks Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer generated random sample.
Allocation concealment (selection bias)	Low risk	Randomization of clusters done in community meeting.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	2% to 3% of person-weeks of observation lost.

Interventions to improve water quality for preventing diarrhoea (Review)

Mäusezhal 2009 BOL

Methods	RCT
Participants	Number: 484 households, 819 children < 5 Inclusion criteria: communities had to have at least 30 children < 5 and rely on contaminated drinking water sources
Interventions	1. SODIS (11 communities, 262 households, 441 children) 2. Primary drinking water supply, unimproved sources (11 communities, 222 households, 378 children)
Outcomes	1. Diarrhoeal episodes for children < 5 2. Dysentery episodes for children < 5
Notes	Location: rural Totora District, Cochabamba Department, Bolivia Length: 12 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random assignment during public event.
Allocation concealment (selection bias)	Low risk	Balls with community codes inscribed on them were drawn from a box; the first ball drawn would be the intervention community.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	21% of person-days of observation missing.

Opryszko 2010a AFG

Methods	RCT
Participants	Number: 553 households, 4507 individuals Inclusion criteria: inadequate access to improved water sources; high areas of diarrhoeal disease according to 2004 census

Interventions to improve water quality for preventing diarrhoea (Review)

Opryszko 2010a AFG (Continued)

Interventions	<ol style="list-style-type: none"> 1. Chlorine disinfection (with improved storage vessel); Improved water supply (tube wells); hygiene promotion (261 households, 1958 individuals) 2. Primary drinking supply (292 households, 2549 individuals)
Outcomes	<ol style="list-style-type: none"> 1. Diarrhoea prevalence 2. Dysentery-diarrhoea prevalence
Notes	Location: rural communities, Wardak province, Afghanistan Length: 16 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomly allocated.
Allocation concealment (selection bias)	Low risk	Randomly allocated by numbered lists.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	10% of households data missing at follow-up.

Opryszko 2010b AFG

Methods	See Opryszko 2010a AFG
Participants	Number: 600 households, 4,966 individuals Inclusion criteria: inadequate access to improved water sources; high areas of diarrhoeal disease according to 2004 census
Interventions	<ol style="list-style-type: none"> 1. Improved water supply (tube wells) 2. Primary drinking supply (292 households, 2549 individuals)
Outcomes	As above

Interventions to improve water quality for preventing diarrhoea (Review)

Opryszko 2010b AFG (Continued)

Notes As above

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomly allocated.
Allocation concealment (selection bias)	Low risk	Randomly allocated by numbered lists.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	10% of households data missing at follow-up.

Opryszko 2010c AFG

 Methods See [Opryszko 2010a AFG](#)

 Participants Number: 591 households, 4575 individuals
 Inclusion criteria: inadequate access to improved water sources; high areas of diarrhoeal disease according to 2004 census

 Interventions

1. Chlorine disinfection (Clorin); Improved storage vessel (299 households, 2026 individuals)
2. Primary drinking supply (292 households, 2549 individuals)

Outcomes As above

Notes As above

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomly allocated.

Interventions to improve water quality for preventing diarrhoea (Review)

Opryszko 2010c AFG (Continued)

Allocation concealment (selection bias)	Low risk	Randomly allocated by numbered lists.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	10% of households data missing at follow-up.

Patel 2012 KEN

Methods	RCT
Participants	Number: 42 schools Inclusion criteria: schools were eligible if they were not near urban centres and did not have pre-existing water-treatment promotion activities
Interventions	1. Chlorine disinfection (WaterGuard); improved vessel (22 schools) 2. Primary drinking supply (20 schools)
Outcomes	1. Student's knowledge and practice of using WaterGuard 2. Any illness 3. Diarrhoeal illness 4. Acute respiratory illness
Notes	Location: rural Nyanza province, Kenya Length: 2 years Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random allocation from census list.
Allocation concealment (selection bias)	Low risk	Random allocation from census list.

Interventions to improve water quality for preventing diarrhoea (Review)

Patel 2012 KEN (Continued)

Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	32% students lost to follow-up.

Peletz 2012 ZMB

Methods	RCT
Participants	Number: 120 households, 599 individuals, 121 children < 2 Inclusion criteria: mothers who disclosed their HIV status, had a child 6-12 months old, and permanently resided in the catchment area
Interventions	1. Filter (LifeStraw® Family); two 5 L storage vessels (61 households, 299 individuals, 61 children < 2) 2. Primary drinking supply (59 households, 300 individuals, 60 children < 2)
Outcomes	1. Use of filter 2. Microbiological water quality 3. Longitudinal diarrhoeal prevalence 4. Weight-for-age Z-scores
Notes	Location: two peri-urban neighbourhoods, Chongwe district, Zambia Length: 12 month Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generator.
Allocation concealment (selection bias)	Low risk	Randomization conducted by person not involved in study.
Comparability of characteristics	Low risk	Irrelevant for study design.

Interventions to improve water quality for preventing diarrhoea (Review)

Peletz 2012 ZMB (Continued)

Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	More than 80% of person-weeks of observation completed.

Quick 1999 BOL

Methods	RCT
Participants	Number: 791 persons of all ages from 127 households Inclusion criteria: all households in the community
Interventions	1. Household level chlorination + vessel + hygiene education (400 people, 64 households) 2. Primary drinking supply (391 people, 63 households)
Outcomes	1. Mean episodes of diarrhoea per person 2. Microbiological water quality
Notes	Location: 2 peri-urban communities in Bolivia Length: 5 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomized by public lottery into two groups.
Allocation concealment (selection bias)	Low risk	Randomized by public lottery into two groups.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.

Interventions to improve water quality for preventing diarrhoea (Review)

Quick 1999 BOL (Continued)

Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Less than 10% of participants lost to follow-up.

Quick 2002 ZMB

Methods	Quasi-RCT
Participants	Number: 1584 persons of all ages from 260 households Inclusion criteria: lack of piped water and presence of health clinic in community
Interventions	1. Household level chlorination + vessel + hygiene education (166 households) 2. Primary drinking supply (94 households)
Outcomes	1. Incidence of diarrhoea 2. Microbiological water quality
Notes	Location: 2 peri-urban communities in Zambia Length: 3 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Irrelevant for study design.
Allocation concealment (selection bias)	Low risk	Irrelevant for study design.
Comparability of characteristics	Low risk	No substantial differences at baseline.
Contemporaneous data collection	Low risk	Data collected at similar points in time.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Irrelevant for study design.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Irrelevant for study design.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Irrelevant for study design.

Interventions to improve water quality for preventing diarrhoea (Review)

Reller 2003a GTM

Methods	RCT
Participants	Number: 492 households Inclusion criteria: household with a child < 12 months or mother in last trimester of pregnancy
Interventions	1. Flocculant-disinfectant (102 households) 2. Primary drinking supply (96 households)
Outcomes	1. Incidence of diarrhoea 2. Intervention knowledge and acceptability 3. Microbiological water quality 4. Intervention utilization
Notes	Location: 12 villages in rural Guatemala Length: 12 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generator assigned eligible households to groups.
Allocation concealment (selection bias)	Low risk	Random number generator assigned eligible households to groups.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	Approximately 13% of participants lost to follow-up.

Reller 2003b GTM

Methods	See Reller 2003a GTM
---------	--------------------------------------

Interventions to improve water quality for preventing diarrhoea (Review)

Reller 2003b GTM (Continued)

Participants	As above
Interventions	1. Bleach only (97 households) 2. Primary drinking supply (as above)
Outcomes	As above
Notes	As above

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generator assigned eligible households to groups.
Allocation concealment (selection bias)	Low risk	Random number generator assigned eligible households to groups.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	Approximately 13% of participants lost to follow-up.

Reller 2003c GTM

Methods	See Reller 2003a GTM
Participants	As above
Interventions	1. Bleach + vessel (97 households) 2. Primary drinking supply (as above)
Outcomes	As above
Notes	As above

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Interventions to improve water quality for preventing diarrhoea (Review)

Reller 2003c GTM (Continued)

Random sequence generation (selection bias)	Low risk	Random number generator assigned eligible households to groups.
Allocation concealment (selection bias)	Low risk	Random number generator assigned eligible households to groups.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	Approximately 13% of participants lost to follow-up.

Reller 2003d GTM

Methods	See Reller 2003a GTM
Participants	As above
Interventions	<ol style="list-style-type: none"> 1. Flocculant-disinfectant + vessel (100 households) 2. Primary drinking supply (as above)
Outcomes	As above
Notes	As above

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generator assigned eligible households to groups.
Allocation concealment (selection bias)	Low risk	Random number generator assigned eligible households to groups.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.

Interventions to improve water quality for preventing diarrhoea (Review)

Reller 2003d GTM (Continued)

Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	Approximately 13% of participants lost to follow-up.

Roberts 2001 MWI

Methods	RCT
Participants	Number: 1160 persons of all ages; of these, 208 were children < 5 years Inclusion criteria: all households in refugee camp
Interventions	1. Improved storage: bucket with spout and narrow opening to limit hand entry (310 people including 51 children, 100 households) 2. Primary drinking supply (850 people including 157 children, 300 households)
Outcomes	1. Incidence of diarrhoea 2. Microbiological water quality 3. Incidence of diarrhoea by selected environmental factors
Notes	Location: Malawi refugee camp Length: 4 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	High risk	"One fourth of the interviewed households were selected at random to receive the improved buckets".
Allocation concealment (selection bias)	High risk	"One fourth of the interviewed households were selected at random to receive the improved buckets".
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.

Interventions to improve water quality for preventing diarrhoea (Review)

Roberts 2001 MWI (Continued)

Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	High risk	88.8% of participants lost to follow-up.

Rodrigo 2011 AUS

Methods	RCT
Participants	Number: 300 households, 1352 individuals, 185 children < 5 Inclusion criteria: households were eligible if they use untreated rainwater as their primary drinking source
Interventions	1. Water filters (Freshwater systems) (152 households, 698 individuals) 2. Sham-water filters (148 households, 654 individuals)
Outcomes	1. Episodes of Highly Credible Gastroenteritis 2. Episodes of diarrhoea
Notes	Location: Adelaide, Australia Length: 12 months Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number sequence by independent researcher.
Allocation concealment (selection bias)	Low risk	Random number sequence by independent researcher.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Sham device (placebo) utilised.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Assessors not blinded.
Incomplete outcome data (attrition bias)	High risk	31% households lost to follow-up.

Interventions to improve water quality for preventing diarrhoea (Review)

Rodrigo 2011 AUS (Continued)

All outcomes

Semenza 1998 UZB

Methods	RCT
Participants	Number and inclusion criteria: 1583 persons of all ages from 240 households, half with access to piped water (first control group) and half without (of which 62 received intervention, and 58 served as a second control group); these included 344 children < 5
Interventions	<ol style="list-style-type: none"> 1. Household level chlorination + vessel + hygiene education 2. Primary drinking supply
Outcomes	<ol style="list-style-type: none"> 1. Incidence of diarrhoea 2. Incidence of diarrhoea by selected household and water management practices
Notes	Location: urban Uzbekistan Length: 9.5 weeks Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Households randomly selected from map of neighbourhoods.
Allocation concealment (selection bias)	Low risk	Households randomly selected from map of neighbourhoods.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Lost to follow-up not discussed.

Interventions to improve water quality for preventing diarrhoea (Review)

Stauber 2009 DOM

Methods	RCT
Participants	Number: 167 households, 907 individuals, 243 children < 5 Inclusion criteria: households were eligible if there was no biosand filter in the house, and there was at least one child < 5
Interventions	1. Biosand filter (81 households, 447 individuals) 2. Primary drinking supply (86 households, 460 individuals)
Outcomes	1. Diarrhoeal incidence 2. Microbiological water quality
Notes	Location: one semi-rural and one urban community, Bona0, Dominican Republic Length: six months follow-up Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generation assigned 50% of households to intervention group.
Allocation concealment (selection bias)	Unclear risk	Households were unaware of whether they would be assigned to the intervention or control group until one week before BSF installation, but it is not clear whether this was foreknowledge of group assignment.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	7% participants lost to follow-up.

Stauber 2012a KHM

Methods	RCT
Participants	Number: 189 households, 1147 individuals, 242 children < 5 Inclusion criteria: households were eligible if there was at least one child < 5

Interventions to improve water quality for preventing diarrhoea (Review)

Stauber 2012a KHM (Continued)

Interventions	<ol style="list-style-type: none"> 1. Plastic Biosand filter (7 villages, 90 households, 546 individuals) 2. Primary drinking supply (6 villages, 99 households, 601 individuals)
Outcomes	<ol style="list-style-type: none"> 1. Diarrhoeal incidence 2. Microbiological water quality
Notes	Location: 13 rural communities, Angk Snoul district, Cambodia Length: four months follow-up Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generation assigned 7 of 13 villages to intervention group.
Allocation concealment (selection bias)	Low risk	All villages were told they would not know to which group they were assigned until halfway through the study (due to surveillance period, pre-intervention).
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	4% of person-observation weeks missing.

Stauber 2012b GHA

Methods	RCT
Participants	Number: 2043 individuals, of which 440 were children < 5, from 260 households Inclusion criteria: households were eligible if there was at least one child < 5
Interventions	<ol style="list-style-type: none"> 1. Plastic Biosand filter (117 households, 1012 individuals) 2. Primary drinking supply (143 households, 1031 individuals)
Outcomes	<ol style="list-style-type: none"> 1. Diarrhoeal incidence 2. Microbiological water quality
Notes	Location: six rural communities, Tamale, Ghana

Interventions to improve water quality for preventing diarrhoea (Review)

Stauber 2012b GHA (Continued)

Length: three months follow-up

Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number generator assigned 3 of the 6 villages to the intervention group.
Allocation concealment (selection bias)	Unclear risk	Not discussed.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Less than 3% of households lost to follow-up.

Tiwari 2009 KEN

Methods	RCT
Participants	Number: 387 individuals, of which 114 were children < 5, from 60 households Inclusion criteria: households were eligible if they had at least one child < 3, used river water as their primary or secondary drinking water source, stable residence for next 12 months, and indicators of lower socio-economic status
Interventions	1. Biosand filter (30 households, 118 children) 2. Primary drinking water supply (30 households, 104 children)
Outcomes	1. Microbiological water quality 2. Diarrhoea prevalence in children
Notes	Location: rural households in River Njoro watershed, Nakuru and Molo districts, Kenya Length: six months Publication status: journal

Risk of bias
Interventions to improve water quality for preventing diarrhoea (Review)

Tiwari 2009 KEN (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	No description of randomization process.
Allocation concealment (selection bias)	Unclear risk	No description of steps to conceal allocation.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	After randomization, 75 (93%) and 79 (92%) of BSF and control households, respectively, completed the study.

Torun 1982 GTM

Methods	Quasi-RCT
Participants	Number: 2103 persons of all ages from 2 villages Inclusion criteria: all households within 2 villages
Interventions	1. Source protection (spring), chlorination facilities, "adequate storage", and water mains with faucets to yards of intervention village (1006 people) 2. Primary drinking supply (1097 people)
Outcomes	1. Incidence of diarrhoea
Notes	Location: 2 small villages in Guatemala Length: 12 months Publication status: book

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Irrelevant for study design.
Allocation concealment (selection bias)	Low risk	Irrelevant for study design.

Interventions to improve water quality for preventing diarrhoea (Review)

Torun 1982 GTM (Continued)

Comparability of characteristics	Low risk	No substantial differences at baseline.
Contemporaneous data collection	Low risk	Data collected at similar points in time.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Irrelevant for study design.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Irrelevant for study design.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Irrelevant for study design.

URL 1995a GTM

Methods	RCT
Participants	<p>Number: 1120 children < 5 years (265 and 289 allocated to the water quality intervention arms, 297 to an education only arm, and 269 to the control arm) from 680 families from three demographic regions</p> <p>Inclusion criteria: households must have children <5 and have indicators of low socio-economic status and microbiological contamination of water source</p>
Interventions	<ol style="list-style-type: none"> 1. Locally fabricated ceramic filters (265 children or 23.6%) 2. Primary drinking supply (269 children)
Outcomes	<ol style="list-style-type: none"> 1. Incidence of diarrhoea 2. Nutritional status (weight/age)
Notes	<p>Location: 3 demographic regions of Guatemala</p> <p>Length: 12 months</p> <p>Publication status: unpublished</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Reported to be randomized, but no description of method of randomization process.
Allocation concealment (selection bias)	Unclear risk	No description of allocation concealment.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.

Interventions to improve water quality for preventing diarrhoea (Review)

URL 1995a GTM *(Continued)*

Blinding of participants and personnel (performance bias) All outcomes	High risk	Participants not blinded.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Not discussed.

URL 1995b GTM

Methods	See URL 1995a GTM
Participants	As above
Interventions	1. Locally fabricated ceramic filters + hygiene education 2. Primary drinking supply (as above)
Outcomes	As above
Notes	As above

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Reported to be randomized, but no description of method of randomization process.
Allocation concealment (selection bias)	Unclear risk	No description of allocation concealment.
Comparability of characteristics	Low risk	Irrelevant for study design.
Contemporaneous data collection	Low risk	Irrelevant for study design.
Blinding of participants and personnel (performance bias) All outcomes	High risk	No placebo used.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Assessors not blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Not discussed.

Interventions to improve water quality for preventing diarrhoea (Review)

Xiao 1997 CHN

Methods	Quasi-RCT
Participants	Number: 4649 people of all ages Inclusion criteria: all households within villages
Interventions	1. Improved water supply + sanitation + hygiene education (2363 people) 2. Primary drinking supply (2286 people)
Outcomes	1. Incidence of diarrhoea
Notes	Location: 2 villages in rural China Length: 3 years Publication status: journal

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Irrelevant for study design.
Allocation concealment (selection bias)	Low risk	Irrelevant for study design.
Comparability of characteristics	Low risk	No substantial differences at baseline.
Contemporaneous data collection	Low risk	Data collected at similar points in time.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Irrelevant for study design.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Irrelevant for study design.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Irrelevant for study design.

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Ahojyo 2011	Allocation was neither randomized nor quasi-randomized.
Aiken 2011	Allocation was neither randomized nor quasi-randomized.

Interventions to improve water quality for preventing diarrhoea (Review)

Study	Reason for exclusion
Alexander 2013	Outcome measures did not include diarrhoea.
Arnold 2009	Allocation was neither randomized nor quasi-randomized.
Arnold 2012a	Comment paper.
Arnold 2013	Design paper.
Asaolu 2002	Allocation was neither randomized nor quasi-randomized; outcome measures did not include diarrhoea.
Aziz 1990 BGD	The intervention included the provision of sanitation facilities.
Azurin 1974	Outcome measures did not include diarrhoea.
Bahl 1976	Allocation was neither randomized nor quasi-randomized.
Bajer 2012	Allocation was neither randomized nor quasi-randomized, outcome not diarrhoea.
Barreto 2007	Allocation was neither randomized nor quasi-randomized.
Barzilay 2011	Allocation was neither randomized nor quasi-randomized.
Bersh 1985	Allocation was neither randomized nor quasi-randomized.
Boubacar 2014	Allocation was neither randomized nor quasi-randomized.
Brown 2012a	Modelling paper.
Capuno 2011	Allocation was neither randomized nor quasi-randomized.
Cavallaro 2011	Allocation was neither randomized nor quasi-randomized, outcomes did not include diarrhoea.
Chang 2012	Outcomes did not include diarrhoea.
Chongsuvivatwong 1994	Allocation was neither randomized nor quasi-randomized.
Christen 2011	Allocation was neither randomized nor quasi-randomized, outcomes did not include diarrhoea.
Clasen 2012	No water quality intervention.
Colford 2005	Outcomes did not include diarrhoea.
Colwell 2003	Outcomes did not include diarrhoea.
Conroy 2001	Outcomes did not include diarrhoea.
Coulliette 2013	Allocation was neither randomized nor quasi-randomized, outcomes did not include diarrhoea.
Crump 2007	Allocation was neither randomized nor quasi-randomized, outcomes did not include diarrhoea.
Davis 2011	Outcomes did not include diarrhoea.
Deb 1986	Outcomes did not include diarrhoea.

Study	Reason for exclusion
Denslow 2010	Allocation was neither randomized nor quasi-randomized.
Devoto 2011	Intervention did not affect water quality.
Dorevitch 2011	Outcomes did not include diarrhoea.
Dreibelbis 2014a KEN	School-based study.
Dreibelbis 2014b KEN	School-based study.
Dreibelbis 2014c KEN	School-based study.
du Preez 2012	Response to comments.
Eisenberg 2006	Study on risk assessment.
Enger 2012	Modelling paper.
Esrey 1988	Allocation was neither randomized nor quasi-randomized.
Fewtrell 1994	Allocation was neither randomized nor quasi-randomized, outcomes did not include diarrhoea.
Fewtrell 1997	Allocation was neither randomized nor quasi-randomized, outcomes did not include diarrhoea.
Firth 2010	Outcomes did not include diarrhoea.
Fisher 2011	Allocation was neither randomized nor quasi-randomized.
Freeman 2012	Outcomes did not include diarrhoea.
Freeman 2014a KEN	School-based study.
Freeman 2014b KEN	School-based study.
Freeman 2014c KEN	School-based study.
Fry 2010	Modelling paper.
Galiani 2009	Allocation was neither randomized nor quasi-randomized
Garrett 2008 KEN	The intervention included the provision of sanitation facilities.
Ghannoum 1981	Allocation was neither randomized nor quasi-randomized, outcomes did not include diarrhoea.
Gorelick 2011	Allocation was neither randomized nor quasi-randomized.
Greene 2012	Outcome not diarrhoea, see Freeman 2012 .
Gómez-Couso 2012	Allocation was neither randomized nor quasi-randomized, outcome not diarrhoea.
Habib 2013	Water quality intervention applied once children had experienced diarrhoea.
Harris 2009	Allocation was neither randomized nor quasi-randomized.
Harshfield 2012	Allocation was neither randomized nor quasi-randomized.

Interventions to improve water quality for preventing diarrhoea (Review)

Study	Reason for exclusion
Hartinger 2011	Design paper.
Hartinger 2012	Outcome measures did not include diarrhoea.
Hellard 2001	Outcome measures did not include diarrhoea.
Hoque 1996	Allocation was neither randomized nor quasi-randomized.
Huda 2012	Allocation was neither randomized nor quasi-randomized.
Hunter 2010	Allocation was neither randomized nor quasi-randomized
Iijima 2001	Allocation was neither randomized nor quasi-randomized.
Islam 2011	Allocation was neither randomized nor quasi-randomized.
Jensen 2002	Outcome not diarrhoea.
Kariuki 2012	Intervention not water.
Karon 2011	Outcome not diarrhoea.
Keraita 2007	Outcome not diarrhoea.
Khan 1984	Outcome not diarrhoea.
Luby 2008	Allocation was neither randomized nor quasi-randomized.
Luoto 2011	Outcome not diarrhoea.
Luoto 2012	Outcome not diarrhoea.
Macy 1998	Allocation was neither randomized nor quasi-randomized; intervention not an improvement in water quality; outcome not diarrhoea.
McCabe 1957	Intervention not an improvement in water quality.
Mertens 1990	Allocation was neither randomized nor quasi-randomized, intervention not an improvement in water quality; outcome not diarrhoea.
Messou 1997	The intervention included the provision of sanitation facilities.
Mäusezahl 2003	Allocation was neither randomized nor quasi-randomized.
Nanan 2003	Allocation was neither randomized nor quasi-randomized.
Nerkar 2014	Allocation was neither randomized nor quasi-randomized.
Nnane 2011	Allocation was neither randomized nor quasi-randomized, no intervention.
Oluyeye 2011	Allocation was neither randomized nor quasi-randomized, no intervention.
Palit 2012	Allocation was neither randomized nor quasi-randomized.
Pavlinac 2014	Allocation was neither randomized nor quasi-randomized.

Interventions to improve water quality for preventing diarrhoea (Review)

Study	Reason for exclusion
Payment 1991a	Allocation was neither randomized nor quasi-randomized, outcomes did not include diarrhoea.
Payment 1991b	Outcomes did not include diarrhoea.
Peletz 2013	Outcomes did not include diarrhoea.
Pinfold 1990	Intervention not an improvement in water quality; outcome not diarrhoea.
Psutka 2012	Allocation was neither randomized nor quasi-randomized, outcomes did not include diarrhoea.
Rosa 2014	Outcomes did not include diarrhoea.
Rose 2006	Allocation was neither randomized nor quasi-randomized.
Rubenstein 1969	Allocation was neither randomized nor quasi-randomized.
Russo 2012	Allocation was neither randomized nor quasi-randomized.
Sathe 1996	Allocation was neither randomized nor quasi-randomized.
Shah 2012	Review paper.
Sharan 2011	Allocation was neither randomized nor quasi-randomized, outcome not diarrhoea.
Sheth 2010	Allocation was neither randomized nor quasi-randomized, outcome not diarrhoea.
Shiffman 1978	Allocation was neither randomized nor quasi-randomized.
Shrestha 2006	Cost-effectiveness paper.
Shum 1971	Allocation was neither randomized nor quasi-randomized, intervention not an improvement in water quality; outcome not diarrhoea.
Sima 2012	Allocation was neither randomized nor quasi-randomized.
Sorvillo 1994	Outcomes did not include diarrhoea.
Stauber 2013	Outcomes did not include diarrhoea.
Sutha 2011	Review paper.
Tonglet 1992	Allocation was neither randomized nor quasi-randomized.
Trivedi 1971	Allocation was neither randomized nor quasi-randomized.
VanDerslice 1995	Allocation was neither randomized nor quasi-randomized, intervention not an improvement in water quality.
Varghese 2002	Allocation was neither randomized nor quasi-randomized.
Wiedenmann 2006	Intervention not an improvement in water quality.
Wolf 2014	Review.
Wood 2012	Qualitative study.

Interventions to improve water quality for preventing diarrhoea (Review)

Study	Reason for exclusion
Wu 2011	Allocation was neither randomized nor quasi-randomized.

Characteristics of ongoing studies [ordered by study ID]

Chlorination, Dhaka

Trial name or title	Impact of Low-Cost In-Line Chlorination Systems in Urban Dhaka on Water Quality and Child Health
Methods	RCT
Participants	All poor households, with at least one child under five, that access one of 160 studied shared water points in Dhaka.
Interventions	In-line chlorination
Outcomes	Water quality, diarrhoea in children under five, weight of children, cost of instilling and maintaining system, hospital visits, health care expenditures, other household expenditures
Starting date	Early 2015
Contact information	
Notes	Funded by SIEF, World Bank

WASH-B, Bangladesh

Trial name or title	WASH Benefits Bangladesh: A Cluster Randomized Controlled Trial of the Benefits of Water, Sanitation, Hygiene Plus Nutrition Interventions on Child Growth
Methods	Parallel, cluster-RCT
Participants	Estimated enrolment: 5040
Interventions	<ol style="list-style-type: none"> 1. Water quality: Storage vessel and chlorine tablets. 2. Sanitation: a) a sani-scoop hoe dedicated to the removal of human and animal faeces from the compound, b) plastic child potties for children ages 6 months and older until they are using the latrine, and c) a new or upgraded dual pit latrine for each household in the compound. The behavior change components of the intervention will emphasize the use of the latrine for defecation and the safe disposal of faeces in the compound courtyard to prevent contact with young children. 3. Handwashing: The hardware components of the Bangladesh handwashing intervention include two handwashing stations. The first station will be located in the kitchen (location of food preparation), and will include a 16 L bucket with a tap fitting, a stool, bowl and soapy water bottle. The second station will be located near the toilet, and will include a 40 L bucket with tap fitting, stool, bowl and soapy water bottle. The study will provide detergent soap to families free of charge to replenish the soapy water bottles. The behavior change component of the intervention will focus messaging for handwashing at two critical times: after defecation and before food preparation. 4. Nutrition: Mothers will be encouraged to exclusively breastfeed their children through age 6 months. When newborns reach 6 months of age, mothers will be encouraged to continue breastfeeding their children until 24 months, and will receive education about supplementing breastfeeding with healthy complementary foods following infant and young child feeding best practice guidelines from Unicef and the WHO. From ages 6 to 24 months, study children will receive a daily

Interventions to improve water quality for preventing diarrhoea (Review)

WASH-B, Bangladesh (Continued)

lipid-based nutritional supplement (LNS) that has been developed and tested through the [iLiNS](#) project.

Outcomes	<ol style="list-style-type: none"> 1. Length-for-Age Z-scores (time frame: measured 24 months after intervention) (Designated as safety issue: no). Child's recumbent length, standardized to Z-scores using the WHO 2006 growth standards. 2. Diarrhoea Prevalence (time frame: measured 12- and 24-months after intervention).
Starting date	May 2012
Contact information	International Centre for Diarrhoeal Disease Research, Bangladesh
Notes	

WASH-B, Kenya

Trial name or title	WASH-Benefits study, Kenya
Methods	Parallel, cluster-RCT
Participants	Estimated: 8000
Interventions	<ol style="list-style-type: none"> 1. Water quality: intervention villages will receive chlorine dispensers at spring water sources. After filling their plastic jerry can of water from the source, users can place the jerry can under the dispenser, and turn a knob to release 3 mL of chlorine. Behavior change messages will focus on the consistent provision of treated water to all children living in the household. 2. Sanitation: a) a sani-scoop hoe dedicated to the removal of human and animal faeces from the compound; b) plastic child potties for children ages 6 months and older until they are using the latrine; and c) a new or upgraded pit latrine for each household in the compound. If participants have a latrine, its structure will be improved if necessary. Plastic slabs will be installed to improve mud or wood floors, and the intervention delivery team will make sure that all latrine structures have walls, doors, roofs that ensure safety and privacy. The behaviour change components of the intervention will emphasize the use of the latrine for defecation and the safe disposal of faeces in the compound courtyard to prevent contact with young children. 3. Handwashing: two handwashing stations in the compound of each respondent, one near the latrine, and one by the cooking area. The handwashing stations are constructed from locally available materials and are of a dual tippy-tap design with independent pedals attached to 5 L jerry cans of clean water and jugs of soapy water. The behavior change component of the intervention will focus messaging for handwashing at two critical times: after defecation and before food preparation. 4. Nutrition: mothers will be encouraged to exclusively breastfeed their children through to 6 months of age. When newborns reach 6 months of age, mothers will be encouraged to continue breastfeeding their children until 24 months, and will receive education about supplementing breastfeeding with healthy complementary foods following infant and young child feeding best practice guidelines from Unicef and WHO. From ages six to 24 months, study children will receive a daily lipid-based nutritional supplement (LNS) that has been developed and tested through the iLiNS project.
Outcomes	<ol style="list-style-type: none"> 1. Length-for-age Z-scores (time frame: measured 24 months after intervention) (designated as safety issue: no). Child's recumbent length, standardized to Z-scores using the WHO 2006 growth standards. 2. Diarrhoea prevalence (time frame: measured 12 and 24 months after intervention)
Starting date	September 2012

Interventions to improve water quality for preventing diarrhoea (Review)

WASH-B, Kenya (Continued)

Contact information Innovations for Poverty Action, Kenya


Notes

DATA AND ANALYSES


Comparison 1. Water quality intervention versus control


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Diarrhoea: all ages	64	81215	Risk Ratio (Random, 95% CI)	0.59 [0.51, 0.69]
1.1 Source water improvement	6	9161	Risk Ratio (Random, 95% CI)	0.76 [0.48, 1.19]
1.2 POU treatment	58	72054	Risk Ratio (Random, 95% CI)	0.58 [0.48, 0.69]
2 Diarrhoea: children < 5 years	49		Risk Ratio (Random, 95% CI)	0.61 [0.49, 0.75]
2.1 Source water improvement	4		Risk Ratio (Random, 95% CI)	0.96 [0.82, 1.12]
2.2 POU treatment	45		Risk Ratio (Random, 95% CI)	0.58 [0.46, 0.73]

Analysis 1.1. Comparison 1 Water quality intervention versus control, Outcome 1 Diarrhoea: all ages.





Interventions to improve water quality for preventing diarrhoea (Review)


Interventions to improve water quality for preventing diarrhoea (Review)



Analysis 1.2. Comparison 1 Water quality intervention versus control, Outcome 2 Diarrhoea: children < 5 years.






Comparison 2. Source: water supply improvement versus control


Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
1 Diarrhoea: CBA studies sub- grouped by age	6		Risk Ratio (Random, 95% CI)	Subtotals only
1.1 Cluster-RCTs	1	3266	Risk Ratio (Random, 95% CI)	1.24 [0.98, 1.57]
1.2 CBA studies	5	5895	Risk Ratio (Random, 95% CI)	0.68 [0.42, 1.09]
2 Diarrhoea: CBA studies sub- grouped by age	5		Risk Ratio (Random, 95% CI)	Subtotals only
2.1 All ages	5	5895	Risk Ratio (Random, 95% CI)	0.68 [0.42, 1.09]


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.2 < 5 years	3	999	Risk Ratio (Random, 95% CI)	0.92 [0.79, 1.07]

Analysis 2.1. Comparison 2 Source: water supply improvement versus control, Outcome 1 Diarrhoea: CBA studies subgrouped by age.



Analysis 2.2. Comparison 2 Source: water supply improvement versus control, Outcome 2 Diarrhoea: CBA studies subgrouped by age.






Comparison 3. POU: water chlorination versus control


Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
1 Diarrhoea: subgrouped by study design	19	34694	Risk Ratio (Random, 95% CI)	0.72 [0.61, 0.84]
1.1 Cluster-RCTs	16	30746	Risk Ratio (Random, 95% CI)	0.77 [0.65, 0.91]
1.2 CBA studies	3	3948	Risk Ratio (Random, 95% CI)	0.51 [0.34, 0.75]
2 Diarrhoea: cluster-RCTs: subgrouped by age	16		Risk Ratio (Random, 95% CI)	Subtotals only
2.1 All ages	16		Risk Ratio (Random, 95% CI)	0.77 [0.65, 0.91]
2.2 < 5 years	15		Risk Ratio (Random, 95% CI)	0.77 [0.64, 0.92]
3 Diarrhoea: cluster-RCTs; subgrouped by adherence	16	30746	Risk Ratio (Random, 95% CI)	0.77 [0.65, 0.91]
3.1 Residual chlorine in 86 to 100% of samples	1	276	Risk Ratio (Random, 95% CI)	0.78 [0.73, 0.83]
3.2 Residual chlorine in 51 to 85% of samples	6	9994	Risk Ratio (Random, 95% CI)	0.60 [0.40, 0.91]
3.3 Residual chlorine in ≤ 50% of samples	4	12613	Risk Ratio (Random, 95% CI)	0.90 [0.76, 1.06]
3.4 Residual chlorine not reported	5	7863	Risk Ratio (Random, 95% CI)	0.85 [0.65, 1.12]
4 Diarrhoea: cluster-RCTs by risk of bias by blinding of participants	16		Risk Ratio (Random, 95% CI)	Subtotals only
4.1 Low risk	5	15867	Risk Ratio (Random, 95% CI)	1.07 [0.97, 1.17]
4.2 High risk	11	14879	Risk Ratio (Random, 95% CI)	0.68 [0.56, 0.83]
5 Diarrhoea: cluster-RCTs; subgrouped by additional water storage intervention	16		Risk Ratio (Random, 95% CI)	0.77 [0.65, 0.91]
5.1 Chlorination kit alone	8		Risk Ratio (Random, 95% CI)	0.75 [0.54, 1.05]
5.2 Chlorination kit plus water storage	8		Risk Ratio (Random, 95% CI)	0.80 [0.66, 0.97]

Interventions to improve water quality for preventing diarrhoea (Review)


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
6 Diarrhoea: cluster-RCTs: subgrouped by sufficiency of water quantity	16	30746	Risk Ratio (Random, 95% CI)	0.77 [0.65, 0.91]
6.1 Sufficient	3	5352	Risk Ratio (Random, 95% CI)	0.90 [0.69, 1.17]
6.2 Insufficient	2	3499	Risk Ratio (Random, 95% CI)	0.91 [0.66, 1.26]
6.3 Unclear	11	21895	Risk Ratio (Random, 95% CI)	0.67 [0.50, 0.88]
7 Diarrhoea: cluster-RCTs: subgrouped by water source	16		Risk Ratio (Random, 95% CI)	Subtotals only
7.1 Improved water source	3	5880	Risk Ratio (Random, 95% CI)	0.82 [0.59, 1.14]
7.2 Unimproved water source	13	24866	Risk Ratio (Random, 95% CI)	0.75 [0.59, 0.93]
8 Diarrhoea: cluster-RCTs: subgrouped by sanitation level	16		Risk Ratio (Random, 95% CI)	Subtotals only
8.1 Improved sanitation	3	4876	Risk Ratio (Random, 95% CI)	0.64 [0.44, 0.92]
8.2 Unimproved sanitation	6	17352	Risk Ratio (Random, 95% CI)	0.81 [0.63, 1.05]
8.3 Unclear	7	8518	Risk Ratio (Random, 95% CI)	0.75 [0.54, 1.05]
9 Diarrhoea: cluster-RCTs; subgrouped by length of follow-up	16		Risk Ratio (Random, 95% CI)	0.77 [0.65, 0.91]
9.1 ≤ 3 months	2		Risk Ratio (Random, 95% CI)	0.42 [0.06, 3.03]
9.2 > 3 to 6 months	7		Risk Ratio (Random, 95% CI)	0.71 [0.51, 0.99]
9.3 > 6 to 12 months	5		Risk Ratio (Random, 95% CI)	0.82 [0.71, 0.96]
9.4 > 12 months	2		Risk Ratio (Random, 95% CI)	0.99 [0.66, 1.48]


Analysis 3.1. Comparison 3 POU: water chlorination versus control, Outcome 1 Diarrhoea: subgrouped by study design.







Analysis 3.2. Comparison 3 POU: water chlorination versus control, Outcome 2 Diarrhoea: cluster-RCTs: subgrouped by age.







Analysis 3.3. Comparison 3 POU: water chlorination versus control, Outcome 3 Diarrhoea: cluster-RCTs; subgrouped by adherence.






Analysis 3.4. Comparison 3 POU: water chlorination versus control, Outcome 4 Diarrhoea: cluster-RCTs by risk of bias by blinding of participants.







Analysis 3.5. Comparison 3 POU: water chlorination versus control, Outcome 5 Diarrhoea: cluster-RCTs; subgrouped by additional water storage intervention.




Analysis 3.6. Comparison 3 POU: water chlorination versus control, Outcome 6 Diarrhoea: cluster-RCTs: subgrouped by sufficiency of water quantity.





Analysis 3.7. Comparison 3 POU: water chlorination versus control, Outcome 7 Diarrhoea: cluster-RCTs: subgrouped by water source.







Analysis 3.8. Comparison 3 POU: water chlorination versus control, Outcome 8 Diarrhoea: cluster-RCTs: subgrouped by sanitation level.





Analysis 3.9. Comparison 3 POU: water chlorination versus control, Outcome 9 Diarrhoea: cluster-RCTs; subgrouped by length of follow-up.






Comparison 4. POU: flocculation and disinfection versus control

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Diarrhoea: cluster-RCTs	7		Risk Ratio (Random, 95% CI)	0.48 [0.20, 1.16]
2 Diarrhoea: cluster-RCTs: subgrouped by age; excluding Doocy 2006 LBR	6		Risk Ratio (Random, 95% CI)	Subtotals only
2.1 All ages	6	11788	Risk Ratio (Random, 95% CI)	0.69 [0.58, 0.82]
2.2 < 5	6	0	Risk Ratio (Random, 95% CI)	0.71 [0.61, 0.84]
3 Diarrhoea: cluster-RCTs: subgrouped by adherence	7		Risk Ratio (Random, 95% CI)	Subtotals only
3.2 Residual chlorine 51 to 85%	1	2191	Risk Ratio (Random, 95% CI)	0.12 [0.11, 0.13]
3.3 Residual chlorine < 50%	4	6914	Risk Ratio (Random, 95% CI)	0.76 [0.67, 0.85]
3.4 Residual chlorine not measured	2	4874	Risk Ratio (Random, 95% CI)	0.41 [0.26, 0.64]
4 Diarrhoea: cluster-RCTs: subgrouped by additional storage container	7		Risk Ratio (Random, 95% CI)	0.48 [0.20, 1.16]
4.1 No storage container	2		Risk Ratio (Random, 95% CI)	0.81 [0.69, 0.95]
4.2 Storage container	5		Risk Ratio (Random, 95% CI)	0.39 [0.14, 1.08]
5 Diarrhoea: cluster-RCTs: subgrouped by sufficiency of water quantity	7		Risk Ratio (Random, 95% CI)	Subtotals only
5.1 Sufficient	1	3401	Risk Ratio (Random, 95% CI)	0.62 [0.47, 0.82]


Interventions to improve water quality for preventing diarrhoea (Review)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
5.2 Insufficient	2	5454	Risk Ratio (Random, 95% CI)	0.31 [0.05, 2.09]
5.3 Unclear	4	5124	Risk Ratio (Random, 95% CI)	0.64 [0.49, 0.85]
6 Diarrhoea: cluster-RCTs: subgrouped by water source	7		Risk Ratio (Random, 95% CI)	Subtotals only
6.1 Improved water source	2	4874	Risk Ratio (Random, 95% CI)	0.41 [0.26, 0.64]
6.2 Unimproved water source	4	5704	Risk Ratio (Random, 95% CI)	0.49 [0.14, 1.68]
6.3 Unclear	1	3401	Risk Ratio (Random, 95% CI)	0.62 [0.47, 0.82]
7 Diarrhoea: cluster-RCTs: subgrouped by sanitation level	7		Risk Ratio (Random, 95% CI)	Subtotals only
7.1 Improved sanitation	2	4874	Risk Ratio (Random, 95% CI)	0.41 [0.26, 0.64]
7.2 Unimproved sanitation	2	5592	Risk Ratio (Random, 95% CI)	0.27 [0.05, 1.36]
7.3 Unclear	3	3513	Risk Ratio (Random, 95% CI)	0.79 [0.69, 0.90]
8 Diarrhoea: cluster-RCTs: subgrouped by length of follow-up	7	13979	Risk Ratio (Random, 95% CI)	0.48 [0.20, 1.16]
8.1 ≤ 3 months	2	5592	Risk Ratio (Random, 95% CI)	0.27 [0.05, 1.36]
8.2 > 3 to 6 months	1	3263	Risk Ratio (Random, 95% CI)	0.83 [0.67, 1.03]
8.3 > 6 to 12 months	4	5124	Risk Ratio (Random, 95% CI)	0.64 [0.49, 0.85]


Analysis 4.1. Comparison 4 POU: flocculation and disinfection versus control, Outcome 1 Diarrhoea: cluster-RCTs.




Analysis 4.2. Comparison 4 POU: flocculation and disinfection versus control, Outcome 2 Diarrhoea: cluster-RCTs: subgrouped by age; excluding Doocy 2006 LBR.




Analysis 4.3. Comparison 4 POU: flocculation and disinfection versus control, Outcome 3 Diarrhoea: cluster-RCTs: subgrouped by adherence.







Analysis 4.4. Comparison 4 POU: flocculation and disinfection versus control, Outcome 4 Diarrhoea: cluster-RCTs: subgrouped by additional storage container.




Analysis 4.5. Comparison 4 POU: flocculation and disinfection versus control, Outcome 5 Diarrhoea: cluster-RCTs: subgrouped by sufficiency of water quantity.







Analysis 4.6. Comparison 4 POU: flocculation and disinfection versus control, Outcome 6 Diarrhoea: cluster-RCTs: subgrouped by water source.




Analysis 4.7. Comparison 4 POU: flocculation and disinfection versus control, Outcome 7 Diarrhoea: cluster-RCTs: subgrouped by sanitation level.



Analysis 4.8. Comparison 4 POU: flocculation and disinfection versus control, Outcome 8 Diarrhoea: cluster-RCTs: subgrouped by length of follow-up.






Comparison 5. POU: filtration versus control


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Diarrhoea: cluster-RCTs: subgrouped by age	23		Risk Ratio (Random, 95% CI)	Subtotals only
1.1 All ages	23		Risk Ratio (Random, 95% CI)	0.48 [0.38, 0.59]
1.2 < 5 years	19		Risk Ratio (Random, 95% CI)	0.49 [0.38, 0.62]
2 Diarrhoea: cluster-RCTs: subgrouped by type of filtration	23		Risk Ratio (Random, 95% CI)	Subtotals only
2.1 Ceramic filter	12	5763	Risk Ratio (Random, 95% CI)	0.39 [0.29, 0.53]
2.2 Sand filtration	5	5504	Risk Ratio (Random, 95% CI)	0.47 [0.39, 0.57]
2.3 LifeStraw®	3	3259	Risk Ratio (Random, 95% CI)	0.69 [0.51, 0.93]
2.4 Plumbed	3	1056	Risk Ratio (Random, 95% CI)	0.73 [0.52, 1.03]
3 Diarrhoea: cluster-RCTs: subgrouped by blinding of participants	23		Risk Ratio (Random, 95% CI)	Subtotals only
3.1 Low risk	5		Risk Ratio (Random, 95% CI)	0.80 [0.68, 0.94]
3.2 High risk	18		Risk Ratio (Random, 95% CI)	0.41 [0.33, 0.52]
4 Diarrhoea: ceramic filter studies subgrouped by water source	12	5763	Risk Ratio (Random, 95% CI)	0.39 [0.29, 0.53]
4.1 Improved water source	8	3607	Risk Ratio (Random, 95% CI)	0.33 [0.23, 0.46]
4.2 Unimproved water source	4	2156	Risk Ratio (Random, 95% CI)	0.54 [0.48, 0.61]
5 Diarrhoea: ceramic filter studies subgrouped by sanitation level	12	5763	Risk Ratio (Random, 95% CI)	0.39 [0.29, 0.53]

Interventions to improve water quality for preventing diarrhoea (Review)


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
5.1 Improved sanitation	7	4198	Risk Ratio (Random, 95% CI)	0.49 [0.38, 0.64]
5.2 Unimproved sanitation	4	1491	Risk Ratio (Random, 95% CI)	0.35 [0.22, 0.56]
5.3 Unclear	1	74	Risk Ratio (Random, 95% CI)	0.21 [0.18, 0.25]
6 Diarrhoea: sand filter studies: subgrouped by water source	5		Risk Ratio (Random, 95% CI)	0.47 [0.39, 0.57]
6.1 Improved water source	2		Risk Ratio (Random, 95% CI)	0.50 [0.33, 0.75]
6.2 Unimproved water source	2		Risk Ratio (Random, 95% CI)	0.44 [0.25, 0.76]
6.3 Unclear	1		Risk Ratio (Random, 95% CI)	0.47 [0.37, 0.60]
7 Diarrhoea: sand filter studies: subgrouped by sanitation level	5		Risk Ratio (Random, 95% CI)	0.47 [0.39, 0.57]
7.1 Improved sanitation	1		Risk Ratio (Random, 95% CI)	0.47 [0.37, 0.60]
7.2 Unimproved sanitation	3		Risk Ratio (Random, 95% CI)	0.48 [0.34, 0.68]
7.3 Unclear	1		Risk Ratio (Random, 95% CI)	0.46 [0.22, 0.96]
8 Diarrhoea: cluster-RCTs: subgrouped by adherence	23		Risk Ratio (Random, 95% CI)	Subtotals only
8.1 86 to 100%	12	7300	Risk Ratio (Random, 95% CI)	0.43 [0.34, 0.55]
8.2 51 to 85%	4	2346	Risk Ratio (Random, 95% CI)	0.56 [0.33, 0.95]
8.3 ≤ 50%	1	1516	Risk Ratio (Random, 95% CI)	0.75 [0.60, 0.94]
8.4 Not reported	6	4420	Risk Ratio (Random, 95% CI)	0.46 [0.28, 0.75]
9 Diarrhoea: cluster-RCTs: subgrouped by additional water storage intervention	19		Risk Ratio (Random, 95% CI)	Subtotals only
9.1 Filtration alone	8		Risk Ratio (Random, 95% CI)	0.60 [0.48, 0.76]
9.2 Filtration plus storage	11		Risk Ratio (Random, 95% CI)	0.38 [0.29, 0.49]
10 Diarrhoea: cluster-RCTs; subgrouped by length of follow-up	23		Risk Ratio (Random, 95% CI)	0.48 [0.38, 0.59]
10.1 ≤ 3 months	3		Risk Ratio (Random, 95% CI)	0.26 [0.20, 0.33]
10.2 > 3 to 6 months	11		Risk Ratio (Random, 95% CI)	0.52 [0.44, 0.60]
10.3 > 6 to 12 months	8		Risk Ratio (Random, 95% CI)	0.51 [0.30, 0.87]
10.4 > 12 months	1		Risk Ratio (Random, 95% CI)	0.87 [0.74, 1.02]


Analysis 5.1. Comparison 5 POU: filtration versus control, Outcome 1 Diarrhoea: cluster-RCTs: subgrouped by age.






Analysis 5.2. Comparison 5 POU: filtration versus control, Outcome 2 Diarrhoea: cluster-RCTs: subgrouped by type of filtration.







Analysis 5.3. Comparison 5 POU: filtration versus control, Outcome 3 Diarrhoea: cluster-RCTs: subgrouped by blinding of participants.




Analysis 5.4. Comparison 5 POU: filtration versus control, Outcome 4 Diarrhoea: ceramic filter studies subgrouped by water source.





Analysis 5.5. Comparison 5 POU: filtration versus control, Outcome 5 Diarrhoea: ceramic filter studies subgrouped by sanitation level.






Analysis 5.6. Comparison 5 POU: filtration versus control, Outcome 6 Diarrhoea: sand filter studies: subgrouped by water source.







Analysis 5.7. Comparison 5 POU: filtration versus control, Outcome 7 Diarrhoea: sand filter studies: subgrouped by sanitation level.





Analysis 5.8. Comparison 5 POU: filtration versus control, Outcome 8 Diarrhoea: cluster-RCTs: subgrouped by adherence.







Analysis 5.9. Comparison 5 POU: filtration versus control, Outcome 9 Diarrhoea: cluster-RCTs: subgrouped by additional water storage intervention.





Analysis 5.10. Comparison 5 POU: filtration versus control, Outcome 10 Diarrhoea: cluster-RCTs; subgrouped by length of follow-up.







Comparison 6. POU: solar disinfection versus control

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Diarrhoea: subgrouped by study design	6		Risk Ratio (Random, 95% CI)	Subtotals only
1.1 Cluster-RCTs	4	3460	Risk Ratio (Random, 95% CI)	0.62 [0.42, 0.94]
1.2 Quasi-RCTs	2	555	Risk Ratio (Random, 95% CI)	0.82 [0.69, 0.97]
2 Diarrhoea: cluster-RCTs; subgrouped by age	4		Risk Ratio (Random, 95% CI)	Subtotals only
2.1 All ages	4		Risk Ratio (Random, 95% CI)	0.62 [0.42, 0.94]


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.2 < 5	3		Risk Ratio (Random, 95% CI)	0.55 [0.34, 0.91]
3 Diarrhoea: cluster-RCTs; subgrouped by adherence	4		Risk Ratio (Random, 95% CI)	Subtotals only
3.1 86 to 100%	1	928	Risk Ratio (Random, 95% CI)	0.37 [0.29, 0.47]
3.2 51 to 85%	0	0	Risk Ratio (Random, 95% CI)	0.0 [0.0, 0.0]
3.3 ≤ 50%	2	1443	Risk Ratio (Random, 95% CI)	0.80 [0.57, 1.11]
3.4 Not reported	1	1089	Risk Ratio (Random, 95% CI)	0.73 [0.63, 0.85]
4 Diarrhoea: cluster-RCTs; subgrouped by sufficiency of water supply level	4	3460	Risk Ratio (Random, 95% CI)	0.62 [0.42, 0.94]
4.1 Sufficient	2	1443	Risk Ratio (Random, 95% CI)	0.80 [0.57, 1.11]
4.3 Unclear	2	2017	Risk Ratio (Random, 95% CI)	0.52 [0.27, 1.02]
5 Diarrhoea: cluster-RCTs; subgrouped by water source	4		Risk Ratio (Random, 95% CI)	Subtotals only
5.1 Improved water source	1	718	Risk Ratio (Random, 95% CI)	0.64 [0.39, 1.05]
5.2 Unimproved water source	3	2742	Risk Ratio (Random, 95% CI)	0.62 [0.38, 1.02]
6 Diarrhoea: cluster-RCTs; subgrouped by sanitation level	4		Risk Ratio (Random, 95% CI)	Subtotals only
6.1 Improved sanitation	0	0	Risk Ratio (Random, 95% CI)	0.0 [0.0, 0.0]
6.2 Unimproved sanitation	2	1653	Risk Ratio (Random, 95% CI)	0.57 [0.24, 1.39]
6.3 Unclear	2	1807	Risk Ratio (Random, 95% CI)	0.72 [0.63, 0.83]
7 Diarrhoea: cluster-RCTs; subgrouped by length of follow-up	4	3460	Risk Ratio (Random, 95% CI)	0.62 [0.42, 0.94]
7.2 > 6 to 12 months	3	2371	Risk Ratio (Random, 95% CI)	0.59 [0.32, 1.09]
7.3 > 12 months	1	1089	Risk Ratio (Random, 95% CI)	0.73 [0.63, 0.85]

Analysis 6.1. Comparison 6 POU: solar disinfection versus control, Outcome 1 Diarrhoea: subgrouped by study design.







Analysis 6.2. Comparison 6 POU: solar disinfection versus control, Outcome 2 Diarrhoea: cluster-RCTs; subgrouped by age.




Analysis 6.3. Comparison 6 POU: solar disinfection versus control, Outcome 3 Diarrhoea: cluster-RCTs; subgrouped by adherence.




Analysis 6.4. Comparison 6 POU: solar disinfection versus control, Outcome 4 Diarrhoea: cluster-RCTs; subgrouped by sufficiency of water supply level.







Analysis 6.5. Comparison 6 POU: solar disinfection versus control, Outcome 5 Diarrhoea: cluster-RCTs; subgrouped by water source.




Analysis 6.6. Comparison 6 POU: solar disinfection versus control, Outcome 6 Diarrhoea: cluster-RCTs; subgrouped by sanitation level.





Analysis 6.7. Comparison 6 POU: solar disinfection versus control, Outcome 7 Diarrhoea: cluster-RCTs; subgrouped by length of follow-up.



Comparison 7. POU: UV disinfection versus control

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Diarrhoea: cluster-RCT	1		Risk Ratio (Random, 95% CI)	Subtotals only

Analysis 7.1. Comparison 7 POU: UV disinfection versus control, Outcome 1 Diarrhoea: cluster-RCT.

Study or subgroup	Inter- vention	Control	log[Risk Ratio] (SE)	Risk Ratio	Weight	Risk Ratio
	N	N		IV, Random, 95% CI		IV, Random, 95% CI
Gruber 2013 MEX	957	956	-0.2 (0.244)		0%	0.79[0.49,1.27]

Favours intervention 0.2 0.5 1 2 5 Favours control

Comparison 8. POU: improved storage versus control

Outcome or subgroup title	No. of studies	No. of partici- pants	Statistical method	Effect size
1 Diarrhoea: cluster-RCTs: subgrouped by age	2		Risk Ratio (Random, 95% CI)	Subtotals only
1.1 All ages	2		Risk Ratio (Random, 95% CI)	0.91 [0.74, 1.11]
1.2 < 5	1		Risk Ratio (Random, 95% CI)	0.69 [0.47, 1.01]

Analysis 8.1. Comparison 8 POU: improved storage versus control, Outcome 1 Diarrhoea: cluster-RCTs: subgrouped by age.

Study or subgroup	Favours in- tervention	Control	log[Risk Ratio] (SE)	Risk Ratio	Weight	Risk Ratio
	N	N		IV, Random, 95% CI		IV, Random, 95% CI
8.1.1 All ages						
Günther 2013 BEN	364	347	-0 (0.076)		63.35%	0.98[0.85,1.14]
Roberts 2001 MWI	310	850	-0.2 (0.135)		36.65%	0.79[0.61,1.03]
Subtotal (95% CI)					100%	0.91[0.74,1.11]
Heterogeneity: Tau ² =0.01; Chi ² =1.95, df=1(P=0.16); I ² =48.59%						
Test for overall effect: Z=0.94(P=0.34)						
8.1.2 < 5						
Roberts 2001 MWI	0	0	-0.4 (0.194)		100%	0.69[0.47,1.01]
Subtotal (95% CI)					100%	0.69[0.47,1.01]
Heterogeneity: Tau ² =0; Chi ² =0, df=0(P<0.0001); I ² =100%						
Test for overall effect: Z=1.91(P=0.06)						

Favours intervention 0.001 0.1 1 10 1000 Favours control

ADDITIONAL TABLES

Table 1. Water quality indicators post-intervention

Trial	Water quality indicator	Water quality post-intervention:	Water quality post intervention:
		Intervention group	Control group

Table 1. Water quality indicators post-intervention (Continued)

Abebe 2014 ZAF	CFUs/100 mL	0	80% of control HHS had 10 to 10000
Austin 1993a GMB	Geometric mean CFUs/100 mL	178	3020
Austin 1993b GMB	Geometric mean CFUs/100 mL	42	3020
Boisson 2009 ETH	Arithmetic mean TTC/100 mL (95% CI)	0	725.7 (621.0 to 830.4)
Boisson 2010 DRC	Geometric mean TTC/100 mL (95% CI)	1.3 (0.9 to 1.7)	173.7 (136.6 to 220.9)
Boisson 2013 IND	Geometric mean TTC/100 mL (95% CI)	50 (44 to 57)	122 (107 to 139)
Brown 2008a KHM	Geometric mean <i>E. coli</i> /100 mL	17	600
Brown 2008b KHM	Geometric mean <i>E. coli</i> /100 mL	15	600
Clasen 2004b BOL	Mean TTC/100 mL	0.13	108
Clasen 2004c BOL	Arithmetic mean TTC/100 mL	100% of intervention households: 0	16% of control households: 0 66% > 10, 34% > 100, and 11% > 1000
Clasen 2005 COL	Arithmetic mean TTC/100 mL (95% CI)	37.3 (6.3 to 48.3)	150.6 (34.8 to 166.4)
Colford 2002 USA; Colford 2005 USA; Colford 2009 USA	All water met FDA requirements	Not measured because of high water quality	Not measured because of high water quality
Crump 2005a KEN	Samples met WHO guidelines for water quality	82%	14%
Crump 2005b KEN	Samples met WHO guidelines for water quality	78%	14%
du Preez 2008 ZAF/ZWE	Samples met WHO guidelines for water quality	57%	30%
du Preez 2010 ZAF	<i>E. coli</i> in concentrations/100 mL	62%	"No significant difference between intervention and control groups"
du Preez 2011 KEN	<i>E. coli</i> in concentrations/100 mL	Storage containers: 0.723 SODIS bottles: -0.727	Not reported
Fabiszewski 2012 HND	Geometric mean <i>E. coli</i> counts per 100 mL (95% CI)	23.4 (20.2 to 27.0)	45.4 (38.6 to 53.4)
Gasana 2002 RWA	Total coliforms/100 mL	Range: 3 to 43	Range: 4 to 1100
Gruber 2013 MEX	Samples with detectable <i>E. coli</i>	43%	59%

Table 1. Water quality indicators post-intervention (Continued)

Günther 2013 BEN	<i>E. coli</i> contamination > 1000 CFU/100 mL	Not reported specifically; findings imply a 70% reduction in <i>E. coli</i> incidence for intervention households	
Handzel 1998 BGD	Stored water samples with <i>E. coli</i> 100 MPN/100 mL	3%	16%
Jain 2010 GHA	Samples with <i>E. coli</i>	8%	54%
Jensen 2003 PAK	Geometric mean <i>E. coli</i> /100 mL	3	49
Kirchhoff 1985 BRA	Mean number of faecal coliforms/dL in the samples	70	16000
Kremer 2011 KEN	Average reduction in log <i>E. coli</i>	-1.07, corresponding to a 66% reduction	
Lule 2005 UGA	Median <i>E. coli</i> CFU/100 mL	23	59
McGuigan 2011 KHM	Geometric mean CFU/100 mL	6.8	48
Mengistie 2013 ETH	Mean <i>E. coli</i>	0	60
Peletz 2012 ZMB	Geometric mean TTC/100 mL	Stored water: 3	Stored water: 181
Quick 1999 BOL	Median <i>E. coli</i> /100 mL	0	6400
Quick 2002 ZMB	Median <i>E. coli</i> /100 mL	0	3
Reller 2003a GTM	Samples with < 1 <i>E. coli</i> /100 mL (flocculant/disinfectant)	40%	7%
Reller 2003b GTM	Samples with < 1 <i>E. coli</i> /100 mL (flocculant/disinfectant+ vessel)	57%	7%
Reller 2003c GTM	Samples with < 1 <i>E. coli</i> /100 mL (bleach)	51%	7%
Reller 2003d GTM	Samples with < 1 <i>E. coli</i> /100 mL (bleach + vessel)	61%	7%
Semenza 1998 UZB	Faecal colonies/100 mL	47	52
Stauber 2009 DOM	<i>E. coli</i> MPN/100 mL	11	19
Stauber 2012a KHM	<i>E. coli</i> CFU/100 mL	2.9	19.7
Stauber 2012b GHA	Geometric mean <i>E. coli</i> MPN/100 mL (95% CI)	Direct filtrate 16 (13 to 20)	490 (426 to 549)
		Stored filtrate: 76 (62 to 91)	
Tiwari 2009 KEN	Geometric mean faecal coliforms/100 mL (95% CI)	30.0 (21.3 to 42.1)	88.9 (58.7 to 135)
URL 1995a GTM	Samples with fecal coliforms	91% had 0 fecal coliforms	Not reported
URL 1995b GTM	Samples with fecal coliforms	91% had 0 fecal coliforms	Not reported

Abbreviations: *E. coli*: *Escherichia coli*; FC: faecal coliform.

Table 2. Studies reporting deaths

Study ID	Intervention		Control		P value	Comment
	Deaths	Partici- pants	Deaths	Partici- pants		
Boisson 2010 DRC	12	546	8	598	0.27	—
Colford 2009 USA	7	385	6	385	> 0.05	—
Crump 2005a KEN	17	2249	28	2277	0.108	—
Crump 2005b KEN	14	2124	28	2277	0.052	—
du Preez 2011 KEN	3	555	3	534	> 0.05	—
Peletz 2012 ZMB	3	300	6	299	0.28	—
Boisson 2013 IND	?	6119	?	5965	—	Only reports total deaths (46)
du Preez 2010 ZAF	?	383	?	335	—	Only reports total deaths (7)
Kremer 2011 KEN	?	—	?	—	—	Reports recording deaths but does not state how many
Boisson 2009 ETH	?	731	?	785	—	Reports recording deaths but does not state how many

Table 3. Summary of findings: improved water source

Improved water source compared with no intervention for preventing diarrhoea in rural settings in low- and middle-income countries					
Patient or population: adults and children					
Settings: low- and middle-income countries in rural areas					
Intervention: water source improvement					
Comparison: no intervention					
Outcomes	Illustrative comparative risks* (95% CI)		Relative ef- fect (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)
	Assumed risk	Corresponding risk			
	No intervention	Water source improvement			
Diarrhoea episodes	3 episodes per person per year	3.7 episodes per person per year (2.9 to 4.7)	RR 1.24 (0.98 to 1.57)	3266 (1 trial)	⊕⊕⊕⊕ very low 1,2,3
Cluster-RCTs					

[Interventions to improve water quality for preventing diarrhoea \(Review\)](#)

Copyright © 2015 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

Table 3. Summary of findings: improved water source (Continued)

Diarrhoea episodes	—	—	—	5895 (5 studies)	⊕○○○ very low 1,4,5
CBA studies					

The basis for the **assumed risk** (for example, the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **RR:** risk ratio.

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: we are very uncertain about the estimate.

The assumed risk is based on 2.9 episodes/child year in 2010 ([Fischer Walker 2012](#)).

¹Downgraded by 1 for serious risk of bias: as diarrhoea episodes were reported by participants this outcome is susceptible to bias from lack of blinding. None of these studies blinded participants and outcome assessors to the treatment allocation.

²No serious inconsistency.

³Downgraded by 2 for serious indirectness: this single RCT from Afghanistan evaluated the provision of protected wells. It is not possible to make broad generalizations to other settings.

⁴Downgraded by 1 for serious inconsistency: statistical heterogeneity was very high (I^2 statistic = 98%), such that the data could not be pooled. Some large and statistically significant effects were seen in some individual trials, but not others.

⁵Downgraded by 1 for serious indirectness: these studies are from a variety of low- and middle-income countries (Bangladesh, Rwanda, Pakistan, South Africa, China). However, as only single trials evaluated each intervention it is not possible to make broad generalizations.

Table 4. Improved water source: description of the interventions

Study ID	Study design	Setting	Incidence of diarrhoea in the control group	Intervention areas		Control areas	
				Water source intervention	Health promotion activities	Water source	Health promotion activities
Opryszko 2010b AFG	Cluster-RCT	Rural villages	3.1 episodes per person per year	One well per 25 households providing 25 litres/person/day	None	35% used unprotected hand dug wells	None
Alam 1989 BGD	CBA	Rural villages	4.1 episodes per child per year	Provision of one hand pump per 4-6 households (3 times as many as control areas)	Female health visitors visited peoples homes and organised group discussion and demonstrations to promote hygienic practices for hand pump use, water storage, child faeces disposal, hand washing.	Shallow, hand-dug wells; some hand pumps	None described
Gasana 2002 RWA	CBA	Rural villages	3 episodes per child per year	Site A: Sedimentation tank/Katadyn filter with communal tap Site B: Gravel-sand-charcoal filter on existing water spring Site C: Protective fence around an existing water spring	None described	An existing water spring	None described
Jensen 2003 PAK	CBA	Rural villages	2.8 episodes per person per year	Chlorination of public water supply	None described	Unchlorinated poorly functioning sand filter system	None described
Majuru 2011 ZAF	CBA	Rural villages	0.6 episodes per person per year	Provision of intermittently operated small community water systems distributing potable water to multiple taps throughout the community	None described	Untreated water from a river and its tributaries	None described
Xiao 1997 CHN	CBA	Rural villages	Not reported	Improved water supply through structural improvements to wells	Hygiene education	Not reported	None described

Table 5. Improved water source: primary drinking water supply and sanitation facilities

Trial	Description	Source ¹	Access to source ²	Quantity available ³	Ambient water quality	Sanitation ⁴
Alam 1989 BGD	Shallow, hand-dug wells; some hand pumps	Unimproved	Unclear	Unclear	Not tested	Unclear
Gasana 2002 RWA	Spring	Unimproved	Unclear	Unclear	Baseline range 4 to 1100 total coliforms/100 mL	Unimproved
Jensen 2003 PAK	Some slow sand filters in poor condition; some household taps; majority used ground water	Improved	Unclear	Unclear	Baseline geometric mean in intervention village: 13.3 <i>E. coli</i> CFU/100 mL; control villages: 137/100 mL	Unclear
Majuru 2011 ZAF	Surface water, boreholes, water tankers	Improved and unimproved	Unclear	Unclear	Not tested	Unclear
Opryszko 2010	35% use unprotected dug wells	Unimproved	Sufficient	Sufficient	Not tested	Unclear
Xiao 1997 CHN	Well water	Unimproved	Unclear	Unclear	Not tested	Unclear

¹'Improved' includes household connection, public standpipe, borehole, protected dug well, protected spring, rainwater collection; 'unimproved' includes unprotected well, unprotected spring, vendor-provided water, bottled water; and 'unclear' means unclear or not reported; definition based on [WHO/UNICEF 2015](#).

²'Sufficient' means located within 500 m, queuing no more than 15 minutes, no more than three minutes to fill 20 L container, and maintained so available consistently; 'insufficient' means that it does not meet any of above; and 'unclear' means unclear or not reported; definition based minimum standards established by The [Sphere Project 2011](#).

³'Sufficient' means a minimum of 15 L/day/person; 'insufficient' means less than 15 L/day/person; and 'unclear' means unclear or not reported; definition based on minimum standards established by The [Sphere Project 2011](#).

⁴'Improved' means connection to a public sewer or septic system, pour flush latrine, simple pit latrine, or ventilated improved pit latrine; 'unimproved' means service or bucket latrine, public latrines, open latrines; and 'unclear' means unclear or not reported; definition based on [WHO/UNICEF 2015](#).

Table 6. POU chlorination: description of the intervention

Trial	Study design	Chlorination product?	Distributed free?	Frequency of distribution?	Storage container also distributed?	Compliance	Additional hygiene promotion
Austin 1993a GMB	Cluster-RCT	Sodium hypochlorite solution	Yes	Fortnightly	No	40% compliance measured by residual chlorine	None
Austin 1993b GMB	Cluster-RCT	Sodium hypochlorite solution	Yes	Fortnightly	No	59% compliance measured by residual chlorine	None
Boisson 2013 IND	Cluster-RCT	Sodim dichloro-iso-cyanurate tablets	Yes	Bimonthly	No	32% compliance measured by residual chlorine	None
Crump 2005a KEN	Cluster-RCT	1% sodium hypochlorite	Yes	Weekly	No	61% compliance during unannounced weekly visits measured by residual chlorine	Use of ORS, treatment seeking for diarrhoea
Handzel 1998 BGD	Cluster-RCT	0.25% to 0.3% chlorine solution	Yes	Weekly	Yes	90% compliance based on residual chlorine measurements	Hygiene and sanitation messages
Jain 2010 GHA	Cluster-RCT	Sodim dichloro-iso-cyanurate tablets	Yes	Twice weekly	Yes	74% to 89% compliance measured by chlorine residual	ORS provided to those with diarrhoea
Kirchhoff 1985 BRA	Cluster-RCT	10% sodium hypochlorite	Yes	Daily	No	Not reported	Chlorination preformed by study staff
Luby 2006a PAK	Cluster-RCT	Sodium hypochlorite solution	Yes	Unclear	Yes	Yes, though rate unclear	Encouraged to only drink treated water
Lule 2005 UGA	Cluster-RCT	0.5% sodium hypochlorite	Yes	Weekly	Yes	Not reported	hygiene education
Mahfouz 1995 KSA	Cluster-RCT	Packets of 50 g calcium hypochloride 70%.	Yes	Unclear	No	Some residual chlorine in all intervention samples	None
Mengistie 2013 ETH	Cluster-RCT	1.25% sodium hypochlorite solution	Yes	Weekly	No	80% compliance measured by chlorine residual	None



Table 6. POU chlorination: description of the intervention (Continued)

Opryszko 2010c AFG	Cluster-RCT	0.05% sodium hypochlorite solution	Yes	Monthly	Yes	78% compliance measured by previous 2 weeks self-report use of chlorine	None
Quick 1999 BOL	Cluster-RCT	MIOX unit electrolytically produced disinfectant with 3% brine solution, hypochlorite, chlorine dioxide, ozone, peroxide and other oxidants.	Yes	Weekly	Yes	63% compliance measured by water in vessel with chlorine residual, average across six rounds	Community health volunteers reinforced messages about proper use of the disinfectant and vessels and of different applications for treated water.
Reller 2003b GTM	Cluster-RCT	Sodium hypochlorite solution (50,000 ppm)	Yes	Monthly	No	36% compliance measure by residual chlorine > 0.1 mg/L on unannounced visits.	Motivational and educational messages about chlorination, use of ORS, care seeking for diarrhoea
Reller 2003c GTM	Cluster-RCT	Sodium hypochlorite solution (50,000 ppm)	Yes	Monthly	Yes	44% compliance measure by residual chlorine > 0.1 mg/L on unannounced visits.	Motivational and educational messages about chlorination, use of ORS, care seeking for diarrhoea
Semenza 1998 UZB	Cluster-RCT	1.5% chlorine solution	Yes	Unclear but households were visited twice weekly	Yes	73% based on residual chlorine levels at time of visit	Only drink chlorinated water and wash all fruit and vegetables with chlorinated water
Luby 2004a PAK	CBA	Bleach (sodium hypochlorite)	Yes	Study workers visited weekly and re-supplied the households with dilute bleach.	Yes	Not reported	Encouraged regular treatment of drinking water
Luby 2004b PAK	CBA	Bleach (sodium hypochlorite)	Yes	Study workers visited weekly and re-supplied the households with dilute bleach.	Yes	Not reported	Encouraged regular treatment of drinking water

Table 6. POU chlorination: description of the intervention *(Continued)*

Quick 2002 ZMB	CBA	0.5% sodium hypochlorite	Yes	Unclear but households were visited once every two weeks	HHs paid for vessel	72% compliance measured by water in vessel with chlorine residual	Community volunteers, gave education about causes and prevention of diarrhoea and safe storage of water and motivated households about the intervention.
---------------------------	-----	--------------------------	-----	--	---------------------	---	--

Table 7. POU chlorination: primary drinking water supply and sanitation facilities

Trial	Description	Source ¹	Access to source ²	Quantity available ³	Ambient water quality	Sanitation ⁴
Austin 1993	Open wells	Unimproved	Sufficient	Unclear	Mean 1871 FC/100 mL in wells; among stored water samples: mean 3358 FC/100 mL in rainy season, 1014 FC/100 mL in dry season	Unclear
Boisson 2013 IND	62% unprotected dug well, 17% tubewell, 14% tap, 5% surface water	Unimproved	Unclear	Unclear	Baseline not reported. Control households: Geometric mean 122 TTC/100 mL	Unimproved
Crump 2005	50% ponds, 49% rivers	Unimproved	Unclear	Insufficient	Baseline mean 98 <i>E. coli</i> /100 mL	Unclear; 33% defecate on ground
Handzel 1998 BGD	48% tap, 52% tubewell; 61% paid for drinking water	Improved	Sufficient	Sufficient	Baseline geometric mean 138.1 faecal coliform counts/100 mL	Unimproved
Jain 2010 GHA	95% of households use tap, 84% surface water, 46% wells, 35% rainwater, 25% borehole	Improved and unimproved	Unclear	Unclear	Baseline: median <i>E. coli</i> MPN 93/100 mL	Unimproved
Kirchhoff 1985 BRA	Pond water stored in clay pots after filtering with cloth	Unimproved	Unclear	Insufficient	Source water: mean 970 faecal coliforms/100 mL	Unimproved
Luby 2004	Tanker trucks, municipal taps (household and community level)	Mostly unimproved	Unclear	Unclear	Baseline: approximately 60% of stored drinking water samples were free of <i>E. coli</i>	Improved
Luby 2006	Tanker trucks, municipal taps (household and community level), water bearer, boreholes	Mostly improved	Unclear	Unclear	Not tested	Improved
Lule 2005 UGA	16% surface or shallow wells, 50% protected springs, 49% boreholes or taps	Unimproved	Sufficient	Sufficient	Source mean <i>E. coli</i> counts: 11/100 mL	Improved
Mahfouz 1995 KSA	Shallow wells	Unimproved	Unclear	Unclear	Source: 92% positive with <i>E. coli</i> ; precise level not reported	Improved
Mengistie 2013 ETH	50% well, 41% spring, 9% river	Unimproved	Unclear	Unclear	Baseline: <i>E. coli</i> MPN 70/100 mL	Unimproved

Table 7. POU chlorination: primary drinking water supply and sanitation facilities (Continued)

Opryszko 2010	35% use unprotected dug wells	Unimproved	Sufficient	Sufficient	Not tested	Unclear
Quick 1999 BOL	Shallow uncovered wells; 38% treated water	Unimproved	Unclear	Unclear	Source water: median colony count <i>E. coli</i> : 57,050/100 mL	Unimproved, but 47% used latrine
Quick 2002 ZMB	Shallow wells; some boiling	Unimproved	Unclear	Unclear	Source water: median colony count <i>E. coli</i> : 34/100 mL	Unclear
Reller 2003	Surface water from shallow wells, rivers and springs	Unimproved	Unclear	Unclear	Baseline drinking water: median colony count <i>E. coli</i> 63/100 mL	Unclear
Semenza 1998 UZB	Households without piped water (procured from street tap, neighbour tap, well, vendor, or river)	Unimproved	Unclear	Unclear	Source water: 54 coliform colonies/100 mL	Unclear

¹'Improved' includes household connection, public standpipe, borehole, protected dug well, protected spring, rainwater collection; 'unimproved' includes unprotected well, unprotected spring, vendor-provided water, bottled water; and 'unclear' means unclear or not reported; definition based on [WHO/UNICEF 2015](#).

²'Sufficient' means located within 500 m, queuing no more than 15 minutes, no more than three minutes to fill 20 L container, and maintained so available consistently; 'insufficient' means that it does not meet any of above; and 'unclear' means unclear or not reported; definition based minimum standards established by The [Sphere Project 2011](#).

³'Sufficient' means a minimum of 15 L/day/person; 'insufficient' means less than 15 L/day/person; and 'unclear' means unclear or not reported; definition based on minimum standards established by The [Sphere Project 2011](#).

⁴'Improved' means connection to a public sewer or septic system, pour flush latrine, simple pit latrine, or ventilated improved pit latrine; 'unimproved' means service or bucket latrine, public latrines, open latrines; and 'unclear' means unclear or not reported; definition based on [WHO/UNICEF 2015](#).

Table 8. Summary of findings: POU chlorination

POU chlorination compared with no intervention for preventing diarrhoea					
Patient or population: adults and children					
Settings: low- and middle-income countries					
Intervention: distribution of chlorine for POU water treatment and instruction on use					
Comparison: no intervention					
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)
	Assumed risk	Corresponding risk			
	No intervention	POU Chlorination			
Diarrhoea episodes cluster-RCTs	3 episodes per person per year	2.3 episodes per year (2.0 to 2.7)	RR 0.77 (0.65 to 0.91)	30,746 (14 trials)	⊕⊕⊕⊕ low 1,2,3,4
Diarrhoea episodes	3 episodes per person per year	1.5 episodes per year (1.0 to 2.3)	RR 0.51 (0.34 to 0.75)	3948 (2 studies)	⊕⊕⊕⊕ very low 5,6,7,8

Interventions to improve water quality for preventing diarrhoea (Review)

Table 8. Summary of findings: POU chlorination (Continued)

CBA studies

The basis for the **assumed risk** is provided in the footnotes. The **corresponding risk** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **RR:** risk ratio.

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: we are very uncertain about the estimate.

The assumed risk is based on 2.9 episodes/child year in 2010 (Fischer Walker 2012).

¹Downgraded by 1 for serious risk of bias: as diarrhoea episodes were reported by participants this outcome is susceptible to bias from lack of blinding. Only two of these studies blinded participants and outcome assessors to the treatment allocation, and these two studies found no evidence of an effect with chlorination.

²Downgraded by 1 for serious inconsistency: statistical heterogeneity was very high (I^2 statistic = 91%). In a subgroup analysis by compliance with the intervention (assessed by measurements of residual chlorine in drinking water) found larger effects in the studies with better compliance.

³No serious indirectness: these studies are mainly from low- and middle-income countries (the Gambia, India, Kenya, Bangladesh, Ghana, Brazil, Pakistan, Uganda, Saudi Arabia, Ethiopia, Afghanistan, Bolivia, Guatemala, and Uzbekistan). The interventions consisted of free distribution of chlorine (every one to four weeks) plus instructions on how to use it. In some cases, the intervention included hygiene education and storage containers in which to treat and store water.

⁴No serious imprecision: the average effect suggests POU chlorination may reduce diarrhoea episodes by about a quarter. The analysis is adequately powered to detect this effect.

⁵Downgraded by 1 for serious risk of bias: as diarrhoea episodes were reported by participants this outcome is susceptible to bias from lack of blinding. None of these studies blinded participants and outcome assessors to the treatment allocation.

⁶Downgraded by 1 for serious inconsistency: statistical heterogeneity was very high (I^2 statistic = 63%).

⁷Downgraded by 1 for serious indirectness: there are only two studies (three comparisons) from Pakistan and Zambia.

⁸No serious imprecision.

Table 9. POU flocculation/disinfection: description of the interventions

Study ID	Study design	Setting	Intervention areas			Control areas	
			Water quality intervention	Health promotion activities	Compliance	Water source	Health promotion activities
Chiller 2006 GTM	Cluster-RCT	Rural villages	Provided households with a large spoon and a wide-mouthed bucket for mixing, a narrow-topped vessel with a lid for storing treated water and provided households with sachets of the flocculant–disinfectant every week	None	44% compliance measured by residual chlorine at week 10 of study	31% tap, 40% river or spring and 25% well.	None
Crump 2005b KEN	Cluster-RCT	Rural villages	Each week households were given sachets of the flocculant–disinfectant	None	44% compliance during unannounced weekly visits measured by residual chlorine	50% pond, 49% river and 2% spring	None
Doocy 2006 LBR	Cluster-RCT	Liberian camps for displaced persons	Households received a bucket and large mixing spoon for preparation, a decanting cloth, a funnel and a storage container with a narrow opening and lid. Each household received a maximum of 21 flocculation–disinfectant packets per week	None	85% compliance based on residual chlorine sampling	Received a funnel and an identical storage container	None
Luby 2006b PAK	Cluster-RCT	Squatter settlements	Provided households with flocculant–disinfectant sachets, a water vessel and soap. Weekly distributions of sachets	Field workers educated neighbourhoods about health problems resulting from hand and water contamination and instructed households on how and when to wash hands	Yes, though rate unclear	Municipal supply at household (33%), at community tap (37%), tanker truck (12%), water bearer (13%) and tube well (5%)	None
Luby 2006c PAK	Cluster-RCT	Squatter settlements	Flocculant–disinfectant and vessel. Weekly distributions of sachets	Field workers educated neighbourhoods about health problems resulting	Yes, though rate unclear	Municipal supply at household (33%), at community tap	None

Table 9. POU flocculation/disinfection: description of the interventions *(Continued)*

				from hand and water contamination		(37%), tanker truck (12%), water bearer (13%) and tube well (5%)	
Reller 2003a GTM	Cluster-RCT	Rural villages	Weekly distribution of flocculant-disinfectant and gave 2 cloths initially, which could be exchanged	Field workers discussed the importance of water treatment and demonstrated the water preparation process	27% compliance measure by residual chlorine > 0.1 mg/L on unannounced visits.	33% tap, 46% river or spring, 21% well.	None
Reller 2003d GTM	Cluster-RCT	Rural villages	Weekly distribution of flocculant-disinfectant and gave 2 cloths initially, which could be exchanged and received a large plastic spoon for stirring, a large-mouthed bucket for mixing, and a vessel with a secure lid and a spigot for storing treated water	Field workers discussed the importance of water treatment and demonstrated the water preparation process	34% compliance measure by residual chlorine > 0.1 mg/L on unannounced visits.	33% tap, 46% river or spring, 21% well.	None

Table 10. POU flocculation/disinfection: primary drinking water supply and sanitation facilities

Trial	Description	Source ¹	Access to source ²	Quantity available ³	Ambient H ₂ O quality	Sanitation ⁴
Chiller 2006 GTM	Rivers, springs, taps, and wells	Unclear	Unclear	Sufficient	98% of source samples contained <i>E. coli</i> ; precise level not reported	Mostly unimproved
Crump 2005b KEN	50% ponds, 49% rivers	Unimproved	Unclear	Insufficient	Baseline mean 98 <i>E. coli</i> /100 mL	Unclear; 33% defecate on ground
Doocy 2006 LBR	Surface sources and some tap stands	Unimproved	Unclear	Insufficient	Source water: 88% samples tested positive for faecal contamination; precise level not reported	Unimproved
Luby 2006b PAK	Tanker trucks, municipal taps (household and community level), water bearer, boreholes	Mostly improved	Unclear	Unclear	Not tested	Improved
Reller 2003a GTM	Surface water from shallow wells, rivers and springs	Unimproved	Unclear	Unclear	Baseline drinking water: median colony count <i>E. coli</i> 63/100 mL	Unclear

¹'Improved' includes household connection, public standpipe, borehole, protected dug well, protected spring, rainwater collection; 'unimproved' includes unprotected well, unprotected spring, vendor-provided water, bottled water; and 'unclear' means unclear or not reported; definition based on [WHO/UNICEF 2015](#).

²'Sufficient' means located within 500 m, queuing no more than 15 minutes, no more than three minutes to fill 20 L container, and maintained so available consistently; 'insufficient' means that it does not meet any of above; and 'unclear' means unclear or not reported; definition based on minimum standards established by The [Sphere Project 2011](#).

³'Sufficient' means a minimum of 15 L/day/person; 'insufficient' means less than 15 L/day/person; and 'unclear' means unclear or not reported; definition based on minimum standards established by The [Sphere Project 2011](#).

⁴'Improved' means connection to a public sewer or septic system, pour flush latrine, simple pit latrine, or ventilated improved pit latrine; 'unimproved' means service or bucket latrine, public latrines, open latrines; and 'unclear' means unclear or not reported; definition based on [WHO/UNICEF 2015](#).

Table 11. Summary of findings: POU flocculation and disinfection

POU water flocculation and disinfection compared with no intervention for preventing diarrhoea					
Patient or population: adults and children					
Settings: low- and middle-income countries					
Intervention: distribution of sachets combining water flocculation and disinfection and instructions on use					
Comparison: no intervention					
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)
	Assumed risk	Corresponding risk			
	No intervention	Water flocculation and disinfection			

Table 11. Summary of findings: POU flocculation and disinfection (Continued)

Diarrhoea episodes	3 episodes per person per year	2.1 episodes per person per year (1.7 to 2.5)	RR 0.69 (0.58 to 0.82)	11,788 (4 trials)	⊕⊕⊕⊖ moderate 1,2,3,4
Cluster-RCTs					

The basis for the **assumed risk** is provided in the footnotes. The **corresponding risk** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **RR:** risk ratio.

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: we are very uncertain about the estimate.

The assumed risk is based on 2.9 episodes/child year in 2010 ([Fischer Walker 2012](#)).

¹Downgraded by 1 for serious risk of bias: as diarrhoea episodes were reported by participants this outcome is susceptible to bias from lack of blinding. None of these studies blinded participants and outcome assessors to the treatment allocation.

²No serious inconsistency: In the complete analysis of five trials statistical heterogeneity was very high (I^2 statistic = 99%). However, this heterogeneity was related to a single trial showing very large effects conducted in an emergency setting in Liberia possibly due to epidemic diarrhoea. When this trial was removed as an outlier, there was a smaller, but more consistent effect.

³No serious indirectness: the studies were conducted in rural areas in Guatemala (two studies), and Kenya (one study), one trial was from a camp for displaced persons in Liberia and one from squatter settlements in Pakistan. Sanitation was improved in only one of these studies.

⁴No serious imprecision: all five studies found benefits with flocculation. The 95% CI of the pooled effect includes the possibility of no effect, but this imprecision is a result of the heterogeneity between studies.

Table 12. POU filtration: description of interventions

Study ID	Intervention sub-group	Study design	Setting	Intervention areas			Control areas	
				Water quality intervention	Health promotion activities	Compliance	Water source	Health promotion activities
Abebe 2014 ZAF	Ceramic filter	Cluster-RCT	Rural	Ceramic water filter impregnated with silver nanoparticles with safe storage containers	Education about safe water and hygiene and information on how to use the filter and maintain it.	Not reported	Personal tap in home (44%), community tap (44%) and river (3%)	Received usual clinical care including education about safe water and hygiene at the clinic
Brown 2008a KHM	Ceramic filter	Cluster-RCT	Rural	CWP (Cambodian Ceramic Water Purifier) including safe storage container.	None	98% compliance measured by self-report	Surface water (55%) and ground water (48%) during the dry season and surface water (45%), ground water (48%) and rain water (73%) during the rainy season	None
Brown 2008b KHM	Ceramic filter	Cluster-RCT	Rural	CWP-Fe (iron-rich ceramic water purifier) including safe storage container.	None	98% compliance measured by self-report	Surface water (55%) and ground water (48%) during the dry season and surface water (45%), ground water (48%) and rain water (73%) during the rainy season	None
Clasen 2004b BOL	Ceramic filter	Cluster-RCT	Rural	Ceramic filters including improved storage	None	67% of households had filters in regular use	68% had taps and 11% boiled water.	None
Clasen 2004c BOL	Ceramic filter	Cluster-RCT	Rural	Ceramic filters including improved storage	None	100% of intervention households' water free of TTC	Water from canal (52%), river (35%) or rainwater (4%)	None
Clasen 2005 COL	Ceramic filter	Cluster-RCT	Rural and urban affected by conflict	Ceramic water filter system including improved storage	None	Not reported	River (27.6%), rainwater (12.1%), yard tap (67.2%). 70.7% claimed to treat water.	None

Table 12. POU filtration: description of interventions (Continued)

du Preez 2008 ZAF/ZWE	Ceramic filter	Cluster-RCT	Rural	Ceramic filters including improved storage	None	55% compliance measured by water quality (approximate compliance across intervention households in Zimbabwe and South Africa).	Protected water source (53.8%) and unprotected water source (46.2%)	None
Lindquist 2014a BOL	Ceramic filter	Cluster-RCT	Peri-urban	Received a PointONE Filter and a 30 L bucket (with lid)	Participants were instructed on diarrhoeal transmission (biological versus cultural beliefs-based), prevention and treatment.	97% compliance based on reported use	83% used water from tanker trucks and 12% from water coolers.	Received weekly messages on life skills and attitudes. Also were instructed on diarrhoeal transmission, prevention and treatment.
Lindquist 2014b BOL	Ceramic filter	Cluster-RCT	Peri-urban	Received a PointONE Filter and a 30-L bucket (with lid) and WASH education	Participants received weekly WASH messages on personal and family hygiene, sanitation, boiling and chlorine-based water treatments (excluding filtration), vitamin A, hygienic food preparation and cleaning, and parasite prevention.	90% compliance based on reported use	83% used water from tanker trucks and 12% from water coolers.	Received weekly messages on life skills and attitudes. Also were instructed on diarrhoeal transmission, prevention and treatment.

Table 12. POU filtration: description of interventions (Continued)

URL 1995a GTM	Ceramic filter	Cluster-RCT	Rural	Handmade ceramic water filter	None	87% to 93% use of filter by children	Majority of households collected water from household tap (not chlorinated)	None
URL 1995b GTM	Ceramic filter	Cluster-RCT	Rural	Handmade ceramic water filter	Education on nutrition (ORS, basic nutrition and maternal and child nutrition), health (hygiene) and family values.	As above	Majority of households collected water from household tap (not chlorinated)	None
Fabiszews-ki 2012 HND	Sand filtration	Cluster-RCT	Rural	Hydradid plastic-housing BioSand filter (BSF) + 20 L water jug	Training for the use and maintenance of the BSF and general education about hygiene and sanitation.	Not reported	Among all study participants- the main source of drinking water were: protected water sources (49% to 69% households per month), protected sources (24% to 50% per month), piped water (1% to 11% per month), and rainwater (0% to 2% per month).	Training for the use and maintenance of the BSF and general education about hygiene and sanitation.
Stauber 2009 DOM	Sand filtration	Cluster-RCT	Semi-rural and urban	Received a biosand filter and safe storage container	Nothing	Water quality testing, however no intervention household level compliance reported	42% reported treating drinking water.	None
Stauber 2012a KHM	Sand filtration	Cluster-RCT	Rural	Plastic biosand filter. HHs were asked to pay USD 10 for the filter.	Health and hygiene education sessions	89% compliance measured by household-reported use at least 3 times per week	Improved water sources during the dry season (7.1%) and during the rainy season (88.9%). 49.5% reported boiling drinking water.	Health and hygiene education sessions
Stauber 2012b GHA	Sand filtration	Cluster-RCT	Rural	Plastic biosand filter	Not specified	97% compliance measured by house-	Use surface water during dry season (95%) and use surface water during rainy season (70.6%). 96.5% re-	nothing

Table 12. POU filtration: description of interventions (Continued)

						hold-reported use	ported sieving drinking water through cloth.	
Tiwari 2009 KEN	Sand filtration	Cluster-RCT	Rural	Provided with the concrete BioSand Filter	At each visit, three oral rehydration packets and instructions were provided.	Not reported	All control houses reported drinking river or unprotected spring water; drink rainwater (96.6%), drink improved source (24.1%). 34.5% reported boiling drinking water.	At each visit, three oral rehydration packets and instructions were provided.
Boisson 2009 ETH	LifeStraw® Personal	Cluster-RCT	Rural	A LifeStraw® personal pipe-style water treatment device was given to each member of the household >6 months and encouraged to use it at home and away from home.	None	13% report use today	The primary drinking water source for 84% was from spring, 12% from rivers, 2.5% from hand dug wells and 4% from communal taps.	None
Boisson 2010 DRC	LifeStraw® Family	Cluster-RCT	Rural	Households received a LifeStraw® Family filters	None	76% compliance measured by self-report use today or yesterday (at 14 month follow-up)	Received a placebo filter.	None
Peletz 2012 ZMB	LifeStraw® Family	Cluster-RCT	Peri-urban	Households received a LifeStraw® Family filter and two 5 L safe storage containers.	None	87% compliance measured by improved water quality	46% use unprotected dug wells, 19% boreholes, 17% public standpipes, 12% protected dug well, 5% piped into home or yard and 2% surface water.	None
Colford 2002 USA	Plumbed in filter	Cluster-RCT	Urban	Installation of water treatment devices to 1 tap in HH that include: a 1-micron absolute prefilter cartridge and a UV lamp.	None	96% compliance measured by not dropping out of study (plumbed-in unit)	Sham device	None
Colford 2005 USA	Plumbed in filter	Cluster-RCT	Urban	Installation of filter (1-micron filter and a UV	All participants received the current CDC	90% compliance measured by not dropping	Sham device	All participants received the

Table 12. POU filtration: description of interventions *(Continued)*

				lamp) to main faucet of household	safe drinking water guidelines for immunocompromised persons	out of study (filter attached to kitchen sink)		current CDC safe drinking water guidelines for immunocompromised persons
Colford 2009 USA	Plumbed in filter	Cluster-RCT	Urban	Installation of filter (1-micron filter and a UV lamp) to main faucet of household	None	83% compliance measured by not dropping out of study (filter attached to kitchen sink)	Sham device	None
Rodrigo 2011 AUS	Ceramic filter/plumbed in	Cluster-RCT	Urban	Bench-top silver impregnated ceramic water treatment units, which required participants to use fill it but then households that had rainwater piped into kitchen were offered an under sink unit	None	Not reported	Sham water treatment unit	None

Table 13. POU filtration: primary drinking water supply and sanitation facilities

Trial	Description	Source ¹	Access to source ²	Quantity available ³	Ambient H ₂ O quality	Sanitation ⁴
Abebe 2014 ZAF	In-home taps or community taps	Improved	Sufficient	Unclear	80% of households had contamination between 10 to 10000 CFUs/100 mL	Unclear
Brown 2008	62% households rely on surface water during dry season and 55% rely on surface water during rainy season	Unimproved	Unclear	Unclear	Baseline not reported. Control households: Geometric mean 600 <i>E. coli</i> /100 mL	Improved
Clasen 2004b BOL	80% yard taps supplied by untreated surface source, 20% directly from untreated surface sources	80% improved, 20% unimproved	Sufficient	Sufficient	Baseline arithmetic mean 86 TTC/100 mL	Unimproved
Clasen 2004c BOL	Irrigation canals and other surface sources	Unimproved	Sufficient	Sufficient	Baseline arithmetic mean 797 TTC/100 mL	Unimproved
Clasen 2005 COL	67% yard tap from municipality (not treated), 28% river, 12% rainwater	Unimproved	Unclear	Unclear	Baseline not reported. Control households: arithmetic mean 151 TTC/100 mL	Mostly improved
du Preez 2008 ZAF/ZWE	Protected wells	Improved	Sufficient	Unclear	Baseline not reported. Control households: 30% samples post-intervention met WHO guidelines for water quality	Improved
Lindquist 2014	Municipal supply	Improved	Sufficient	Unclear	Not tested	Unimproved
URL 1995	Household tap (27%), public tap (21%), well (23%)	Improved	Unclear	Unclear	Range 5 to 260; average 106 faecal coliforms/100 mL across three sites.	Improved
Fabiszewski 2012 HND	49% to 69% households use unprotected sources, 24% to 50% use protected sources, 1% to 11% piped water, 0% to 2% rainwater	Improved and unimproved	Unclear	Unclear	Geometric mean <i>E. coli</i> concentrations of both unprotected and protected sources were > 100 MPN/100 mL	Unimproved
Stauber 2009 DOM	Unclear	Unclear	Unclear	Unclear	Baseline: geometric mean 21 MPN <i>E. coli</i> /100 mL	Improved
Stauber 2012a KHM	77% used improved water source during dry season, 89% during rainy season	Improved	Unclear	Unclear	Baseline: geometric mean 27.5 CFU/100 mL	Unimproved
Stauber 2012b GHA	Surface water 70% in dry season, 95% in rainy season	Unimproved	Unclear	Unclear	Baseline: geometric mean 792 or 832 <i>E. coli</i> /100 mL	Unimproved

Table 13. POU filtration: primary drinking water supply and sanitation facilities *(Continued)*
 for control and intervention households, respectively

Tiwari 2009 KEN	Primarily river water; 27% drink protected sources	Unimproved	Unclear	Unclear	Baseline not reported. Control households: 88.9 faecal coliforms/100 mL	Unclear
Boisson 2009 ETH	84% springs, 12% river, 2% handdug well, 4% communal tap	Unimproved	Unclear	Unclear	Baseline arithmetic mean 449 TTC/100 mL	Unimproved
Boisson 2010 DRC	97% surface water, 38% rainwater, 16% springs	Unimproved	Unclear	Unclear	Source drinking water: 75% of household samples > 1000 TTC/100 mL	Unimproved
Peletz 2012 ZMB	46% unprotected dug wells, 22% taps, 16% borehole or protected dug well, 2% surface water	Improved and unimproved	Unclear	Unclear	Unfiltered water: Geometric mean 190 TTC/100 mL	Unimproved
Colford 2002 USA	Household taps supplied by municipal water treatment	Improved	Sufficient	Sufficient	Data from water treatment plant: met US federal and California drinking water standards	Improved
Colford 2005 USA	Household taps supplied by municipal water treatment	Improved	Sufficient	Sufficient	Data from water treatment plant: met US federal drinking water standards	Improved
Colford 2009 USA	Household taps supplied by municipal water treatment	Improved	Sufficient	Sufficient	Data from water treatment plant: met US federal drinking water standards	Improved
Rodrigo 2011 AUS	Untreated rainwater	Improved	Sufficient	Sufficient	Not tested	Improved

Abbreviations: TTC: thermotolerant coliforms, MPN: most probable number, CFU: colony-forming units

¹'Improved' includes household connection, public standpipe, borehole, protected dug well, protected spring, rainwater collection; 'unimproved' includes unprotected well, unprotected spring, vendor-provided water, bottled water; and 'unclear' means unclear or not reported; definition based on [WHO/UNICEF 2015](#).

²'Sufficient' means located within 500 m, queuing no more than 15 minutes, no more than three minutes to fill 20 L container, and maintained so available consistently; 'insufficient' means that it does not meet any of above; and 'unclear' means unclear or not reported; definition based minimum standards established by The [Sphere Project 2011](#).

³'Sufficient' means a minimum of 15 L/day/person; 'insufficient' means less than 15 L/day/person; and 'unclear' means unclear or not reported; definition based on minimum standards established by The [Sphere Project 2011](#).

⁴'Improved' means connection to a public sewer or septic system, pour flush latrine, simple pit latrine, or ventilated improved pit latrine; 'unimproved' means service or bucket latrine, public latrines, open latrines; and 'unclear' means unclear or not reported; definition based on [WHO/UNICEF 2015](#).

Table 14. Summary of findings: POU filtration
POU filtration compared with no intervention for preventing diarrhoea

Patient or population: adults and children

Settings: low-, middle- and high-income countries

Interventions to improve water quality for preventing diarrhoea (Review)

Copyright © 2015 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

Table 14. Summary of findings: POU filtration (Continued)

Intervention: distribution of water filters and instructions on use

Comparison: no intervention

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)
	Assumed risk	Corresponding risk			
	No intervention	Water filtration			
Diarrhoea episodes Cluster-RCTs	3 episodes per person per year	All filters	RR 0.48	15,582	⊕⊕⊕⊙
		1.4 episodes per person per year (1.1 to 1.8)	(0.38 to 0.59)	(18 trials)	moderate ^{1,2,3,4}
	3 episodes per person per year	Ceramic filters	RR 0.39 (0.29 to 0.53)	5763	⊕⊕⊕⊙
		1.1 episodes per person per year (0.8 to 1.5)		(8 trials)	moderate ^{2,4,5,6}
		Biosand filters	RR 0.47	5504	⊕⊕⊕⊙
		1.4 episodes per person per year (1.2 to 1.7)	(0.39 to 0.57)	(4 trials)	moderate ^{4,7,8,9}
	LifeStraw® filters	RR 0.69	3259	⊕⊕⊙⊙	
	2.1 episodes per person per year (1.5 to 2.8)	(0.51 to 0.93)	(3 trials)	low ^{2,4,10,11}	
	Plumbed filters	RR 0.73	1056	⊕⊕⊕⊙	
	2.2 episodes per person per year (1.6 to 3.1)	(0.52 to 1.03)	(3 trials)	moderate ^{2,4,12,13}	

The basis for the **assumed risk** is provided in the footnotes. The **corresponding risk** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **RR:** risk ratio.

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: we are very uncertain about the estimate.

The assumed risk is based on 2.9 episodes/child year in 2010 (Fischer Walker 2012).

¹Downgraded by 1 for serious risk of bias: as diarrhoea episodes were reported by participants, this outcome is susceptible to bias from lack of blinding. Only five studies blinded participants and outcome assessors to the treatment allocation and only one found an effect of the intervention.

²No serious inconsistency: statistical heterogeneity was very high, however there is consistency in the direction of the effect.

³No serious indirectness: these studies are from a variety of low-, middle-, and high-income countries (South Africa, Ethiopia, Democratic Republic of Congo, Cambodia, Bolivia, Colombia, USA, Australia, Honduras, Zimbabwe, Zambia, Dominican Republic, Ghana, Kenya and Guatemala).

⁴No serious imprecision.

⁵Downgraded by 1 for serious risk of bias: as diarrhoea episodes were reported by participants, this outcome is susceptible to bias from lack of blinding. Only one of these studies, [Rodrigo 2011 AUS](#), blinded participants and outcome assessors to the treatment allocation.

⁶No serious indirectness: these studies are from a variety of low-, middle-, and high-income countries (South Africa, Cambodia, Bolivia, Colombia, Zimbabwe, Guatemala and Australia). The interventions consisted of distribution of water filters (which included a safe storage chamber) plus instructions on how to use them. In some cases, the intervention included hygiene education.

⁷Downgraded by 1 for serious risk of bias: as diarrhoea episodes were reported by participants, this outcome is susceptible to bias from lack of blinding. None these studies blinded participants and outcome assessors to the treatment allocation.

⁸No serious inconsistency: there was no statistical heterogeneity between studies, I^2 statistic = 0%.

⁹No serious indirectness: the studies were conducted in a variety of rural and urban settings in a variety of low- and middle-income countries (Honduras, Dominican Republic, Cambodia, Ghana and Kenya). The interventions consisted of distribution of water filters plus instructions on how to use them. In some cases, the intervention included hygiene education and a separate storage vessel.

¹⁰Downgraded by 1 for serious risk of bias: as diarrhoea episodes were reported by participants this outcome is susceptible to bias from lack of blinding. Only one of these studies, [Boisson 2010 DRC](#), blinded participants and outcome assessors to the treatment allocation and found no evidence of effect of the filter.

¹¹Downgraded by 1 for some indirectness, the studies were only performed in three sub-Saharan African countries (Ethiopia, Democratic Republic of Congo, and Zambia).

¹²No serious risk of bias: the three studies blinded participants and outcome assessors to the treatment allocation.

¹³Downgraded by 1 for some indirectness, the three studies were only performed in the USA in water conditions that presumed to meet US EPA standards.

Table 15. POU solar disinfection (SODIS): description of the interventions

Study ID	Study design	Setting	Intervention areas			Control areas	
			Water quality intervention	Health promotion activities	Compliance	Water source	Health promotion activities
Conroy 1996 KEN	Quasi-RCT	Rural	Children were given two 1.5 L plastic bottles and told to keep the bottles on the roof of the hut throughout the day in full sunlight	None	100%- random checks by project workers uncovered no evidence of non-compliance	Children were given two 1.5 L plastic bottles and told to keep the bottles indoors	None
Conroy 1999 KEN	Quasi-RCT	Rural	Mothers were given plastic bottles and told to keep the bottles on the roof of the hut throughout the day in full sunlight	None	Not reported	Mothers were given plastic bottles and told to keep the bottles indoors	None
du Preez 2010 ZAF	Cluster-RCT	Peri urban	Received two 2 L polyethylene terephthalate (PET) bottles for each child. Carers were instructed to fill one bottle and place it in full, unobscured sunlight for a minimum of 6 h every day.	None	25% compliance measured by participants filling out diarrhoeal diaries at least 75% of the time	No SODIS bottles and maintain their usual practices	None
du Preez 2011 KEN	Cluster-RCT	Peri urban and rural	Received two 2 L PET bottles for each child. Carers were instructed to fill one bottle and place it in full, unobscured sunlight for a minimum of 6 h every day.	None	Not specified.	No SODIS bottles and maintain their usual practices	None
Mäusezhal 2009 BOL	Cluster-RCT	Rural	Households were supplied regularly with clean, PET bottles. They were instructed to expose the water-filled bottles for at least 6 h to the sun.	Households were taught about the importance and benefits of drinking only treated water, the germ-disease concept, and promoted hygiene measures such as	32% compliance measured by observation	Drinking water from spring (48.1%), tap (51.9%), river (22.1%), rain (14.9%) and dug well (14.9%)	None

Table 15. POU solar disinfection (SODIS): description of the interventions *(Continued)*

				safe drinking water storage and hand washing.			
McGuigan 2011 KHM	Cluster-RCT	Rural	Households were provided with two transparent 2 L plastic bottles for each child and a sheet of corrugated iron on which to place the bottles to expose them to sunlight. Carers were instructed to fill one bottle and place it in full, unobscured sunlight for a minimum of 6 h every day.	The parents or carers were given verbal and written information on the disease concept and a simple explanation of the solar disinfection process and its effect on the microbial quality of their drinking water and subsequently the health of their children	90% (5% of children having < 10 months of follow-up and 2.3% having < 6 months)	Almost all of the households (97%) obtained water from unprotected boreholes. An important subgroup of these, 25%, drew water from shallow tube wells fitted with hand pumps. The remainder used unprotected wells or surface ponds	None

Table 16. POU solar disinfection (SODIS): primary drinking water supply and sanitation facilities

Trial	Description	Source ¹	Access to source ²	Quantity available ³	Ambient H ₂ O quality	Sanitation ⁴
Conroy 1996 KEN	Open water holes, tank fed by untreated piped water supply.	Unimproved	Unclear	Unclear	Source water: 10 ³ CFU/100 mL	Unclear
Conroy 1999 KEN	Open water holes, tank fed by untreated piped water supply.	Unimproved	Unclear	Unclear	Source water: 10 ³ CFU/100 mL	Unclear
du Preez 2010 ZAF	39% standpipes, 28% protected borehole, 10% unprotected boreholes, protected springs	Mostly improved	Sufficient	Sufficient	Baseline not reported. Intervention households: 62% of samples met WHO guidelines for water quality; no significant difference from control households	Unclear
du Preez 2011 KEN	Spring, protected and unprotected dug wells protected, canals, other	Mostly unimproved	Unclear	Unclear	50% of samples from stored water had 10 CFU/100 mL or less; no significant difference for intervention and controls	Unclear
Mäusezhal 2009 BOL	48% spring, 52% tap, 22% river, 15% rain, 15% dug well	Improved and unimproved	Sufficient	Sufficient	Not tested	Unimproved
McGuigan 2011 KHM	97% households use unprotected sources: unprotected wells, surface ponds	Unimproved	Unclear	Unclear	Baseline not reported. Control households: geometric mean 48 CFU/100 mL	Unimproved

¹'Improved' includes household connection, public standpipe, borehole, protected dug well, protected spring, rainwater collection; 'unimproved' includes unprotected well, unprotected spring, vendor-provided water, bottled water; and 'unclear' means unclear or not reported; definition based on [WHO/UNICEF 2015](#).

²'Sufficient' means located within 500 m, queuing no more than 15 minutes, no more than three minutes to fill 20 L container, and maintained so available consistently; 'insufficient' means that it does not meet any of above; and 'unclear' means unclear or not reported; definition based minimum standards established by The [Sphere Project 2011](#).

³'Sufficient' means a minimum of 15 L/day/person; 'insufficient' means less than 15 L/day/person; and 'unclear' means unclear or not reported; definition based on minimum standards established by The [Sphere Project 2011](#).

⁴'Improved' means connection to a public sewer or septic system, pour flush latrine, simple pit latrine, or ventilated improved pit latrine; 'unimproved' means service or bucket latrine, public latrines, open latrines; and 'unclear' means unclear or not reported; definition based on [WHO/UNICEF 2015](#).

Table 17. Summary of findings: POU solar disinfection (SODIS)

POU solar disinfection (SODIS) of water compared with no intervention for preventing diarrhoea
Patient or population: adults and children
Settings: low- and middle-income countries
Intervention: distribution of plastic bottles with instructions on using them to treat water using the SODIS method.

Interventions to improve water quality for preventing diarrhoea (Review)

Copyright © 2015 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

Table 17. Summary of findings: POU solar disinfection (SODIS) (Continued)

Comparison: no intervention

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)
	Assumed risk	Corresponding risk			
	No intervention	SODIS			
Diarrhoea episodes Cluster-RCTs	3 episodes per person per year	1.9 episodes per person per year (1.3 to 2.8)	RR 0.62 (0.42 to 0.94)	3460 (4 trials)	⊕⊕⊕⊖ moderate 1,2,3,4
Diarrhoea episodes Quasi-RCTs	3 episodes per person per year	2.5 episodes per person per year (2.1 to 2.9)	RR 0.82 (0.69 to 0.97)	555 (2 studies)	⊕⊕⊖⊖ low ^{1,5,6,7}

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: we are very uncertain about the estimate.

The assumed risk is based on 2.9 episodes/child year in 2010 (Fischer Walker 2012).

¹Downgraded by 1 for serious risk of bias: as diarrhoea episodes were reported by participants this outcome is susceptible to bias from lack of blinding. None of these studies blinded participants and outcome assessors to the treatment allocation.

²No serious inconsistency: statistical heterogeneity was very high (I^2 statistic = 89%), however there is consistency in the direction of the effect. This heterogeneity may relate to differences in compliance across the studies, however compliance was not measured in the same way across studies.

³No serious indirectness: the studies were conducted in peri-urban South Africa (one study), peri-urban and rural Kenya (one study), rural Bolivia (one study) and rural Cambodia (one study).

⁴No serious imprecision: the average effect suggests that the intervention may reduce diarrhoea episodes by about one third.

⁵No serious inconsistency: statistical heterogeneity was low (I^2 statistic = 0%).

⁶Downgraded by 1 for serious indirectness: there are only two studies and both were conducted in the same province in Kenya (one study included children five to 16 years old and the other included children younger than six years old).

⁷No serious imprecision.

Table 18. POU UV: description of the interventions

Study ID	Study design	Setting	Intervention areas			Control areas	
			Water quality intervention	Health promotion activities	Compliance	Water source	Health promotion activities
Gruber 2013 MEX	Cluster-RCT	Rural	Promotion of the UV Tube disinfection technology and safe storage	Unclear	51% compliance measured by access to treatment device	Unclear	None

Table 19. POU UV: primary drinking water supply and sanitation facilities

Trial	Description	Source ¹	Access to source ²	Quantity available ³	Ambient H ₂ O quality	Sanitation ⁴
Gruber 2013 MEX	Unclear	Unclear	Unclear	Unclear	Baseline: 60% of samples with detectable <i>E. coli</i>	Improved

¹'Improved' includes household connection, public standpipe, borehole, protected dug well, protected spring, rainwater collection; 'unimproved' includes unprotected well, unprotected spring, vendor-provided water, bottled water; and 'unclear' means unclear or not reported; definition based on [WHO/UNICEF 2015](#).

²'Sufficient' means located within 500 m, queuing no more than 15 minutes, no more than three minutes to fill 20 L container, and maintained so available consistently; 'insufficient' means that it does not meet any of above; and 'unclear' means unclear or not reported; definition based minimum standards established by The [Sphere Project 2011](#).

³'Sufficient' means a minimum of 15 L/day/person; 'insufficient' means less than 15 L/day/person; and 'unclear' means unclear or not reported; definition based on minimum standards established by The [Sphere Project 2011](#).

⁴'Improved' means connection to a public sewer or septic system, pour flush latrine, simple pit latrine, or ventilated improved pit latrine; 'unimproved' means service or bucket latrine, public latrines, open latrines; and 'unclear' means unclear or not reported; definition based on [WHO/UNICEF 2015](#).

Table 20. POU Improved storage: description of the interventions

Study ID	Study design	Setting	Intervention areas			Control areas	
			Water quality intervention	Health promotion activities	Compliance	Water source	Health promotion activities
Günther 2013 BEN	Cluster-RCT	Rural	Provided households with a new 30 L household water storage with a tap at the bottom, a new plastic container to transport water from the water source to the household and a sign attached to the transport and storage containers which emphasized the importance of avoiding hand-contact with the water and to only use water from an improved water source.	None	After 7 months, 88% of households were still using the improved storage containers	68% only consume improved water source	None
Roberts 2001 MWI	Cluster-RCT	Refugee camp	All of the participating household's water collection vessels were exchanged for improved buckets (20 L with a narrow opening to limit hand entry). Households were offered 1 improved bucket in exchange for 1 vessel, 2 for 2, and 3 improved buckets for any number of containers > 2. Households were asked never to put their hands in the improved buckets and were shown how to rinse the bucket without hand entry.	None	Intervention householders received buckets; actual use was not reported	Provided with 20 L standard ration bucket	None

Table 21. POU Improved storage: primary drinking water supply and sanitation facilities

Trial	Description	Source ¹	Access to source ²	Quantity available ³	Ambient H ₂ O quality	Sanitation ⁴
Günther 2013 BEN	Public tap or pump	Improved	Sufficient	Unclear	12% source water contaminated (≥ 1000 CFU per 100 mL)	Unclear
Roberts 2001 MWI	Traditional pots or standard ration buckets filled at refugee camp water point	Improved	Unclear	Unclear	Source water: 71% of samples had ≤ 1 faecal coliform/100 mL	Unclear

¹'Improved' includes household connection, public standpipe, borehole, protected dug well, protected spring, rainwater collection; 'unimproved' includes unprotected well, unprotected spring, vendor-provided water, bottled water; and 'unclear' means unclear or not reported; definition based on [WHO/UNICEF 2015](#).

²'Sufficient' means located within 500 m, queuing no more than 15 minutes, no more than three minutes to fill 20 L container, and maintained so available consistently; 'insufficient' means that it does not meet any of above; and 'unclear' means unclear or not reported; definition based minimum standards established by The [Sphere Project 2011](#).

³'Sufficient' means a minimum of 15 L/day/person; 'insufficient' means less than 15 L/day/person; and 'unclear' means unclear or not reported; definition based on minimum standards established by The [Sphere Project 2011](#).

⁴'Improved' means connection to a public sewer or septic system, pour flush latrine, simple pit latrine, or ventilated improved pit latrine; 'unimproved' means service or bucket latrine, public latrines, open latrines; and 'unclear' means unclear or not reported; definition based on [WHO/UNICEF 2015](#).

Table 22. Summary of findings: POU improved water storage
Improved water storage compared with no intervention for preventing diarrhoea

Patient or population: adults and children in sub-Saharan Africa

Settings: areas with improved water sources

Intervention: distribution of improved water containers

Comparison: no intervention

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	Number of participants (studies)	Quality of the evidence (GRADE)
	Assumed risk	Corresponding risk			
	No intervention	Water storage			
Diarrhoea episodes	3 episodes per person per year	2.7 episodes per person per year	RR 0.91 (0.74 to 1.11)	1871 (2 trials)	⊕⊕⊕⊕ low 1,2,3,4
Cluster-RCTs		(2.2 to 3.3)			

The basis for the **assumed risk** is the median control group risk across studies. The **corresponding risk** (and its 95% CI) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: confidence interval; **RR:** risk ratio.

GRADE Working Group grades of evidence

High quality: further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Interventions to improve water quality for preventing diarrhoea (Review)

Table 22. Summary of findings: POU improved water storage (Continued)

Very low quality: we are very uncertain about the estimate.

The assumed risk is based on 2.9 episodes/child year in 2010 (Fischer Walker 2012).

¹Downgraded by 1 for serious risk of bias: as diarrhoea episodes were reported by participants this outcome is susceptible to bias from lack of blinding. None of these studies blinded participants and outcome assessors to the treatment allocation.

²No serious inconsistency.

³Downgraded by 1 for indirectness: only 2 studies, from rural Benin and a refugee camp in Malawi, have been conducted to assess improved water storage.

⁴No serious imprecision.

Table 23. Estimates of household-level interventions after adjustment for non-blinding

POU intervention	Number of comparisons	Not adjusted for non-blinding		Adjusted for non-blinding	
		RR	95% CI	RR	95% CI
All	55	0.56	(0.46 to 0.68)	0.70	(0.64 to 0.77)
Chlorination	19	0.72	(0.61 to 0.84)	0.80	(0.69 to 0.92)
Filtration	23	0.48	(0.38 to 0.59)	0.62	(0.55 to 0.70)
Flocculation and disinfection	7	0.48	(0.20 to 1.16)	0.65	(0.40 to 1.09)
SODIS	6	0.68	(0.53 to 0.89)	0.80	(0.60 to 1.01)

Abbreviation: SODIS: solar disinfection; CI: confidence interval.

Table 24. Potential reasons for finding of no-effect in trials with adequate blinding

Study	Risk from ambient water quality	Compliance	Other issues
Colford 2002 USA	Very low (USA)	High (Sham filter)	None
Colford 2005 USA	Very low (USA)	High (Sham filter)	None
Colford 2009 USA	Very low (USA)	High (Sham filter)	None
Rodrigo 2011 AUS	Very low (Australia)	Not reported	None
Jain 2010 GHA	Low (11 CFU/100 mL)	High (RFC)	Control group received jerry can; 13 week follow-up
Kirchhoff 1985 BRA	Very high (mean 16000 FC/dL)	Not reported	Only 112 persons from 16 households; 18 week trial
Austin 1993	High (1871 FC/100 mL)	Low ("50% to 60%")	No test of blinding; not peer reviewed
Boisson 2010 DRC	High (75% of samples > 1000 TTC/100 mL)	High, but 73% of adults and 95% of children drank from untreated sources	"Placebo" removed > 90% of TTC in control arm

Table 24. Potential reasons for finding of no-effect in trials with adequate blinding (Continued)

Boisson 2013 IND	Moderate (mean 122 TTC/100 mL)	Low and inconsistent (32% of samples positive for RFC)	None
-------------------------	--------------------------------	--	------

Abbreviations: TTC: thermotolerant coliforms, CFU: colony-forming units, FC: faecal coliforms, RFC: residual free chlorine.

APPENDICES

Appendix 1. Search methods: detailed search strategies

Search set	CIDG SR ^a	CENTRAL	MEDLINE ^b	EMBASE ^b	LILACS ^b
1	water	WATER PURIFICATION	WATER PURIFICATION	WATER PURIFICATION	water
2	purification OR treatment OR chlorination OR decontamination OR filtration OR supply OR storage OR consumption	WATER MICROBIOLOGY	WATER MICROBIOLOGY	WATER MICROBIOLOGY	purification OR treatment OR chlorination OR decontamination OR filtration OR supply OR storage OR consumption
3	diarrhea	1 OR 2	1 OR 2	1 OR 2	diarrhea
4	1 AND 2 AND 3	water	water	water	1 AND 2 AND 3
5	—	purification OR treatment OR chlorination OR decontamination OR filtration OR supply OR storage OR consumption OR drink*	purification OR treatment OR chlorination OR decontamination OR filtration OR supply OR storage OR consumption OR drink*	purification OR treatment OR chlorination OR decontamination OR filtration OR supply OR storage OR consumption OR drink\$	—
6	—	4 AND 5	4 AND 5	4 AND 5	—
7	—	3 OR 6	3 OR 6	3 OR 6	—
8	—	DIARRHEA/EPIDEMIOL- OGY	DIARRHEA/EPIDEMIOL- OGY	DIAR- RHEA/EPIDEMIOL- OGY	—
9	—	DIARRHEA/MICROBIOLO- OGY	DIARRHEA/MICROBIOLO- OGY	DIARRHEA/PREVEN- TION	—
10	—	DIARRHEA/PREVENTION AND CONTROL	DIARRHEA/PREVENTION AND CONTROL	waterborne infection\$	—
11	—	waterborne infection*	waterborne infection*	cholera OR shigell\$ OR dysenter\$ OR crypt- osporidi\$ OR giardia\$ OR Escherichia coli OR clostridium	—

(Continued)

12	—	INTESTINAL DISEASES	INTESTINAL DISEASES	ENTEROBACTERIACEAE	—
13	—	cholera OR shigell* OR dysenter* OR cryptosporidi* OR giardia* OR Escherichia coli OR clostridium	cholera OR shigell* OR dysenter* OR cryptosporidi* OR giardia* OR Escherichia coli OR clostridium	8-12/OR	—
14	—	ENTEROBACTERIACEAE	ENTEROBACTERIACEAE	7 AND 13	—
15	—	8-14/OR	8-14/OR	LIMIT 14 TO HUMAN	—
16	—	7 AND 15	7 AND 15	—	—
17	—	—	LIMIT 16 TO HUMAN	—	—

^aCochrane Infectious Diseases Group Specialized Register.

^bSearch terms used in combination with the search strategy for retrieving trials developed by Cochrane ([Higgins 2005](#)); upper case: MeSH or Emtree heading; lower case: free text term.

Appendix 2. Data extracted from included studies

Type	Fields
Trial data	Country and setting (urban, rural)
	Number of participants/groups
	Unit of randomization, and whether measurement of effect adjusts for clustering where randomization is other than individual
	Definition and practices of control group
	Type and details of water quality intervention (filtration, flocculation, chemical disinfection, heat, or UV radiation)
	Other components of intervention (hygiene message, improved supply, improved sanitation, improved storage)
	Whether water protected to POU (i.e. by pipe, residual disinfection, or safe storage)
	Case definition of diarrhoea
	Method for diarrhoea assessment (self-reported, observed, or clinically confirmed)
	Where self reported, recall period used
	Study duration; Adherence rates
	Publication status
	Prescribed criteria of methodological quality

Interventions to improve water quality for preventing diarrhoea (Review)

Copyright © 2015 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

(Continued)

Individual characteristics	Age group
	Type and description of water source
	Level of faecal contamination of control water (low (< 100 thermotolerant coliforms (TTC)/100 mL), medium (100 to 1000 TTC/100 mL), and high (> 1000 TTC/100 mL))
	Causative agents identified (yes or no)
	Water collection, storage, and drawing practices
	Distance to and other constraints regarding water supply
	Sanitation facilities (improved or unimproved)
	Hygiene practices
	Outcomes
	Diarrhoea morbidity and 95% CI for each age group reported
	Manner of measuring diarrhoea morbidity
	Mortality attributed to diarrhoea
	Rate of utilization of intervention and manner of assessing same

Abbreviations: POU: point of use; CI: confidence interval; UV: ultraviolet.

WHAT'S NEW

Date	Event	Description
21 October 2015	Amended	Amended author affiliations.

HISTORY

Protocol first published: Issue 2, 2004

Review first published: Issue 3, 2006

Date	Event	Description
15 October 2015	New search has been performed	The review authors updated the review, and included several new studies, a 'Summary of findings' table, and 'Risk of bias' assessments.
15 October 2015	New citation required and conclusions have changed	The review authors performed an updated literature search, reapplied the inclusion criteria, repeated data extraction, added new studies, and used the GRADE approach to assess the quality of the evidence. They also applied statistical methods to uni-

Interventions to improve water quality for preventing diarrhoea (Review)

Copyright © 2015 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

Date	Event	Description
		fy the measures of effect and applied additional criteria for sub-grouping based on study design, setting, and length of follow-up.

CONTRIBUTIONS OF AUTHORS

TC and SC conceived the review. TC coordinated the review. TC, KA, SB, RP, HC, and SC designed the review. TC and authors of the initial review drafted the protocol. SB and Cochrane Infectious Diseases Group (CIDG) performed the search strategy. SB and RP screened search results. KA, SB, and RP retrieved papers. SB and RP applied inclusion criteria. KA, SB, and RP extracted data. KA, SB, RP, HC, and FM computed estimates of effect. KA, TC, FM, and DS applied quality criteria. KA contacted study authors for additional information. TC, KA, HC, DS, and CIDG addressed statistical issues. KA entered data into [Review Manager \(RevMan\)](#). TC, KA, and DS drafted the review. SB, RP, HC, and SC commented on the review. TC, KA, HC, FM, and DS prepared tables. KA prepared figures. TC is guarantor of this Cochrane Review.

DECLARATIONS OF INTEREST

TC, KA, SB, and SC have provided research or consulting services for Unilever, Ltd., Medentech, Ltd., DelAgua Health and Science, Ltd., and Vestergaard-Frandsen SA who manufacture or sell household-based water treatment devices.

SOURCES OF SUPPORT

Internal sources

- Liverpool School of Tropical Medicine, UK.

External sources

- Department for International Development (DFID), UK.

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

Risk of bias has been assessed using GRADE rather than the original methods expressed in the protocol. Statistical methods have been used to pool odds ratios, rate ratios, RRs and longitudinal prevalence ratios. Subgrouping has been done separately for each water quality intervention, and additional subgrouping has been conducted based on study design and length of follow up. Data has been provided on adjustment of studies for non-blinding.

INDEX TERMS

Medical Subject Headings (MeSH)

Controlled Before-After Studies; Diarrhea [*prevention & control]; Drinking Water [standards]; Randomized Controlled Trials as Topic; Water Purification [*methods] [standards]; Water Supply [*standards]

MeSH check words

Adult; Child; Child, Preschool; Humans; Infant