

POSTER PRESENTATION

Open Access

Pitfalls of whole exome-sequencing: hidden *DYNC2H1* mutations in patients with Jeune asphyxiating thoracic dystrophy

H Arts^{1*}, M Schmidts^{2†}, EMHF Bongers¹, MM Oud¹, LEM Duijkers¹, Z Yap², J Stalker³, JL Yntema¹, A Hoischen¹, C Gilissen¹, JA Veltman¹, A Kutkowska-Kaźmierczak⁴, EJ Kamsteeg¹, PJ Scambler², PL Beales², NVAM Knoers^{5†}, R Roepman^{1†}, HM Mitchison^{2†}

From First International Cilia in Development and Disease Scientific Conference (2012) London, UK. 16-18 May 2012

In recent years whole-exome sequencing has been developed, a technique by which all exons of the genome (all the protein-coding DNA) can be sequenced at once. Here we show that whole-exome sequencing, using either 35 or 50 Mb Agilent kits for exome capture, was insufficient to detect pathogenic *DYNC2H1* variants in patients with Jeune asphyxiating thoracic dystrophy (ATD; MIM208500). Jeune syndrome is a rare inherited ciliopathy involving chondrodysplasia characterized by shortened ribs and long bones, and polydactyly, progressive kidney and liver disease as well as retinitis pigmentosa. Reduced thoracic capacity causes approximately 60% early lethality. *DYNC2H1* encodes a subunit of the dynein 1B motor that drives tip-to-base ciliary intraflagellar transport, and mutations have previously been associated both with embryonically lethal short rib-polydactyly and the milder, but overlapping Jeune asphyxiating thoracic dystrophy. Although the *DYNC2H1* gene was targeted in our whole-exome experiments many sequence reads were not properly aligned, resulting in 30-70% of the gene not being covered. Only a combination of whole-exome sequencing and a candidate gene approach (i.e. analysis of non-covered *DYNC2H1* exons using Sanger sequencing) enabled us to detect the missing *DYNC2H1* mutations. Whole-exome data analysis of the 90 exon *DYNC2H1* gene is therefore comparable to playing 'hide and seek', whereby certain mutations are easier to find than others according to their relative coverage. In conclusion, although whole-exome sequencing has revolutionized the

field of human genetics, our findings emphasize that next-generation sequencing also presents significant challenges for gene identification and for implementation of this technique in DNA diagnostics.

Author details

¹Department of Genetics, Radboud University Nijmegen Medical Centre, the Netherlands. ²Molecular Genetics Molecular Medicine Unit, UCL Institute of Child Health, UK. ³UK10K, Wellcome Trust Genome Center, Cambridge, UK. ⁴Department of Medical Genetics, Institute of Mother and Child, Poland. ⁵Department of Medical Genetics, University Medical Centre Utrecht, the Netherlands.

Published: 16 November 2012

doi:10.1186/2046-2530-1-S1-P80

Cite this article as: Arts et al.: Pitfalls of whole exome-sequencing: hidden *DYNC2H1* mutations in patients with Jeune asphyxiating thoracic dystrophy. *Cilia* 2012 1(Suppl 1):P80.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: h.arts@antrg.umcn.nl

† Contributed equally

¹Department of Genetics, Radboud University Nijmegen Medical Centre, the Netherlands

Full list of author information is available at the end of the article