

Available online at www.sciencedirect.com**ScienceDirect****Biomedical Journal**journal homepage: www.elsevier.com/locate/bj

Highlights

Tainted air: The link between pollution and Alzheimer's disease

Emma Louise Walton*

Staff Writer at the Biomedical Journal, 56 Dronningens Gate, 7012 Trondheim, Norway

ARTICLE INFO

Article history:

Available online 25 July 2018

Keywords:

Air pollution
Alzheimer's disease
Fusobacterium
Periodontitis

ABSTRACT

In this issue of the *Biomedical Journal*, we learn how air pollution may contribute to cognitive decline and even increase risk for Alzheimer's disease. We also highlight original research documenting the body's response to infection with a common oral pathogen. Finally, we learn how a cellular antioxidant protein protects against mitochondrial dysfunction in Parkinson's disease.

Spotlight on reviews

Tainted air: the link between air pollution and Alzheimer's disease

Without prevention, the incidence of Alzheimer's disease (AD) is expected to triple by the year 2050 [1]. The insurmountable pressure that this would place on health care systems has intensified the search for major preventable risk factors for AD. In this issue of the *Biomedical Journal*, Kilian and Kitazawa [2] discuss how one such risk factor for AD may literally be in the air we breath, and describe the potential mechanisms linking pollution to decline in brain function and AD.

Chronic exposure to polluted air is a major health issue worldwide, with the WHO estimating that a staggering 91% of the world's population lives in places where air pollution exceeds recommended health guidelines [3]. Polluted air

contains particulate matter (PM) of various sizes along with noxious compounds such as nitrogen and sulfur oxide species, carbon monoxide, metals and other inorganic compounds. PM contains sulfates, nitrates, ammonium, chlorides, carbon, and other biological material and dust, and is divided according to size: "ultrafine" ($PM < 100$ nm, $PM_{0.1}$), "fine" ($PM < 2.5$ μm , $PM_{2.5}$), and "coarse" ($PM < 10 \mu m$, PM_{10}) [4]. Once inhaled, fine and ultrafine PM are capable of crossing into the bloodstream where they are taken up by cells leading to oxidative stress and mitochondrial damage [5]. Ultrafine PM in particular is considered the most toxic form of PM, and may even be able to penetrate the brain directly through the olfactory nerve [6].

It has long been known that exposure to polluted air affects respiratory health [7], but could the effects be more widespread to extend even to the brain? Many epidemiological studies have focused on vehicle exhaust as a source of PM, measuring how urban vs. rural or distance to roadway affects

* Corresponding author.

E-mail address: ewalton86@gmail.com.

Peer review under responsibility of Chang Gung University.

<https://doi.org/10.1016/j.bj.2018.06.004>

2319-4170/© 2018 Chang Gung University. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1 The link between PM exposure and AD. PM and other neurotoxic substances are capable of infiltrating the brain where they lead to inflammation, oxidative stress, and ultimately the neurotoxicity underlying the cognitive effects in AD. Figure kindly provided by Kilian et al. [2].

brain function and cognition. Overall, these studies reveal an alarming trend: exposure to PM at any stage of life negatively impacts cognition. In childhood, exposure to PM negatively affects performance in multiple intelligence subscales [8] and is associated with poor motor coordination and response time [9]. Similarly, elderly residents living in highly polluted areas perform poorly in cognitive tests [10] and it is estimated that every $10 \mu\text{g}/\text{m}^3$ increase in exposure to black carbon is equivalent to 2 years cognitive decline by ageing [11]. Moreover, several studies support the notion that these cognitive effects translate into increased risk of dementia. For example, in a Canadian cohort of 2.2 million residents aged over 55 years, the incidence of dementia was significantly higher for those living within 50 m of a major roadway [12], and exposure to various pollutants including ozone [13] and nitrogen oxide [14] is associated with an increased risk of dementia. The exact risk seems to be modified by other environmental factors [15] along with genetic predisposition, since possessing copies of the APOE gene variant linked to dementia appears to exacerbate the effect of pollution [16].

These epidemiological studies are backed up by human pathological studies and animal studies revealing that PM exposure affects molecular pathways linked to AD. Individuals living in highly polluted areas accumulate higher amounts of $\text{A}\beta_{42}$ [17], a toxic form of β -amyloid, and show hyperphosphorylated tau pre-tangles in the olfactory bulb and hippocampus [18]. Likewise, in a mouse model of AD, exposure to ultrafine PM led to an increase in the expected amount of $\text{A}\beta$ plaques and reduced neuron density in the hippocampus [16]. These findings suggest that PM exposure can affect

amyloid processing in the brain. The mechanism by which this occurs is likely to involve oxidative stress and an inflammatory response gone haywire, characterized by the massive release of pro-inflammatory cytokines from glial cells, generation of reactive oxygen species and increase in cellular antioxidant proteins [19] [Fig. 1].

Still, many questions remain unanswered regarding the link between PM and AD. The reliance of epidemiological studies on proxy measures of exposure like distance to roadway means that it is difficult to determine whether any specific air pollutants increase AD risk. If this were indeed the case, it would be arguably easier to enact policy to limit the release of these compounds into the environment. There is also a need for epidemiological studies over longer time courses, since the time window for the development of AD is still unknown. Despite these unknowns, evidence is mounting that make an “airtight” case against pollution in AD, with a call to clean up our cities and urban environments of the future.

Spotlight on original articles

Tooth loss in periodontitis: the role of *Fusobacterium* and inflammation

We all know the importance of a daily routine of brushing and flossing, or we may otherwise suffer the consequences of gingivitis, an inflammation of the gums that is caused by the build-up of bacteria. If left untreated, gingivitis can lead to periodontitis, a serious gum infection that damages soft tissue

and destroys the bone supporting teeth. In this issue of the *Biomedical Journal*, Johnson et al. [20] investigate the *in vivo* responses to infection with one of the leading perpetrators of periodontitis, *Fusobacterium nucleatum*.

The epithelium surrounding our teeth is a haven for anaerobic bacteria like *F. nucleatum*, which can grow in biofilms to form dental plaque [21]. To defend against this colonization, the oral epithelium secretes antimicrobial peptides, which both penetrate the bacterial membrane and act as chemoattractants to recruit immune cells to the site of infection [22]. Gingival epithelial cells (GECs) themselves also alerted to the infection when bacterial molecules bind to host pathogen recognition receptors on their surface. As a result, GECs activate NF- κ B and express cytokines and chemokines to call in for reinforcement innate and adaptive immune cells [23]. Some oral bacteria, like *Porphyromonas gingivalis*, are able to evade these host responses by negatively regulating the production of pro-inflammatory cytokines in GECs [24]. *F. nucleatum* on the other hand, triggers inflammatory responses [25], but infection with this bacteria in isolation had not been studied before in mice until now.

To investigate responses to *F. nucleatum*, Johnson et al. [20] infected BALB/c mice and collected maxillas at several time points after infection. Consistent with previous findings, *F. nucleatum* infection caused a strong pro-inflammatory response characterized by the upregulation of several cytokines, including IL-1 β and IFN- γ . This response was accompanied by the infiltration of immune cells, notably macrophages, into the maxilla. Although the goal of inflammation is to resolve the infection, pro-inflammatory cytokines like IL-1 β and IFN- γ can activate osteoclasts, which break down bone tissue [26]. Using immunohistochemistry, Johnson et al. detected bone resorption pits in the alveolar bone seven days after infection, which were accompanied by a striking loss in bone elasticity as measured with an atomic force microscope, as early as one day after infection.

Overall, these findings reveal that the oral cavity launches a strong pro-inflammatory attack when challenged with *F. nucleatum*. Unlike many other pathogenic oral bacteria, *F. nucleatum* does little to hide from these host defenses and as such, may inadvertently promote bone resorption and tooth loss through macrophage recruitment and osteoclast activation, very early during the infection process.

Also in this issue

Review

Autophagy inhibition drives liver cancer

In this short review, Chen et al. [27] discuss how the inhibition of autophagy drives the progression of liver cancer and describe the molecular mechanisms in the tumor microenvironment underpinning this effect.

Original articles

Nrf2 protects against mitochondrial dysfunction in Parkinson's disease

It has been known for many years that defects in mitochondria play a pivotal role in Parkinson's disease [28]. Here, Fu

et al. [29] examine the relationship between mitochondrial dysfunction and the accumulation of α -synuclein (SNCA) into Lewy bodies, the pathological hallmark of Parkinson's disease. Using a SNCA mutant that is prone to aggregation, they find that the accumulation of SNCA is accompanied by an increase in reactive oxygen species and decrease in both mitochondrial fitness and mitochondrial DNA copy number. Interestingly, these effects could be reversed by activating nuclear factor/erythroid-derived 2-like 2 (Nrf2), a transcription factor that regulates many anti-oxidant proteins [30].

Antimicrobial activity discovered in medicinal plant

Recent findings of an entirely new class of antibiotics in soil [31] suggest that many more antimicrobials are awaiting discovery in the environment. Here, da Silva et al. [32] test the oral toxicity and antimicrobial activity of the South American medicinal plant *Baccharis trimera*. Not only was the plant well tolerated in rats, it also inhibited the growth of Gram positive and Gram negative bacteria and limited biofilm production. These promising findings suggest that the hunt for new antimicrobials may prove fruitful by taking a "leaf" out of the book of nature.

Risk factors for hip fracture among elderly Taiwanese women
Taiwan has one of the highest incidences of hip fracture in the world [33], in particular among postmenopausal women. In this case-control study, Chen et al. [34] investigate the factors that put elderly Taiwanese women at risk of hip fracture and find that education level and total hip bone mineral density were the strongest predictors of first incident hip fracture.

Correspondence

Kaposi sarcoma-like lesions found in other skin conditions

In Kaposi's sarcoma, patches of abnormal tissue growing under the skin that appear as multi-colored areas or the so-called "rainbow pattern" are considered a hallmark of the disease. However, in this correspondence, Kelati and Mernissi [35] report finding these Kaposi-like lesions in various dermatological conditions, suggesting that they are more widespread than previously thought.

Conflicts of interest

The author declares no conflicts of interest.

REFERENCES

- [1] <https://www.alzinfo.org/articles/alzheimers-cases-triple-2050/>. [Accessed 25 June 2018].
- [2] Kilian J, Kitazawa M. The emerging risk of exposure to air pollution on cognitive decline and Alzheimer's disease – evidence from epidemiological and animal studies. *Biomed J* 2018;41:141–62.
- [3] World Health Organization. Air pollution, <http://www.who.int/airpollution/en/>. [Accessed 25 June 2018].

- [4] U.S. EPA. Air quality criteria for particulate matter (final report, 2004). Washington, DC: U.S. Environmental Protection Agency; 2004. EPA 600/P-99/002aF-bF.
- [5] Li N, Sioutas C, Cho A, Schmitz D, Misra C, Sempf J, et al. Ultrafine particulate pollutants induce oxidative stress and mitochondrial damage. *Environ Health Perspect* 2003;111:455–60.
- [6] Block ML, Calderon-Garcidueñas L. Air pollution: mechanisms of neuroinflammation & CNS disease. *Trends Neurosci* 2009;32:506–16.
- [7] Khafaei MA, Yajnik CS, Salvi SS, Ojha A. Critical review of air pollution health effects with special concern on respiratory health. *J Air Pollut Health* 2016;1:123–36.
- [8] Calderon-Garcidueñas L, Engle R, Antonieta Mora-Tiscareño AM, Styner M, Gomez-Garza G, Zhu H, et al. Exposure to severe urban air pollution influences cognitive outcomes, brain volume and systemic inflammation in clinically healthy children. *Brain Cognit* 2011;77:345–55.
- [9] Wang S, Zhang J, Zeng X, Zeng Y, Wang S, Chen S. Association of traffic-related air pollution with children's neurobehavioral functions in Quanzhou, China. *Environ Health Perspect* 2009;117:1612–8.
- [10] Sánchez-Rodríguez MA, Santiago E, Arronte-Rosales A, Vargas-Guadarrama LA, Mendoza-Nuñes VM. Relationship between oxidative stress and cognitive impairment in the elderly of rural vs. urban communities. *Life Sci* 2006;78:1682–7.
- [11] Power MC, Weisskopf MG, Alexeeff SE, Coull BA, Avron S, Schwartz J. Traffic-related air pollution and cognitive function in a cohort of older men. *Environ Health Perspect* 2011;119:682–7.
- [12] Chen H, Kwong JC, Copes R, Tu K, Villeneuve PJ, van Donkelaar A, et al. Living near major roads and the incidence of dementia, Parkinson's disease, and multiple sclerosis: a population-based cohort study. *Lancet* 2017;389:718–26.
- [13] Jung CR, Lin YT, Hwang BF. Ozone, particulate matter, and newly diagnosed Alzheimer's disease: a population-based cohort study in Taiwan. *J Alzheim Dis* 2015;44:573–84.
- [14] Oudin A, Forsberg B, Adolfsson AN, Lind N, Modig L, Nordin M, et al. Traffic-related air pollution and dementia incidence in Northern Sweden: a longitudinal study. *Environ Health Perspect* 2016;124:306–12.
- [15] Chen JC, Schwartz J. Neurobehavioral effects of ambient air pollution on cognitive performance in US adults. *Neurotoxicology* 2009;30:231–9.
- [16] Cacciottolo M, Wang X, Driscoll I, Woodward N, Saffari A, Reyes J, et al. Particulate air pollutants, APOE alleles and their contributions to cognitive impairment in older women and to amyloidogenesis in experimental models. *Transl Psychiatry* 2017;7:e1022.
- [17] Calderón-Garcidueñas L, Reed W, Maronpot RR, Henríquez-Roldán C, Delgado-Chavez R, Calderón-Garcidueñas A, et al. Brain inflammation and Alzheimer's-like pathology in individuals exposed to severe air pollution. *Toxicol Pathol* 2004;32:650–8.
- [18] Calderón-Garcidueñas L, Kavanaugh M, Block ML, D'Angilli A, Delgado-Chavez R, Torres-Jardón R, et al. Neuroinflammation, hyperphosphorylated tau, diffuse amyloid plaques and down-regulation of the cellular prion protein in air pollution exposed children and adults. *J Alzheim Dis* 2012;28:93–107.
- [19] Guerra R, Vera-Aguilar E, Uribe-Ramirez M, Gookin G, Camacho J, Osornio-Vargas AR, et al. Exposure to inhaled particulate matter activates early markers of oxidative stress, inflammation and unfolded protein response in rat striatum. *Toxicol Lett* 2013;222:146–54.
- [20] Johnson L, Almeida-da-Silva CLC, Takiya CM, Figliuolo V, Rocha GM, Weissmüller G, et al. Oral infection of mice with *Fusobacterium nucleatum* results in macrophage recruitment to the dental pulp and bone resorption. *Biomed J* 2018;41:184–93.
- [21] Hasan A, Palmer RM. A clinical guide to periodontology: pathology of periodontal disease. *Br Dent J* 2014;216:457–61.
- [22] Diamond G, Ryan L. Beta-defensins: what are they really doing in the oral cavity? *Oral Dis* 2011;17:628–35.
- [23] Laube DM, Dongari-Bagtzoglou A, Kashleva H, Eskdale J, Gallagher G, Diamond G. Differential regulation of innate immune response genes in gingival epithelial cells stimulated with Aggregatibacter actinomycetemcomitans. *J Periodontal Res* 2008;43:116–23.
- [24] Johnson L, Atanasova KR, Bui PQ, Lee J, Hung SC, Yilmaz O, et al. *Porphyromonas gingivalis* attenuates ATP-mediated inflammasome activation and HMGB1 release through expression of a nucleoside-diphosphate kinase. *Microb Infect* 2015;17:369–77.
- [25] Bui FQ, Johnson L, Roberts J, Hung SC, Lee J, Atanasova KR, et al. *Fusobacterium nucleatum* infection of gingival epithelial cells leads to NLRP3 inflammasome-dependent secretion of IL-1beta and the danger signals ASC and HMGB1. *Cell Microbiol* 2015;18:970–81.
- [26] Zupan J, Jeras M, Marc J. Osteoimmunology and the influence of pro-inflammatory cytokines on osteoclasts. *Biochem Med* 2013;23:43–63.
- [27] Chen KD, Lin CC, Tsai MC, Huang KT, Chiu KW. Tumor microenvironment mediated by suppression of autophagic flux drives liver cancer malignancy. *Biomed J* 2018;41:163–8.
- [28] Winklhofer K, Haass C. Mitochondrial dysfunction in Parkinson's disease. *Biochim Biophys Acta* 2010;1802:29–44.
- [29] Fu Mh, Wu CW, Lee YC, Hung CY, Chen IC, Wu KLH. Nrf2 activation attenuates the early suppression of mitochondrial respiration due to the α -synuclein overexpression. *Biomed J* 2018;41:169–83.
- [30] Ishii T, Itoh K, Takahashi S, Sato H, Yanagawa T, Katoh Y, et al. Transcription factor Nrf2 coordinately regulates a group of oxidative stress-inducible genes in macrophages. *J Biol Chem* 2000;275:16023–9.
- [31] Hover BM, Kim SH, Katz M, Charlop-Powers Z, Owen JG, Ternei MA, et al. Culture-independent discovery of the malacidins as calcium-dependent antibiotics with activity against multidrug-resistant Gram-positive pathogens. *Nat Microbiol* 2018;3:415–22.
- [32] Da Silva ARH, Lopes LQS, Cassanego GB, de Jesus PR, Figueiredo KC, Santos RCV, et al. Acute toxicity and antimicrobial activity of leaf tincture *Baccharis trimera* (Less.). *Biomed J* 2018;41:194–201.
- [33] Kanis JA, Oden A, McCloskey EV, Johansson H, Wahl DA, Cooper C, IOF Working Group on Epidemiology and Quality of Life. A systematic review of hip fracture incidence and probability of fracture worldwide. *Osteoporos Int* 2012;23:2239–56.
- [34] Chen FP, Fu TS, Lin YC, Fan CM. Risk factors and quality of life for the occurrence of hip fracture in postmenopausal women. *Biomed J* 2018;41:202–8.
- [35] Kelati A, Mernissi FZ. The rainbow pattern in dermoscopy: a zoom on nonkaposi sarcoma skin diseases. *Biomed J* 2018;41:209–10.