

## CORRECTION

# Correction: Dihydroxyacetone of wheat root exudates serves as an attractant for *Heterodera avenae*


Gaofeng Wang, Yunhe Wang, Hazem Abdelnabby, Xueqiong Xiao, Wenkun Huang, Deliang Peng, Yannong Xiao

An additional affiliation is missing for the third author. Hazem Abdelnabby is also affiliated with Department of Plant Protection, College of Agriculture, Benha University, Qaliubia, Egypt.

The following information is missing from the Funding statement: Yannong Xiao is supported by the National Key R & D Program of China (2018YFD0200500).

## Reference

1. Wang G, Wang Y, Abdelnabby H, Xiao X, Huang W, Peng D, et al. (2020) Dihydroxyacetone of wheat root exudates serves as an attractant for *Heterodera avenae*. PLoS ONE 15(7): e0236317. <https://doi.org/10.1371/journal.pone.0236317> PMID: 32702002



## OPEN ACCESS

**Citation:** Wang G, Wang Y, Abdelnabby H, Xiao X, Huang W, Peng D, et al. (2021) Correction: Dihydroxyacetone of wheat root exudates serves as an attractant for *Heterodera avenae*. PLoS ONE 16(5): e0251537. <https://doi.org/10.1371/journal.pone.0251537>

**Published:** May 6, 2021

**Copyright:** © 2021 Wang et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.