

ERRATUM Open Access



Erratum to: A zeta potential value determines the aggregate's size of penta-substituted [60]fullerene derivatives in aqueous suspension whereas positive charge is required for toxicity against bacterial cells

Dmitry G. Deryabin^{1*}, Ludmila V. Efremova^{1,4}, Alexey S. Vasilchenko^{1,2}, Evgeniya V. Saidakova³, Elena A. Sizova^{1,4}, Pavel A. Troshin⁵, Alexander V. Zhilenkov⁵ and Ekaterina A. Khakina⁵

Erratum to: J Nanobiotechnol (2015) 13:50 DOI 10.1186/s12951-015-0112-6

Unfortunately, the original version of this article [1] contained an error. In the author list, Ekaterina A. Khakina was misspelt as Ekaterina E. Khakina. This has now been corrected in the original version.

Author details

- ¹ Department of Microbiology, Orenburg State University, Orenburg, Russia.
- ² Institute of Cellular and Intracellular Symbiosis, RAS, Orenburg, Russia.
- ³ Institute for Ecology and Genetics of Microorganisms, RAS, Perm, Russia.
- ⁴ All-Russia Research Institute of Beef Cattle Breeding, RAS, Orenburg, Russia.
- ⁵ Institute for Problems of Chemical Physics of RAS, Chernogolovka, Russia.

The online version of the original article can be found under doi:10.1186/s12951-015-0112-6.

Received: 28 September 2015 Accepted: 28 September 2015 Published online: 13 October 2015

Doforonce

 Deryabin DG, Efremova LV, Vasilchenko AS, Saidakova EV, Sizova EA, Troshin PA, Zhilenkov AV, Khakina EE. A zeta potential value determines the aggregate's size of penta-substituted [60] fullerene derivatives in aqueous suspension whereas positive charge is required for toxicity against bacterial cells. J Nanobiotechnol. 2015;13:50. doi:10.1186/ s12951-015-0112-6.

Full list of author information is available at the end of the article



^{*}Correspondence: dgderyabin@yandex.ru

¹ Department of Microbiology, Orenburg State University, Orenburg,