

Poster presentation

Open Access

## A new role of the HIV-1 nucleocapsid in the spatiotemporal control of the reverse transcription throughout the virus replication cycle

Bing Yu<sup>\*1,2</sup>, Laurent Houzet<sup>1,3</sup>, Ludovic Didierlaurent<sup>1</sup>, Célia Chamontin<sup>1</sup>, Zakia Morichaud<sup>1</sup>, Jean Luc Darlix<sup>4</sup> and Marylène Mougel<sup>1</sup>

Address: <sup>1</sup>CNRS UMR 5236-UMI/UMII, CPBS - Equipe «Assemblage et Réplication des Rétrovirus», Institut de Biologie, Montpellier, France, <sup>2</sup>Department of Pathogen Biology, Tongji Medical College of Huazhong University of Science and Technology, Wuhan, PR China, <sup>3</sup>Molecular Virology Section, Laboratory of Molecular Microbiology National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland 20892, USA and <sup>4</sup>LaboRetro, Unité de Virologie humaine INSERM U758, IFR128, ENS, 46 allée d'Italie, Lyon, France

\* Corresponding author

from *Frontiers of Retrovirology: Complex retroviruses, retroelements and their hosts* Montpellier, France. 21-23 September 2009

Published: 24 September 2009

*Retrovirology* 2009, **6**(Suppl 2):P14 doi:10.1186/1742-4690-6-S2-P14

This abstract is available from: <http://www.retrovirology.com/content/6/S2/P14>

© 2009 Yu et al; licensee BioMed Central Ltd.

### Background

Retroviral nucleocapsid (NC) is multifunctional in that it acts throughout the virus replication cycle via a number of molecular interactions. During the early stage, mature NC molecules extensively interact with the viral genome and reverse transcriptase to chaperone proviral DNA synthesis. At the late stage, NC as part of Gag, selects and dimerizes the genomic RNA, which is thought to start the Gag assembly process in infected cells. Interestingly, the RT reaction appears to be tightly controlled during the late steps of HIV-1 replication since the viral DNA synthesis is completed only after virions infect target cells [1]. How this is regulated is yet poorly understood and we hypothesized that the NC might be involved in the timing of RT. To function, NC needs its two conserved CCHC zinc fingers and the flanking basic residues. Therefore, we investigated their role in the temporal control of the RT.

### Results

We undertook a detailed quantitative analysis of the viral nucleic acid production throughout the replication cycle by qPCR and qRT-PCR. We measured the effects of NC zinc finger and basic residue deletions and mutations on the conversion of both the genomic and spliced RNA species into DNA. We discovered that viral particles released from the cells expressing HIV-1 NC mutants, contained a high level of DNA (up to 100-fold as compared with wild-type HIV-1) [2,3]. This unexpected accumulation of DNA

in NC mutant virions was also independently reported by Thomas et al [4]. Furthermore, we reported for the first time that intravirion DNA presence did not result from natural endogenous reverse transcriptase activity (NERT), but rather from the activation of the RT in the virus producer cells [2,3].

### Conclusion

These results provide the first example of RT during the late steps of HIV-1 replication and could bring an alternative explanation for the presence of viral DNA in HIV-1 particles isolated from the peripheral blood and semen of HIV-1-infected patients [5]. The occurrence of late RT inside producer cells is also a property of the foamy viruses that release viral DNA-containing particles and whose Gag domain naturally lacks NC zinc-fingers. It is of great interest to determine whether such timing RT control is a characteristic of the HIV-1 NC or rather a common characteristic among all retroviruses. These investigations are in progress.

Last, these new findings on the role of HIV-1 NC emphasize the fact that the conserved zinc-finger motifs should be viewed as a major target for new drugs against HIV-1.

## Acknowledgements

This work was supported by grants from ANRS, SIDACTION and CNRS. YB was supported by RTRS, LH by SIDACTION, and LD by ANRS. We thank Emilie Dumas for support of our current investigation.

## References

1. Mougel M, Houzet L, Darlix JL: **When is it time for reverse transcription to start and go?** *Retrovirology* 2009, **6**.
2. Houzet L, Morichaud Z, Didierlaurent L, Muriaux D, Darlix JL, Mougel M: **Nucleocapsid mutations turn HIV-1 into a DNA-containing virus.** *Nucleic Acids Res* 2008, **36**:2311-9.
3. Didierlaurent L, Houzet L, Morichaud Z, Darlix JL, Mougel M: **The conserved N-terminal basic residues and zinc-finger motifs of HIV-1 nucleocapsid restrict the viral cDNA synthesis during virus formation and maturation.** *Nucleic Acids Res* 2008, **36**:4745-53.
4. Thomas JA, Bosche WJ, Shatzer TL, Johnson DG, Gorelick RJ: **Mutations in human immunodeficiency virus type 1 nucleocapsid protein zinc fingers cause premature reverse transcription.** *J Virol* 2008, **82**:9318-28.
5. Zhang H, Dornadula G, Pomerantz RJ: **Endogenous reverse transcription of human immunodeficiency virus type 1 in physiological microenvironments: an important stage for viral infection of nondividing cells.** *J Virol* 1996, **70**:2809-2824.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

