


# Sharpening focus through wider collaboration: evolving heterogeneity in the bi-directional relationship between cardiovascular disease and COVID-19

**Amit Kaura** <sup>1,2</sup>, **Jamil Mayet** <sup>1,2</sup>, and **Charlotte Manisty** <sup>3,4\*</sup>

<sup>1</sup>National Institute for Health Research Imperial Biomedical Research Centre, Imperial College London and Imperial College Healthcare NHS Trust, London, UK; <sup>2</sup>National Heart and Lung Institute, Imperial College London, London, UK; <sup>3</sup>Institute of Cardiovascular Science, University College London, Gower Street, London, UK; and <sup>4</sup>Barts Heart Centre, St Bartholomew's Hospital, West Smithfield, London, UK

Online publish-ahead-of-print 4 November 2021

This editorial refers to 'Clinical presentation, disease course, and outcome of COVID-19 in hospitalized patients with and without pre-existing cardiac disease: a cohort study across 18 countries', by The CAPACITY-COVID Collaborative Consortium and LEOSS Study Group, <https://doi.org/10.1093/eurheartj/ehab656>.



**Graphical Abstract** The bidirectional relationship between cardiovascular disease and COVID-19 infection. ECMO, extracorporeal membrane oxygenation.

The recent outbreak of a novel coronavirus illness (coronavirus disease 2019; COVID-19) has emerged as a public health crisis of global proportions. The cardiology community has been captivated by COVID-19, perhaps more than any other systemic disease in history, due to its apparent links with cardiovascular disease (CVD). Researchers rapidly pivoted attention to concentrate on the disease, with initial case series and single-centre studies suggesting poor outcomes in patients with prior cardiac conditions and, soon after, reports of a high prevalence of cardiovascular complications. However, now, more than a year since the pandemic's onset, larger studies have emerged that are beginning to refine our understanding of the complex interplay between COVID-19 and CVD.

Previous COVID-19 studies have predominantly evaluated the association between pre-existing chronic cardiac disease and COVID-19-related mortality, where all cardiac conditions are combined and analysed together.<sup>1–3</sup> In this issue of the *European Heart Journal*, the CAPACITY-COVID and LEOSS (Lean European Open Survey on SARS-CoV-2 Infected Patients) study groups report on their collaborative retrospective study using coordinated large-scale data collection to evaluate heterogeneity in associations between various heart disease subtypes and in-hospital mortality.<sup>4</sup> The authors describe and compare the disease course and outcomes from >16 000 hospitalized COVID-19 patients across 18 countries admitted between March and May 2021, including more than one-third of patients with pre-existing cardiac disease. Although overall crude mortality was almost twice as high in those with prior CVD when compared with those without, after multivariable adjustment this difference was no longer significant. Furthermore, although for patients with heart failure the relative risk (RR) of death in hospital was significantly higher and related to severity (NYHA III/IV, RR 1.41, 1.20–1.64), for the majority of cardiovascular disorders, including ischaemic heart disease and atrial fibrillation, outcomes were similar to those of patients without CVD. This heterogeneity in outcomes between different cardiovascular conditions with COVID-19 provides support for a more stratified public health approach during the pandemic. Examples might include prioritizing those with heart failure at highest risk for early vaccine administration or reducing shielding advice for those considered at lower risk.

The contrast in these findings when compared with those from prior smaller studies highlights the importance firstly of robust adjustment for co-variants when considering the association of CVD with outcomes. Secondly, and perhaps more importantly, they demonstrate that 'prior CVD' should no longer be considered a single uniform risk factor but should be disaggregated into heart disease subtypes to better stratify risk and inform decision-making. This is not only relevant for COVID-19 but should be considered as an approach across other clinical scenarios when analysing the impact of cardiac conditions on outcomes. Unfortunately, although intuitive, in practice this presents challenges, with study design requiring significantly larger sample sizes and multicentre support for delivery.

Here perhaps COVID-19 has provided a unique opportunity for progress. The urgent public health crisis has led to international focus on prioritizing large-scale collaborative studies to answer pertinent clinical questions in a timely manner. Both LEOSS and CAPACITY-COVID are examples of such efforts, having received support from national and international professional organizations including endorsement by the European Society of Cardiology (ESC) for

international enrolment in CAPACITY-COVID. Similarly, combining results from independent (potentially competing) clinical studies to accelerate delivery of results was rare prior to COVID-19; however, the expert committees of LEOSS and CAPACITY-COVID should be congratulated for adopting this approach in the current study.

The study also assessed the prevalence of cardiovascular complications of severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) infection, given the presumed bi-directional relationship between CVD and COVID-19. SARS-CoV-2 binds to the angiotensin-converting enzyme 2 (ACE2) receptor expressed in cardiac myocytes, has been found in autopsy specimens, and hence could potentially cause myocarditis.<sup>5</sup> Elevated troponin levels are both common and prognostic in patients hospitalized with COVID-19, and asymptomatic abnormalities have been reported on cardiovascular imaging studies following severe disease.<sup>6</sup> However, despite targeted data collection for cardiovascular complications, the current study found clinical cardiac adverse effects to be uncommon, with an overall prevalence of <2%. In contrast, venous thrombo-embolism was common despite the enhanced antithrombotic prophylactic strategies instituted after the first wave of the pandemic, with pulmonary emboli found in >1 in 10 patients admitted to intensive care and with overall prevalence 3–4 times higher when compared with seasonal influenza.<sup>7</sup> The multicentre ISARIC study, which prospectively collected in-hospital COVID-19 complications from >80 000 patients in the UK starting early during the pandemic, has recently reported cardiac complication rates almost twice higher than those found in the current study,<sup>3</sup> despite lower rates of venous thrombo-embolism. One common finding, however, was that thrombo-embolic complications are unexpectedly less commonly reported among patients with known cardiac disease, possibly related to pre-admission anticoagulant or antiplatelet prescription.

The reasons behind the differences between the current study and ISARIC remain unclear, particularly given the similarities in clinical and demographic characteristics of the study cohorts. The one major difference is in the timing of data collection—ISARIC reported findings from patients hospitalized between January and August 2020, whereas the current study included patients hospitalized between March and May 2021. In the interval between these periods, the findings of randomized clinical trials and the inevitable growth in clinical experience resulted in changes to the management of patients with severe COVID-19 (including dexamethasone administration,<sup>8</sup> judicious fluid management, and empirical anticoagulation for those at highest risk<sup>9</sup>), any of which may have impacted cardiac complication rates. This highlights the changing pattern of patient demographics, clinical manifestations, and outcomes as the pandemic progresses—with the additional impact of vaccination likely to result in widening of the geographical and socio-economic differences in COVID-19. The bi-directional relationship between COVID-19 and CVD is therefore likely to evolve, with increasing heterogeneity as time progresses (*Graphical Abstract*).

There are several considerations left to be answered in future studies. Despite collaboration between LEOSS and CAPACITY-COVID, the combined dataset lacked sufficient power to draw conclusions on the interaction of age in the association between COVID-19 and CVD. Second, whilst ethnicity is known to be relevant to SARS-CoV-2 infection, the demographic of this current cohort is narrow (84.5% white), with no adjustment for ethnicity in

multivariable models. It is also important to recognize that the scope of this study is relatively narrow, with focus solely on in-hospital survival status and cardiovascular complications following severe COVID-19. Understanding the effect of both severe and (the more prevalent) community COVID-19 on late cardiovascular complications and overall functional recovery requires further investigation, which will demand careful data collection across both hospital and primary care data. Recently resources linking individual person-level data across healthcare settings have been developed to address this issue,<sup>10</sup> and the results which will emerge over the coming months are likely to improve our understanding of this complex inter-relationship. Currently, however, the authors are to be congratulated for coordinating data from such a large number of centres over 18 countries during a pandemic. Their study serves to emphasize the strength of scientific collaboration across Europe and beyond, and has highlighted that a 'one-size-fits-all' approach to patients with CVD in the pandemic may no longer be appropriate.

## Funding

C.M. is supported directly and indirectly from the NIHR Biomedical Research Centres at University College London Hospitals and Barts Health NHS Trusts. A.K. is funded by a British Heart Foundation clinical research training fellowship (FS/20/18/34972). J.M. is supported by the BHF Imperial Centre for Research Excellence (RE/18/4/34215).

**Conflict of interest:** none declared.

## References

- Williamson EJ, Walker AJ, Bhaskaran K, Bacon S, Bates C, Morton CE, Curtis HJ, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature* 2020;**584**:430–436.
- Docherty AB, Harrison EM, Green CA, Hardwick HE, Pius R, Norman L, Holden KA, et al. Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: prospective observational cohort study. *BMJ* 2020;**369**:m1985.
- Drake TM, Riad AM, Fairfield CJ, Egan C, Knight SR, Pius R, Hardwick HE, Norman L, Shaw CA, McLean KA, Thompson AAR, Ho A, Swann OV, Sullivan M, Soares F, Holden KA, Merson L, Plotkin D, Sigfrid L, de Silva TI, Girvan M, Jackson C, Russell CD, Dunning J, Solomon T, Carson G, Oliaro P, Nguyen-Van-Tam JS, Turtle L, Docherty AB, Openshaw PJ, Baillie JK, Harrison EM, Semple MG; ISARIC4C investigators. Characterisation of in-hospital complications associated with COVID-19 using the ISARIC WHO Clinical Characterisation Protocol UK: a prospective, multicentre cohort study. *Lancet* 2021;**398**:223–237.
- Gitt AK, Bernhardt A, Zahn R, Zeymer U, Grau A, Beutel ME, Werdan K. The COVID-19 Registry in Rhineland-Palatinate in the context of international registry activities documenting COVID-19 outcomes. *Herz* 2020;**45**:316–318.
- CAPACITY-COVID collaborative consortium and LEOSS Study Group. Clinical presentation, disease course and 1 outcome of COVID-19 in hospitalized patients with and without pre-existing cardiac disease: a cohort study across 18 countries. *Eur Heart J* 2022;**43**:1104–1120.
- Wichmann D, Sperhake JP, Lutgehetmann M, Steurer S, Edler C, Heinemann A, Wichmann D, Sperhake JP, Lutgehetmann M, Steurer S, Edler C, Heinemann A. Autopsy findings and venous thromboembolism in patients with COVID-19: a prospective cohort study. *Ann Intern Med* 2020;**173**:268277.
- Kotecha T, Knight DS, Razvi Y, Kumar K, Vimalasvaran K, Thornton G, Patel R, Chacko L, Brown JT, Coyle C, Leith D, Shetye A, Ariff B, Bell R, Captur G, Coleman M, Goldring J, Gopalan D, Heightman M, Hillman T, Howard L, Jacobs M, Jeetley PS, Kanagaratnam P, Kon OM, Lamb LE, Manisty CH, Mathurdas P, Mayet J, Negus R, Patel N, Pierce I, Russell G, Wolff A, Xue H, Kellman P, Moon JC, Treibel TA, Cole GD, Fontana M. Patterns of myocardial injury in recovered troponin-positive COVID-19 patients assessed by cardiovascular magnetic resonance. *Eur Heart J* 2021;**42**:1866–1878.
- Piroth L, Cottenet J, Mariet AS, Bonniaud P, Blot M, Tubert-Bitter P, Quantin C. Comparison of the characteristics, morbidity, and mortality of COVID-19 and seasonal influenza: a nationwide, population based retrospective cohort study. *Lancet Respir Med* 2020;**9**:251–259.
- RECOVERY Collaborative Group, Horby P, Lim WS, Emberson JR, Mafham M, Bell JL, Linsell L, Dexamethasone in hospitalized patients with Covid-19. *N Engl J Med* 2021;**384**:693–704.
- Poissy J, Goutay J, Caplan M, Parmentier E, Duburcq T, Lassalle F, Jeanpierre E, Rauch A, Labreuche J, Susen S; Lille ICU Haemostasis COVID-19 Group. Pulmonary embolism in patients with COVID-19: awareness of an increased prevalence. *Circulation* 2020;**142**:184–186.
- Wood A, Denholm R, Hollings S, Cooper J, Ip S, Walker V, Denaxas S, Akbari A, Banerjee A, Whiteley W, Lai A, Sterne J, Sudlow C; CVD-COVID-UK consortium. Linked electronic health records for research on a nationwide cohort of more than 54 million people in England: data resource. *BMJ* 2021;**373**:n826.