RESEARCH ARTICLE



REVISED A patient satisfaction survey and educational package

to improve the care of people hospitalised with COVID-19: a

quality improvement project, Liverpool, UK [version 2; peer

review: 2 approved]

Muhammad Shamsher Ahmad^{1,2}, Scott Rory Hicks¹, Rebecca Watson¹, Rajia Akter Ahmed¹, Lewis Jones¹, Marcella Vaselli¹, Meng-San Wu¹, Fatima Hayat¹, Libuse Ratcliffe¹, Mark McKenna¹, Paul Hine^{1,3}, Sylviane Defres¹⁻³, Tom Wingfield¹⁻⁴

¹Tropical and Infectious Disease Unit, Liverpool University Hospital NHS Foundation Trust, Liverpool, Merseyside, L7 8XP, UK ²Clinical Infection Microbiology and Immunology, Institute of Infection and Global Health, Liverpool, UK ³Clinical Sciences and International Public Health, Liverpool School of Tropical Medicine, Liverpool, Merseyside, L3 5QA, UK ⁴WHO Collaborating Centre for Social Medicine and Tuberculosis, Department of Global Public Health Sciences, Karolinska Institute, Stockholm, Sweden

V2 First published: 03 Sep 2021, 6:222 https://doi.org/10.12688/wellcomeopenres.171		
	Latest published: 25 Jan 2022, 6:222 https://doi.org/10.12688/wellcomeopenres.17163.2	

Abstract

Background: The perspectives and experiences of people hospitalised with COVID-19 have been under-reported during the coronavirus pandemic. We developed and conducted a COVID-19 patient satisfaction survey in a large university-affiliated secondary healthcare centre in Liverpool, UK, during Europe's first coronavirus wave (April-June 2020). The survey found that care was rated highly, including among people of Black Asian and Minority Ethnic (BAME) backgrounds. However, sleep-quality and communication about medications and discharge-planning were identified as areas for improvement.

Methods: To improve care for people with COVID-19 admitted to our centre, we designed an educational package for healthcare professionals working on COVID-19 wards. The package, implemented in August 2020, included healthcare worker training sessions on providing holistic care and placement of "Practice Pointers" posters. Patient satisfaction was re-evaluated during the second/third COVID-19 waves in Liverpool (September 2020 - February 2021).

Results: Across waves, most (95%) respondents reported that they would recommend our hospital to friends and/or family and rated overall care highly. Comparison of the responses of second/third-



1. Katherine Ajdukiewicz 🛄, North

Manchester General Hospital, Manchester, UK

2. Vishesh Paul (D), Carle Foundation Hospital, Urbana, USA

Any reports and responses or comments on the article can be found at the end of the article.

wave respondents (n=101) with first-wave respondents (n=94) suggested improved patient satisfaction across most care domains but especially those related to having worries and fears addressed and being consulted about medications and their side-effects.

Conclusions: People admitted with COVID-19 to our centre in Liverpool, including those from BAME backgrounds, rated the care they received highly. A simple education package improved the feedback on care received by respondents between the first and second/third waves. These UK-first findings are informing regional strategies to improve person-centred care of hospitalised people with COVID-19.

Keywords

Coronavirus; Covid-19; patient satisfaction; patient perspective; patient feedback; patient experience; quality improvement project

Corresponding author: Tom Wingfield (tom.wingfield@lstmed.ac.uk)

Author roles: Ahmad MS: Data Curation, Formal Analysis, Investigation, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; Hicks SR: Data Curation, Investigation, Project Administration, Validation, Writing – Review & Editing; Watson R: Data Curation, Investigation, Project Administration, Writing – Review & Editing; Ahmed RA: Investigation, Project Administration, Writing – Review & Editing; Jones L: Investigation, Writing – Review & Editing; Vaselli M: Investigation, Project Administration, Writing – Review & Editing; Wu MS: Conceptualization, Data Curation, Investigation, Methodology, Project Administration, Writing – Review & Editing; Hayat F: Investigation, Project Administration, Writing – Review & Editing; Ratcliffe L: Investigation, Methodology, Writing – Review & Editing; McKenna M: Methodology, Validation, Writing – Review & Editing; Hine P: Investigation, Methodology, Writing – Review & Editing; Defres S: Investigation, Methodology, Writing – Review & Editing; Wingfield T: Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This work was supported by Wellcome [209075, https://doi.org/10.35802/209075]. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Copyright: © 2022 Ahmad MS *et al*. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Ahmad MS, Hicks SR, Watson R *et al.* A patient satisfaction survey and educational package to improve the care of people hospitalised with COVID-19: a quality improvement project, Liverpool, UK [version 2; peer review: 2 approved] Wellcome Open Research 2022, 6:222 https://doi.org/10.12688/wellcomeopenres.17163.2

First published: 03 Sep 2021, 6:222 https://doi.org/10.12688/wellcomeopenres.17163.1

REVISED Amendments from Version 1

We have updated the article as per reviewer's minor comments. This includes:

1. Changing "BAME background" to "BAME backgrounds"

2. Including explanation of the abbreviation LUHFT (Liverpool University Hospitals NHS Foundation Trust)

Any further responses from the reviewers can be found at the end of the article

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has infected over 171 million people worldwide and caused over 3.5 million deaths¹. In the UK, there have been in excess of 4.5 million confirmed cases of whom over 10% have required admission to hospital and more than 152,000 have died².

Despite extensive research into COVID-19 vaccines, diagnostics, and biomedical treatments, evidence concerning the perspectives of people with COVID-19, especially from vulnerable and Black Asian Minority Ethnic (BAME) groups, is negligible³. To address this, we designed and implemented a satisfaction survey of people with COVID-19 admitted during the first wave of Covid-19 infections (March-June 2020) to our large university hospital in Liverpool, UK. The survey findings (published in Future Healthcare Journal⁴) showed that nursing and medical care was rated highly and most respondents (96%) reported that they would recommend our hospital to friends or family. However, the survey also highlighted potential areas for improvement including communication about medications and their side effects and informing patients about discharge plans⁴. To address these shortcomings and improve holistic COVID-19 hospital care, we subsequently implemented a package of complementary interventions on COVID-19 wards in our centre

Here we report the satisfaction survey findings from the second/third COVID-19 waves in Liverpool (September 2020 - February 2021) and compare them to those of the first wave.

Methods

This was an unpowered before-and-after observational quality improvement project (QIP) registered with the local Clinical Effectiveness Department. The COVID-19 patient satisfaction survey was developed in collaboration with our centre's 'Patient Experience' team to ensure it was patientfriendly and suitable for people with learning and reading difficulties. According to the policy activities that constitute research at the Liverpool University Hospitals NHS Trust (LUHFT), this Quality Improvement Project met criteria for operational improvement activities exempt from ethics review. All respondents were consented verbally but, in line with local QIP policy, written consent was not deemed to be required⁴. The survey was adapted from existing patient satisfaction surveys and integrated with our centre's "Friends and Family Test" questions. An open-access version of the survey itself and accompanying standard operating procedure can be found in 5.

The survey was implemented during the first COVID-19 wave (15th March 2020 to 15th June 2020) and second/third waves (21st September 2020 to 6th February 2021). Between these time periods, the first-wave survey findings were presented at Tropical Infectious Disease Unit QIP meetings and centre-wide Patient Experience meetings. A package of complementary interventions was designed and implemented in August-September 2021. The package consisted of: concise feedback and training sessions for healthcare workers on holistic care on the COVID-19 wards; updated COVID-19 patient information leaflets for admission and discharge; and a 'COVID-19 Practice Pointers Poster' (Figure 1), which was also placed in visible, shared ward areas.

Descriptive analysis summarised overall responses, compared second/third vs first wave responses, and further compared responses by BAME, age, and gender.

An earlier version of this article can be found on medRxiv (doi: 10.1101/2021.03.23.21253630).

Results

Surveys from 195 respondents, 94 (48%) from the first wave and 101 (52%) from the second/third wave, were collated (Table 1). Compared to the first wave, median age of respondents in the second/third wave was higher (64 vs 59 years) and there were more respondents who were female (58% vs 48%), obese (48% vs 38%), active/ex-smokers (49% vs 33%), and had at least one chronic comorbidity (70% vs 67%). The proportion of BAME respondents was the same across both cohorts (10%).

Patient satisfaction was high with overall care rated 4.8/5 on average. Nearly all (95%) respondents reported that they would recommend our hospital to friends and/or family (Figure 2).

Compared to first wave respondents, second/third wave patient satisfaction increased across multiple domains of care but, most notably, being involved in care decisions (81% to 95%), able to share worries and fears (77% to 93%), and communication about medications and side effects (63% to 85%, Figure 2). Satisfaction decreased with relation to being kept informed of discharge plans (84% to 75%) and sleep environment (77% to 73%, Figure 2).

Reported patient satisfaction was higher amongst BAME than non-BAME respondents across all domains except discussion about medications and side-effects (70% vs 75%, Figure 3). Responses were similar by gender and age.

Comparison of monthly average "Friends and Family" test responses performed routinely by our centre among all

Liverpool University Hospitals

COVID-19 How can we improve the patient experience?

Between March and June 2020, 98 people with COVID-19 admitted to Liverpool University Hospitals NHS Foundation Trust completed a satisfaction survey. Respondents rated the care they had received highly (on average 4.7 out of 5 stars) and made suggestions of how their care could have been improved. We are grateful for their feedback, we created the 'Practice Pointers' below, which we hope will lead to improved patient care.

Communication



- Clearly communicate to new patients their diagnosis of COVID-19.
- Daily updates on ward rounds about test results including COVID-19 swabs.
- Explain new medications and possible side effects (e.g. remdesivir, steroids, enoxaparin) and ask about need for pain relief.
- The use of face masks can make communication difficult, especially for people who are hearing impaired.
- Where needed use masks with window adaptations, increase speaking volume, and use visual aids.
- **Communicate sensitively** about treatment escalation plans.

Food and sleep

- · Ensure meal times are protected.
- Offer snacks and hot drinks in between meals.
- Involve dietitians if you are concerned about poor nutrition.
- To improve sleep, avoid all unnecessary night time disruptions.
- Be aware of daytime noise levels on the ward, which some patients may find stressful.
- Explain to patients as early as possible the day-to-day routine of ward activities that they can expect.
- Consider morning dosing of enoxaparin to allow day-time Xa levels.

Family



- Try to facilitate phone or video calls wherever possible.
- Ask for a nominated family member or next of kin and update them regularly.
- Visitors may be permitted for those approaching end of life, and under certain circumstances for those experiencing distress due to dementia or a mental health issue - please see Trust intranet for the latest guidance.

Additional information

- Provide written information on admission and discharge - available from the Trust intranet or PHE website.
- Discuss additional treatments available through clinical trials to allow patients to consider all of their options.
- Advise patients of plans for follow up in those with chest X-ray changes, a chest X-ray is recommended approximately three months following discharge.

This poster is an aide memoire for holistic care of people with COVID-19 admitted to our hospital. It is currently only for use on the wards on which the original COVID-19 patient satisfaction survey took place. For more information see Wu et al, BMJ, 2020: https://www.bmj.com/content/370/bmj.m3496

Figure 1. COVID-19 practice pointer poster. Figure 1 showing the COVID-19 practice pointer poster introduced after the patient satisfaction survey during the first wave of pandemic to improve the holistic care services.

Variables	Respondents (%)			
	First wave (n=94)	Second/Third wave (n=101)	All (n=195)	
Male	49 (52)	42 (42)	91 (47)	
Age, median (IQR)	59 (46-72)	64 (54-73)	62 (50-73)	
BMI, median (IQR) (n=84/94, 86/101, 170/195)*	28 (25-32)	29 (24-34)	29 (25-33)	
Obesity (BMI ≥ 30 kg/m2, n=84/94, 86/101, 170/195)*	32 (38)	41 (48)	73 (43)	
Black, Asian and Minority Ethnicity	10 (11)	10 (10)	20 (10)	
Smoker / ex-smoker	31 (33)	49 (49)	80 (41)	
Health characteristics				
Chronic lung diseases**	36 (38)	12 (12)	48 (25)	
Hypertension	29 (31)	40 (40)	69 (35)	
Chronic cardiovascular diseases***	15 (16)	22 (22)	37 (19)	
Diabetes	17 (18)	32 (32)	49 (25)	
Chronic Kidney Disease	13 (14)	16 (16)	29 (15)	
Non-HIV immunosuppression	9 (10)	7 (7)	16 (8)	
HIV positive	2 (2)	1 (1)	3 (2)	
History of DVT / PE	1 (1)	11 (11)	12 (6)	
Number of Comorbidities				
0	31 (33)	31 (31)	62 (32)	
1	34 (36)	20 (20)	54 (28)	
2	14 (15)	34 (34)	48 (25)	
3 or more	15 (16)	16 (16)	31 (16)	

 Table 1. Sociodemographic and clinical characteristics of survey respondents (n=195).

Legend: This table shows the number of respondents (%) in the first (n=94), second/third wave (n=101), and overall (n=195). *BMI was only available or calculable for 170/195 respondents because height and/ or weight was not documented for 10 respondents in the first wave (n=84/94) and 15 respondents in the second/third wave (n=86/101). **Chronic lung diseases include asthma, COPD, bronchiectasis, interstitial lung diseases, lung cancer. ***Chronic cardiovascular diseases include ischaemic heart disease, heart failure, cerebrovascular disease, and peripheral vascular disease. Abbreviations: IQR interquartile range, BMI body mass index, HIV human immunodeficiency virus, DVT deep vein thrombosis, PE pulmonary embolus

admissions showed that people with COVID-19 were more likely to recommend our centre to friends and family than other patients apart from in October 2020 and January 2021 (Figure 4). When asked to rate the overall services received out of five, 80% of Covid-19 survey respondents rated it 5/5 and 20% rated 4/5 (Figure 5).

Free text comments on care received were mostly positive with negative comments focusing on noise level, sleep disruption, and response time when asking for help or pressing patient buzzer (Table 2).

Discussion

In the UK to date, nearly half a million people have been hospitalised with COVID-19². In addition to physical symptoms of COVID-19, those hospitalised can experience negative psychosocial consequences: isolation, including related to infection prevention and control policies; lack of contact with family and friends unable to visit hospital; and uncertainty related to their prognosis⁶. This may be compounded by constrained communication, trust and rapport with healthcare professionals, particularly for people with hearing impairment due to personal protective equipment including masks^{3,5,7}.



Figure 2. Responses from patients admitted with COVID-19. Figure 2 showing patient satisfaction survey responses (n=195) during the first (n=94) and second-third (n=101) waves of COVID-19 in Liverpool with an overall improvement of satisfaction in most domains after introduction of educational package.



Figure 3. Comparison between BAME and non-BAME Respondents. Figure 3 comparing the patient satisfaction survey responses between the BAME (n=20 and non-BAME (n=175) respondents showing a comparable or even better satisfaction among the BAME patients admitted with COVID-19.



Figure 4. Patient satisfaction between COVID-19 And overall admission. Figure 4 is the month-wise comparison between the overall hospital admissions and the survey participants with COVID-19 admissions during first and second-third wave of COVID-19 pandemic in Liverpool; showing overall better satisfaction in COVID-19 admissions with higher percentage of patients recommending our services in most of the months apart from the two months in the height of second and third waves.



Figure 5. Overall rating of the Hospital Services. Figure 5 showing the rating given to our services by the patients admitted with COVID-19 with nearly all of them rating it above four out of five.

Despite these challenges, our COVID-19 patient satisfaction survey showed that the quality of care at our centre was rated highly throughout the COVID-19 pandemic even during the second/third wave when our local health system was under significant operating pressures. While this is the first peerreviewed COVID-19 patient satisfaction survey of its kind in the UK, a national satisfaction survey of the 'UK Patient Association' has been published against which we can compare our findings⁸. Our survey showed a higher percentage of satisfaction for quality of care and being involved in care decisions. Notably, the 75% of respondents reporting satisfaction regarding discharge planning was consistent with the national survey. Unlike the national survey, the overall satisfaction in patients admitted with COVID-19 remained higher than the general admissions, apart from the dip during the height of the second (October 2020) and third waves (January 2021) in our centre. Keeping in touch with friends and family was identified as a key determinant of inpatient satisfaction in the 'UK Patient Association' survey⁹. Indeed, this was a priority in our Covid-19 Practice Pointers poster (Figure 1) and the overall

Table 2. Patients free text comments during the second-third waves (n=101).				
	Excellent standard of care by all staff.			
	Always someone within seconds when I rang for help.			
	Outstanding staff in all roles.			
	Regular updates to family			
	Allowing family to bring food is very helpful, mentally and physically			
	Couldn't ask for better			
Things that went well	Conversation with consultants			
	Most of the staff were great - some exceptional.			
	Food - delicious. Cleaners very thorough at all times.			
	All staff including the cleaners and porters are credit to the NHS			
	Staff are attentive, caring, explained everything nicely			
	Never been in hospital care. Amazing - You are all so good			
	I couldn't ask for better staff, love them all, made me relaxed away from home			
	Don't leave people alone when they are so helpless			
Things that could be	Ward was very noisy - doors and bins banging, phone ringing unanswered.			
Things that could be improved	Had trouble sleeping for outside noise one night			
	Give patients something to pass time (books, TV etc.)			
	Not being able to cope with the pain of the illness			

Table 2. Patients' free text comments during the second-third waves (n=101).

improvement of patient satisfaction during the second/third wave in our centre may reflect the positive impact of this practice. Outside of the UK, our findings are broadly consistent with a COVID-19 patient survey carried out in Saudi Arabia¹⁰.

Our finding of higher satisfaction amongst BAME respondents across nearly all care domains was encouraging. Compared to non-BAME, people of BAME backgrounds have higher rates of severe COVID-19 disease and death¹¹ and restricted healthcare access¹² in the UK. This serves as a reminder of the importance of addressing widening health inequalities related to socioeconomic status and ethnicity in the UK¹¹.

Despite our package of interventions improving some aspects of care⁵, sleep quality was rated low across both waves. These findings align with pre-COVID surveys and are a persistent issue in hospital care¹³. There is ongoing work within our centre to address this including noise monitoring and designated "noise free" hours.

Our survey suggested that more respondents with COVID-19 would recommend our centre's services to friends and family than people admitted for other reasons except in October 2020 and January 2021 – the peak of second and third waves of admissions when our centre was it under the most pressure. However, the overall trend of general patient satisfaction in our centre remained stable during the COVID pandemic as reported in other UK patient experience surveys⁸.

Almost all of the patients in our survey rated the care they had received during the admission highly. Indeed, COVID-19 respondents in our centre appeared to rate the care they received more highly than those in the UK Patient Association national survey⁹.

Although not necessarily reflecting the actual quality or standard of clinical services, patient satisfaction is a popular tool to measure quality of health care services^{14,15}. A patient survey in the emergency department in New York revealed generally higher level of satisfaction among their attendees during the COVID pandemic compared to the preceding years¹⁶. This was attributed in free text responses to gratitude towards the healthcare staff for efforts and sacrifices during the pandemic. The appreciation for the NHS could also be a contributing factor for the positive responses received in our survey.

Limitations of this study

This was a single-centre, opportunistic, non-randomised survey of a small sample of clinically stable patients, which limits generalisability. Despite this, this peer-reviewed study was a UK first and its findings important when considering person-centred COVID-19 care strategies.

Conclusions

In a cohort of people hospitalised with COVID-19 in Liverpool, hospital care was rated highly including by those of BAME background. Implementation of an education and training package between COVID-19 waves was associated with improved patient feedback, particularly regarding involvement in and communication about care. The survey and package are being expanded locally to further improve care of people with COVID-19 and other conditions.

Data availability

Open Science Framework. A patient satisfaction survey and educational package to improve the care of people hospitalised with COVID-19: a quality improvement project, Liverpool, UK.

DOI: https://doi.org/10.17605/OSF.IO/7MBJQ¹⁶

This project contains the following data:

 Deidentified COVID patient satisfaction survey data 23082021 OSF.xlsx

Data are available under the terms of the Creative Commons Zero "No rights reserved" data waiver (CC BY 4.0 Public domain dedication).

References

- European Centre for Disease Prevention and Control: COVID-19 situation update worldwide. accessed 09/06/2021. Reference Source
- GOV.UK, Coronavirus (COVID-19) in the UK: Patients admitted in hospital. Healthcare in the United Kingdom, 2021; accessed 09/06/2021. Reference Source
- Wu MS, Hayat F, Ratcliffe L, et al.: What do people hospitalised with covid-19 think about the care they received? BMJ. 2020; 370: m3496.
 PubMed Abstract | Publisher Full Text
- British Medical Journal Quality & Safety: Journal Policy on ethics review for quality improvement reports. accessed 25/08/2021. Reference Source
- Wu MS, Watson R, Hayat F, et al.: What do people hospitalised with COVID-19 think about their care? Results of a satisfaction survey during the first wave of COVID-19 in Liverpool. Future Healthc J. 2021; 8(1): e70-e75. PubMed Abstract | Publisher Full Text | Free Full Text
- Millstein JH, Kindt S: Reimagining the Patient Experience During the Covid-19 Pandemic. Nejm Catalyst Innovations in Care Delivery. 2020. Reference Source
- The Patient Associations: Pandemic Patient Experience UK patient experience of health, care and other support during the COVID-19 pandemic. accessed 09/06/2021. Reference Source
- NHS Patient Survey Programme: Inpatient experience during the COVID-19 pandemic: National report. accessed 09/06/2021. Reference Source
- 9. Bin Traiki TA, AlShammari SA, AlAli MN, et al.: Impact of COVID-19 pandemic on patient satisfaction and surgical outcomes: A retrospective and cross

sectional study. Ann Med Surg (Lond). 2020; 58: 14–19. PubMed Abstract | Publisher Full Text | Free Full Text

- Apea VJ, Wan YI, Dhairyawan R, et al.: Ethnicity and outcomes in patients hospitalised with COVID-19 infection in East London: an observational cohort study. BMJ Open. 2021; 11(1): e042140.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Razai MS, Kankam H, Majeed A, et al.: Mitigating ethnic disparities in covid-19 and beyond. *BMJ*. 2021; 372: m4921.
 PubMed Abstract | Publisher Full Text
- Wesselius HM, van den Ende ES, Alsma J, et al.: Quality and Quantity of Sleep and Factors Associated With Sleep Disturbance in Hospitalized Patients. JAMA Intern Med. 2018; 178(9): 1201–1208.
 PubMed Abstract | Publisher Full Text | Free Full Text
- Abidova A, da Silva PA, Moreira S: Predictors of Patient Satisfaction and the Perceived Quality of Healthcare in an Emergency Department in Portugal. West J Emerg Med. 2020; 21(2): 391–403. PubMed Abstract | Publisher Full Text | Free Full Text
- Farley H, Enguidanos ER, Coletti CM, et al.: Patient Satisfaction Surveys and Quality of Care: An Information Paper. Ann Emerg Med. 2014; 64(4): 351–357. PubMed Abstract | Publisher Full Text
- Grissom MO, Farra M, Cruzen ES, et al.: What can COVID-19 teach us about patient satisfaction in the emergency department? A mixed-methods approach. J Am Coll Emerg Physicians Open. 2021; 2(2): e12436. PubMed Abstract | Publisher Full Text | Free Full Text
- Wingfield T: A patient satisfaction survey and educational package to improve the care of people hospitalised with COVID-19: a quality improvement project, Liverpool, UK. 2021. http://www.doi.org/10.17605/OSF.IO/7MBJQ

Open Peer Review

Current Peer Review Status:

Version 2

Reviewer Report 07 February 2022

https://doi.org/10.21956/wellcomeopenres.19534.r48262

© **2022 Paul V.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Vishesh Paul 匝

Pulmonary and Critical Care Medicine, Carle Foundation Hospital, Urbana, IL, USA

Looks good. Strongly consider for indexing.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Proning in COVID procedural complications in the ICU

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 24 January 2022

https://doi.org/10.21956/wellcomeopenres.18957.r48161

© **2022 Paul V.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Vishesh Paul ២

Pulmonary and Critical Care Medicine, Carle Foundation Hospital, Urbana, IL, USA

I enjoyed reading this manuscript and highly recommend it for indexing. As the article discusses, the majority of the medical community's focus has been on the biomedical treatments of COVID-19. In this, the psychosocial impact of this illness on hospitalised patients may have taken a backseat. Identifying areas of improvement through patient satisfaction surveys such as this one should help us provide more comprehensive and fulfilling care. The points that the team identified, such as discussion about medications and side effects, discharge planning, are being observed globally. Improvement projects such as this one can help guide many centers and providers, especially regarding the care of minorities who often get suboptimal care.

I have a couple of suggestions:

- 1. In the 'conclusion' section of the abstract, it should be BAME backgrounds (not background).
- 2. Figure 4: Please explain what 'LUFTH' stands for. Readers from outside institutions might not know it.

Overall, I feel the article is very well written, is succinct, and should be of interest to health care providers and centers worldwide.

Is the work clearly and accurately presented and does it cite the current literature? $\ensuremath{\mathsf{Yes}}$

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others? $\ensuremath{\mathsf{Yes}}$

If applicable, is the statistical analysis and its interpretation appropriate? Yes

Are all the source data underlying the results available to ensure full reproducibility? $\ensuremath{\mathsf{Yes}}$

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Proning in COVID procedural complications in the ICU

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 24 Jan 2022

Tom Wingfield, Liverpool School of Tropical Medicine, UK

Dear Dr Paul,

Thank you for your constructive and supportive comments.

We have made the updates suggested.

Yours sincerely,

Tom Wingfield (on behalf of the co-authors)

Competing Interests: None

Reviewer Report 11 November 2021

https://doi.org/10.21956/wellcomeopenres.18957.r46861

© **2021 Ajdukiewicz K.** This is an open access peer review report distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Katherine Ajdukiewicz 匝

Department of Infectious Diseases & Tropical Medicine, North Manchester General Hospital, Manchester, UK

Well written and well executed study. Most limitations identified - excludes those who were extremely ill and either died or went to critical care; pre-dates vaccination which may alter future similar studies (with use of ronapreve for example) in spike protein antibody negative patients. Relevant study with relevant findings and easily reproducible in other settings with realistic interventions. It would be interesting to see if these outcomes are maintained over time and in non-COVID patients.

Is the work clearly and accurately presented and does it cite the current literature? $\ensuremath{\mathsf{Yes}}$

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others? $\ensuremath{\mathsf{Yes}}$

If applicable, is the statistical analysis and its interpretation appropriate? $\ensuremath{\mathsf{Yes}}$

Are all the source data underlying the results available to ensure full reproducibility? $\ensuremath{\mathsf{Yes}}$

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Infectious diseases clinician treating patients with COVID-19 since 2020 (in addition to patients with other infections). Most research is on other areas - bacterial meningitis RCT with adjuvant therapy, CSF and plasma cytokine studies, perinatal HIV cohort studies

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.