

First survey of Interstitial molluscs from Cayo Nuevo, Campeche Bank, Gulf of Mexico

Deneb Ortigosa¹, Nancy Yolimar Suárez-Mozo¹, Noe C. Barrera², Nuno Simões^{1,3,4}

1 Unidad Multidisciplinaria de Docencia e Investigación Sisal (UMDI-SISAL), Facultad de Ciencias, Universidad Nacional Autónoma de México, Puerto de abrigo s/n, Sisal, CP 97356 Yucatán, Mexico **2** Harte Research Institute, Texas A&M University-Corpus Christi, 300 Ocean Dr., Unit 5869, Corpus Christi, Texas 78412-5869, U.S.A. **3** Laboratorio Nacional de Resiliencia Costera Laboratorios Nacionales, CONACYT, Mexico City, Mexico **4** International Chair for Coastal and Marine Studies, Harte Research Institute for Gulf of Mexico Studies, Texas A&M University – Corpus Christi, Corpus Christi, Texas, USA

Corresponding author: Nuno Simões (ns@ciencias.unam.mx)

Academic editor: N. Yonow | Received 20 February 2018 | Accepted 9 July 2018 | Published 2 August 2018

<http://zoobank.org/CE8D6959-7966-480C-ADF6-D514071AAF29>

Citation: Ortigosa D, Suárez-Mozo NY, Barrera NC, Simões N (2018) First survey of Interstitial molluscs from Cayo Nuevo, Campeche Bank, Gulf of Mexico. ZooKeys 779: 1–17. <https://doi.org/10.3897/zookeys.778.24562>

Abstract

Six sediment samples weighing between 224–735 g were collected in June of 2016 from Cayo Nuevo reef, located at the Campeche Bank, southern Gulf of Mexico. Samples were collected by SCUBA diving, from were two stations at depths of 7.6 and 18.2 m. Sediment was sieved and molluscs (adults and micromolluscs ≤ 10 mm) were sorted, examined, and identified to the lowest taxonomic level. A total of 1,347 specimens was found, of which 224 shells were dead and 1,123 were alive. Thirty-four families, 53 genera, and 67 species were identified. The most abundant families were Chamidae and Arcidae for the Bivalvia class, and Caecidae and Tornidae for the Gastropoda class. The vertical range of *Bentharca* sp. was extended.

Keywords

baseline, inventory, Campeche Bank, Gulf of Mexico, micromolluscs

Introduction

Frequently, species molluscan biodiversity accounts are incomplete because of the lack of some groups such as sea slugs and micromolluscs. Compared with macromolluscs, the study of micromolluscs is still in its infancy, which is probably due not to the difficulty involved in obtaining samples, but difficulties in identification of such small animals and the time-consuming process required to separate specimens from sand or other substrates (e.g., algae or rocks), and photography. In order to get a more realistic picture of the biodiversity for different habitats, micromolluscs should be incorporated into the different studies (Sasaki 2008, and pers. obs.).

The term micromollusc has been applied in arbitrary and non-standardised ways. Micromolluscs are molluscs not visible without some type of artificial assistance, such as a microscope or magnifying glass. The most restrictive definition, or *sensu stricto*, stated that micromollusc size should be less than 5 mm as an adult (Narciso 2005, Geiger et al. 2007). Other authors considered micromolluscs as specimens smaller than 10 mm as an adult (Barrera and Tunnell 2001). Finally, the wider definition of micromollusc, or *sensu lato*, includes molluscs whose size is typically less than 10 mm as an adult and also included juvenile representative of macromolluscs (Moore 1964; García-Cubas 1970; Tunnell 1974; Kay 1980; Vokes and Vokes 1983).

Barrera (2001) stated that within Texas and Mexico, the majority of the studies involved macromolluscs. In Mexico, more than 4,643 species of marine molluscs have been recorded, and approximately 2,067 of them inhabit the Gulf of Mexico (GoMx) and the Mexican Caribbean Sea (Castillo-Rodríguez 2014). Unfortunately, Castillo-Rodríguez did not state which of them were micromolluscs. Important molluscan compilations focusing on these groups include publications by García-Cubas and Reguero (2007) and Vokes and Vokes (1983), although only a few publications have addressed the micromolluscan fauna specifically such as García-Cubas (1963, 1970 and 1971) for lagoons in the Gulf of Mexico and Hicks et al. (2001) at Alacranes reef.

The Campeche Bank is located at the southern GoMx and is composed of several emergent and submerged coral reefs (Tunnell et al. 2010). One of the smaller and most remote reefs is Cayo Nuevo, located between Arenas and Triángulos reefs (190 km offshore) in the GoMx (Fig. 1). Studies on this reef are almost non-existent with the exception of polychaetes (Granados-Barba et al. 2003).

The present work focuses on the molluscs of the Cayo Nuevo sandy bottoms, Gulf of Mexico. In this substrate we could find micromolluscs *sensu stricto* and juveniles of macromolluscs species that inhabit the interstitial as but also empty shells that could be carried by external factors such wind and currents.

Materials and methods

Using SCUBA gear, six sediment samples of coarse sand to fine gravel weighing 224–735 g each were collected by hand at 7.6 and 18.2 m on 19 June 2017 at Cayo Nuevo




Figure 1. Map of the Gulf of Mexico, with the largest reef in the GoMx and sampling locations where micromolluscs have been documented in the literature and this study (adapted from Felder and Camp 2009).

Table 1. Sampling stations and coordinates at Cayo Nuevo on 18 June 2016.

Station	Depth (m)	Latitude (N)	Longitude (W)	Sample
1	18.2	21°49'40.32"	92°4'37.62"	GoMex-001
				GoMex-002
				GoMex-002
				GoMex-004
2	7.6	21°49'47.82"	92°4'34.32"	GoMex-005
				GoMex-006
				GoMex-006

reef (Table 1). Each sample was sieved by pouring water through six differently sized sieves (2 mm, 1.4 mm, 1 mm; 710 μ m, 500 μ m, and 250 μ m) (Table 2) and sorted dry using a dissecting Nikon SMZ800 microscope. Specimens were picked out using soft forceps and 000 fine paint brushes. Molluscs were placed into 2 mm tubes and micromolluscs were placed into PCR tubes, both with 70% ethanol for long-term storage. Identification of specimens to species level was based on Abbott (1974), Tunnell et al. (2010) and Redfern (2013). Whenever possible, at least one specimen of each species

Table 2. Weight of each sample per sieve size in grams.

Sieve	GoMex-001	GoMex-002	GoMex-002	GoMex-004	GoMex-005	GoMex-006
2 mm	86.89	335.85	372.2	645.4	679.1	615.19
1.4 mm	46.94	60.04	21.7	46	80.7	91.53
1 mm	34.01	28.43	5.7	27.3	14.8	22.96
710 µm	27.1	11.13	1.3	28.5	2.3	5.32
500 µm	16.04	3.07	0.3	16.7	0.7	0.82
250 µm	13.86	1.2	0.07	2.6	0.6	0.004
Total	224.84	439.72	401.27	766.5	778.2	735.824

or morphotype was photographed. All the specimens were deposited at the “Colección de Moluscos de la Península de Yucatán” (CMPY), Unidad Multidisciplinaria de Docencia e Investigación Campus Sisal, Universidad Nacional Autónoma de México.

The nomenclature of the species listed was assigned according to Bouchet and Rocroi (2010) for Bivalvia, Bouchet et al. (2017) for Gastropoda, and Kaas and Van Belle (1985) for Polyplacophora, due to the variability of some categories we only present the Linnaean ones. Abundance categories were assigned following Hicks et al. (2001): Abundant ≥ 50 (A); Common = 6–49 (C); Uncommon = 2–5 (UC), and Rare = 1 (R). Juvenile species are denoted by an asterisk (*).

Results

The results from the analysis of the sediment from Cayo Nuevo reef revealed 67 species of molluscs, from which 50 species are gastropods, 14 species are bivalves and three are chitons. These species belong to 38 different families.

Phylum Mollusca Linnaeus, 1758

Class Polyplacophora Gray, 1821

Polyplacophora sp. 1 * (R) (Fig. 2-1)

Polyplacophora sp. 2 * (R) (Fig. 2-2)

Order Chitonina Thiele, 1909

Family Chitonoidea Rafinesque, 1815

Genus *Ischnochiton* Gray, 1847

***Ischnochiton* sp.** * (UC) (Fig. 2-3)

Class Bivalvia Linnaeus, 1758

Order Arcida Stoliczka, 1871

Family Arcidae Lamarck, 1809

Arcidae sp. (UC) (Fig. 2-4)

Genus *Arca* Linnaeus, 1758

***Arca imbricata* Bruguière, 1789** * (UC) (Fig. 2-5a, b)




Figure 2. Polyplacophora 1–3: **1** Polyplacophora sp. 1, dorsal view, scale bar 0.5 mm **2** Polyplacophora sp. 2, dorsal view, scale bar 1 mm **3** *Ischnochiton* sp., dorsal view, scale bar 0.5 mm. Bivalvia 4–15: **4** Arcidae sp., dorsal view, scale bar 0.5 mm **5** *Arca imbricata* **5a** ventral view **5b** dorsal view, scale bar 1 mm **6** *Barbatia domingensis* **6a** ventral view **6b** dorsal view, scale bar 1 mm **7** *Barbatia* sp. **7a** ventral view **7b** dorsal view, scale bar 1 mm **8** *Bentharca* sp. **8a** ventral view **8b** dorsal view, scale bar 1 mm **9** *Anomia* sp. **9a** ventral view **9b** dorsal view, scale bar 1 mm **10** *Carditopsis smithii* **10a** ventral view **10b** dorsal view, scale bar 1 mm, **11** Lucinidae sp., ventral view, scale bar 0.5 mm **12** *Chama sinuosa* **12a** ventral view **12b** dorsal view, scale bar 0.5 mm **13** Chamidae sp. **13a** ventral view **13b** dorsal view, scale bar 1 mm, **14** *Crassinella lunulata* **14a** ventral view **14b** dorsal view, scale bar 1 mm **15** *Semele bellastrata* **15a** dorsal view **15b** lateral view, scale bar 1 mm. Gastropoda 16–23: **16** Gastropoda sp., ventral view, scale bar 0.5 mm, **17** *Diodora minuta*, ventral view, scale bar 0.25, **18** *Diodora listeri* **18a** ventral view **18b** dorsal view, scale bar 1 mm, **19** *Scissurella redfernii* **19a** ventral view **19b** dorsal view, scale bar 1 mm **20** *Synaptocoblea picta* **20a** ventral view **20b** dorsal view, scale bar 1 mm **21** *Lodderena ornata* **21a** ventral view **21b** Apical view, scale bar 0.5 mm **22** *Cerithium* sp. 1 **22a** ventral view **22b** dorsal view, 1 scale bar 1 mm **23** *Cerithium* sp. 2 **23a** ventral view **23b** dorsal view, scale bar 1 mm.

Genus *Barbatia* Gray, 1842

Barbatia domingensis * (Lamarck, 1819) (UC) (Fig. 2-6a, b)

***Barbatia* sp.** * (C) (Fig. 2-7a, b)

Genus *Bentharca* Verrill & Bush, 1898

***Bentharca* sp.** (R) (Fig. 2-8a, b)

Order Mytilida Férussac, 1822

Family Mytilidae Rafinesque, 1815

Genus *Crenella* T. Brown, 1827

***Crenella* sp.** (R)

Order Pectinida Gray, 1854

Family Anomiidae Rafinesque, 1815

Genus *Anomia* Linnaeus, 1758

Anomia* sp. (R) (Fig. 2-9a, b)

Order Cardita Bruguière, 1792

Family Carditidae Férussac, 1822

Genus *Carditopsis* E. A. Smith, 1881

***Carditopsis smithii* (Dall, 1896)** (C) (Fig. 2-10a, b)

Order Lucinida Gray, 1854

Family Lucinidae J. Fleming, 1828

Lucinidae sp. * (UC) (Fig. 2-11)

Order Venerida Gray, 1854

Family Chamidae Lamarck, 1809

Chamidae sp. (UC) (Fig. 2-12a, b)

Genus *Chama* Linnaeus, 1758

***Chama sinuosa* Broderip, 1835** * (A) (Fig. 2-13a, b)

Family Galeommatidae Gray, 1840

Galeommatidae sp. (R)

Order Carditida Dall, 1889

Family Crassatellidae Férussac, 1822

Genus *Crassinella* Guppy, 1874

***Crassinella lunulata* (Conrad, 1834)** * (R) (Fig. 2-14a, b)

Order Cardiida Férussac, 1822

Family Semelidae Stoliczka, 1870 (1825)

Genus *Semele* Schumacher, 1817

***Semele bellastrata* (Conrad, 1837)** * (UC) (Fig. 2-15a, b)

Class Gastropoda Cuvier, 1795

Gastropoda sp. * (R) (Fig. 2-16)

Order Lepetelloidea Dall, 1882

Family Fissurellidae Fleming, 1822

Genus *Diodora* Gray, 1821

***Diodora minuta* (Lamarck, 1822)** * (UC) (Fig. 2-17)

***Diodora listeri* (d'Orbigny, 1847)** (R) (Fig. 2-18a, b)

Family Scissurellidae Gray, 1847

Genus *Scissurella* d'Orbigny, 1824

***Scissurella redferni* (Rolán, 1996)** (C) (Fig. 2-19a, b)

Order Trochida Rafinesque, 1815

Family Trochidae Rafinesque, 1815

Genus *Synaptocochlea* Pilsbry, 1890

***Synaptocochlea picta* (d'Orbigny, 1847)** (A) (Fig. 2-20a, b)

Family Skeneidae W. Clark, 1851

Genus *Lodderena* Iredale, 1924

***Lodderena ornata* (Olsson & McGinty, 1958)** (A) (Fig. 2-21a, b)

Family Cerithiidae Fleming, 1822

Genus *Cerithium* Bruguière, 1789

Cerithium sp. 1 (R) (Fig. 2-22a, 22b)

Cerithium sp. 2 (UC) (Fig. 2-23a, 23b)

***Cerithium atratum* (Borns, 1778)** (R) (Fig. 3-1a, b)

Family Litiopidae Gray, 1847

Genus *Alaba* H. Adams & A. Adams, 1853

***Alaba incerta* (d'Orbigny, 1841)** (C) (Fig. 3-2a, b)

Family Scaliolidae Jousseaume, 1912

Finella sp. (UC) (Fig. 3-3a, 3b)

Family Pickworthiidae Iredale, 1917

Subfamily Pickworthiinae Iredale, 1917

Genus *Sansonia* Jousseaume, 1892

***Sansonia tuberculata* (Watson, 1886)** (R) (Fig. 3-4a, b)

Family Hypponicidae Troschel, 1861

Hipponix sp. (C) (Fig. 3-5a, 5b)




Figure 3. Gastropoda 1–26: **1** *Cerithium atratum* **1a** ventral view **1b** dorsal view, scale bar 1 mm **2** *Alaba incerta* **2a** ventral view **2b** dorsal view, scale bar 0.25 mm **3** *Finella* sp. **3a** ventral view **3b** dorsal view, scale bar 1 mm **4** *Sansonia tuberculata* **4a** ventral view **4b** dorsal view, scale bar 1 mm **5** *Hipponix* sp. **5a** ventral view **5b** dorsal view, scale bar 0.1 mm **6** *Iniforis turristhormae* **6a** ventral view **6b** dorsal view, scale bar 1 mm **7** *Metaxia rugulosa* **7a** ventral view **7b** dorsal view, scale bar 0.5 mm **8** *Cerithiopsis* sp. **8a** ventral view **8b** dorsal view, scale bar 1 mm **9** *Cerithiopsis* cf. *iuxtafuniculata* **9a** ventral view **9b** dorsal view, scale bar 1 mm **10** Vermetidae *incertae sedis* irregularis scale bar 1 mm **11** *Dendropoma corrodens* scale bar 1 mm **12** Vermetid sp. C, ventral view, scale bar 0.25 mm **13** *Petaloconchus mcgintyi*, ventral view, scale bar 0.25 mm **14** *Thylacodes* sp. scale bar 1 mm **15** *Simulamereлина caribaea* **15a** ventral view **15b** dorsal view, scale bar 1 mm **16** *Schwartziella fischeri* **16a** ventral view **16b** dorsal view, scale bar 1 mm **17** *Zebina* sp. 2 **17a** ventral view **17b** dorsal view, scale bar 1 mm **18** *Zebina* sp. 2 **18a** ventral view **18b** dorsal view, scale bar 0.5 mm **19** *Truncatella* sp. **19a** ventral view **19b** dorsal view, scale bar 0.5 mm **20** *Caecum circumvolutum*, lateral view, scale bar 0.2 mm **21** *Caecum donmoorei* **21a** stage 1, lateral view, scale bar 0.2 mm **21b** adult, lateral view, scale bar 0.2 mm **22** *Caecum johnsoni*, lateral view, scale bar 0.2 mm **23** *Caecum pulchellum*, lateral view, scale bar 0.2 mm **24** *Caecum textile*, lateral view, scale bar 0.2 mm **25** *Caecum* sp. B, lateral view, scale bar 0.2 mm **26** *Meioceras nitidum*, lateral view, scale bar 0.2 mm.

Family Triphoridae Gray, 1847

Genus *Iniforis* Jousseaume, 1884

Iniforis turristhormae (Holten, 1802) (UC) (Fig. 3-6a, b)

Genus *Metaxia* Monterosato, 1884

***Metaxia rugulosa* (C. B. Adams, 1850) (R)** (Fig. 3-7a, b)

Family Cerithiopsidae H. Adams & A. Adams, 1853

Genus *Cerithiopsis* Forbes & Hanley, 1850

***Cerithiopsis* sp.** (R) (Fig. 3-8a, 8b)

***Cerithiopsis* cf. *iuxtafuniculata* Rolán, Espinosa & Fernández-Garcés, 2007 (R)**
(Fig. 3-9a, b)

Family Vermetidae Rafinesque, 1815

Vermetidae *incertae sedis irregularis* d'Orbigny, 1841 (Fig. 3-10)

Genus *Dendropoma* Mörch, 1861

***Dendropoma corrodens* (d'Orbigny, 1841) (R)** (Fig. 3-11)

Vermetid sp. C Redfern 2013 (A) (Fig. 3-12)

Genus *Petalococonchus* Lea, 1843

***Petalococonchus mcgintyi* (Olsson & Harbison, 1953) * (C)** (Fig. 3-13)

Genus *Thylacodes* Guettard, 1770

***Thylacodes* sp.** * (R) (Fig. 3-14)

Family Rissoidae Gray, 1847

Genus *Simulamereлина* Ponder, 1985

***Simulamereлина caribaea* (d'Orbigny, 1842) (UC)** (Fig. 3-15a, b)

Family Zebinidae Coan, 1964

Genus *Schwartzziella* G. Nevill, 1881

***Schwartzziella fischeri* (Desjardin, 1949) (UC)** (Fig. 3-16a, b)

Genus *Zebina* H. Adams & A. Adams, 1854

***Zebina* sp. 1** (A) (Fig. 3-17a, 17b)

***Zebina* sp. 2** (C) (Fig. 3-18a, 18b)

Family Truncatellidae Gray, 1840

Genus *Truncatella* Risso, 1826

***Truncatella* sp.** (R) (Fig. 4-19a, 19b)

Family Caecidae Gray, 1850

Genus *Caecum* Fleming, 1813

***Caecum circumvolutum* de Folin, 1867 (C)** (Fig. 3-20)

***Caecum donmoorei* Mitchell-Tapping, 1979 (C)** (Fig. 3-21a, b)

***Caecum johnsoni* Winkley, 1908 (A)** (Fig. 3-22)




Figure 4. Gastropoda 1–18: **1** *Parviturboides* sp. **1a** ventral view **1b** dorsal view, scale bar 1 mm **2** *Vitrinella* sp. **2a** ventral view **2b** dorsal view, scale bar 0.5 mm **3** *Gibberula lavalleeana* **3a** ventral view **3b** dorsal view, scale bar 1 mm **4** *Volvarina* sp. **4a** ventral view **4b** dorsal view, scale bar 1 mm **5** *Volvarina* sp. **2** **5a** ventral view **5b** dorsal view, scale bar 1 mm **6** Columbellidae sp. **1** **6a** ventral view **6b** dorsal view, scale bar 0.5 mm **7** Columbellidae sp. **2** **7a** ventral view **7b** dorsal view, scale bar 0.5 mm **8** *Steironepion moniferum* **8a** ventral view **8b** dorsal view, scale bar 1 mm **9** *Phrontis* sp. **9a** ventral view **9b** dorsal view, scale bar 0.5 mm **10** *Trachypollia* sp. **10a** ventral view **10b** dorsal view, scale bar 1 mm **11** Turritidae sp. **1** **11a** ventral view **11b** dorsal view, scale bar 1 mm **12** *Ammonicera lineofuscata*, Apical view, scale bar 0.25 mm **13** *Ammonicera minorialis* **13a** Apical view **13b** ventral view, scale bar 0.1 mm **14** *Rissoella galba* **14a** ventral view **14b** dorsal view, scale bar 0.1 mm **15** *Pseudoscilla* aff. *babylonia* **15a** ventral view **15b** dorsal view, scale bar 1 mm.

Caecum pulbellum Stimpson, 1851 (C) (Fig. 3-23)

Caecum textile de Folin, 1867 (UC) (Fig. 3-24)

Caecum sp. **B sensu** Redfern 2013 (C) (Fig. 3-25)

Genus *Meioceras* Carpenter, 1859

Meioceras nitidum (Stimpson, 1851) (UC) (Fig. 3-26)

Family Tornidae Sacco, 1896 (1884)

Genus *Parviturboides* Pilsbry & McGinty, 1949

Parviturboides sp. (C) (Fig. 4-1a, b)

Genus *Vitrinella* C. B. Adams, 1850

Vitrinella sp. (A) (Fig. 4-2a, b)

Family Cystiscidae Stimpson, 1865

Genus *Gibberula* Swainson, 1840

***Gibberula lavalleana* (d'Orbigny, 1824)** (UC) (Fig. 4-3a, b)

Family Marginellidae Fleming, 1828

Genus *Volvarina* Hinds, 1844

***Volvarina* sp. 1** (UC) (Fig. 4-4a, 4b)

***Volvarina* sp. 2** (R) (Fig. 4-5a, 5b)

Family Columbelloidea Swainson, 1840

Columbellidae sp. 1 (UC) (Fig. 4-6a, b)

Columbellidae sp. 2 (R) (Fig. 4-7a, 7b)

Genus *Steironepion* Pilsbry & H. N. Lowe, 1932

***Steironepion moniliferum* (G. B. Sowerby I, 1844)** (UC) (Fig. 4-8a, b)

Family Nassariidae Iredale, 1916 (1835)

Genus *Phrontis* H. Adams & A. Adams, 1853

***Phrontis* sp.** (UC) (Fig. 4-9a, b)

Family Muricidae Rafinesque, 1815

Genus *Trachypollia* Woodring, 1928

***Trachypollia* sp.** (R) (Fig. 4-10a, b)

Family "Turridae" H. Adams & A. Adams, 1853 (1838)

Turridae sp. 1 (R) (Fig. 4-11a, 11b)

Family Omalogyridae G.O. Sars, 1878

Genus *Ammonicera* Vayssière, 1893

***Ammonicera lineofuscata* Rolán, 1992** (A) (Fig. 4-12)

***Ammonicera minortalis* Rolán, 1992** (A) (Fig. 4-13a, b)

Family Rissoellidae Gray, 1850

Genus *Rissoella* Gray, 1847

***Rissoella galba* Robertson, 1961** (R) (Fig. 4-14a, b)

Order Aplysiida

Family Aplysiidae Lamarck, 1809

Genus *Aplysia* Linnaeus, 1767

***Aplysia* sp.** (R)

Order Siphonarimorpha

Family Pyramidellidae Gray, 1840

Genus *Pseudoscilla* Boettger, 1901

***Pseudoscilla* aff. *babylonia* (C. B. Adams, 1845)** (R) (Fig. 4-15a, 15b)

Discussion

The most abundant families of gastropods were the Caecidae (456 specimens, seven species), Tornidae (221 specimens, two species), and Omalogyridae (132 specimens, two species). The most abundant families of bivalves were Arcidae (40 specimens, five species) and Chamidae (59 specimens, two species). The most abundant gastropod species were *Caecum johnsoni* (310 specimens), *Vitrinella* sp. (208 specimens), *Vermetid* sp. C (91 specimens), *Lodderena ornata* (71 specimens), and *Caecum donmoorei* (147 specimens). For the Bivalvia the most abundant species were *Chama sinuosa* (57 specimens), *Barbatia domingensis* (57 specimens) and *Carditopsis smithii* (12 specimens).

From the six sediment samples, the most commonly found molluscs were *Lodderena ornata*, *Caecum johnsoni* and *Ammonicera lineofuscata*, while other species appeared only once: *Leptochiton* sp., Arcidae sp., *Bentharca* sp., *Crenella* sp., *Anomia* sp., *Chama* sp., Galeommatidae sp., *Chione elevata*, *Semele bellastrata*, *Cerithium* sp. 1, Gastropoda sp., *Diodora listeri*, *Cerithium atratum*, *Sansonia tuberculata*, *Iniforis turristhormae*, *Metaxia rugulosa*, *Cerithiopsis* cf. *iuxtafuniculata*, *Cerithiopsis* sp., *Vermetidae incertae sedis* irregularis, *Dendropoma corrondens*, *Thylacodes* sp., *Finella* sp., *Caecum textile*, *Hipponix* sp., *Volvarina* sp. 2, Columbellidae sp. 2, *Aplysia* sp., and *Pseudoscilla* aff. *babylonia*.

This new data becomes a taxonomic reference list for the molluscs that inhabit Cayo Nuevo, GoMx, including micromolluscs as well as juvenile macromolluscs. To place it within a useful context we mention other inventories made in this area: Felder and Camp (2009) recorded some 5,517 species of invertebrates in the GoMx, of which 2,455 were marine molluscs (Moretzsohn et al. 2009). González et al. (1991) recorded 298 species of molluscs and included 33 localities distributed around the coasts of the Yucatan Peninsula and adjacent coral reefs but did not mention Cayo Nuevo. García-Cubas et al. (1999) recorded 110 species of gastropods in the northern and northeastern regions of the Yucatan Peninsula. Rice and Kornicker (1962) recorded 130 species for Alacranes reef in the Campeche Bank and later, Hicks et al. (2001) recorded 215 species of molluscs on the same reef. Although earlier articles include reefs or sampling locations within the Bank of Campeche (e.g., Rehder and Abbott 1951, Springer and Bullis 1956, Kornicker et al. 1959), no mention of molluscs from Cayo Nuevo were found. Only Barrera (2001) study focuses on reef micromolluscs, recording 131 species from the East and West Flower Garden Banks (FGB).

The molluscan assemblage at Cayo Nuevo shares many species also present at the FGB (Barrera 2001) and Alacranes Reef (Hicks et al. 2001) (19 families/21 genera and 21 families/22 genera, respectively) (Table 3). The most diverse families recorded by Barrera (2001) and Hicks et al. (2001) were Caecidae (six genera and ten species) and Rissoidae (five genera and seven species) for Gastropoda and Arcidae (four genera and seven species) for Bivalvia.

Barrera (2001) most abundant gastropod species were *Amphithalamus vallei* (672 individuals) and cf. *Vitrinella* sp. (534 individuals); however, at Cayo Nuevo, 208 individuals of sf. *Vitrinella* sp., were found. Differences in the numbers of collected individ-

Table 3. Comparison of molluscs recorded at Cayo Nuevo (present study) and other interstitial records at the GoMx; Madre: García-Cubas (1970), Tamiahua: García-Cubas (1971), Términos: García-Cubas (1963, 1981), East Flower Garden: Barrera and Tunnell (2001), Alacranes: Hicks et al. (2001). The maximum recorded size is provided: Key: a) García-Cubas and Reguero (2004), b) Tunnell et al. (2010), c) Redfern (2013).

Systematics	Lagoons			Reefs			Maximum recorded size (mm)
	Madre	Tamiahua	Términos	East Flower Garden	Alacranes	Cayo Nuevo	
Class Bivalvia							
<i>Arca imbricata</i>			*		*	*	48 ^c
<i>Barbatia domingensis</i>				*	*	*	30 ^b
<i>Carditopsis smithii</i>				*	*	*	1.5 ^c
<i>Chama sinuosa</i>					*	*	76 ^b
<i>Crassinella lunulata</i>		*				*	8 ^b
<i>Semele bellastrata</i>						*	14 ^c
Class Gastropoda							
<i>Diodora minuta</i>						*	10.5 ^c
<i>Diodora listeri</i>					*	*	45 ^b
<i>Scissurella redferni</i>						*	1 ^c
<i>Synaptocochlea picta</i>					*	*	3.5 ^c
<i>Lodderena ornata</i>				*	*	*	0.8 ^c
<i>Cerithium atratum</i>						*	
<i>Alaba incerta</i>						*	
<i>Sansonia tuberculata</i>						*	1.5 ^c
<i>Iniforis turrishomae</i>				*	*	*	6 ^c
<i>Metaxia rugulosa</i>						*	7 ^c
<i>Cerithiopsis</i> cf. <i>iuxtafuniculata</i>						*	3 ^c
Vermetidae <i>incertae sedis irregularis</i>						*	6 ^b
<i>Dendropoma corrodens</i>					*	*	10 ^b
<i>Petalocochus mcgintyi</i>				*	*	*	35 ^b
<i>Simulamerelina caribaea</i>						*	3 ^b
<i>Schwartziella fischeri</i>						*	3.5 ^c
<i>Caecum circumvolutum</i>						*	4 ^b
<i>Caecum donmoorei</i>						*	2 ^c
<i>Caecum johnsoni</i>			*	*		*	5 ^b
<i>Caecum pulchellum</i>	*	*	*	*	*	*	2 ^a
<i>Caecum textile</i>					*	*	2 ^b
<i>Meioceras nitidum</i>		*	*			*	3 ^b
<i>Gibberula lavalleana</i>		*				*	4 ^b
<i>Steironepion moniliferum</i>						*	
<i>Ammonicera lineofuscata</i>						*	0.6 ^c
<i>Ammonicera minortalis</i>				*		*	0.5 ^c
<i>Rissoella galba</i>				*		*	1 ^c
<i>Pseudoscilla</i> aff. <i>babylonia</i>				*	*	*	3 ^b

uals can be explained by geography, but also by differences in the quantity of sediment collected and processed, fifteen sites with 300 ml sediment sampled at FGB and six samples of 224–735 g at Cayo Nuevo. *Vitrinella* sp. could not be identified to specific

level due to the low similarity of characters shown with other described western Atlantic species. Barrera (2001) previously suggested that it could be an undescribed new species, but further detailed studies are required to establish its identity.

Regarding bivalves, the most abundant species reported by Barrera (2001) for the FGB were *Gregariella coralliophaga* (145 individuals) (summing nine identified as *Barbatia domingensis* (102 individuals) and eleven identified as *Barbatia cancellaria* (68 individuals, currently a synonym), and *Carditopsis smithii* (51 individuals). In comparison, 37 individuals of *Barbatia domingensis* and 12 *Carditopsis smithii* were collected at Cayo Nuevo and these were not the most abundant species.

It should come as no surprise that many organisms were not identified to species level (e.g., *Leptochiton* sp., *Bentharca* sp., *Crenella* sp., *Anomia* sp., *Lottia* sp., *Diodora* sp., *Cerithium* sp., *Cerithiopsis* sp., *Thylacodes* sp., *Zebina* sp. 1, *Zebina* sp. 2, and *Phrontis* sp.). We relied on regional and local literature that in fact was scarce. In the case of juveniles, shells within a genus are similar because they share many characters and the differential characters are difficult to discern even as adults and almost impossible in juveniles. Our specimens identified under the name of *Gibberula lavalleeana* could be considered as a species complex, due to the evidence and description of new species in Cuban waters (Espinosa and Ortea 2007).

These faunistic results from Cayo Nuevo represent the first inventory of molluscs from this remote reef. These findings contribute to record expansions for the southern GoMx of *Bentharca* sp. This contribution highlights the importance of conserving small areas that can harbour a considerable diversity of organisms. Seasonal changes on the mollusc community assemblages were not evaluated but would be an interesting future project, as would collecting growth series of species to assist in confirming identifications. In 2004, González and Torruco stated the importance of the Campeche Bank's reefs and proposed a marine reserve for the reefs located within this area, Cayo Nuevo included. However, this proposal never materialized and, up to now, only Alacranes reef has governmental protection under the status of marine reserve. Species checklists of micromolluscs, as well as other faunal groups, are of vital importance to serve as a baseline data set, due to the proximity to Mexico's offshore oil production area within the GoMx. The soft benthic interstitial mollusc communities are diverse, and their monitoring could well represent ecological indicators of ecosystem health, especially in the light of potential future oil-spills.

Acknowledgements

We give our gratitude to the researchers from the Yucatan Marine Biodiversity Team (BDMY) who participated in the expedition on the boat "Isla Mujeres", especially to P. Tapia Díaz who collected the samples. Thanks to M. Badillo Alemán and C. Galindo de Santiago for their general laboratory technical assistance and to B. Rodrigo Cobarrubias Contreras for his technical assistance with the granulometry analysis, all at the Unidad Académica Sisal, UNAM. Professor Fabio Moretzsohn kindly provided the

base map. Financial support provided by the Harte Charitable Foundation, through the Harte Research Institute (Biodiversity of the southern Gulf of Mexico) and by the CONABIO NE018 Project (Update on the knowledge of the biodiversity of shallow water benthic marine invertebrate species (<50m) from the southern GoMx). First author received a postdoctoral grant from DGAPA-UNAM 2018-2019. We are grateful to Editor Nathalie Yonow for comments in the manuscript. We also thank the reviewer Dr. Henk Dekker and two other anonymous reviewers for their constructive comments and suggestions. We dedicate this work to our malacologist colleague, Dr. Wes Tunnell, who just left us. Dr. Tunnell was always curious for the Gulf of Mexico invertebrates and tireless promoter of marine biodiversity knowledge in the region.

References

- Abbott RT (1974) American Seashells. The marine Mollusca of the Atlantic and Pacific Coast of North America. Van Nostrand Reinhold, New York, 663 pp.
- Albano PG, Sabelli B, Bouchet P (2011) The challenge of small and rare species in marine biodiversity surveys: Microgastropod diversity in a complex tropical coastal environment. *Biodiversity and Conservation* 20: 3223–3237. <https://doi.org/10.1007/s10531-011-0117-x>
- Barrera N, Tunnell JW (2001) Micromolluscan faunal survey of the East and West Flower Garden Banks. Long-term monitoring of the East and West Flower Garden Banks, 1998–1999. *EUA*, New Orleans, Louisiana, 99–108.
- Barrera N (2001) Micromolluscan Assemblages on the Flower Garden Banks, Northwestern Gulf of Mexico. Texas A&M University, Corpus Christi, 90 pp.
- Bouchet P, Lozouet P, Maestrati P, Heros V (2002) Assessing the magnitude of species richness in tropical marine environments: Exceptionally high numbers of molluscs at a New Caledonia site. *Biological Journal of the Linnean Society* 75: 421–436. <https://doi.org/10.1046/j.1095-8312.2002.00052.x>
- Bouchet P, Rocroi J-P, Bieler R, Carter JG, Coan EV (2010) Nomenclator of Bivalve Families with a Classification of Bivalve Families. *Malacologia* 52: 1–184. <https://doi.org/10.4002/040.052.0201>
- Bouchet P, Rocroi J-P, Hausdorf B, Kaim A, Kano Y, Nützel A, Parkhaev P, Schrödl M, Strong EE (2017) Revised Classification, Nomenclator and Typification of Gastropod and Monoplacophoran Families. *Malacologia* 61: 1–526. <https://doi.org/10.4002/040.061.0201>
- Castillo-Rodríguez ZG (2014) Biodiversidad de moluscos marinos en México. *Revista mexicana de Biodiversidad* 85: 410–430. <https://doi.org/10.7550/rmb.33003>
- Espinosa J, Ortea J (2007) El género *Gibberula* Swainson, 1840 (Mollusca: Neogastropoda: Cystiscidae) en Cuba, con la descripción de nuevas especies. *Avicennia* 19: 99–120.
- Felder DL, Camp DK (2009) Gulf of Mexico: origin, waters, and biota. Volume 1. Biodiversity. Texas A&M University Press, Texas, 1312 pp.
- García-Cubas A (1963) Sistemática y distribución de los micromoluscos recientes de la laguna de Terminos, Campeche, Mexico. *Universidad Nacional Autónoma de México. Instituto de Geología* 67: 1–55.

- García-Cubas A, Escobar F, Reguero M (1999) Gastrópodos marinos de la Península de Yucatán, México. *Revista de la Sociedad Mexicana de Historia Natural* 49: 127–146.
- García-Cubas A (1970) Ecología y distribución de los micromoluscos recientes de la Laguna Madre, Tamaulipas, México. *Instituto de Geología, Boletín* 86: 1–44.
- García-Cubas A (1971) Ecología y distribución de los micromoluscos de la Laguna de Tamiahua, Veracruz, Mexico. *Universidad Nacional Autónoma de Mexico* 91: 1–53.
- García-Cubas A (1981) Moluscos de un sistema lagunar tropical en el sur del Golfo de México (Laguna de Términos, Campeche). *Instituto de Ciencias del Mar y Limnología. Universidad Nacional Autónoma de México. Publicación especial* 5: 182.
- García-Cubas A, Reguero M (2007) Catálogo ilustrado de moluscos bivalvos del Golfo de México y Mar Caribe. *Instituto de Ciencias del Mar y Limnología. Universidad Nacional Autónoma de México, México*, 94 pp.
- Geiger DL, Marshall BA, Ponder WF, Sasaki T, Warén A (2007) Techniques for collecting, handling, preparing, storing and examining small molluscan specimens. *Molluscan Research* 27: 1–50.
- González MA, Chávez E, de la Cruz G, Torruco D (1991) Patrones de distribución de gasterópodos y bivalvos en la Península de Yucatán, México. *Ciencias Marinas* 17: 147–172. <https://doi.org/10.7773/cm.v17i3.810>
- González MA, Torruco D (2004) Diagnóstico para la creación de una reserva de la biosfera en los arrecifes de Campeche, México; Diagnosis for creation of a biosphere reserve in the reefs of Campeche, Mexico. *Proceedings of the Gulf and Caribbean Fisheries Institute* 55: 685–692.
- Granados-Barba A, Solís-Weiss V, Tovar-Hernández MA, Ochoa-Rivera V (2003) Distribution and diversity of the Syllidae (Annelida: Polychaeta) from Mexican Gulf of Mexico and Caribbean. *Hydrobiologia* 496: 337–345. <https://doi.org/10.1023/A:1026177722364>
- Hicks DW, Barrera N, Tunnell JW (2001) Ecological distribution of shallow-water Mollusca on Alacran Reef, Campeche Bank, Yucatan, Mexico. *Texas Conchologist* 38: 7–30.
- Kaas P, Bell RA van, Strack HL (1985) Monograph of living chitons (Mollusca: Polyplacophora). E.J. Brill/Dr. W. Backhuys, Leiden, 240 pp.
- Kay EA (1980) Micromollusks: Techniques and patterns in benthic marine communities. In: *Environmental Survey Techniques for Coastal Water Conference proceedings. Water Resources Research Center, University of Hawaii at Manoa, Honolulu, and Hawaii Water Pollution Control Association*, 93–112.
- Kornicker S, Bonet F, Ross C, Charles H (1959) Alacran Reef, Campeche Bank, Mexico. *Publications of the Institute of Marine Science, University of Texas* 6: 1–22.
- Moore DR (1964) The family Vitrinellidae in south Florida and the Gulf of Mexico. PhD diss., University of Miami Coral Gables.
- Moretzsohn F, Tunnell Jr W, Lyons WG, Baqueiro E, Barrera N, Espinosa E, García EF, Ortea J, Regueiro M (2009) Mollusca: Introduction. In: *Felder DL, Camp DK (Eds) Gulf of Mexico Origin, Waters, and Biota. Volume 1, Biodiversity. Texas A&M University Press, Texas*, 559–564.
- Redfern C (2013) Bahamian seashells: 1161 species from Abaco, Bahamas. *Bahamian Seashells, Inc., Florida, USA*, 501 pp.

- Rehder HA, Abbott RT (1951) Some new and interesting mollusks from the deeper waters of the Gulf of Mexico. *Revista de la Sociedad Malacológica "Carlos de la Torre"* 8: 53–66.
- Rice WH, Kornicker S (1962) Mollusks of Alacran Reef, Campeche Bank, México. *Publications of the Institute of Marine Science* 8: 366–403.
- Sasaki T (2008) Micromolluscs in Japan: taxonomic composition, habitats, and future topics. *Zoosymposia* 1: 147–232. <https://doi.org/10.11646/zoosymposia.1.1.12>
- Springer S, Bullis HR (1956) Collections by the Oregon in the Gulf of Mexico. List of crustaceans, mollusks and fishes. Identified from collections by the exploratory fishing vessel Oregon in the Gulf of Mexico and adjacent seas 1950 through 1955. United States Fish and Wildlife Service, Special Scientific Report. Fisheries 196: 1–134.
- Tunnell JW (1974) Ecological and geographical distribution of Mollusca of Lobos and Enmedio coral reefs, Southwestern Gulf of Mexico. PhD Thesis, Texas A&M University, College Station.
- Tunnell JW, Felder DL, Earle SA, Camp DK (2009) Mollusca of the Gulf of Mexico. In: Felder DL, Camp DK (Eds) *Gulf of Mexico Origin, Waters, and Biota*. Volume 1, Biodiversity. Texas A&M University Press, Texas, 559–749.
- Tunnell JW, Hicks DW, Withers K (2010) *Encyclopedia of Texas seashells: identification, ecology, distribution, and history*. Texas A&M University Press, Texas, 512 pp.
- Vokes HE, Vokes EH (1983) Distribution of shallow-water marine mollusca, Yucatan Peninsula, Mexico. Mesoamerican Ecology Institute, Middle American Research Institute, Tulane University, New Orleans, 183 pp.

Supplementary material I

Species abundance per sample

Authors: Deneb Ortigosa, Nancy Yolimar Suárez-Mozo, Noe C. Barrera, Nuno Simões

Data type: occurrences

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.779.24562.suppl1>