

HLA-Cw6 increases the risk of psoriasis and early onset before twenty-seven years of age among the Vietnamese population

Nhi Thi Uyen Pham,^{1,2} Thuong Van Nguyen,³ Hao Trong Nguyen¹

¹Ho Chi Minh City Hospital of Dermato-Venereology, Ho Chi Minh City; ²Hanoi Medical University, Hanoi; ³National Hospital of Dermatology and Venereology, Hanoi, Vietnam

Correspondence: Hao Trong Nguyen, Ho Chi Minh City Hospital of Dermato-Venereology, 2 Nguyen Thong, Ward Vo Thi Sau, District 3, Ho Chi Minh City, Vietnam.
Tel.: +84.903639234.
E-mail: bshao312@yahoo.com

Key words: psoriasis; polymorphism; HLA-Cw6; psoriasis susceptibility; early onset.

Acknowledgments: the authors thank the Center for Molecular Biomedicine, University of Medicine, and Pharmacy at Ho Chi Minh City for supporting the simulated gene sequencing assays for psoriasis. We also thank our colleagues from the Psoriasis Clinic of Ho Chi Minh City Hospital of Dermato-Venereology for collecting the data.

Contributions: NTUP, TVN, data acquisition; NTUP, TVN, HTN, analysis of data, and publishing edition finalization; HTN, manuscript drafting. All the authors approved the final version to be published.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: none.

Ethical approval and consent to participate: this study exhibited scrupulous adherence to the ethical guidelines in biomedical research. All laboratory examinations were meticulously conducted strictly following established protocols and testing regulations. This study was approved by the Ethics Council for Biomedical Research at Hanoi Medical University under reference number 66/GCN-HDDNCCBH HMU, issued on March 16th, 2020. The participants were clearly explained the study's objectives and protocols and written informed consent was obtained prior to study enrollment.

Availability of data and material: all data used to support the findings of this study are included within the article. Moreover, the Excel data used to support the findings of this study are available from the corresponding author upon request.

Received: 18 September 2023.

Accepted: 5 October 2023.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0).

©Copyright: the Author(s), 2023

Licensee PAGEPress, Italy

Dermatology Reports 2024; 16:9854

doi:10.4081/dr.2023.9854

Publisher's note: all claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Abstract

Psoriasis is a chronic inflammatory skin disease that affects people all over the world. It is linked to the HLA-Cw6 allele, which is more common in Caucasians than in Asians and varies across ethnic groups. We investigated the association between the disease severity and the onset age of HLA-Cw6 prevalence in Vietnamese psoriasis patients. In 121 psoriasis patients and 30 healthy controls, we looked at the relationship between HLA-Cw6 and clinical features. We found that patients with psoriasis had significantly higher levels of HLA-Cw6 (64.5%) than controls (26.7%) ($p=0.0001$), with an odds ratio of 4.98 (2.04-12.15). Positive HLA-Cw6 patients had a significantly lower mean age of psoriasis onset than negative HLA-Cw6 patients. Patients with mild psoriasis (100%) were more likely to have the AA genotype, while patients with moderate to severe psoriasis (47.2% and 59.0%, respectively) and those with high PASI scores (55.1% and 54.1%, respectively) were more likely to have the TA genotype. Thus, HLA-Cw6 is a major genetic risk factor for psoriasis in Vietnamese patients, especially early-onset cases. Variations in HLA-Cw6 genotypes also affect disease severity.

Introduction

Psoriasis is a chronic inflammatory skin disorder that recurs frequently and affects more than 3% of the United States adult population.¹ Besides causing skin lesions, psoriasis profoundly impacts the overall well-being of patients and their quality of life in comparison with internal medical conditions, such as cardiovascular diseases and cancer.² The pathogenesis of psoriasis primarily involves a complex interplay of genetics, the immune system, and environmental factors, leading to diverse clinical manifestations, prognoses, and comorbidities among patients. Among the genetic factors, the HLA-Cw6 allele has emerged as the most extensively studied, with associations ranging from disease pathogenesis, epidemiology, and clinical manifestations to comorbidities and treatment responses.

HLA-Cw6 is considered to be one of the most susceptible alleles to psoriasis. The frequency of the HLA-Cw6 allele in psoriasis varies across different ethnicities worldwide, ranging from 10.5 to 77.2%.³ The prevalence of HLA-Cw6 positivity among patients with psoriasis is generally lower in Asian countries than in Caucasians. In China, the rate ranges from 32.5 to 50.8%,^{4,5} while in Japan, it fluctuates between 10.2-12.0%.^{6,7} In contrast, in South Korea, the prevalence is relatively high at 76.1%.⁸ This indicates that there is indeed variability in the prevalence of HLA-Cw6, even within the Asian population. However, these rates were consistently higher than those observed in the control groups in the aforementioned studies. This suggests a correlation between the pres-

ence of HLA-Cw6 and psoriasis, confirming its significant role in the development of this condition. HLA-Cw6 has been extensively associated with early-onset psoriasis. Previous studies have established an association between HLA-Cw6 and the age at which psoriasis first appears, indicating a heightened susceptibility of individuals carrying the HLA-Cw6 allele to develop psoriasis at a younger age. Queiro *et al.* observed a significant difference in the average age of onset between individuals with a positive HLA-Cw6 (23±12 years) and those with a negative HLA-Cw6 (32±12 years).⁹ Enerbäck *et al.* revealed that individuals with an onset age between 30 and 35 had a similar prevalence of HLA-Cw6 compared with the control group. In contrast, those who developed psoriasis before the age of 21 years showed a significantly higher frequency of HLA-Cw6.¹⁰ These findings support the association between the HLA-Cw6 allele and early onset of psoriasis. Furthermore, there is evidence from a few studies that mention the correlation between the HLA-Cw6 allele and the severity of psoriasis. The clinical phenotype and severity of the disease may be associated with certain genetic factors that may be predicted in advance.¹¹ Individuals who carry the HLA-Cw6 allele are likely to have a higher prevalence of extensive plaques on their arms, legs, and trunk.³ However, previous studies have not elucidated the association between HLA-Cw6 and disease severity. This novel study investigated the prevalence of the HLA-Cw6 allele in Vietnamese patients with psoriasis and its potential correlation with clinical characteristics, such as age at onset and disease severity.

Materials and Methods

Patients and controls

A total of 121 patients with psoriasis and 30 healthy controls were enrolled. The inclusion criteria were patients diagnosed with psoriasis based on the 2016 World Health Organization criteria at Ho Chi Minh City Hospital of Dermato-Venereology, aged ≥18, having both parents of Vietnamese descent, and volunteering to participate. Exclusion criteria included patients with severe acute illnesses; direct family members with autoimmune, hereditary, or other immunological disorders; and pregnant or breastfeeding women. The control group comprised healthy individuals without a personal or family history of autoimmune diseases, hereditary disorders, or psoriasis.

HLA-Cw6 sequencing

Sample collection and DNA extraction

2 mL of peripheral blood were collected into ethylene diamine tetra acetic acid anticoagulant tubes, refrigerated, and processed using the ReliaPrep™ Blood gDNA Miniprep System kit (Promega, USA).

Polymerase chain reaction primer design

The Oligo 4.1 software was used based on the standard sequence of the HLA-C gene with accession number NG_029422 in GenBank. Exon 3 of the HLA-C gene was amplified using the primer pair HLA-Cg3F (5'-AGTCGCCTTTACCCGGTTTC-3') and HLA-Cg3R (5'-AGTGCTGATCCCATTTCCT-3'), resulting in a polymerase chain reaction (PCR) product of 472 base pairs. Integrated DNA Technologies (Park Road, Singapore) performed primer synthesis.

Polymerase chain reaction response

The 25 µl mixture included 2.5 µL of 10X PCR buffer, 2.5 µL of dNTPs (250 µM each), 2.5 µL of forward and reverse primers (0.5 µM each), 0.25 µl of TaKaRa Taq™ HotStart Polymerase (Takara Bio, Japan), and 50-100 ng of DNA. Thermal cycling conditions were 98°C for 3 minutes, followed by 35 cycles of 98°C for 10 seconds, 60°C for 30 seconds, and 72°C for 40 seconds, with a final extension at 72°C for 2 minutes. The PCR product was analyzed on a 1.5% agarose gel using the Geldoc-It™ gel documentation system (UVP, USA) and purified with ExoSAP-IT® PCR Product Cleanup kit (Thermo Scientific) following the instructions.

Sequencing reaction

Purified PCR products were cycle-sequenced in both directions using BigDye V3.1 (Applied Biosystems, USA). DNA sequences were determined using an ABI 3500 Genetic Analyzer. Sequencer software (version 5.0) was used to analyze the results of the HLA-C gene sequence (NG_029422, GenBank).

Statistical analyses

Excel was used for data entry and R for processing. T-test, ANOVA, and Kruskal-Wallis assessed mean differences, while χ^2 and Fisher's exact tests analyzed proportions. The correlation used an odds




Figure 1. HLA-Cw6 sequencing.

ratio with a 95% confidence interval, providing valuable insights into the research variables. Statistical significance was set at $p < 0.05$.

Results

We collected and analyzed the genotype of HLA-Cw6 in 121 patients with psoriasis and 30 controls at the Ho Chi Minh City Hospital of Dermato-Venereology.

Characteristics of the study group

All basic characteristics of the study group are presented in Table 1. The mean age of the patients was 42.5 ± 12.4 years (ranging from 21 to 73 years). Male patients were predominantly in the psoriasis group, accounting for 63.3% of the total patients.

The duration of psoriasis in patients was 12.22 ± 10.22 years, with the shortest duration being < 1 year and the longest duration reaching up to 69 years. The mean age of onset for the psoriasis patient group was 27.79 ± 10.94 , ranging from as young as four years old to as old as 60. Among patients with psoriasis, 25.6% had a family history. The average body mass index was recorded as 23.39 ± 5.33 , while the mean psoriasis area and severity index (PASI) score was reported as 20.31 ± 12.70 .

The association between HLA-Cw6 and clinical features of psoriasis

Our study involved sequencing the HLA-Cw6 gene in wells 49: AA, 50: TT, and 51: TA (Figure 1).

Sequencing of 151 cases, including the disease and control groups, revealed that HLA-Cw6 was positive in 57% of the total samples and negative in 43%. Specifically, patients with psoriasis showed a significantly higher proportion of HLA-Cw6 positivity (64.5%), whereas in the control group, the positivity rate for HLA-Cw6 was only 26.7%. The difference between the two groups was significant ($P = 0.0001$). Those with a positive HLA-Cw6 had 4.98 times (95% CI: 2.04-12.15, $p < 0.01$) higher odds of having psoriasis than those with a negative HLA-Cw6. On analysis of the allele genotypes of HLA-Cw6, the results reveal that patients with psoriasis had HLA-Cw6 genotypes in a heterogeneous (TA) form. Homogeneous (TT) genotypes comprised 14.1% of cases, whereas genotype AA represented 35.5%. The proportion of genotype AA in the control group was higher in the control group than that in the psoriasis group (73.3% vs. 35.5%). This difference was significant ($p = 0.001$). The details of this analysis are presented in Table 2. Our study observed that the group of psoriasis patients with positive HLA-Cw6 had a mean onset age of 26.1 ± 10.9 years, which was significantly different from the group of patients with

Table 1. Basic characteristics.

Characteristics	Total n (%) [*]	Study group		p
		Psoriasis (n=121) n (%) [*]	Control (n=30) n (%) [*]	
Age (mean)	42.5±12.3 (21-73)	42.5±12.4 (21-73)	42.4±12.2 (24-67)	1
Ethnicity				0.48
Kinh (Vietnamese)	148 (98.0)	119 (98.4)	29 (96.7)	
Ethnic minorities of Vietnamese	3 (2.0)	2 (1.6)	1 (3.3)	
Sex			0.71	
Male	95 (62.9)	77 (63.6)	18 (60.0)	
Female	56 (37.1)	44 (36.4)	12 (40.0)	
Address			-	-
Ho Chi Minh City	64 (52.9)	64 (52.9)		
Outside Ho Chi Minh City	57 (47.1)	57 (47.1)		
Education			-	-
Primary school	3 (2.5)	3 (2.5)		
Secondary school	10 (8.3)	10 (8.3)		
High school	21 (17.4)	21 (17.4)		
University and postgraduate	44 (36.3)	44 (36.3)		
Career			-	-
Office worker	33 (27.3)	33 (27.3)		
Unskilled laborer	23 (29.7)	23 (29.7)		
Self-employed	11 (9.1)	11 (9.1)		
Student	10 (8.3)	10 (8.3)		
Laborer	4 (3.3)	4 (3.3)		
Homemaker	4 (3.3)	4 (3.3)		
Others	36 (29.7)	36 (29.7)		
Disease duration, years (mean)	12.22±10.22 (0-69)	12.22±10.22 (0-69)	-	-
Age of onset, years (mean)	27.79±10.94 (4-60)	27.79±10.94 (4-60)	-	-
Family history of psoriasis			-	-
Yes	31 (25.6)	31 (25.6)		
No	90 (74.4)	90 (74.4)		
Body mass index (mean)	23.39±5.33 (15.43-37.04)	23.39±5.33 (15.43-37.04)	-	-
Psoriasis area and severity index (mean)	20.31±12.70 (1.8-61.6)	20.31±12.70 (1.8-61.6)	-	-

SD, standard deviation. ^{*}Results are expressed as mean ± SD (range).

psoriasis with negative HLA-Cw6 (31.9±10.3 years) (p=0.002). There was a significant difference between the HLA-Cw6 gene and the age of onset of the patients, with the TT genotype showing the earliest onset, followed by the TA and AA genotypes (P=0.01) (Table 3). Our results revealed that the AA phenotype was commonly observed in mild cases according to the body surface area (BSA) classification (100%). The TA phenotype was more prevalent in the moderate and severe cases (47.2% and 59.0%, respectively). This difference was significant (p=0.01). Similarly, based on the BSA classification, the AA phenotype was commonly found in mild cases according to the PASI classifications (73.3%). The TA phenotype was more prevalent in the moderate and severe

cases (55.1% and 54.1%, respectively). This difference was significant (P=0.03). However, there was no significant difference between the HLA-Cw6 genotypes and the dermatology life quality index classification (P=0.28) (Table 4).

Discussion

The HLA-Cw6 gene is a major risk factor for psoriasis and is present in people of all races and ethnicities.¹² The global prevalence of the HLA-Cw6 allele in patients with psoriasis ranges from 10.5-77.2%, with significant variation across different ethnic

Table 2. HLA-Cw6 allele genotypes (n=151).

HLA-Cw6	Total n (%)	Study group		p	OR (95% CI)
		Psoriasis n (%)	Control n (%)		
HLA-Cw6, n	151	121	30	0.0001	
Positive	86 (57.0)	78(64.5)	8 (26.7)		4.98 (2.04-12.15)
Negative	65 (43.0)	43 (35.5)	22 (73.3)		
HLA-Cw6 genotypes, n	151	121	30	0.001	-
TA	66 (43.7)	61 (50.4)	5 (16.7)		
TT	20 (13.3)	17 (14.1)	3 (10.0)		
AA	65 (43.0)	43 (35.5)	22 (73.3)		

OR, odds ratio; CI, confidence interval; TA, heterogeneous; TT, homogeneous; AA, negative.

Table 3. HLA-Cw6 and age of onset (n=121).

	N	Age of onset, mean ± SD	p
HLA-Cw6			
Positive	78	26.1±10.9	0.002
Negative	43	31.9±10.3	
HLA-Cw6 genotypes			
TA	61	26.4±10.6	0.01
TT	17	24±12.1	
AA	43	31.9±10.3	

SD, standard deviation; TA, heterogeneous; TT, homogeneous; AA, negative.

Table 4. HLA-Cw6 genotypes and clinical characteristics of the psoriasis group (n=121).

Gen HLA-Cw6	AA n (%)	TA n (%)	TT n (%)	p
Body surface area (BSA)	N=43	N=61	N=17	
Mild (BSA <10)	7 (100.0)	0 (0.0)	0 (0.0)	0.01
Moderate (10≤BSA ≤30)	18 (33.9)	25 (47.2)	10 (18.9)	
Severe (BSA >30)	18 (29.5)	36 (59.0)	7 (11.5)	
Psoriasis area and severity index (PASI)				
Mild (PASI <10)	11 (73.3)	3 (20.0)	1 (6.7)	0.03
Moderate (10≤PASI <20)	21 (30.4)	38 (55.1)	10 (14.5)	
Severe (PASI ≥20)	11 (29.7)	20 (54.1)	6 (16.2)	
Dermatology life quality index (effect on patient's life)				
No effect	0 (0.0)	1 (100.0)	0 (0.0)	0.28
Small effect	10 (55.6)	5 (27.8)	3 (16.6)	
Moderate effect	11 (39.3)	12 (42.9)	5 (17.8)	
Very large effect	15 (33.3)	26 (57.8)	4 (8.9)	
Extremely large effect	7 (24.1)	17 (58.6)	5 (17.3)	

AA, negative; TA, heterogeneous; TT, homogeneous.

groups.³ Our study discovered that the overall frequency of HLA-Cw6 was 57%. A review by Chen *et al.* showed that the prevalence of HLA-Cw6 in patients with psoriasis is higher in Caucasians than in Asians.³ However, the prevalence varies among countries and ethnicities. The prevalence in our study of Vietnamese patients was 64.5%, which is higher than that reported in certain European studies, such as Croatia (59.3%),¹³ Spain (39.0-49.4%),^{14,15} and Iceland (64.2%).¹⁶ A few Asian countries have a high prevalence of HLA-Cw6, such as South Korea (76.1%),⁸ and India (67.6-71.1%),¹⁷⁻¹⁹ which is higher than that in our study. Meanwhile, our study is more prevalent than in other Asian countries, such as China (32.5-50.8%),^{4,5} Japan (10.2-12.0%),^{6,7} and Taiwan (12.1-32.8%).²⁰⁻²² These findings suggest that genetic characteristics may vary among different populations, and not only due to geographical location. Studies on the genetic characteristics of Vietnamese people have contributed to global gene mapping. Our study revealed that the proportion of patients with psoriasis was significantly higher (64.5%) than that of the controls (26.7%) ($P=0.0001$). This result correlates with those of studies conducted in other countries around the world. Although the prevalence of HLA-Cw6 positivity varies among different races and nationalities,³ it is clear that the prevalence in patients with psoriasis is always significantly higher than that in controls. This finding indicated that individuals with psoriasis were more likely to carry the HLA-Cw6 allele than those without psoriasis in Vietnamese people. The odds ratio of having psoriasis for people with HLA-Cw6 was 4.98. This indicates that people with HLA-Cw6 were 4.98 times more likely to develop psoriasis than those without HLA-Cw6. Our study confirms that HLA-Cw6 is a genetic factor associated with psoriasis in the Vietnamese population. Individuals with HLA-Cw6 have a significantly higher risk of developing psoriasis than those without HLA-Cw6. Therefore, HLA-Cw6 may play a role in the development of psoriasis. Our study also revealed that the most common genotype in patients with psoriasis was TA (50.4%), followed by TT (14.1%), and AA (35.5%). This finding shows that the TA genotype is the most common in patients with psoriasis. In contrast, the AA genotype was the most common in the control group (73.3%). Genotypes TA and TT were associated with a higher risk of the disease, whereas genotype AA was associated with a lower risk. The clear difference between the two groups suggests that HLA-Cw6 polymorphisms are associated with the risk of psoriasis.

Our study revealed that individuals with HLA-Cw6 had a significantly earlier onset of psoriasis than those without the gene. The results demonstrated a significantly lower mean age of onset of psoriasis in patients with positive HLA-Cw6 (26.1 ± 10.9 years) than those with negative HLA-Cw6 (31.9 ± 10.3 years). Furthermore, our study is consistent with the findings of Queiro *et al.*, who noted a significant difference in the average age of onset of psoriasis between people with the HLA-Cw6 gene (23 ± 12 years) and those without the gene (32 ± 12 years).⁹ Previous studies have shown that HLA-Cw6 is associated with early-onset psoriasis. Additionally, a study by Henseler *et al.* suggested that the prevalence of HLA-Cw6 positivity was 85.3% in patients with early-onset psoriasis.²³ Enerbäck *et al.* observed that the frequency of HLA-Cw6 was significantly higher in patients with psoriasis that developed before age 21 than in patients with psoriasis that developed at age ≥ 21 .¹⁰ Our findings suggest that individuals with the HLA-Cw6 allele are more likely to develop psoriasis at an earlier age. The difference in the age at onset was significant ($p=0.002$). The TT genotype showed the earliest onset, at 24 ± 12.1 years, while the TA and AA genotypes had a later onset, at 27 ± 10.3 years and 30 ± 9.5 years, respectively. This difference in the age at onset was significant ($p=0.01$). Additionally, our study

indicates that HLA-Cw6 may affect the age of psoriasis onset. We observed that patients with psoriasis in Vietnam with HLA-Cw6 positivity had a higher risk of early onset psoriasis, with the onset age possibly <27 years of age. This is an important finding because it may help doctors identify individuals who are at an increased risk of developing psoriasis at an early age.

Recent studies have suggested a possible association between the HLA-Cw6 allele and the severity of psoriasis. This suggests that people who carry the HLA-Cw6 allele may be more likely to experience severe psoriasis, with more extensive plaques on their arms, legs, and trunk.³ The clinical phenotype and severity of the disease may be associated with certain genetic factors, which may be predicted in advance.¹¹ The study results indicate that the AA phenotype was more common in mild cases of psoriasis, while the TA phenotype was more common in moderate and severe cases. This difference was significant for both BSA ($P=0.01$) and PASI classifications ($p=0.03$). The AA phenotype is the most common genotype in the general population (73.3%), however, it is less common in people with psoriasis (35.5%). This suggests that the AA genotype is associated with milder cases of psoriasis. In contrast, the TA phenotype was less common in the general population (19.3%) but more common in patients with psoriasis (50.4%). This suggests that the TA genotype may be associated with severe psoriasis. Our findings insinuate that the HLA-Cw6 genotype may contribute to the severity of psoriasis. However, further studies are required to confirm these findings and understand the exact mechanism of this association.

Limitations and recommendations

This study has a few limitations. However, these limitations do not detract from the significance of our findings. The study was conducted at a single center, however, the sample size was large enough to be considered significant. This study did not assess environmental factors, which is a common limitation in genetic studies. Overall, this study provides valuable insights into the genetic basis of psoriasis in the Vietnamese population. Future studies are needed to confirm these findings and identify other genetic and environmental factors that may contribute to the development of psoriasis.

Conclusions

In conclusion, the findings of this study suggest that HLA-Cw6 is a major genetic risk factor for psoriasis in the Vietnamese population. This is supported by the higher prevalence of HLA-Cw6 in patients with early-onset disease, and the association between different HLA-Cw6 genotypes and disease severity.

References

1. Armstrong AW, Mehta MD, Schupp CW, et al. Psoriasis prevalence in adults in the United States. *JAMA Dermatol* 2021;157:940-6.
2. Rapp SR, Feldman SR, Exum ML, et al. Psoriasis causes as much disability as other major medical diseases. *J Am Acad Dermatol* 1999;41:401-7.
3. Chen L, Tsai TF. HLA-Cw6 and psoriasis. *Br J Dermatol* 2018;178:854-62.
4. Zhang XJ, Zhang AP, Yang S, et al. Association of HLA class I alleles with psoriasis vulgaris in southeastern Chinese Hans.

- J Dermatol Sci 2003;33:1-6.
5. Fan X, Yang S, Sun LD, et al. Comparison of clinical features of HLA-Cw*0602-positive and -negative psoriasis patients in a Han Chinese population. *Acta Derm Venereol* 2007;87:335-40.
 6. Asahina A, Akazaki S, Nakagawa H, et al. Specific nucleotide sequence of HLA-C is strongly associated with psoriasis vulgaris. *J Invest Dermatol* 1990;97:254-58.
 7. Nakagawa H, Asahina A, Akazaki S, et al. Association of Cw11 in Japanese patients with psoriasis vulgaris. *Tissue Antigens* 1990;36:241-2.
 8. Kim TG, Lee HJ, Youn JI, et al. The association of psoriasis with human leukocyte antigens in the Korean population and the influence of age of onset and sex. *J Invest Dermatol* 2000;114:309-13.
 9. Queiro R, Torre JC, González S, et al. HLA antigens may influence the age of onset of psoriasis and psoriatic arthritis. *J Rheumatol* 2003;30:505-7.
 10. Enerbäck C, Martinsson T, Inerot A, et al. Evidence that HLA-Cw6 determines early onset of psoriasis, obtained using sequence-specific primers (PCR-SSP). *Acta Derm Venereol* 1997;77:273-6.
 11. Christensen TE, Callis KP, Papenfuss J, et al. Observations of psoriasis in the absence of therapeutic intervention identifies two unappreciated morphologic variants, thin-plaque and thick-plaque psoriasis, and their associated phenotypes. *J Invest Dermatol* 2006;126:2397-403.
 12. Nair RP, Stuart PE, Nistor I, et al. Sequence and haplotype analysis supports HLA-C as the psoriasis susceptibility 1 gene. *Am J Hum Genet* 2006;78:827-51.
 13. Kastelan M, Gruber F, Cecuk E, et al. Analysis of HLA antigens in Croatian patients with psoriasis. *Acta Derm Venereol Suppl (Stockh)* 2000;211:12-3.
 14. Gonzalez S, Martinez-Borra J, Torre-Alonso JC, et al. The MICA-A9 triplet repeat polymorphism in the transmembrane region confers additional susceptibility to the development of psoriatic arthritis and is independent of the association of Cw*0602 in psoriasis. *Arthritis Rheum* 1999;42:1010-6.
 15. Martinez-Borra J, Brautbar C, Gonzalez S, et al. The region of 150 kb telomeric to HLA-C is associated with psoriasis in the Jewish population. *J Invest Dermatol* 2005;125:928-32.
 16. Gudjonsson JE, Karason A, Antonsdottir A, et al. Psoriasis patients who are homozygous for the HLA-Cw*0602 allele have a 2.5-fold increased risk of developing psoriasis compared with Cw6 heterozygotes. *Br J Dermatol* 2003;148:233-5.
 17. Rani R, Narayan R, Fernandez-Vina MA, et al. Role of HLA-B and C alleles in development of psoriasis in patients from North India. *Tissue Antigens* 1998;51:618-22.
 18. Chandra A, Lahiri A, Senapati S, et al. Increased risk of psoriasis due to combined effect of HLA-Cw6 and LCE3 risk alleles in Indian population. *Sci Rep* 2016;6:24059.
 19. Indhumathi S, Rajappa M, Chandrashekar L, et al. The HLA-C*06 allele as a possible genetic predisposing factor to psoriasis in South Indian Tamils. *Arch Dermatol Res* 2016;308:193-9.
 20. Tsai TF, Hu CY, Tsai WL, et al. HLA-Cw6 specificity and polymorphic residues are associated with susceptibility among Chinese psoriatics in Taiwan. *Arch Dermatol Res* 2002;294:214-20.
 21. Chang YT, Tsai SF, Lee DD, et al. A study of candidate genes for psoriasis near HLAC in Chinese patients with psoriasis. *Br J Dermatol* 2003;148:418-23.
 22. Chang YT, Tsai SF, Lin MW, et al. SPR1 gene near HLA-C is unlikely to be a psoriasis susceptibility gene. *Exp Dermatol* 2003;12:307-14.
 23. Henseler T, Christophers E. Psoriasis of early and late onset: characterization of two types of psoriasis vulgaris. *J Am Acad Dermatol* 1985;13:450-6.