

RESEARCH ARTICLE

Bottom-up and top-down effects on phytoplankton communities in two freshwater lakes

Yanran Li¹, Jiao Meng¹, Chao Zhang¹, Shuping Ji¹, Qiang Kong², Renqing Wang³, Jian Liu^{1*}

1 Environment Research Institute, Shandong University, Qingdao, China, **2** College of Geography and Environment, Shandong Normal University, Jinan, China, **3** School of Life Sciences, Shandong University, Qingdao, China

* ecology@sdu.edu.cn



OPEN ACCESS

Citation: Li Y, Meng J, Zhang C, Ji S, Kong Q, Wang R, et al. (2020) Bottom-up and top-down effects on phytoplankton communities in two freshwater lakes. *PLoS ONE* 15(4): e0231357. <https://doi.org/10.1371/journal.pone.0231357>

Editor: Xiao Guo, Qingdao Agricultural University, CHINA

Received: December 24, 2019

Accepted: March 21, 2020

Published: April 9, 2020

Copyright: © 2020 Li et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: JL is supported by the China Major Science and Technology Program for Water Pollution Control and Treatment under contract No.2017ZX07101003 and the Natural Science Foundation of Shandong Province, China under contract No. ZR2017MC013. The funders have no role in study design, data collection and analysis, decision to publish or preparation of manuscript.

Abstract

The relative importance of bottom-up versus top-down effects in aquatic ecosystems remains a longstanding and ongoing controversy. To investigate these effects on phytoplankton communities in freshwater lakes, phytoplankton and zooplankton were sampled, and physical-chemical variables were measured during spring and summer in two important freshwater lakes in northern China: Nansi Lake and Dongping Lake. The redundancy analysis results showed that phytoplankton density and biomass were regulated by physical-chemical variables (bottom-up effects) and predation (top-down effects) together, and the former was more prominent in both lakes. However, the correlation analysis indicated that the top-down effects of zooplankton on phytoplankton were not significant in spring and summer in both lakes, while the bottom-up regulation of physical-chemical variables on phytoplankton had different patterns in the two lakes. In Nansi Lake, the bottom-up effects of physical-chemical variables on phytoplankton were weaker in summer than that in spring due to the abundant nutrients in summer. In Dongping Lake, the bottom-up effects of physical-chemical on phytoplankton were significant both in spring and summer, and the dominant bottom-up control factor shifted from total nitrogen in spring to total phosphorus in summer, with an increased ratio of nitrogen to phosphorus due to changes in limiting factors. In the two studied lakes, with fish culture, the bottom-up effects of phytoplankton on zooplankton were more important than the top-down effects of zooplankton on phytoplankton. These results demonstrate the interactions between phytoplankton and zooplankton and highlight the importance of phytoplankton regulation in freshwater lakes, which has implications for the effective management of freshwater lake ecosystems.

Introduction

Phytoplankton and zooplankton not only play important roles in aquatic ecosystems but also serve as key indicators for water quality assessment [1–4]. Phytoplankton is the primary

Competing interests: The authors have declared that no competing interests exist.

producer in lake ecosystems, producing oxygen and organic matter through photosynthesis. Zooplankton is a predator of phytoplankton, connecting primary producers with more advanced consumers in the biological chain and providing a crucial link in the aquatic food web. Seasonal changes in plankton communities have been found [5], which are performed not only as changes in species number, density, biomass and diversity but also as seasonal changes in community structure [6–8]. Moreover, the dominant species of phytoplankton may also change between different seasons within a year [9]. These changes are mainly influenced by physical-chemical (bottom-up effects) and predation (top-down effects) through the aquatic food web [10,11].

The bottom-up effect means that a lower trophic level in the biological network affects the community structure of higher trophic levels by means of resource restriction [12]. The top-down effect refers to a higher trophic level influences the community structure of a lower trophic level through predation [10]. McQueen et al. [13] proposed that the bottom-up effect is strongest at the bottom of the food web and weakens further up the trophic levels, while the top-down effect is strongest at the top of the food web and weakens further down the trophic levels. Some studies have indicated that nutrients influence the density and species composition of phytoplankton through bottom-up effects while predation by zooplankton (top-down effects) controlled the size, distribution, and abundance of phytoplankton [14–16]. Zhang et al. [17] and Song et al. [18] found that phytoplankton was more influenced by the bottom-up effects of environmental factors, while Severiano et al. [19] indicated that the top-down effects of zooplankton had a more significant influence on phytoplankton. Experimental studies have suggested that phytoplankton is controlled more by the bottom-up effects of nutrients than the top-down effects of zooplankton when zooplankton are under strong predation pressure from fish [20,21]. However, few studies have addressed the seasonal dynamics of bottom-up and top-down effects, which have an important influence on phytoplankton succession in freshwater [22]. It is necessary to explore the bottom-up and top-down effects in phytoplankton regulation and the seasonal dynamics of these influences, as the dynamics and structure of phytoplankton communities play an important role in aquatic ecosystems. Understanding the influence of bottom-up and top-down effects in freshwater lakes would provide meaningful evidence for better management.

Nansi Lake and Dongping Lake are two major freshwater lakes in Shandong Province in northern China that are important diversion lakes on the eastern route of the South-to-North Water Diversion Project. In preparation for the water diversion project, water quality and biodiversity of both lakes were restored while they were used for economic development, such as aquaculture. Therefore, these two lakes are suitable for studying the different responses of phytoplankton and zooplankton in lakes in which water quality has been improving in recent years. Based on the measurement of physical-chemical variables, phytoplankton and zooplankton in Nansi Lake and Dongping Lake, we tested the following hypotheses: (i) the bottom-up effects of physical-chemical have more significant regulatory effects on phytoplankton than the top-down effects of zooplankton, and (ii) different patterns in the seasonal regulation of bottom-up and top-down effects exist in different lakes.

Materials and methods

Ethics statement

No specific permissions were required to collect the samples in our study. We confirm that the field studies did not involve endangered or protected species.

Study area and sampling sites

Nansi Lake (116° 34' E– 117° 21' E, 34° 27' N– 35° 20' N) is the largest freshwater lake in northern China, with a total area of 1266 km². The lake consists of four connected lakes, which are Nanyang, Dushan, Zhaoyang and Weishan. There are about fifty-three inflowing rivers and three outflowing rivers. Nansi Lake plays a vital role in the east route of the South-to-North Water Diversion Project, which is the largest project to solve the demand for water resources in northern China. With the warm-temperate monsoon climate, Nansi Lake has an annual temperature of 13.7°C and a total capacity of 6.37×10⁹ m³. Dongping Lake (116° 00' E– 116° 30' E, 35° 30' N– 36° 20' N) is located in Tai'an in the middle of Shandong Province, China. The total area of Dongping Lake is 632 km², and the average water depth of the lake is 2.5 m. Dongping Lake is the catchment for the tributaries of Dahan River in the Yellow River Basin. It is also one of the storage lakes on the east route of the South-to-North Water Diversion Project, and it also has a warm-temperate monsoon climate. Well distributed sampling sites were established across Nansi Lake (n = 10) and Dongping Lake (n = 8) (Fig 1), and samples were collected in spring (April and May) and summer (July and August) from both lakes in 2015.

Sampling methods

Plankton samples were collected at each site with three replications by two different sized plankton nets, with mesh sizes of 112 and 64 μm. The qualitative samples, which were collected by the plankton nets with mesh sizes of 112 μm (macro plankton samples) and 64 μm (micro plankton samples), were classified under a light microscope [23,24]. The quantitative samples were collected using different approaches. Micro plankton samples, which were collected with a 1 L plankton sampler, were kept in brown bottles and fixed with Lugol's iodine solution for microscopic enumeration. Macro plankton samples, which were collected with 40 L water samples and concentrated to 30 mL through a plankton net with a mesh size of 112 μm, were fixed with formaldehyde solution for microscopic enumeration. The biomass and density of plankton were calculated following the Handbook of fishery natural resource investigation in the inside water area [25]. The biomass calculations, which used the specific gravity and body length-weight regression equation, were converted, taking sample volume into account.




Fig 1. Location and sampling sites of Nansi Lake and Dongping Lake, China. (Software of ArcGIS 10.2 and Adobe Photoshop CS6 were used in drawing the figure. The outline of study area and two lakes was drawn by using ArcGIS (version 10.2) and referring to the map from <http://www.dsac.cn/>).

<https://doi.org/10.1371/journal.pone.0231357.g001>

Water samples were collected along with biodiversity samples *in situ* at a depth of 0.5 m using a 1 L Ruttner water sampler. There were three replications for each sample and the samples were carried back to the laboratory as soon as possible under low-temperature conditions for total nitrogen (TN) and total phosphorus (TP) measurement using a 5B-3B(V8) multi-parameter water quality meter and LH-3BNT total nitrogen analyzer. Water temperature (WT) and chlorophyll a were measured in the field using a thermometer and a portable chlorophyll a meter. Dissolved oxygen, pH and water transparency were also measured in the field using a portable dissolved oxygen meter, a pH meter and a Secchi disk. Chemical oxygen demand (COD_{Cr}) was measured in the laboratory using 5B-1 COD rapid monitor, and chemiluminescence detection of permanganate index (COD_{Mn}) and NH₃-N were measured using acidic potassium permanganate method and spectrophotometry.

Statistical analysis

Redundancy analysis (RDA) was used to test the variables with a significant top-down or bottom-up influence on phytoplankton in Nansi Lake and Dongping Lake as the detrended correspondence analysis (DCA) revealed that the gradient length of the response data was less than three. In addition, Pearson correlations among phytoplankton density, phytoplankton biomass, different physical-chemical variables, zooplankton (especially Crustacea) density, and zooplankton biomass were performed comparing spring and summer periods. The RDA highlighted the influence of water temperature, TN, and TP on phytoplankton. Therefore, correlation analysis among these variables was performed for further study. To explore the top-down effects, correlation analysis between phytoplankton and zooplankton was performed across the seasonal factor and biotic indices of density and biomass. Four different patterns were assessed: the same indices for phytoplankton and zooplankton in the same season, the same indices for phytoplankton and zooplankton in different seasons, different indices for phytoplankton and zooplankton in the same season, and different indices for phytoplankton and zooplankton in different seasons. SPSS 22.0 and CANOCO for windows were used to implement the aforementioned analyses.

Results

Abiotic and biological variables in Nansi and Dongping lakes

In 2015, the average water temperature in summer increased by about 12°C compared with that in spring (Table 1), with decreased transparency and dissolved oxygen compared with that in spring. Water quality deteriorated in summer in both Nansi and Dongping lakes, with

Table 1. Physical-chemical variables in Nansi Lake and Dongping Lake in 2015 in different seasons.

Variable	Nansi Lake		Dongping Lake	
	Spring	Summer	Spring	Summer
Water temperature (°C)	19.9±1.06	30.86±1.52	17.39±2.95	30.98±0.76
pH	8.29±0.91	8.16±0.45	8.54±0.65	8.45±0.50
Water transparency (m)	0.54±0.33	0.19±0.13	0.58±0.36	0.36±0.18
Dissolved oxygen (mg·L ⁻¹)	10.57±1.69	5.96±1.97	9.81±1.59	8.35±1.74
Chlorophyll a (mg·L ⁻¹)	12.23±6.52	23.63±12.25	9.97±6.85	27.81±26.43
Total phosphorus (mg·L ⁻¹)	0.04±0.02	0.10±0.04	0.18±0.21	0.14±0.08
Permanganate index (mg·L ⁻¹)	5.17±1.20	5.59±1.33	4.89±0.75	5.67±1.01
Chemical oxygen demand (mg·L ⁻¹)	20.25±9.89	32.51±7.77	34.61±21.07	28.03±7.10
NH ₃ -N (mg·L ⁻¹)	0.75±0.20	0.36±0.28	0.48±0.27	0.20±0.08
Total nitrogen (mg·L ⁻¹)	1.99±1.86	2.81±1.82	0.93±0.38	1.35±0.50

<https://doi.org/10.1371/journal.pone.0231357.t001>

changes in different indices. The concentrations of chlorophyll a, TP, TN, COD_{Mn}, and COD_{Cr} increased in Nansi Lake in summer, while the concentrations of chlorophyll a, TN, and COD_{Mn} increased in Dongping Lake in summer. The density and biomass of phytoplankton and the density of zooplankton were higher in summer than that in spring in both lakes. However, the biomass of zooplankton increased in summer in Dongping Lake, whereas, the biomass of zooplankton showed a higher value in spring in Nansi Lake. The dominant algae in the two lakes were Bacillariophyta with seasonal changes in their composition in spring and summer (Fig 2). The sub-dominant algae in Nansi Lake changed from Euglenophyta in spring to Cyanophyta in summer. In Dongping Lake, the sub-dominant algae changed from Chlorophyta in spring to Cyanophyta in summer. For zooplankton, Crustacea was dominant in Nansi Lake in both spring and summer, while Protozoa was the dominant in Dongping Lake.

The response of phytoplankton variables to bottom-up and top-down effects

The RDA (Fig 3A) demonstrated the ranking of phytoplankton density, zooplankton density, and physical-chemical factors in Nansi Lake, of which the first two axes accounted for 69.72%. Euglenophyta density and Cyanophyta density showed a positive correlation with TP. In addition, Euglenophyta density was positively correlated with water temperature, while Pyrrophyta density was positively influenced by Crustacea density. There were no significant correlations between the density of all phyla of phytoplankton in Nansi Lake and the densities of zooplankton and Crustacea. For biomass, the RDA ranking of phytoplankton biomass, zooplankton biomass, and physical-chemical factors in Nansi Lake showed that the first two axes accounted for 58.24% of the total variation (Fig 3B). The biomass of phyla of phytoplankton was mainly related to physical-chemical variables, and the Cyanophyta biomass presented a significant positive relationship with water temperature. The biomass of Pyrrophyta was significantly affected by the biomass of zooplankton and Crustacea, indicating the predation effects on Pyrrophyta in Nansi Lake.

The RDA results for the phytoplankton variables and physical-chemical factors in Dongping Lake showed that the first two axes accounted for 63.82% of the total variation in phytoplankton density (Fig 4A) and 64.32% of the total variation in phytoplankton biomass (Fig 4B). The density and biomass of phytoplankton were mainly related to water variables that differed with phylum and physical-chemical factors. There was no significant correlation between phytoplankton and zooplankton in Dongping Lake, suggesting that weak top-down effects of zooplankton and Crustacea existed.

Seasonal patterns of bottom-up and top-down effects

The relationships between water quality variables and phytoplankton density and biomass differed between spring and summer in Nansi and Dongping lakes (Tables 2 and 3). In Nansi Lake in spring, the densities of Euglenophyta and Bacillariophyta were positively related to TP, and the biomass of Euglenophyta was positively influenced by water temperature and TN. The phytoplankton showed no significant relationship with water temperature, TP, or TN in summer in Nansi Lake. However, there were different patterns in the bottom-up effects of physical-chemical on phytoplankton in Dongping Lake during seasonal changes. Most of the phytoplankton, except Cyanophyta, Pyrrophyta, and Chlorophyta, were significantly influenced by water temperature and TN in spring, and most of the phytoplankton, except Pyrrophyta and Xanthophyta, were influenced by TP in summer in Dongping Lake. These seasonal changes showed the strong bottom-up effects of physical-chemical on phytoplankton in Dongping Lake in both spring and summer.




Fig 2. Percentage composition of phytoplankton (a) and zooplankton (b). Phytoplankton taxa are Chlorophyta (Chl), Bacillariophyta (Bac), Cryptophyta (Cry), Xanthophyta (Xan), Pyrrophyta (Pyr), Euglenophyta (Eug), Cyanophyta (Cya); zooplankton taxa are Crustacea (Cru), Rotifera (Rot), Protozoa (Pro).

<https://doi.org/10.1371/journal.pone.0231357.g002>

The top-down effects of zooplankton on phytoplankton were unclear in both lakes. Phytoplankton were positively correlated with zooplankton in Nansi Lake, indicating possible bottom-up effects of phytoplankton on zooplankton (Table 4). However, the top-down effects of zooplankton on phytoplankton were not significant in the two lakes. In Nansi Lake, Cyanophyta strongly regulated zooplankton in spring, and Cryptophyta did so in summer. During the seasonal changes, the density of Bacillariophyta in spring was positively correlated with zooplankton in summer. In Dongping Lake, the regulation of phytoplankton on zooplankton mainly occurred between Pyrrophyta and zooplankton (Table 5). In particular, the analyses indicated a significant relationship between Pyrrophyta and Crustacea. Most phytoplankton taxa variables in spring had a positive influence on the total density of zooplankton in summer. The results showed that the biomass accumulation of zooplankton and Crustacea mainly depended on phytoplankton to provide rich sources of food in spring, while the predation of zooplankton and Crustacea had weak top-down control on phytoplankton.

Discussion

The response of phytoplankton to bottom-up and top-down effects

Our study showed that the bottom-up effects of physical-chemical variables on phytoplankton were stronger than the top-down effects of zooplankton in both Nansi and Dongping lakes, which is consistent with previous studies [17,18,26]. The growth and reproduction of phytoplankton were influenced by physical-chemical factors in the water [27,28], among which water temperature and nutrient concentration contributed more than the other factors. The rate of photosynthesis of phytoplankton is probably promoted by the rising water temperature, which promotes the accumulation of biomass. In addition, seasonal shifts provide suitable living conditions for phytoplankton, such as promoting the absorption of nutrients at night by phytoplankton due to the increased daily minimum temperature in summer [29,30]. Nitrogen




Fig 3. Redundancy analysis (RDA) plots for different phytoplankton phyla, zooplankton and water physicochemical parameters in Nansi Lake. ((a) density; (b) biomass).

<https://doi.org/10.1371/journal.pone.0231357.g003>

and phosphorus are essential nutrients for the growth of phytoplankton, and at a particular concentration, they can also limit the growth of phytoplankton [31,32]. In an applicable concentration range of nitrogen and phosphorus, increasing concentrations can promote phytoplankton growth [33,34]. Therefore, nutrients factors in water bodies can effectively control the density and biomass of phytoplankton through bottom-up effects.

Zooplankton, especially Crustacea, can directly affect the density and biomass of phytoplankton through predation, while zooplankton can also be influenced by fish predation. Previous studies have indicated that zooplankton are mainly controlled by top-down effects [35],




Fig 4. Redundancy analysis (RDA) plots for different phytoplankton phyla, zooplankton and water physicochemical parameters in Dongping Lake. ((a) density; (b) biomass).

<https://doi.org/10.1371/journal.pone.0231357.g004>

and they deliver top-down effects on phytoplankton through food chains. The predation pressure of fish on zooplankton changes the biomass and density of zooplankton, which further influences the top-down effects of zooplankton on phytoplankton. Fish that feed on zooplankton prefer large Cladocera, followed by Copepods, which have a strong ability to escape, and finally, the smaller species of Rotifera [36,37]. In lakes with aquaculture, the community structure of Crustacea is mainly influenced by predation by fish, which is not significant enough to affect the biomass of phytoplankton [38]. Cage aquaculture in Nansi and Dongping lakes

Table 2. The Pearson correlations between water temperature, total phosphorus, total nitrogen and phytoplankton of Nansi Lake in different seasons.

	Spring			Summer		
	WT	TN	TP	WT	TN	TP
Density _{TPh}	-0.499	-0.286	-0.233	0.541	-0.183	0.222
Density _{Cya}	0.325	-0.406	0.313	0.6	-0.149	0.294
Density _{Eug}	0.246	0.477	0.869**	0.606	-0.128	0.499
Density _{Pyr}	0.474	0.249	0.07	0.331	-0.487	-0.489
Density _{Xan}	-0.147	-0.125	-0.493	0.278	0.231	0.027
Density _{Cry}	-0.403	-0.172	-0.11	0.437	-0.238	-0.001
Density _{Bac}	-0.145	-0.353	0.746*	0.443	-0.168	0.158
Density _{Chl}	-0.46	-0.278	0.098	0.568	-0.222	0.291
Biomass _{TPh}	0.399	0.23	-0.169	0.503	-0.175	0.201
Biomass _{Cya}	0.342	-0.391	0.323	0.6	-0.149	0.294
Biomass _{Eug}	0.815**	0.837**	0.622	0.606	-0.128	0.499
Biomass _{Pyr}	0.472	0.25	0.066	0.19	-0.331	-0.303
Biomass _{Xan}	-0.147	-0.125	-0.492	0.286	0.226	0.033
Biomass _{Cry}	-0.262	-0.08	0.107	0.412	-0.281	-0.238
Biomass _{Bac}	-0.133	-0.355	-0.573	0.443	-0.168	0.158
Biomass _{Chl}	-0.46	-0.279	0.097	0.568	-0.223	0.291

WT = water temperature; TN = total nitrogen; TP = total phosphorus; TPh = total phytoplankton; Cya = Cyanophyta; Eug = Euglenophyta; Pyr = Pyrrophyta; Xan = Xanthophyta; Cry = Cryptophyta; Bac = Bacillariophyta; Chl = Chlorophyta

*p<0.05

**p<0.01.

<https://doi.org/10.1371/journal.pone.0231357.t002>

Table 3. The Pearson correlations between water temperature, total phosphorus, total nitrogen and phytoplankton of Dongping Lake in different seasons.

	Spring			Summer		
	WT	TN	TP	WT	TN	TP
Density _{TPh}	0.787*	0.559	-0.578	-0.153	0.185	0.972**
Density _{Cya}	0.014	-0.043	-0.289	-0.226	0.231	0.897**
Density _{Eug}	0.807*	0.912**	-0.272	-0.175	0.157	0.993**
Density _{Pyr}	-0.518	-0.191	-0.03	-0.257	-0.22	0.011
Density _{Xan}	0.773*	0.942**	-0.326	--	--	--
Density _{Cry}	0.821*	0.895**	-0.242	-0.171	0.262	0.824*
Density _{Bac}	0.887**	0.39	-0.553	-0.115	0.162	0.939**
Density _{Chl}	0.359	0.395	-0.413	0.033	0.297	0.870**
Biomass _{TPh}	0.850**	0.591	-0.581	-0.147	0.171	0.971**
Biomass _{Cya}	0.013	-0.043	-0.288	-0.226	0.231	0.897**
Biomass _{Eug}	0.807*	0.912**	-0.272	-0.175	0.157	0.993**
Biomass _{Pyr}	-0.56	-0.22	0.028	-0.257	-0.22	0.011
Biomass _{Xan}	0.772*	0.942**	-0.328	--	--	--
Biomass _{Cry}	0.826*	0.891**	-0.243	-0.172	0.262	0.825*
Biomass _{Bac}	0.887**	0.39	-0.553	-0.115	0.162	0.939**
Biomass _{Chl}	0.359	0.395	-0.413	0.033	0.297	0.870**

WT = water temperature; TN = total nitrogen; TP = total phosphorus; TPh = total phytoplankton; Cya = Cyanophyta; Eug = Euglenophyta; Pyr = Pyrrophyta; Xan = Xanthophyta; Cry = Cryptophyta; Bac = Bacillariophyta; Chl = Chlorophyta

*p<0.05

**p<0.01.

<https://doi.org/10.1371/journal.pone.0231357.t003>

Table 4. The Pearson correlations between phytoplankton and zooplankton variables of Nansi Lake in different seasons.

		Spring				Summer			
		Density _{TZ}	Density _{Cru}	Biomass _{TZ}	Biomass _{Cru}	Density _{TZ}	Density _{Cru}	Biomass _{TZ}	Biomass _{Cru}
Spring	Density _{Cya}	0.650	0.805**	0.815**	0.805**	0.757*	0.252	0.261	0.248
	Density _{Bac}	-0.143	0.446	0.426	0.446	0.295	0.789*	0.788*	0.790*
	Biomass _{Cya}	0.660	0.803**	0.814**	0.803**	0.755*	0.242	0.251	0.239
Summer	Density _{Cry}	0.065	0.771*	0.751*	0.771*	0.488	0.730*	0.731*	0.725*
	Biomass _{Cry}	0.099	0.685*	0.670*	0.685*	0.503	0.665	0.667*	0.660

TZ = total zooplankton; Cru = Crustacea; Cya = Cyanophyta; Cry = Cryptophyta; Bac = Bacillariophyta

*p<0.05

**p<0.01; Complete data is attached in Supporting information.

<https://doi.org/10.1371/journal.pone.0231357.t004>

increased the feeding pressure on zooplankton due to the increasing fish stock, resulting in less zooplankton, especially Crustacea. However, the reduced zooplankton density in our study did not show measurable top-down effects of zooplankton on phytoplankton. Although some studies have shown that phytoplankton were under the top-down regulation of zooplankton, these studies were normally conducted in eutrophic ecosystems, such as eutrophic reservoirs, lakes, and bays [19,39]. Under field studies on bottom-up and top-down effects, the environmental complexity obscured the effects between phytoplankton and zooplankton with the participation of higher-level predators.

Seasonal patterns of bottom-up and top-down effects

The bottom-up effects of physical-chemical variables were stronger than the top-down effects of zooplankton on phytoplankton in both Nansi and Dongping lakes in spring and summer.

Table 5. The Pearson correlations between phytoplankton and zooplankton variables of Dongping Lake in different seasons.

		Spring				Summer			
		Density _{TZ}	Density _{Cru}	Biomass _{TZ}	Biomass _{Cru}	Density _{TZ}	Density _{Cru}	Biomass _{TZ}	Biomass _{Cru}
Spring	Density _{TPh}	-0.051	-0.371	-0.026	-0.032	0.835**	-0.317	0.347	-0.246
	Density _{Eug}	-0.160	-0.122	-0.206	-0.259	0.853**	-0.160	0.486	-0.086
	Density _{Pyr}	-0.339	0.895**	-0.229	0.351	-0.419	-0.230	-0.468	-0.260
	Density _{Xan}	-0.278	-0.001	-0.228	-0.003	0.728*	-0.260	0.295	-0.224
	Density _{Cry}	-0.095	-0.220	-0.108	-0.171	0.877**	-0.149	0.561	-0.013
	Density _{Bac}	0.322	-0.261	0.336	-0.079	0.812*	-0.032	0.401	-0.147
	Biomass _{TPh}	-0.005	-0.355	-0.010	-0.119	0.886**	-0.261	0.396	-0.224
	Biomass _{Eug}	-0.160	-0.122	-0.206	-0.259	0.853**	-0.160	0.486	-0.086
	Biomass _{Pyr}	-0.331	0.857**	-0.225	0.314	-0.416	-0.256	-0.484	-0.282
	Biomass _{Xan}	-0.280	-0.001	-0.226	0.007	0.725*	-0.263	0.290	-0.227
	Biomass _{Cry}	-0.086	-0.227	-0.103	-0.181	0.885**	-0.147	0.566	-0.013
	Biomass _{Bac}	0.322	-0.262	0.336	-0.079	0.812*	-0.032	0.402	-0.147
Summer	Density _{Pyr}	0.588	-0.366	0.450	-0.371	0.036	0.859**	0.779*	0.961**
	Biomass _{Pyr}	0.450	-0.371	0.588	-0.366	0.035	0.859**	0.779*	0.962**

TZ = total zooplankton; Cru = Crustacea; TPh = total phytoplankton; Eug = Euglenophyta; Pyr = Pyrrophyta; Xan = Xanthophyta; Cry = Cryptophyta; Bac = Bacillariophyta

*p<0.05

**p<0.01; Complete data is attached in Supporting information.

<https://doi.org/10.1371/journal.pone.0231357.t005>

However, the seasonal bottom-up effects of physical-chemical factors and the top-down effects of zooplankton on phytoplankton were different between Nansi and Dongping lakes. The results showed a significant positive correlation between phytoplankton and physical-chemical factors and no significant negative correlation between phytoplankton and zooplankton in either lake. However, there was a strong correlation showing the bottom-up effects of physical-chemical on phytoplankton. In Nansi Lake, the bottom-up effects of physical-chemical on phytoplankton were stronger in spring than in summer, as there was no significant correlation in summer. The density and biomass of phytoplankton in summer increased significantly as nutrient concentrations and water temperature increased. The same conclusions can be drawn from Dongping Lake. Our results implied that the bottom-up effects of nutrients on phytoplankton also changed between seasons. The control of bottom-up effects on phytoplankton altered between nitrogen and phosphorus in Dongping Lake. Chai et al. [40] and Zhou and Liu [41] also found that these nutrient elements changed, which affected phytoplankton between seasons. The primary nutrients, which influenced phytoplankton in Dongping Lake, changed from TN in spring to TP in summer. The growth of phytoplankton needs a suitable N/P ratio [42], and a previous study found that the N/P ratios in Dongping Lake changed from 5.17 to 9.64 [43].

Frau et al. [44] pointed out that the control patterns of bottom-up and top-down effects on phytoplankton were different during different hydrological periods. The top-down effects on phytoplankton were stronger in the dry season, while the bottom-up effects were stronger in the wet season. Neither sunlight nor nutrients are limiting factors of phytoplankton biomass in summer, and it follows that a short-term top-down effect appears in the lake [45]. In Nansi and Dongping lakes, fish culture influenced the density of Crustacea through predation pressure, which led to no measurable top-down control of zooplankton on phytoplankton. However, the bottom-up effects of phytoplankton on zooplankton were significant in spring and summer. The growth of zooplankton biomass can also be affected by the biomass of phytoplankton due to predator-prey relationships [46]. As temperature rises in summer, higher predation pressure of fish leads to a decrease in the density of *Daphnia* [47,48], which in turn results in a decreased intensity of predation on phytoplankton. Our results showed reductions in both Cladocera and Copepoda, leading to weak top-down effects of zooplankton on phytoplankton in summer.

Contrary to expectations, the zooplankton did not significantly decrease the density and biomass of phytoplankton, which corresponds with a previous study [49]. Gliwicz [50] indicated that fish play an important role in the composition and distribution of the zooplankton community, which indirectly influences the top-down effects of zooplankton on phytoplankton. Fish culture blocks the top-down effects of zooplankton on phytoplankton in these kinds of aquatic ecosystems. Von Ruckert and Giani [51] suggested that fish regulate phytoplankton more than zooplankton in certain systems. As the phytoplankton composition changed from spring to summer and there were no significant top-down effects of zooplankton on phytoplankton, we also suggest the plant defense hypothesis, which states that herbivores control the plant species composition rather than plant biomass [49,52,53]. The diversity of phytoplankton possibly made the phytoplankton resilient to predation by zooplankton and fish, with changes in composition but not biomass. The top-down effects of zooplankton on phytoplankton were not significant, but they might contribute to shaping the community composition of phytoplankton [54–56].

Conclusions

Our study showed that the bottom-up effects of physical-chemical on phytoplankton were weaker in summer than in spring in Nansi Lake, and there was a shift in the bottom-up effects of nutrients on phytoplankton in Dongping Lake. The control nutrients of bottom-up effects on phytoplankton altered from TN in spring to TP in summer. While the bottom-up effects of physical-chemical on phytoplankton are clear, the top-down effects of zooplankton on phytoplankton, which may have been regulated indirectly by fish, are more difficult to predict in aquatic management. Conclusions should be drawn cautiously because more data are needed for a thorough analysis of the two effects. Our results can serve as a basis for identifying how phytoplankton are influenced, which have implications for developing sustainable management strategies and conserving services in freshwater lake ecosystems.

Supporting information

S1 Table. The Pearson correlations between phytoplankton and zooplankton variables of Nansi Lake in different seasons.

(DOCX)

S2 Table. The Pearson correlations between phytoplankton and zooplankton variables of Dongping Lake in different seasons.

(DOCX)

Acknowledgments

We thank Xudong Yao for his help in the fieldwork. Thanks to professional editors at Editage for their English language editing.

Author Contributions

Conceptualization: Jiao Meng, Jian Liu.

Data curation: Yanran Li, Jiao Meng, Qiang Kong.

Formal analysis: Yanran Li, Jiao Meng.

Funding acquisition: Jian Liu.

Investigation: Yanran Li, Jiao Meng, Qiang Kong.

Software: Yanran Li, Jiao Meng.

Writing – original draft: Yanran Li, Jian Liu.

Writing – review & editing: Yanran Li, Chao Zhang, Shuping Ji, Renqing Wang, Jian Liu.

References

1. Cadotte MW, Carscadden K, Mirotchnick N. Beyond species: functional diversity and the maintenance of ecological processes and services. *J Appl Ecol.* 2011; 48: 1079–1087. <https://doi.org/10.1111/j.1365-2664.2011.02048.x>
2. Jiang Y, He W, Liu W, Qin N, Ouyang H, Wang Q, et al. The seasonal and spatial variations of phytoplankton community and their correlation with environmental factors in a large eutrophic Chinese lake (Lake Chaohu). *Ecol Indic.* 2014; 40: 58–67. <https://doi.org/10.1016/j.ecolind.2014.01.006>
3. Stevenson J. Ecological assessments with algae: a review and synthesis. Graham L, editor. *J Phycol.* 2014; 50: 437–461. <https://doi.org/10.1111/jpy.12189> PMID: 26988318

4. Liu B, Stevenson RJ. Improving assessment accuracy for lake biological condition by classifying lakes with diatom typology, varying metrics and modeling multimetric indices. *Sci Total Environ*. 2017; 609: 263–271. <https://doi.org/10.1016/j.scitotenv.2017.07.152> PMID: 28750229
5. Lv J, Wu H, Chen M. Effects of nitrogen and phosphorus on phytoplankton composition and biomass in 15 subtropical, urban shallow lakes in Wuhan, China. *Limnologica*. 2011; 41: 48–56. <https://doi.org/10.1016/j.limno.2010.03.003>
6. Dooley Á, Isbell F, Kirwan L, Connolly J, Finn JA, Brophy C. Testing the effects of diversity on ecosystem multifunctionality using a multivariate model. Buckley Y, editor. *Ecol Lett*. 2015; 18: 1242–1251. <https://doi.org/10.1111/ele.12504>
7. Vallina SM, Cermenó P, Dutkiewicz S, Loreau M, Montoya JM. Phytoplankton functional diversity increases ecosystem productivity and stability. *Ecol Modell*. 2017; 361: 184–196. <https://doi.org/10.1016/j.ecolmodel.2017.06.020>
8. Ke Z, Xie P, Guo L. Controlling factors of spring–summer phytoplankton succession in Lake Taihu (Meiliang Bay, China). *Hydrobiologia*. 2008; 607: 41–49. <https://doi.org/10.1007/s10750-008-9365-5>
9. Ye R, Qian X, Shan K, Gao H. Temporal distribution patterns of phytoplankton community structure in a large shallow lake, Lake Taihu, China. *Adv Mater Res*. 2014; 955–959: 1363–1367. <https://doi.org/10.4028/www.scientific.net/AMR.955-959.1363>
10. Carpenter SR, Kitchell JF, Hodgson JR. Cascading trophic interactions and lake productivity. *Bioscience*. 1985; 35: 634–639. <https://doi.org/10.2307/1309989>
11. Doi H, Chang KH, Nishibe Y, Imai H, Nakano S. Lack of congruence in species diversity indices and community structures of planktonic groups based on local environmental factors. Convertino M, editor. *PLoS One*. 2013; 8: e69594. <https://doi.org/10.1371/journal.pone.0069594> PMID: 23936054
12. White TCR. The importance of a relative shortage of food in animal ecology. *Oecologia*. 1978; 33: 71–86. <https://doi.org/10.1007/BF00376997> PMID: 28309267
13. McQueen DJ, Post JR, Mills EL. Trophic relationships in freshwater pelagic ecosystems. *Can J Fish Aquat Sci*. 1986; 43: 1571–1581. <https://doi.org/10.1139/f86-195>
14. Carignan R, Neiff JJ. Nutrient dynamics in the floodplain ponds of the Paraná River (Argentina) dominated by the water hyacinth *Eichhornia crassipes*. *Biogeochemistry*. 1992; 17: <https://doi.org/10.1007/BF00002642>
15. Mayora G, Devercelli M, Giri F. Spatial variability of chlorophyll-a and abiotic variables in a river–floodplain system during different hydrological phases. *Hydrobiologia*. 2013; 717: 51–63. <https://doi.org/10.1007/s10750-013-1566-x>
16. Nicolle A, Hansson LA, Brodersen J, Nilsson PA, Brönmark C. Interactions between predation and resources shape zooplankton population dynamics. Perc M, editor. *PLoS One*. 2011; 6: e16534. <https://doi.org/10.1371/journal.pone.0016534> PMID: 21304980
17. Zhang X, Xie P, Chen F, Li S, Qin J. Driving forces shaping phytoplankton assemblages in two subtropical plateau lakes with contrasting trophic status. *Freshw Biol*. 2007; 52: 1463–1475. <https://doi.org/10.1111/j.1365-2427.2007.01776.x>
18. Song H, Zhang X, Wang B, Sun X, Wang X, Xin M. Bottom-up and top-down controls of the phytoplankton standing stock off the Changjiang Estuary. *Acta Oceanol Sin (in Chinese)*. 2014; 36: 91–100. <https://doi.org/10.3969/j>
19. Severiano J dos S, Moura A do N, Magalhães EM de M, Almeida VL dos S. Study about top-down and bottom-up controls in regulating the phytoplankton biomass in a eutrophic reservoir in northeastern Brazil. *J Water Resour Prot*. 2012; 04: 616–621. <https://doi.org/10.4236/jwarp.2012.48071>
20. Sinistro R. Top-down and bottom-up regulation of planktonic communities in a warm temperate wetland. *J Plankton Res*. 2010; 32: 209–220. <https://doi.org/10.1093/plankt/fbp114>
21. Sinistro R, Sánchez ML, Marinone MC, Izaguirre I. Experimental study of the zooplankton impact on the trophic structure of phytoplankton and the microbial assemblages in a temperate wetland (Argentina). *Limnologica*. 2007; 37: 88–99. <https://doi.org/10.1016/j.limno.2006.09.001>
22. Sommer U, Adrian R, De Senerpont Domis L, Elser JJ, Gaedke U, Ibelings B, et al. Beyond the Plankton Ecology Group (PEG) Model: Mechanisms Driving Plankton Succession. *Annu Rev Ecol Evol Syst*. 2012; 43: 429–448. <https://doi.org/10.1146/annurev-ecolsys-110411-160251>
23. Arhonditsis GB, Winder M, Brett MT, Schindler DE. Patterns and mechanisms of phytoplankton variability in Lake Washington (USA). *Water Res*. 2004; 38: 4013–4027. <https://doi.org/10.1016/j.watres.2004.06.030> PMID: 15380991
24. Lansac-Tôha F, Bonecker C, Velho L, Simões N, Dias J, Alves G, et al. Biodiversity of zooplankton communities in the Upper Paraná River floodplain: interannual variation from long-term studies. *Brazilian J Biol*. 2009; 69: 539–549. <https://doi.org/10.1590/S1519-69842009000300009> PMID: 19738961

25. Zhang J, He Z. Handbook of fishery natural resource investigation in inside water area. Beijing, China: Agriculture Press; 1991.
26. Du X, García-Berthou E, Wang Q, Liu J, Zhang T, Li Z. Analyzing the importance of top-down and bottom-up controls in food webs of Chinese lakes through structural equation modeling. *Aquat Ecol*. 2015; 49: 199–210. <https://doi.org/10.1007/s10452-015-9518-3>
27. Xu H, Paerl HW, Qin B, Zhu G, Gao G. Nitrogen and phosphorus inputs control phytoplankton growth in eutrophic Lake Taihu, China. *Limnol Oceanogr*. 2010; 55: 420–432. <https://doi.org/10.4319/lo.2010.55.1.0420>
28. Elliott JA, Jones ID, Thackeray SJ. Testing the sensitivity of phytoplankton communities to changes in water temperature and nutrient load, in a temperate lake. *Hydrobiologia*. 2006; 559: 401–411. <https://doi.org/10.1007/s10750-005-1233-y>
29. Walther GR. Adapted behaviour and shifting ranges of species—a result of recent climate warming? “Fingerprints” of Climate Change. Boston, MA: Springer US; 2001. pp. 1–15. https://doi.org/10.1007/978-1-4419-8692-4_1
30. Straub C, Quillardet P, Vergalli J, de Marsac NT, Humbert JF. A day in the life of microcystis aeruginosa strain pcc 7806 as revealed by a transcriptomic analysis. Schönbach C, editor. *PLoS One*. 2011; 6: e16208. <https://doi.org/10.1371/journal.pone.0016208> PMID: 21283831
31. Elmgren R, Larsson U. Nitrogen and the Baltic Sea: managing nitrogen in relation to phosphorus. *Sci World J*. 2001; 1: 371–377. <https://doi.org/10.1100/tsw.2001.291> PMID: 12805876
32. Smith VH. Eutrophication of freshwater and coastal marine ecosystems a global problem. *Environ Sci Pollut Res*. 2003; 10: 126–139. <https://doi.org/10.1065/espr2002.12.142> PMID: 12729046
33. Conley DJ, Paerl HW, Howarth RW, Boesch DF, Seitzinger SP, Havens KE, et al. Ecology: controlling eutrophication: nitrogen and phosphorus. *Science*. 2009; 323: 1014–1015. <https://doi.org/10.1126/science.1167755> PMID: 19229022
34. Sun X, Dong S, Tang Z. Influences of nutrients and illuminance on phytoplankton community structure. *South China Fish Sci*. 2008; 4: 5–13.
35. Rejas D, Declerck S, Auwerkerken J, Tak P, Meester L De. Plankton dynamics in a tropical floodplain lake: fish, nutrients, and the relative importance of bottom-up and top-down control. *Freshw Biol*. 2005; 50: 52–69. <https://doi.org/10.1111/j.1365-2427.2004.01306.x>
36. Winfield IJ, Peirson G, Cryer M, Townsend CR. The behavioural basis of prey selection by underyearling bream (*Abramis brama* (L.)) and roach (*Rutilus rutilus* (L.)). *Freshw Biol*. 1983; 13: 139–149. <https://doi.org/10.1111/j.1365-2427.1983.tb00666.x>
37. Rajasilta M, Vuorinen I. A field study of prey selection in planktivorous fish larvae. *Oecologia*. 1983; 59: 65–68. <https://doi.org/10.1007/BF00388074> PMID: 25024149
38. Wang S, Xie P, Wu S, Wang H. Crustacean zooplankton size structure in aquaculture lakes: is larger size structure always associated with higher grazing pressure? *Hydrobiologia*. 2007; 575: 203–209. <https://doi.org/10.1007/s10750-006-0394-7>
39. Nagdali SS, Gupta PK. Impact of mass mortality of a mosquito fish, *Gambusia affinis* on the ecology of a fresh water eutrophic lake (Lake Naini Tal, India). *Hydrobiologia*. 2002; 468: 45–52. <https://doi.org/10.1023/A:1015270206187>
40. Chai Y, Peng T, Guo K, He Y, Yang D, Luo J. Community characteristics of phytoplankton in Lake Changhu and relationships with environmental factors in the summer of 2012. *Chinese J Plant Ecol*. 2014; 38: 857–867. <https://doi.org/10.3724/SP.J.1258.2014.00080>
41. Zhou M, Liu Z. Nitrogen and phosphorus control the cyanobacteria of Huizhou West Lake, a tropical shallow lake. *Ecol Sci*. 2012; 31: 115–120. <https://doi.org/10.3969/j.issn>
42. Schanz F, Juon H. Two different methods of evaluating nutrient limitations of periphyton bioassays, using water from the River Rhine and eight of its tributaries. *Hydrobiologia*. 1983; 102: 187–195. <https://doi.org/10.1007/BF00006346>
43. Meng J, Yu Z, Miao M, Kong Q, Zhang Y, Liu J. Differentiated responses of plankton and zoobenthos to water quality based on annual and seasonal analysis in a freshwater lake. *Polish J Environ Stud*. 2017; 26: 755–764. <https://doi.org/10.15244/pjoes/66713>
44. Frau D, Devercelli M, José de Paggi S, Scarabotti P, Mayora G, Battauz Y, et al. Can top-down and bottom-up forces explain phytoplankton structure in a subtropical and shallow groundwater-connected lake? *Mar Freshw Res*. 2015; 66: 1106. <https://doi.org/10.1071/MF14177>
45. Mataloni G, Tesolín G, Sacullo F, Tell G. Factors regulating summer phytoplankton in a highly eutrophic Antarctic lake. *Hydrobiologia*. 2000; 432: 65–72. <https://doi.org/10.1023/A:1004045219437>
46. Li J, Duan C, Yang F. Relationship between phytoplankton and nutrient dynamics in the enclosed water-body, Dianchi Lake. *Adv Mater Res*. 2013; 779–780: 1514–1517. <https://doi.org/10.4028/www.scientific.net/AMR.779-780.1514>

47. Geller W, Müller H. The filtration apparatus of Cladocera: Filter mesh-sizes and their implications on food selectivity. *Oecologia*. 1981; 49: 316–321. <https://doi.org/10.1007/BF00347591> PMID: 28309989
48. Jiang X, Han B, Lin Q. Effect of farming fish density on zooplankton community of reservoirs. *J Hydroecology*. 2013; 34: 30–36. <https://doi.org/10.15928/j.1674-3075.2013.01.002>
49. Sommer U, Sommer F, Santer B, Jamieson C, Boersma M, Becker C, et al. Complementary impact of copepods and cladocerans on phytoplankton. *Ecol Lett*. 2001; 4: 545–550. <https://doi.org/10.1046/j.1461-0248.2001.00263.x>
50. Gliwicz ZM. Between hazards of starvation and risk of predation: the ecology of offshore animals. Kinne O, editor. Oldendorf/Luhe, Germany: International Ecology Institute; 2003.
51. von Ruckert G, Giani A. Biological interactions in the plankton community of a tropical eutrophic reservoir: is the phytoplankton controlled by zooplankton? *J Plankton Res*. 2008; 30: 1157–1168. <https://doi.org/10.1093/plankt/fbn065>
52. Pace ML, Cole JJ, Carpenter SR, Kitchell JF. Trophic cascades revealed in diverse ecosystems. *Trends Ecol Evol*. 1999; 14: 483–488. [https://doi.org/10.1016/s0169-5347\(99\)01723-1](https://doi.org/10.1016/s0169-5347(99)01723-1) PMID: 10542455
53. Power ME. Top-down and bottom-up forces in food webs: do plants have primacy. *Ecology*. 1992; 73: 733–746. <https://doi.org/10.2307/1940153>
54. Denno RF, Gratton C, Döbel H, Finke DL. Predation risk affects relative strength of top-down and bottom-up impacts on insect herbivores. *Ecology*. 2003; 84: 1032–1044. [https://doi.org/10.1890/0012-9658\(2003\)084\[1032:PRARSO\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2003)084[1032:PRARSO]2.0.CO;2)
55. Gripenberg S, Roslin T. Up or down in space? Uniting the bottom-up versus top-down paradigm and spatial ecology. *Oikos*. 2007; 116: 181–188. <https://doi.org/10.1111/j.2006.0030-1299.15266.x>
56. Hunter MD, Price PW. Playing chutes and ladders: heterogeneity and the relative roles of bottom-up and top-down forces in natural communities. *Ecology*. 1992; 73: 724–732. <https://doi.org/10.2307/1940152>