

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2,4-Dinitrobenzaldehyde hydrazone

Nan Zhang,^a Ruji Wang,^b Chunyan Tan,^c Yuyang Jiang^{d*} and Yufen Zhao^a

^aKey Laboratory of Bioorganic Phosphorus Chemistry & Chemical Biology, Ministry of Education, Department of Chemistry, Tsinghua University, Beijing 100084, People's Republic of China, ^bDepartment of Chemistry, Tsinghua University, Beijing 100084, People's Republic of China, ^cKey Laboratory of Chemical Biology, Guangdong Province, Graduate School at Shenzhen, Tsinghua University, Shenzhen 518055, People's Republic of China, and ^dSchool of Medicine, Tsinghua University, Beijing 100084, People's Republic of China

Correspondence e-mail: jiangyy@sz.tsinghua.edu.cn


Received 17 March 2008; accepted 19 March 2008

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.049; wR factor = 0.116; data-to-parameter ratio = 11.9.

The title compound, $\text{C}_7\text{H}_6\text{N}_4\text{O}_4$, plays an important role in the synthesis of biologically active compounds. The planar hydrazone group is oriented at a dihedral angle of $8.27(3)^\circ$ with respect to the benzene ring. In the crystal structure, intermolecular $\text{N}-\text{H}\cdots\text{O}$ and $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds link the molecules.

Related literature

For related literature, see: Allen *et al.* (1987); Chaulk *et al.* (2007); Kawakami *et al.* (2000); Moreno-Mañas *et al.* (2001).



Experimental

Crystal data

$\text{C}_7\text{H}_6\text{N}_4\text{O}_4$
 $M_r = 210.16$

Triclinic, $P\bar{1}$
 $a = 4.5839(7)$ Å

$b = 9.6840(16)$ Å
 $c = 9.9287(15)$ Å
 $\alpha = 90.785(12)^\circ$
 $\beta = 96.149(11)^\circ$
 $\gamma = 98.955(13)^\circ$
 $V = 432.66(12)$ Å³

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.14$ mm⁻¹
 $T = 295(2)$ K
 $0.4 \times 0.3 \times 0.2$ mm

Data collection

Bruker P4 diffractometer
Absorption correction: none
2238 measured reflections
1616 independent reflections
1160 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$
3 standard reflections
every 97 reflections
intensity decay: none

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.116$
 $S = 1.07$
1616 reflections

136 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.16$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.18$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1B}\cdots\text{O1}^{\text{i}}$	0.90	2.52	3.305 (3)	146
$\text{N1}-\text{H1C}\cdots\text{N2}^{\text{ii}}$	0.90	2.34	3.123 (4)	146

Symmetry codes: (i) $-x + 2, -y, -z + 1$; (ii) $-x + 2, -y, -z$.

Data collection: *XSCANS* (Bruker, 1997); cell refinement: *XSCANS*; data reduction: *XSCANS*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The authors acknowledge financial support from the Ministry of Science and Technology of China (2005CCA03400, 2007 A A02Z160), the Chinese National Natural Science Foundation (20572060, 20472043), and the Department of Science and Technology of Guangdong Province (2005 A11601008).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2106).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
Bruker (1997). *XSCANS*. Bruker AXS Inc., Madison, Wisconsin, USA.
Chaulk, S. G. & MacMillan, A. M. (2007). *Nat. Protoc.* **2**, 1052–1058.
Kawakami, T., Uehata, K. & Suzuki, H. (2000). *Org. Lett.* **2**, 413–415.
Moreno-Mañas, M., Pleixats, R., Andreu, R., Garín, J., Orduna, J., Villacampa, B., Levillain, E. & Sallé, M. (2001). *J. Mater. Chem.* **11**, 374–380.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supplementary materials

Acta Cryst. (2008). E64, o745 [doi:10.1107/S1600536808007514]

2,4-Dinitrobenzaldehyde hydrazone

N. Zhang, R. Wang, C. Tan, Y. Jiang and Y. Zhao

Comment

Benzaldehyde hydrazone and its analogues are important intermediates in heterocyclic chemistry, and they have been widely used for the synthesis of biologically active compounds such as [1,2,4]triazino[6,5-*f*]quinolines, pyrazolo[3,4-*f*]quinolines (Kawakami *et al.*, 2000), 1,3-dithiol-2-ylidene derivatives (Moreno-Mañas *et al.*, 2001), and oligo-RNAs with photocaged adenosine 2'-hydroxyls (Chaulk *et al.*, 2007). Here we report the synthesis and crystal structure of a nitro-analogue: 2,4-dinitrobenzaldehyde hydrazone. The molecule of the title compound (Fig. 1) contains a benzene ring, a hydrazone chain and two nitril groups. Most of the bond lengths and angles are within normal ranges (Allen *et al.*, 1987). Because of the pi-pi conjugation and two nitril groups electron withdrawing effect, the distance of C=N bond (1.282 (3) Å) is obviously shorter than that of the normal range (1.34–1.38 Å). The molecule is essentially planar, with a dihedral angle of 8.27° between the hydrazone group and the benzene ring. In the crystal structure, the molecules are linked by intermolecular N—H···O and N—H···N hydrogen bonds (Table 1, Fig. 2), which seem to be effective in the stabilization of the structure.

Experimental

2,4-Dinitrobenzaldehyde (1.96 g, 10 mmol) was dissolved in 100 ml absolute ethanol, after which hydrazine hydrate (0.96 ml, 20 mmol) was added. The mixture was stirred at about 353 K for 5 h. The solution was cooled and kept at about 279 K overnight. Brown powder was collected by filtration (1.41 g, yield 67%) and then single crystals suitable for X-ray measurements were obtained by recrystallization from ethanol.

Refinement

All non-H atoms were refined anisotropically. All H atoms were placed in calculated positions, with N—H = 0.9 Å and C—H = 0.93 Å. Final difference Fourier maps showed the highest and lowest electron densities of 0.160 and -0.177 e Å⁻³, respectively.

Figures




Fig. 1. Perspective drawing of the title compound, with the atomic numbering scheme. Displacement ellipsoids are shown at the 35% probability level.




Fig. 2. The unit cell packing of the title compound, viewed along the a direction.

2,4-Dinitrobenzaldehyde hydrazone

Crystal data

$C_7H_6N_4O_4$

$M_r = 210.16$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 4.5839$ (7) Å

$b = 9.6840$ (16) Å

$c = 9.9287$ (15) Å

$\alpha = 90.785$ (12)°

$\beta = 96.149$ (11)°

$\gamma = 98.955$ (13)°

$V = 432.66$ (12) Å³

$Z = 2$

$F_{000} = 216$

$D_x = 1.613$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 39 reflections

$\theta = 5.9$ – 12.5 °

$\mu = 0.14$ mm⁻¹

$T = 295$ (2) K

Prism, yellow

$0.4 \times 0.3 \times 0.2$ mm

Data collection

Bruker P4
diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 295$ (2) K

ω scans

Absorption correction: none

2238 measured reflections

1616 independent reflections

1160 reflections with $I > 2\sigma(I)$

$R_{int} = 0.027$

$\theta_{max} = 25.5$ °

$\theta_{min} = 2.1$ °

$h = -5 \rightarrow 1$

$k = -11 \rightarrow 11$

$l = -11 \rightarrow 11$

3 standard reflections

every 97 reflections

intensity decay: none

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.048$

$wR(F^2) = 0.116$

$S = 1.07$

1616 reflections

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.001P)^2 + 0.38P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{max} < 0.001$

$\Delta\rho_{max} = 0.16$ e Å⁻³

136 parameters

$$\Delta\rho_{\min} = -0.18 \text{ e } \text{\AA}^{-3}$$

Primary atom site location: structure-invariant direct methods

Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.7840 (6)	0.1234 (3)	0.5319 (2)	0.0887 (8)
O2	0.4576 (6)	0.2314 (2)	0.6030 (2)	0.0798 (7)
O3	0.0003 (6)	0.5548 (2)	0.3362 (3)	0.0872 (8)
O4	0.1241 (6)	0.5961 (3)	0.1351 (3)	0.0946 (9)
N1	0.9783 (6)	-0.0870 (3)	0.1463 (3)	0.0758 (8)
H1B	1.0167	-0.1364	0.2204	0.091*
H1C	1.0581	-0.1043	0.0700	0.091*
N2	0.8489 (5)	0.0275 (2)	0.1456 (2)	0.0578 (6)
N3	0.5907 (6)	0.1955 (2)	0.5116 (2)	0.0569 (6)
N4	0.1273 (6)	0.5291 (3)	0.2381 (3)	0.0675 (7)
C1	0.7626 (6)	0.0654 (3)	0.2571 (3)	0.0520 (7)
H1A	0.7984	0.0169	0.3358	0.062*
C2	0.6075 (6)	0.1858 (3)	0.2585 (3)	0.0471 (6)
C3	0.5165 (6)	0.2445 (3)	0.3752 (3)	0.0470 (6)
C4	0.3552 (6)	0.3540 (3)	0.3687 (3)	0.0518 (7)
H4A	0.2918	0.3885	0.4466	0.062*
C5	0.2906 (6)	0.4105 (3)	0.2464 (3)	0.0534 (7)
C6	0.3755 (6)	0.3582 (3)	0.1277 (3)	0.0580 (7)
H6A	0.3302	0.3978	0.0448	0.070*
C7	0.5269 (6)	0.2473 (3)	0.1360 (3)	0.0561 (7)
H7A	0.5791	0.2108	0.0565	0.067*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.117 (2)	0.1053 (19)	0.0558 (13)	0.0634 (17)	-0.0041 (13)	-0.0040 (12)
O2	0.1006 (18)	0.0891 (16)	0.0580 (13)	0.0258 (14)	0.0307 (12)	0.0011 (11)
O3	0.0851 (17)	0.0709 (15)	0.115 (2)	0.0314 (13)	0.0264 (15)	-0.0126 (14)
O4	0.136 (2)	0.0714 (16)	0.0829 (17)	0.0524 (16)	-0.0107 (16)	0.0009 (13)

supplementary materials

N1	0.109 (2)	0.0739 (17)	0.0583 (15)	0.0542 (17)	0.0153 (15)	0.0001 (13)
N2	0.0690 (15)	0.0583 (14)	0.0517 (14)	0.0267 (12)	0.0088 (11)	-0.0026 (11)
N3	0.0658 (15)	0.0538 (14)	0.0513 (14)	0.0095 (12)	0.0085 (12)	-0.0042 (11)
N4	0.0686 (17)	0.0521 (15)	0.083 (2)	0.0200 (13)	-0.0023 (15)	-0.0111 (14)
C1	0.0609 (17)	0.0507 (15)	0.0484 (15)	0.0176 (13)	0.0106 (13)	0.0035 (12)
C2	0.0447 (14)	0.0465 (14)	0.0510 (15)	0.0082 (11)	0.0085 (12)	-0.0016 (11)
C3	0.0482 (15)	0.0464 (14)	0.0459 (15)	0.0056 (12)	0.0063 (11)	-0.0001 (11)
C4	0.0494 (15)	0.0486 (15)	0.0578 (17)	0.0080 (12)	0.0091 (13)	-0.0096 (12)
C5	0.0505 (15)	0.0449 (15)	0.0661 (18)	0.0140 (12)	0.0029 (13)	-0.0031 (13)
C6	0.0649 (18)	0.0569 (17)	0.0542 (17)	0.0183 (14)	0.0027 (14)	0.0034 (13)
C7	0.0661 (18)	0.0586 (17)	0.0473 (15)	0.0211 (14)	0.0073 (13)	-0.0027 (12)

Geometric parameters (\AA , $^\circ$)

O1—N3	1.214 (3)	C1—H1A	0.9300
O2—N3	1.221 (3)	C2—C7	1.400 (4)
O3—N4	1.229 (3)	C2—C3	1.414 (3)
O4—N4	1.219 (3)	C3—C4	1.382 (3)
N1—N2	1.336 (3)	C4—C5	1.361 (4)
N1—H1B	0.8999	C4—H4A	0.9300
N1—H1C	0.9000	C5—C6	1.393 (4)
N2—C1	1.282 (3)	C6—C7	1.365 (4)
N3—C3	1.464 (3)	C6—H6A	0.9300
N4—C5	1.463 (3)	C7—H7A	0.9300
C1—C2	1.458 (3)		
N2—N1—H1B	124.3	C4—C3—C2	122.2 (2)
N2—N1—H1C	115.3	C4—C3—N3	115.3 (2)
H1B—N1—H1C	119.5	C2—C3—N3	122.5 (2)
C1—N2—N1	117.5 (2)	C5—C4—C3	118.9 (2)
O1—N3—O2	121.8 (3)	C5—C4—H4A	120.6
O1—N3—C3	119.9 (2)	C3—C4—H4A	120.6
O2—N3—C3	118.3 (2)	C4—C5—C6	121.7 (3)
O4—N4—O3	123.8 (3)	C4—C5—N4	119.6 (3)
O4—N4—C5	118.6 (3)	C6—C5—N4	118.7 (3)
O3—N4—C5	117.6 (3)	C7—C6—C5	118.4 (3)
N2—C1—C2	118.9 (2)	C7—C6—H6A	120.8
N2—C1—H1A	120.5	C5—C6—H6A	120.8
C2—C1—H1A	120.5	C6—C7—C2	123.2 (3)
C7—C2—C3	115.5 (2)	C6—C7—H7A	118.4
C7—C2—C1	119.4 (2)	C2—C7—H7A	118.4
C3—C2—C1	125.0 (2)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1B \cdots O1 ⁱ	0.90	2.52	3.305 (3)	146
N1—H1C \cdots N2 ⁱⁱ	0.90	2.34	3.123 (4)	146

Symmetry codes: (i) $-x+2, -y, -z+1$; (ii) $-x+2, -y, -z$.

Fig. 1




Fig. 2

