

Published in final edited form as:

Nat Ecol Evol. 2018 May ; 2(5): 800–809. doi:10.1038/s41559-018-0518-2.

***Homo sapiens* in Arabia by 85,000 years ago**

Huw S. Groucutt^{1,2,*}, Rainer Grün^{3,4}, Iyad S.A. Zalmout⁵, Nick A. Drake⁶, Simon J. Armitage^{7,8}, Ian Candy⁷, Richard Clark-Wilson⁷, Julien Louys³, Paul S. Breeze⁶, Mathieu Duval^{3,9}, Laura T. Buck^{10,11}, Tracy L. Kivell^{12,13}, Emma Pomeroy^{10,14}, Nicholas B. Stephens¹³, Jay T. Stock^{10,15}, Mathew Stewart¹⁶, Gilbert J. Price¹⁷, Leslie Kinsley⁴, Wing Wai Sung¹⁸, Abdullah Alsharekh¹⁹, Abdulaziz Al-Omari²⁰, Muhammad Zahir²¹, Abdullah M. Memesh⁵, Ammar J. Abdulshakoor⁵, Abdu M. Al-Masari⁵, Ahmed A. Bahameem⁵, Khaled S.M. Al Murayyi²⁰, Badr Zahrani²⁰, Eleanor M.L. Scerri^{1,2}, and Michael D. Petraglia^{2,22,*}

¹School of Archaeology, Research Laboratory for Archaeology and the History of Art, Hayes House, 75 George Street, University of Oxford, Oxford, OX1 2BQ, UK

²Department of Archaeology, Max Planck Institute for the Science of Human History, Kahlaische Stasse 10, D-07743, Jena, Germany

³Australian Research Centre for Human Evolution (ARCHE), Environmental Futures Research Institute, Griffith University, Nathan, QLD 4111, Australia

⁴Research School of Earth Sciences, The Australian National University, Canberra ACT, 0200, Australia

⁵Saudi Geological Survey, Sedimentary Rocks and Palaeontology Department, Jeddah 21514, Saudi Arabia

⁶Department of Geography, King's College London, Strand, London WC2R 2LS, UK

⁷Department of Geography, Royal Holloway, University of London, TW20 0EX UK

⁸SFF Centre for Early Sapiens Behaviour (SapienCE), University of Bergen, Post Box 7805, 5020, Bergen, Norway

⁹Geochronology, Centro Nacional de Investigación sobre la Evolución (CENIEH), Paseo Sierra de Atapuerca, 3, 09002 Burgos, Spain

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

*Correspondence and requests for materials should be addressed to HSG (huw.groucutt@rlaha.ox.ac.uk) and MDP (petraglia@shh.mpg.de).

Author Contributions H.S.G. and M.D.P. designed, coordinated and supervised the study. H.S.G., I.S.Z., N.D., S.A., I.C., R.C.-W., J.L., P.S.B., M.S., G.J.P., A.A., A.A.-O., A.M. B.A., E.M.L.S. and M.D.P. conducted excavation, survey and multidisciplinary sampling at Al Wusta. L.T.B., T.L.K., E.P., N.B.S. and J.T.S. conducted the morphological analysis and comparative study of the AW-1 phalanx. R.G., M.D. and L.K. carried out the U-series and ESR analyses. S.J.A. and R.C.W. carried out the OSL dating. I.C. and R.C.W. conducted the stratigraphic and sedimentological analysis of the site, with input from N.D., J.L. and G.J.P. W.W.S. analysed the diatoms. M.S. and J.L. analysed the vertebrate fossils, with input from G.J.P. Lithic analysis was conducted by H.S.G. and E.M.L.S. Spatial analyses were conducted by P.S.B. All authors helped to write the paper.

Author Information The authors declare no competing financial interests.

Readers are welcome to comment on the online version of the paper.

Data availability statement. Authors can confirm that all relevant data are included in the paper and/or its supplementary information files.

¹⁰PAVE Research Group, Dept. of Archaeology, University of Cambridge, Cambridge, CB2 3QG, UK

¹¹Earth Sciences Department, Natural History Museum, Cromwell Road, London, SW7 5BD, UK

¹²Skeletal Biology Research Centre, School of Anthropology and Conservation, University of Kent, Canterbury, CT2 7NR, UK

¹³Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, 04103, Leipzig, Germany

¹⁴School of Natural Sciences and Psychology, Liverpool John Moores University, Liverpool, L3 3AF, UK

¹⁵Department of Anthropology, University of Western Ontario, London, Ontario, N6A 3K7, Canada

¹⁶Palaeontology, Geobiology and Earth Archives Research Centre, School of Biological, Earth and Environmental Science, University of New South Wales, Sydney 2052, Australia

¹⁷School of Earth and Environmental Sciences, The University of Queensland, St Lucia 4072, Queensland, Australia

¹⁸Department of Life Sciences, The Natural History Museum, Cromwell Road, London SW7 5BD, UK

¹⁹Department of Archaeology, King Saud University, Riyadh, 12372 Saudi Arabia

²⁰Saudi Commission for Tourism and National Heritage, Riyadh 11586, Saudi Arabia

²¹Department of Archaeology, Hazara University, Mansehra, 21300, Pakistan

²²Human Origins Program, National Museum of Natural History, Smithsonian Institution, Washington, DC, 20560, USA

Abstract

Understanding the timing and character of *Homo sapiens* expansion out of Africa is critical for inferring the colonisation and admixture processes that underpin global population history. It has been argued that dispersal out of Africa had an early phase, particularly ~130-90 thousand years ago (ka), that only reached the East Mediterranean Levant, and a later phase, ~60-50 ka, that extended across the diverse environments of Eurasia to Sahul. However, recent findings from East Asia and Sahul challenge this model. Here we show that *H. sapiens* was in the Arabian Peninsula before 85 ka. We describe the Al Wusta-1 (AW-1) intermediate phalanx from the site of Al Wusta in the Nefud Desert, Saudi Arabia. AW-1 is the oldest directly dated fossil of our species outside Africa and the Levant. The palaeoenvironmental context of Al Wusta demonstrates that *H. sapiens* using Middle Palaeolithic stone tools dispersed into Arabia during a phase of increased precipitation driven by orbital forcing, in association with a primarily African fauna. A Bayesian model incorporating independent chronometric age estimates indicates a chronology for Al Wusta of ~95-86 ka, which we correlate with a humid episode in the later part of Marine Isotope Stage 5 known from various regional records. Al Wusta shows that early dispersals were more spatially and temporally extensive than previously thought. Early *H. sapiens* dispersals out of Africa were not limited to winter rainfall-fed Levantine Mediterranean woodlands immediately adjacent to

Africa, but extended deep into the semi-arid grasslands of Arabia, facilitated by periods of enhanced monsoonal rainfall.

Background

Homo sapiens evolved in Africa in the late Middle Pleistocene¹. Early dispersals out of Africa are evidenced at the Levantine site of Misliya at ~194-177 ka², followed by Skhul and Qafzeh, where *H. sapiens* fossils have been dated to ~130-100 and ~100-90 ka respectively³. While the Levantine fossil evidence has been viewed as the onset of a much broader dispersal into Asia^{4–6}, it has generally been seen as representing short-lived incursions into the woodlands of the Levant immediately adjacent to Africa, where relatively high precipitation is produced by winter storms tracking across the Mediterranean^{7,8}. While the Levantine record indicates the subsequent local replacement of early *H. sapiens* by Neanderthals, the failure of early dispersals to extend beyond the Levant is largely inferred from interpretations of genetic data⁹. Genetic studies have suggested that recent non-African populations stem largely¹⁰, if not entirely⁹, from an expansion ~60-50 ka, but this model remains debated. The absence of low latitude Pleistocene human DNA and uncertainties regarding ancient population structure undermine conclusions drawn from genetic studies alone. The paucity of securely dated archaeological, palaeontological and ancient DNA data - particularly across southern Asia - has made testing dispersal hypotheses challenging^{4,7,11}.

Recent fossil discoveries in East Asia indicate that the early (particularly Marine Isotope Stage 5) dispersals of *Homo sapiens* extended across much of southern Asia. At Tam Pa Ling in Laos, *Homo sapiens* fossils date to between 70 and 46 ka¹². Teeth assigned to *Homo sapiens* from Lida Ajer cave, Sumatra, were recovered from a breccia dating to 68 ± 5 ka, with fauna from the site dating to 75 ± 5 ka¹³. Several sites in China have produced fossil material claimed to represent early *Homo sapiens*¹⁴. These include teeth from Fuyan Cave argued to be older than 80 ka based on the dating of an overlying speleothem a few metres from the fossils¹⁵, and teeth from Luna Cave that were found in a layer dating to between 129.9 ± 1.5 ka and 70.2 ± 1.4 ka¹⁶. Teeth and a mandible from Zhiren Cave, China, date to at least 100 ka and have been argued to represent *Homo sapiens*, but other species attributions are possible¹⁷. The recent documentation of a human presence in Australia from ~65 ka is consistent with these findings¹⁸. Likewise, some interpretations of genetic data are consistent with an early spread of *Homo sapiens* across southern Asia¹⁰. These discoveries are leading to a radical revision of our understanding of the dispersal of *Homo sapiens*, yet there remain stratigraphic and taxonomic uncertainties for many of the east Asian fossils^{14,19}, and thousands of kilometers separate these findings from Africa.

The Arabian Peninsula is a vast landmass at the crossroads of Africa and Eurasia. Growing archaeological evidence demonstrates repeated hominin occupations of Arabia^{20,21} each associated with a strengthened summer monsoon which led to the re-activation of lakes and rivers^{22–24}, as it did in North Africa²⁵. Here we report the discovery of the first pre-Holocene human fossil in Arabia, Al Wusta-1 (AW-1), as well as the age, stratigraphy,

vertebrate fossils and stone tools at the Al Wusta site (Fig. 1, see also Supplementary Information).

Results

AW-1 is an intermediate manual phalanx, most likely from the 3rd ray (Fig. 2a, Supplementary Information 1: see below for detail on siding and species identification). It is generally well-preserved, although there is some erosion of the cortical/subchondral bone, and minor pathological bone formation (likely an enthesophyte) affecting part of the diaphysis (Supplementary Information 1). The phalanx measures 32.3 mm in proximo-distal length, and 8.7 mm and 8.5 mm in radio-ulnar breadth of the proximal base and midshaft, respectively (Supplementary Table 1).

AW-1 is more gracile than the robust intermediate phalanges of Neanderthals^{26–28}, which are broader radio-ulnarly relative to their length and have a more 'flared' base. AW-1's proximal radio-ulnar maximum breadth is 14.98 mm, which provides an intermediate phalanx breadth-length index (proximal radio-ulnar maximum breadth relative to articular length) of 49.6. This is very similar to the mean (\pm SD) for the Skhul and Qafzeh *H. sapiens* of 49.7 (\pm 4.1) and 49.1 (\pm 4.0) for Upper Palaeolithic Europeans, but 1.89 standard deviations below the Neanderthal mean of 58.3 (\pm 4.6)²⁹.

To provide a broad interpretive context for the Al Wusta phalanx, we conducted linear and geometric morphometric (GMM) landmark analyses (Supplementary Information 1) on phalanges from non-human primates, fossil hominins and geographically widespread recent *H. sapiens*. Comparative linear analyses (Supplementary Information 1, Supplementary Tables 2 and 3, Supplementary Figure 1) reveal that there is substantial overlap across most taxa for all shape ratios, so AW-1 falls within the range of variation of *H. sapiens*, cercopiths, *Gorilla*, *Australopithecus afarensis*, *A. sediba* and Neanderthals. However, AW-1 is most similar to the median value or falls within the range of variation of recent and early *H. sapiens* for all shape ratios.

Geometric morphometric (GMM) analyses of AW-1 and various primate groups including hominins (see Supplementary Table 4 and Supplementary Figure 2 for landmarks, and Supplementary Table 5 for sample) are illustrated in Figure 3 and Supplementary Figure 3. PC1 and PC2 together account for 61% of group variance in shape. AW-1 is separated on these two shape vectors from the non-human primates and most of the Neanderthals. AW-1 falls closest to the recent and early *H. sapiens* and is clearly differentiated from all non-human primates. This is also shown by the Procrustes distances from AW-1 to the mean shapes of each taxonomic group (Supplementary Table 6).

Three of the Neanderthal phalanges (from Kebara 2 and Tabun C1) are quite disparate from the main Neanderthal cluster and fall closer to the *H. sapiens* and Al Wusta cluster on PC1 and 2 (Figure 3 and Supplementary Figure 3). Having established the hominin affinity of AW-1, shape was analysed in more detail using a smaller hominin sample for which ray number and side were known, which included Kebara 2 and Tabun C1. The broader primate sample used in the first GMM analysis was not used for the more detailed shape analysis, as

the initial comparisons show clearly that AW-1 is not a non-human primate and including this level of variation could potentially mask more subtle shape differences between hominins. The side and ray are also not known for most of the Neanderthal and non-human primate samples, meaning it would be impossible to evaluate the effect of these factors using this sample.

The more in-depth shape comparison and modelling using the hominin sample of phalanges of known ray and side (Supplementary Table 7) demonstrates that the long and slender morphology of AW-1 falls just outside the range of variation of comparative Middle Palaeolithic modern humans, but that its affinity is clearly with *H. sapiens* rather than Neanderthals (Fig. 4, Supplementary Table 8). Although both Pleistocene *H. sapiens* and Neanderthal landmark configurations fall almost completely inside the scatter for the Holocene *H. sapiens* sample in the principal components analysis (Figure 4), AW-1 is closest to Holocene *H. sapiens* 3rd intermediate phalanges. AW-1 overlaps with the Holocene *H. sapiens* sample, but is separated from the Pleistocene *H. sapiens* specimens by a higher score on PC2 and from the Neanderthal group by a simultaneously higher score on PC1 and PC2. The Procrustes distances (Supplementary Table 8), also show that AW-1 is most distinct from the Neanderthal phalanges, which fall towards the lower ends of both PCs and are characterised by shorter and broader dimensions. PC1 and PC2 in this analysis show that AW-1 is taller and narrower (in all directions: dorso-palmarly, proximo-distally and radio-ulnarly) than almost all the phalanges in the comparative sample and is particularly distinct from most of the Neanderthal phalanges. In this analysis AW-1 is closest in shape to 3rd phalanges of individuals from (in descending order of proximity) Egyptian Nubia, and Medieval Canterbury (UK), and Maiden Castle (Iron Age Dorset, UK) (Supplementary Table 9), although there is not a great difference in its distance to any of these specimens. These analyses suggest that the AW-1 phalanx is likely to be a 3rd intermediate phalanx from a *H. sapiens* individual.

The third ray is the most symmetrical ray in the hand and is therefore difficult to side, particularly when not all of the phalanges of a particular individual are present. Comparing AW-1 separately to right and to left phalanges (Supplementary Information 1.4) gives results which are very similar to the pooled sample, such that AW-1 is closest to Holocene *H. sapiens* 3rd rays for both right and left hands (Supplementary Figure 4, Supplementary Table 10). There is little difference in morphological closeness between AW-1 and its nearest neighbour in the samples of right and left bones (Supplementary Table 11), reflecting the lack of difference in morphology between the sides. It is therefore not possible to suggest whether AW-1 comes from a right or a left hand using these analyses.

AW-1 is unusual in its more circular midshaft cross-sectional shape (Fig. 2B), which is confirmed by cross-sectional geometric analyses (Supplementary Information 1.5). This may reflect the pronounced palmar median bar that makes the palmar surface slightly convex at the midshaft rather than flat, the latter being typical of most later *Homo* intermediate phalanges. However, more circular shafts may reflect greater loading of the bone in multiple directions and enthesophytes are a common response to stress from high levels of physical activity³⁰. This morphology may reflect high and varied loading of the fingers during intense manual activity.

To determine the age of AW-1, and associated sediments and fossils, we used a combination of uranium series (U-series), electron spin resonance (ESR) and optically stimulated luminescence (OSL) dating (Methods, Supplementary Information 2 and 3). U-series ages were produced for AW-1 itself (87.6 ± 2.5 ka) and hippopotamus dental tissues (WU1601), which yielded ages of 83.5 ± 8.1 ka (enamel) and 65.0 ± 2.1 ka (dentine). They should be regarded as minimum estimates for the age of the fossils. In addition, a combined U-series-ESR age calculation for WU1601 yielded an age of $103 +10/-9$ ka. AW-1 was found on an exposure of Unit 3b, and WU1601 excavated from Unit 3a, one metre away (Fig 1b). Unit 1 yielded OSL ages of 85.3 ± 5.6 ka (PD17), 92.2 ± 6.8 ka (PD41) and 92.0 ± 6.3 ka (PD15), while Unit 3a yielded an OSL age of 98.6 ± 7.0 ka (PD40). The OSL age estimates agree within error with the US-ESR age obtained for WU-1601 and the minimum age of ~ 88 ka obtained for AW-1. These data were incorporated into a Bayesian sequential phase model³¹ which indicates that deposition of Unit 1 ceased 93.1 ± 2.6 ka (Phase 1: PD15, 17, 41) and that Units 2 and 3 and all associated fossils were deposited between 92.2 ± 2.6 ka and 90.4 ± 3.9 ka (Phase 2: all other ages) (Supplementary Information 4, Supplementary Figure 11).

This ~ 95 - 86 ka timeframe is slightly earlier than most other records of increased humidity in the region in late MIS 5^{32,33}, which correlate with a strengthened summer monsoon associated with an insolation peak at 84 ka (Fig. 6). The underlying (Unit 3) aeolian sand layer at Al Wusta correlates with an insolation minimum at the end of MIS 5c. The chronometric age estimates for the site suggest that lake formation and the associated fauna and human occupation occurred shortly after this in time. Regional indications of increased humidity around the 84 ka insolation peak include speleothem formation at ~ 88 ka in the Negev³⁴, and the formation of sapropel S3 beginning ~ 86 ka³⁵. In both the Levant and Arabia, records are consistent with this switch from aridity to humidity around this time³²⁻⁴⁰. Precisely reconstructing regional palaeoclimate at this time and relating it to human demographic and behavioural change has proved challenging. This reflects both rapid changes in climate, as well as the complexities involved in dating relevant deposits⁴¹. In summary, combining chronological data (Supplementary sections 2-4), interpretation of the sedimentary sequence (described below), and the regional setting of Al Wusta, we conclude that lake formation and associated finds such as the AW-1 phalanx relate to the late MIS 5 humid period associated with the 84 ka insolation peak.

The sedimentary sequence at Al Wusta consists of a basin-like deposit of exposed carbonate-rich sediments (Unit 2, 0.4-0.8 m thick), underlain by wind-blown sand (Unit 1) and overlain by water-lain sands (Unit 3). The carbonate rich sediments of Unit 2 are interpreted as lacustrine marl deposits on the basis of their sedimentology, geochemistry, and diatom palaeoecology (Figure 1c, Methods, Supplementary Information 5). At both the macro- and micro-scale, these beds are relatively massive and comprise fine-grained calcite, typical of material precipitating and accumulating in a still-water lacustrine environment⁴². At the micro-scale there is no evidence for the desiccation or fluctuation of water levels typical of palustrine/wetland environments⁴², implying that the lake body was perennial. The diatom flora support this, containing species such as *Aulacoseira italica* and *Aulacoseira granulata* throughout the sequences, indicating an alkaline lake a few metres deep. The water was fresh, not saline or brackish, since saline tolerant species and evaporitic minerals are absent throughout. While $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of continental carbonates are controlled by a wide-

range of variables, the values derived from the Al Wusta marl beds are compatible with the suggestion of marls precipitated in a perennial lake basin. The Al Wusta carbonate beds therefore indicate a perennial lake body a few metres in depth. The existence of a marl precipitating lake basin implies that this system was groundwater fed (to allow for sufficient dissolved mineral material to be present in the lake waters). Although the Al Wusta sequence represents a single lake basin, the development of such a feature over highly permeable aeolian sands in a region where no lake systems exist at the present day implies a local increase in water table that would require an increase in mean annual rainfall. Consequently, the Al Wusta sequence represents the occurrence of a humid interval at this time. The Unit 2 marl is overlain by a medium-coarse sand (Unit 3) with crude horizontal laminations, occasional clasts, fragments of ripped up marl and shells of *Melanoides tuberculata* and *Planorbis* sp. While some vertebrate fossils and lithics were found in the upper part of Unit 2, most were found in or on the surface of Unit 3. Unit 3a sands are waterlain and represent the encroachment of fluvial sediment as the lake environment shallowed and contracted. Unit 3b represents a winnowed lag formed by aeolian deflation of 3a. The sequence is capped by a dense network of calcitic rhizoliths marking the onset of fully terrestrial conditions.

A total of 860 vertebrate fossils were excavated from Unit 3 and the top of Unit 2 (n=371) and systematically surface collected (n=489). These include specimens attributed to Reptilia, Aves, and Mammalia (Supplementary Table 19, Methods, Supplementary Information 6). Notable taxa now extinct in Arabia are predominately grazers and include *Hippopotamus*, *Pelorovis*, and *Kobus*. The faunal community demonstrates a clear preference for temperate to semi-arid grasslands, and the presence of *Hippopotamus* and *Kobus* indicate permanent muddy, fluvial, or lacustrine conditions⁴³ not currently found in the Nefud Desert, but consistent with the geological evidence from the site. The faunal assemblages show a strong affinity to African fauna, particularly *Hippopotamus*, *Pelorovis*, and *Kobus*⁴⁴. Many large tooth pits on fossils indicate that large carnivores played a role in the accumulation of the deposit. Long bone circumference, completeness and numbers of green fractures suggests modification of bones by bone-breaking agents such as large carnivores or hominins (Supplementary Information 6). However, no evidence of cut-marks or hammerstone damage to the bones was observed.

An assemblage of 380 lithic artefacts (stone tools) was recovered from the excavation of upper Unit 2 and Unit 3 and systematic surface collection (Methods, Figure 5, Supplementary Information 7). They are of Middle Palaeolithic character and most are chert and quartzite. The assemblage demonstrates a focus on centripetal Levallois reduction, and is similar to other late Marine Isotope Stage 5 assemblages in the west and north of Arabia⁴⁵, and contemporaneous assemblages in east (e.g. Aduma, BNS at Omo Kibish) and northeast Africa (e.g. Bir Tarfawi), as well as those from the Levant (e.g. Qafzeh)¹¹ (Fig. 5).

Discussion

Al Wusta-1 is the oldest directly dated *H. sapiens* fossil outside Africa and the Levant. It joins a small but growing corpus of evidence that the early dispersal of *H. sapiens* into Eurasia was much more widespread than previously thought. The site of Al Wusta is located

in the Nefud desert more than 650 km southeast of Skhul and Qafzeh (Fig. 1A). This site establishes that *H. sapiens* were in Arabia in late MIS 5, rather than being restricted to Africa and the Levant as suggested by traditional models (Fig. 6). With Skhul dating to ~130-100 ka, Qafzeh to ~100-90 ka^{3,46} and Al Wusta to ~95-85 ka it is currently unclear if the southwest Asian record reflects multiple early dispersals out of Africa or a long occupation during MIS 5. The association of the Al Wusta site with a late MIS 5 humid phase (Fig. 6), suggests that significant aspects of this dispersal process were facilitated by enhanced monsoonal rainfall. While changes in behaviour and demography are crucial to understanding the dispersal process, climatic windows of opportunity were also key in allowing *H. sapiens* to cross the Saharo-Arabian arid belt, which often constituted a formidable barrier^{24,25}.

Conclusion

Al Wusta shows that the early, Marine Isotope Stage 5, dispersals of *H. sapiens* out of Africa were not limited to the Levantine woodlands sustained by winter rainfall, but extended deep into the Arabian interior where enhanced summer rainfall created semi-arid grasslands containing abundant fauna and perennial lakes. After long being isolated in Africa^{1,47,48}, the Late Pleistocene saw the expansion of our species out of Africa and into the diverse ecologies of Eurasia. Within a few thousand years of spreading into Eurasia our species was occupying rainforest environments and making long sea crossings to remote islands^{13,18}. Adapting to the semi-arid conditions of the Saharo-Arabian arid belt represented a crucial step on this pathway to global success and the Al Wusta *Homo sapiens* fossil demonstrates this early ability to occupy diverse ecologies which led to us becoming a cosmopolitan species.

Methods

Site identification, survey and excavation

The site of Al Wusta (field code WNEF16_30) was discovered in 2014 as part of a programme of joint survey fieldwork of the Palaeodeserts Project, the Saudi Commission for Tourism and National Heritage, and the Saudi Geological Survey. It is located in the western Nefud desert, a few kilometres from the Middle Pleistocene fossil locality of Ti's al Ghaddah⁴⁹. The locations of all materials of interest (fossils, stone tools, geomorphological features, excavations and sample points) were recorded using a high-precision Trimble XRS Pro Differential GPS system and a total station, and entered into a GIS (Fig. 1). Elevation data (masl) were recorded as a series of transects across the site, and a digital elevation model (DEM) and contours interpolated (Spline) from all data with precisions of better than 10 cm in all (x,y,z) dimensions (22,047 points). This allowed visualisation and recording of the spatial relationships between materials in three dimensions (Fig. 1). Eight trenches were excavated into the fossil and artefact bearing deposits. These trenches revealed vertebrate remains and lithics, but no further human fossils were recovered.

Morphological analysis of AI Wusta-1 phalanx

The phalanx was scanned using micro-computed tomography (micro-CT) on the Nikon Metrology XT H 225 ST High Resolution scanner and X-Tek software (Nikon Metrology, Tring, UK) housed in the Cambridge Biotomography Centre, University of Cambridge, UK. Scan parameters were: a tungsten target; 0.5 mm copper filter; 150 kV; 210 mA; 1080 projections with 1000 ms exposure, and resulted in a voxel size of 0.02 mm³. The micro-CT data were reconstructed using CT-PRO 3D software (Nikon Metrology) and exported as an image (.tif) stack. Other CT data were obtained from the institutions cited in Supplementary Table 5 with permissions following the memoranda of understanding with each institution.

3D landmarks and semilandmarks were chosen to best describe the overall shape of the morphology of the AW-1 phalanx (Supplementary Table 4, Supplementary Figure 2), and were digitised on virtual reconstructions of phalanges created from micro-CT data in AVIZO 8 and 9.1 (FEI Software, Burlington, Mass.). Landmark coordinates were exported for use in Morphologika50. In Morphologika, generalized Procrustes analyses were performed to superimpose landmark coordinate data, and principal components analyses (PCA) were run to investigate similarities in shape between specimens. Shape differences along principal components were visualised and wireframes were produced in Morphologika, PC scores were exported to create graphs in R51. Procrustes distances between specimens were calculated using MorphoJ52. To avoid representing the same phalanges from different sides of a single individual as independent data points and to maximise sample sizes in pooled analyses, right phalanges were used in cases where the phalanges from both sides were present. Where only the left was present, this was used and ‘reflected’ (i.e. mirrored) in Morphologika to generate landmark configurations consistent with right phalanges.

U-series and combined US-ESR dating of fossil bone and teeth

The AW-1 phalanx (lab number 3675) and a hippopotamus tooth fragment (lab number WU1601) were collected from Trench 1 (Fig.1) for U-series and combined US-ESR dating, respectively. The external dose rate utilised the data of OSL sample PD40, which was collected in an equivalent position within unit 3a.

U-series analysis—U-series analyses were conducted at the Research School of Earth Sciences, The Australian National University, Canberra. The experimental setup for the U-series analysis of the phalanx was described in detail by Grün and colleagues⁵³ (Supplementary Figures 2 and 3, Supplementary Information 2). Laser ablation (LA) was used to drill a number of holes into AW-1 following the approach of Benson and colleagues⁵⁴. After a cleaning run with the laser set at a diameter of 460 µm, seven holes were drilled for 1000 s with the laser set at 330 µm. The isotopic data streams were converted into ²³⁰Th/²³⁴U and ²³⁴U/²³⁸U activity ratios and apparent Th/U age estimates and subsequently binned into 30 successive sections (each containing 33 cycles) for the calculation of average isotopic ratios and ages. A similar experimental setup and methodology were employed for the LA U-series analysis of tooth sample WU1601. The whole closed system U-series analytical datasets of the enamel and dentine sections were integrated to provide the data input for the ESR age calculations.

Combined US-ESR dating of the fossil tooth: ESR dose evaluation—The ESR dose evaluation of the hippo tooth was carried out at CENIEH, Burgos, Spain, following a similar procedure to that described in Stimpson and colleagues⁴⁹. Enamel was collected from WU1601 and powdered <200 μm . The sample was then divided into 11 aliquots and gamma irradiated with a Gammacell-1000 Cs-137 source to increasing doses until 3.4 kGy. ESR measurements were carried out at room temperature with an EMXmicro 6/1 Bruker ESR spectrometer coupled to a standard rectangular ER 4102ST cavity. ESR intensities were extracted from T1-B2 peak-to-peak amplitudes of the ESR signal of enamel. Fitting procedures were carried out with a single saturating exponential (SSE) function through the pooled ESR experimental data derived from the repeated measurements, with data weighting by the inverse of the squared ESR intensity ($1/I^2$) and following the recommendations by Duval and Grün⁵⁵. Full details about the experimental conditions and analytical procedure may be found in Supplementary Information 2.

Combined US-ESR dating of the fossil tooth: Dose rate evaluation and age calculations—The combined US-ESR age of WU1601 was calculated with the DATA programme⁵⁶ using the US model defined by Grün and colleagues⁵⁷. The following parameters were used for the dose rate evaluation: an alpha efficiency of 0.13 ± 0.0258 , Monte-Carlo beta attenuation factors from Marsh⁵⁹, dose-rate conversion factors from Guerin and colleagues⁶⁰, external sediment (beta and gamma) dose rate from the OSL sample PD40, a depth of 25 ± 10 cm, resulting in an age of $103 \pm 10/-9$ ka.

Optically Stimulated Luminescence Dating

Three samples (PD15, PD17 and PD41) were collected from the aeolian sands (Unit 1) underlying the southern marl outcrop (Unit 2, Fig 1B). A fourth sample (PD40) was taken from the main fossil bearing bed (Unit 3). Individual quartz grains were measured on a Risø TL/OSL-DA-15 instrument using the single-aliquot regenerative-dose (SAR) method⁶¹. The burial dose for each sample (D_b) was calculated using the central age model (CAM)⁶².

Environmental dose rates were determined using a Risø GM-25-5 low-level beta counting system⁶³ (beta dose rate), field gamma spectrometry (gamma dose rate), and an estimate of the cosmic dose rate derived using site location and present day sediment burial depths⁶⁴. Full optically stimulated luminescence dating methods and results are presented in Supplementary Information Section 3. All analyses were carried out in the Royal Holloway Luminescence Laboratory by SA and R C-W.

Age modelling

Chronometric ages for samples from the Al Wusta site were incorporated into a Bayesian sequential phase model implemented in OxCal v4.231 (Supplementary Information 4; Supplementary Figure 11). The model consists of two discrete phases separated by a hiatus. Phase 1 was defined by the three OSL ages (PD15, 17 and 41) for samples from the aeolian sands (Unit 1) underlying the lacustrine marls (Unit 2). Phase 2 was defined by the ages for the sand (PD40) and fossils (AW-1 and WU1601) from the waterlain sediments (Unit 3) overlying Unit 2. U-series ages for WU1601 and AW-1 were treated as minimum age estimates, whereas PD40 and the combined U-series-ESR age on WU1601 were treated as

finite age estimates. Since the Al Wusta sequence accumulated over a short period of time, and contains only five finite ages (and three minimum ages), the General Outlier Model³¹ was unable to function, and instead a simpler model using agreement indices was employed. This analysis yielded Amodel (76) and Aoverall (79) values well in excess of the generally accepted threshold (6031), with only one age yielding an individual agreement index below this threshold (PD17, 51). These data indicate that no ages should be excluded from the model, and that the age model itself is robust. The Bayesian sequential model yielded an age for the end of Phase 1 of 93.1 ± 2.6 ka (1 σ uncertainties), while Phase 2 yielded start and end dates of 92.2 ± 2.6 ka and 90.4 ± 3.9 ka respectively. The end date for phase 2 should be treated as a maximum value since no overlying material is present, precluding the possibility of further constraining the end of this phase.

Stratigraphy and sedimentology

Sediment analysis—Bulk samples (in the form of coherent blocks) were taken at 10 cm intervals through each of the marl beds in four sections (Fig. 1C and Supplementary Figures 13 and 14). Each block was air-dried and subsamples (ca 0.5 g) were removed, powdered and analysed for percentage carbonate content using Bascomb calcimetry, which measures the volume of carbon dioxide liberated from a known sample mass during reaction with 10% HCl⁶⁵. Thin sections were prepared from fresh sediment blocks. The sediments did not require acetone treatment as they were already dry and, due to their permeability, were impregnated with a bonding resin. Standard thin section preparation was then carried out using techniques developed in the Centre for Micromorphology at Royal Holloway, University of London⁶⁶. Thin sections were analysed using an Olympus BX-50 microscope with magnifications from 20x to 200x and photomicrographs were captured with a Pixera Penguin 600es camera. A point-count approach was used to produce semi-quantified data from the thin sections, based on counting micro-features at 3 mm intervals along linear transects 1 cm apart. Kemp⁶⁷, Stoops⁶⁸ and Alonso-Zarza⁴² were referred to when identifying features. X-ray diffraction analysis (XRD) was carried out in the Department of Earth Sciences (Royal Holloway, University of London). Powdered samples were analysed on a Philips PW1830/3020 spectrometer with copper K α X-rays. Mineral peaks were identified manually from the ICDD Powder Diffraction File (PDF) database. The methods and results are described further in Supplementary Information 5.

Diatoms—*Sample preparation*. Samples were analysed using the standard method of Renberg⁶⁹ (Supplementary Information 5). Thus, all samples were treated with 30% H₂O₂ and 5% HCl to digest organic material and remove calcium carbonate. Distilled water was added to dilute the samples after heating, which were then stored in the refrigerator for four days to minimise further chemical reactions. The samples were rinsed daily and allowed to settle overnight. A known volume of microspheres was added to the supernatant after the last rinse to enable calculation of the diatom concentration⁷⁰. The slides were air-dried at room temperature in a dust free environment before mounting with Naphrax diatom mountant. Diatom taxonomy followed Krammer and Lange-Bertalot^{71–73} and taxonomic revisions^{74,75} with at least 300 valves enumerated for a representative sample at x1000 magnification.

Numerical analysis. Prevalent trends in the diatom assemblage were explored using ordination analyses using CANOCO 4.5 of ter Braak and Šmilauer⁷⁶. Detrended Correspondence Analysis (DCA⁷⁷) with detrending by segments and down-weighting of rare species was used to investigate taxonomic variations within each site and to determine whether linear or unimodal models should be used for further analyses. If the gradient length of the first axis is <1.5 SD units, linear methods (Principle Component Analysis, PCA) should be used; however, if the gradient length is >1.5 SD units, unimodal methods (Correspondence Analysis) should be used⁷⁸. Detrended Canonical Correspondence Analysis (DCCA⁷⁹) was also used to show changes in compositional turnover scaled in SD units. Therefore, variations in the down-core DCCA first axis sample scores show an estimate of the compositional change between samples along an environmental or temporal gradient. Depth was used as the sole constraint as the samples in each site are in a known temporal order⁸⁰. The dataset was square-root transformed to normalise the distribution prior to analyses. Optimal sum-of-squares partitioning⁸¹ with the program ZONE⁸² and comparison of the zones with the Broken-stick model using the program BSTICK⁸³ were used to determine significant zones. The planktonic: benthic ratio, habitat summary, concentration and the F index (a dissolution index⁸⁴) were calculated for all the samples.

Stable isotopes—It is common practice, when analysing the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of lacustrine/palustrine carbonates to either: 1) sieve the sediment and analyse the <63 μm fraction, or 2) use the microstructure of the sample, as identified under thin section, to identify pure, unaltered fabrics, which can then be drilled out and analysed⁸⁵. The former procedure ensures that the analysed fraction comprises pure authigenic marl (rather than a mixture of ostracod, mollusc, chara and marl components that will contain different isotopic values). The latter is done to ensure that any carbonate that has been affected by diagenesis is sampled. Neither of these approaches were carried out here as; 1) microfabric analysis showed no evidence for diagenesis (although some of the samples are cemented the cement makes a negligible component of sample mass), and 2) some of the samples have incipient cementation, which means that they cannot be sieved. Bulk carbonate powders were consequently analysed for $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$. To show that the analysis of bulk samples had no impact on the derived isotopic data, samples that were friable enough to be sieved were treated with sodium hexametaphosphate to disaggregate them and then homogenised and separated into two subsamples for isotopic analysis; (1) a sieved <63 μm fraction and (2) a homogenised bulk sample. The resulting isotopic data showed no difference between the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of the sieved and bulk samples (Supplementary Figure 13b), highlighting that the homogenous and unaltered nature of the material results in bulk carbonate isotopic analysis generating valid data. Two samples were taken from different locations of each sampled block to generate a larger dataset of independent samples. The $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values of each samples were determined by analysing CO_2 liberated from the reaction of the sample with phosphoric acid at 90°C using a VG PRISM series 2 mass spectrometer in the Earth Sciences Department at Royal Holloway. Internal (RHBNC) and external (NBS19, LSVEC) standards were run every 4 and 18 samples respectively. 1σ uncertainties are 0.04‰ ($\delta^{18}\text{O}$) and 0.02‰ ($\delta^{13}\text{C}$). All isotope data presented in this study are quoted against the Vienna Pee Dee Belemnite (VPDB) standard.

Vertebrate fossil analyses

Each fossil specimen was identified to lowest taxonomic and anatomical level possible (Supplementary Figure 20, Supplementary Table 19 and Supplementary Information 6). Taxonomic identification and skeletal element portions were determined based on anatomical landmarks, and facilitated by comparisons with the Australian National University Archaeology and Natural History reference collection (Canberra), unregistered biological collections held at the University of New South Wales (Sydney), and the large mammal collections of the Zoologische Staatssammlung München (Munich). Each specimen was assigned a size category (small, medium, and large) following Dominguez-Rodrigo and colleagues⁸⁶, and corresponding to the five size classes described in Bunn⁸⁷, where small, medium and large denote size classes 1-2, 3A-3B and 4-6, respectively. Element abundance is reported as Number of Identified Specimens (NISP).

Each specimen was examined for modification by eye and hand-lens (10x) under both natural and high-incidence light, and examined at different angles to assist identification of fine-scale surface modifications. Where required, further examination and photography was carried out using a digital microscope (Model: Dino-lite, AM7013MZ). Morphometric data (length, breadth and width) was measured using digital callipers (Model: Mitutoyo Corp, CD-8"PMX), and specimen weights using a digital scale. Bone surface modifications were identified and recorded following standard methodologies: butchery and tooth marks^{88–94}, burning^{95–96}, rodent gnawing^{97,98}, weathering⁹⁹ and trampling¹⁰⁰. Carnivore damage was categorized as pit, score, furrow or puncture, and the location noted⁹⁴. Tooth mark morphometric data – short and long axes – was also recorded. Any additional modifications, i.e. polish, manganese staining, and root etching, were also reported and described. Bone breakage was recorded as green, dry, or both, following Villa and Mahieu¹⁰¹. Long bone circumference completeness was recorded using the three categories described by Bunn¹⁰²: type 1 (<1/2), type 2 (>1/2 but < complete) and type 3 (complete).

Lithic analysis

Lithics were systematically collected during pedestrian transects and excavations of Al Wusta. This produced a total studied assemblage of 380 lithics (Supplementary Information 7). Further lithics extended for a considerable distance to the north, seeming to track the outlines of the palaeolake, but we only conducted detailed analysis on lithics from the southern part of the site, close to AW-1 and the sedimentary ridge on which it was found (i.e. south of the Holocene playa). These were analysed using the methodology described in Scerri and colleagues^{25,103,104} and Groucutt and colleagues^{45,105}. As well as qualitative analysis of technological features indicating particular techniques and methods of reduction, a variety of quantitative features such as dimensions, the number of scars and % of cortex were recorded. Informative examples were selected for photography and illustration. This approach allows both a characterisation and description of the assemblage and broad comparison with other assemblages from surrounding regions.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank HRH Prince Sultan bin Salman bin Abdulaziz Al-Saud, President of the Saudi Commission for Tourism and National Heritage (SCTH), and Prof. Ali Ghabban, Vice President of the SCTH for permission to carry out this study. Dr Zohair Nawab, President of the Saudi Geological Survey, provided research support and logistics. Fieldwork and analyses were funded by the European Research Council (no. 295719, to MDP and 617627, to JTS), the SCTH, the British Academy (HSG and EMLS), The Leverhulme Trust, the Australian Research Council (DP110101415 to RG, FT150100215 and TF15010025 to MD, and FT160100450 to JL), and the Research Council of Norway (SFF Centre for Early Sapiens Behaviour, 262618). We thank Patrick Cuthbertson, Klint Janulis, Marco Bernal, Salih Al-Soubhi, Mohamad Haptari, Adel Matari, and Yahya Al-Mufarreh for assistance in the field. We thank Ian Cartwright (Institute of Archaeology, University of Oxford) for the photographs of AW-1 (Fig. 2a), Ian Matthews (RHUL) for producing the Bayesian age model, and Michelle O'Reilly (MPI-SHH) for assistance with the preparation of figures. We acknowledge the Max Planck Society for supporting us with comparative fossil data, and we thank curators for access to comparative extant and fossil material in their care (Supplementary Tables 5 and 7).

References

1. Stringer C. The origin and evolution of *Homo sapiens*. *Philos Trans R Soc B Biol Sci*. 2016; 371:20150237–20150237.
2. Hershovitz I, et al. The earliest modern humans outside Africa. *Science*. 2018; 359:456–459. [PubMed: 29371468]
3. Grün R, et al. U-series and ESR analyses of bones and teeth relating to the human burials from Skhul. *J Hum Evol*. 2005; 49:316–334. [PubMed: 15970310]
4. Groucutt HS, et al. Rethinking the dispersal of *Homo sapiens* out of Africa. *Evol Anthropol*. 2015; 24:149–164. [PubMed: 26267436]
5. Petraglia MD, et al. Middle Paleolithic assemblages from the Indian subcontinent before and after the Toba super-eruption. *Science*. 2007; 317:114–116. [PubMed: 17615356]
6. Bae CJ, Douka K, Petraglia MD. On the origin of modern humans: Asian perspectives. *Science*. 2017; 358doi: 10.1126/science.aai9067
7. Mellars P, Gori KC, Carr M, Soares PA, Richards MB. Genetic and archaeological perspectives on the initial modern human colonization of southern Asia. *Proc Natl Acad Sci USA*. 2013; 110:10699–10704. [PubMed: 23754394]
8. Shea JJ. Transitions or turnovers? Climatically-forced extinctions of *Homo sapiens* and Neanderthals in the east Mediterranean Levant. *Quatern Sci Rev*. 2008; 27:2253–2270.
9. Mallick S, et al. The Simons Genome Diversity Project: 300 genomes from 142 diverse populations. *Nature*. 2016; 538:201–206. [PubMed: 27654912]
10. Pagani L, et al. Genomic analyses inform on migration events during the peopling of Eurasia. *Nature*. 2016; 538:238–242. [PubMed: 27654910]
11. Groucutt HS, et al. Stone tool assemblages and models for the dispersal of *Homo sapiens* out of Africa. *Quatern Int*. 2015; 382:8–30.
12. Demeter F, et al. Early Modern Humans from Tam Pà Ling, Laos. *Fossil Review and Perspectives. Curr Anthropol*. 2017; 57:S17. doi: 10.1086/694192
13. Westaway KE, et al. An early modern human presence in Sumatra 73,000–63,000 years ago. *Nature*. 2017; 548:322–325. [PubMed: 28792933]
14. Michel V, et al. The earliest modern *Homo sapiens* in China? *J Hum Evol*. 2016; 101:101–104. [PubMed: 27586079]
15. Liu W, et al. The early unequivocally modern humans in southern China? *Nature*. 2015; 526:696–699. [PubMed: 26466566]
16. Bae C, et al. Modern human teeth from Late Pleistocene Luna Cave (Guangxi, China). *Quatern Int*. 2015; 354:169–183.
17. Liu W, et al. Human remains from Zhirendong, South China, and modern human emergence in East Asia. *Proc Natl Acad Sci USA*. 2010; 107:19201–19206. [PubMed: 20974952]
18. Clarkson C, et al. Human occupation of northern Australia by 65,000 years ago. *Nature*. 2017; 547:306–310. [PubMed: 28726833]

19. Martínón-Torres M, Wu X, de Castro JMB, Xing S, Liu W. *Homo sapiens* in the Eastern Asian Late Pleistocene. *Curr Anthropol*. 58:S17.
20. Groucutt HS, Petraglia MD. The prehistory of Arabia: Deserts, dispersals and demography. *Evol Anthropol*. 2012; 21:113–125. [PubMed: 22718479]
21. Petraglia MD, Groucutt HS, Parton A, Alsharekh A. Green Arabia: Human prehistory at the Cross-Roads of continents. *Quatern Int*. 2015; 382:1–7.
22. Jennings RP. The greening of Arabia: Multiple opportunities for human occupation in the Arabian Peninsula during the Late Pleistocene inferred from an ensemble of climate model simulations. *Quatern Int*. 2015; 205:181–199.
23. Rosenberg TM, et al. Middle and Late Pleistocene humid periods recorded in palaeolake deposits in the Nafud desert, Saudi Arabia. *Quatern Sci Rev*. 2013; 70:109–123.
24. Breeze PS, et al. Palaeohydrological corridors for hominin dispersals in the Middle East ~ 250-70,000 years ago. *Quatern Sci Rev*. 2016; 11:155–185.
25. Scerri EML, Drake NA, Jennings R, Groucutt HS. Earliest evidence for the structure of *Homo sapiens* populations in Africa. *Quatern Sci Rev*. 2014; 101:207–216.
26. Trinkaus, E. *The Shanidar Neandertals*. Academic Press; New York: 1981.
27. McCown, TD., Keith, A. *The Stone Age of Mount Carmel Volume 2: The fossil human remains from the Levallouiso-Mousterian*. Clarendon Press; Oxford: 1939.
28. Vandermeersch, B. *Les hommes fossiles de Qafzeh (Israel)*. CNRS; Paris: 1981.
29. Walker MJ, Ortega J, López MV, Parmová K, Trinkaus E. Neanderthal postcranial remains from the Sima de las Palomas del Cabezo Gordo, Murcia, Southeastern Spain. *Am J Phys Anthropol*. 2011; 144:505–515. [PubMed: 21404228]
30. Benjamin M, et al. Where tendons and ligaments meet bone: attachment sites ('entheses') in relation to exercise and/or mechanical load. *J Anat*. 208:471–490.
31. Bronk Ramsey C. Bayesian analysis of radiocarbon dates. *Radiocarbon*. 2009; 51:337–360.
32. Drake NA, Breeze P, Parker A. Palaeoclimate in the Saharan and Arabian Deserts during the Middle Palaeolithic and the potential for hominin dispersals. *Quatern Int*. 2013; 300:48–61.
33. Parton A, et al. Orbital-scale climate variability in Arabia as a potential motor for human dispersals. *Quatern Int*. 2015; 382:82–97.
34. Vaks A, Bar-Matthews M, Matthews A, Ayalon A, Frumkin A. Middle-Late Quaternary paleoclimate of northern margins of the Saharan-Arabian Desert: reconstruction from speleothems of Negev Desert, Israel. *Quatern Sci Rev*. 2010; 29:2647–2662.
35. Grant KM, et al. The timing of Mediterranean sapropel deposition relative to insolation, sea-level and African monsoon changes. *Quatern Sci Rev*. 2016; 140:125–141.
36. Bar-Matthews M, Ayalon A, Gilmour M, Matthews A, Hawkesworth CJ. Sea-land oxygen isotopic relationships from planktonic foraminifera and speleothems in the Eastern Mediterranean region and their implication for paleorainfall during interglacial intervals. *Geochim Cosmochim Acta*. 2003; 67:3181–3199.
37. Lisiecki LE, Raymo ME. A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}O$ records. *Paleoceanography*. 2005; 20:1–17.
38. Berger A, Loutre MF. Insolation values for the climate of the last 10 million years. *Quatern Sci Rev*. 1991; 10:297–317.
39. Fleitmann D, Burns SJ, Neff U, Mangini A, Matter A. Changing moisture sources over the last 333,000 years in Northern Oman from fluid-inclusion evidence in speleothems. *Quatern Res*. 2003; 60:223–232.
40. Rosenberg TM, et al. Humid periods in southern Arabia: Windows of opportunity for modern human dispersal. *Geology*. 2011; 39:1115–1118.
41. Clark-Balzan L, Parton A, Breeze PS, Groucutt HS, Petraglia MD. Resolving problematic luminescence chronologies for carbonate- and evaporite-rich sediments spanning multiple humid periods in the Jubba Basin, Saudi Arabia. *Quatern Geochron*. 2018; doi: 10.1016/j.quageo.2017.06.002
42. Alonso-Zarza AM. Palaeoenvironmental significance of palustrine carbonates and calcretes in the geological record. *Earth Sci Rev*. 2003; 60:261–298.

43. Estes, RD. *The Behaviour Guide to African Mammals*. University of California Press; Berkeley: 1991.
44. O'Regan HJ, Turner A, Bishop LC, Elton S, Lamb AL. Hominins without fellow travellers? First appearances and inferred dispersals of Afro-Eurasian large-mammals in the Plio-Pleistocene. *Quatern Sci Rev*. 2011; 30:1343–1352.
45. Groucutt HS, et al. Human occupation of the Arabian Empty Quarter during MIS 5: evidence from Mundafan al-Buhayrah. *Quatern Sci Rev*. 2015; 119:116–135.
46. Millard AR. A critique of the chronometric evidence for hominid fossils: I. Africa and the Near East 500–50 ka. *J Hum Evol*. 2008; 54:848–874. [PubMed: 18201747]
47. Hublin JJ, et al. New fossils from Jebel Irhoud, Morocco and the pan-African origin of *Homo sapiens*. *Nature*. 2017; 546:289–292. [PubMed: 28593953]
48. Richter D, et al. The age of the hominin fossils from Jebel Irhoud, Morocco, and the origins of the Middle Stone Age. *Nature*. 2017; 546:293–296. [PubMed: 28593967]

Methods References

49. Stimpson C, et al. Middle Pleistocene vertebrate fossils from the Nefud Desert, Saudi Arabia: Implications for biogeography and palaeoecology. *Quatern Sci Rev*. 2016; 143:13–36.
50. O'Higgins P, Jones N. Facial growth in *Cercocebus torquatus*: an application of three-dimensional geometric morphometric techniques to the study of morphological variation. *J Anat*. 1998; 193:251–72. [PubMed: 9827641]
51. R Core Team. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing; 2015. <http://www.R-project.org>
52. Klingenberg CP. MorphoJ: an integrated software package for geometric morphometrics. *Mol Ecol Resour*. 2011; 11:353–7. [PubMed: 21429143]
53. Grün R, Eggins S, Kinsley L, Mosely H, Sambridge M. Laser ablation U-series analysis of fossil bones and teeth. *Palaeogeogr, Palaeoclimatol, Palaeoecol*. 2014; 416:150–167.
54. Benson A, Kinsley L, Defleur A, Kokkonen H, Mussi M, Grün R. Laser ablation depth profiling of U-series and Sr isotopes in human fossils. *J Arch Sci*. 2013; 40:2991–3000.
55. Duval M, Grün R. Are published ESR dose assessments on fossil tooth enamel reliable? *Quat Geochron*. 2016; 31:19–27.
56. Grün R. The DATA program for the calculation of ESR age estimates on tooth enamel. *Quatern Geochron*. 2009; 4:231–232.
57. Grün R, Schwarcz HP, Chadam J. ESR dating of tooth enamel: Coupled correction for U-uptake and U-series disequilibrium. *Int J Radiat Appl Instrum Nucl Tracks Radiat Meas*. 1988; 14:237–241.
58. Grün R, Katzenberger-Apel O. An alpha irradiator for ESR dating. *Ancient TL*. 1994; 12:35–38.
59. Marsh, RE. *Beta-gradient Isochrons Using Electron Paramagnetic Resonance: Towards a New Dating Method in Archaeology*. MSc thesis, McMaster University; Hamilton: 1999.
60. Guérin G, Mercier N, Adamiec G. Dose-rate conversion factors: update. *Ancient TL*. 2011; 29(1): 5–8.
61. Murray AS, Wintle AG. Luminescence dating of quartz using an improved single-aliquot regenerative-dose protocol. *Radiat Meas*. 2000; 32:57–73.
62. Galbraith RF, Roberts RG, Laslett GM, Yoshida H, Olley JM. Optical dating of single and multiple grains of quartz from Jinnium rock shelter, northern Australia: Part I, experimental design and statistical models. *Archaeometry*. 1999; 41:339–364.
63. Bøtter-Jensen L, Mejdahl V. Assessment of beta dose-rate using a GM multiscaler system. *Int J Rad Appl Instrum B*. 1988; 14:187–191.
64. Prescott JR, Hutton JT. Cosmic ray and gamma ray dosimetry for TL and ESR. *Int J Rad Apple Instrum*. 1988; 14:223–227.
65. Gale, S., Hoare, P. *Quaternary Sediments: Petrographic Methods for the Study of Unlithified Rocks*. Belhaven and Halsted Press; 1991.

66. Palmer, AP., Lee, JA., Kemp, RA., Carr, SJ. Revised laboratory procedures for the preparation of thin sections from unconsolidated sediments. Centre for micromorphology publication, Royal Holloway, University of London; 2008.
67. Kemp, RA. Soil micromorphology and the Quaternary. Quaternary Research Association; 1985.
68. Stoops, G. Interpretation of micromorphological features of soils and regoliths. Elsevier; 2010.
69. Rengberg I. A procedure for preparing large sets of diatom sets from sediment. *J Palaeolimnol.* 1990; 4:87–90.
70. Battarbee RW, Knen MJ. The use of electronically counter microspheres in absolute diatom analysis. *Limnol Oceanogr.* 1982; 27:184–188.
71. Krammer, K., Lange-Bertalot, H. Bacillariophyceae 2. Teil Epithemiaceae, Suiellaceae. Gustav-Fisher Verlag; 1988.
72. Krammer, K., Lange-Bertalot, H. Bacillariophyceae 3. Teil Centrales, Fragicariaceae, Eunotiaceae. Gustav-Fisher Verlag; 1991a.
73. Krammer, K., Lange-Bertalot, H. Bacillariophyceae 4. Teil Achnanthaceae, Kritshe Ergänzungen zu Navicula (Lineolate) und Gomphonema. Gustav-Fisher Verlag; 1991b.
74. Crawford RM, Likhoshway YV, Jahn R. Morphology and identity of *Aulacoseira italic* and typification of *Aulacoseira* (Bacillariophyta). *Diatom Research.* 2003; 18:1–19.
75. Navok T, Guillory WX, Julius ML, Theriort EC, Alverson AJ. Towards a phylogenetic classification of species belonging to the diatom genus *Cyclotella* (Bacillariophyceae): Transfer of species formerly placed in *Puncticulata*, *Handmannia*, *Pliocaenicus* and *Cyclotella* to the genus *Lindavia*. *Phytotaxa.* 2015; 217:249–264.
76. Ter Braak, C.J.F., Šmilauer, P. CANOCO reference manual and CanoDraw for Windows user's guide: software for canonical community ordination (version 4.5). Microcomputer Power; 2002.
77. Hill MO, Gauch HG. Detrended correspondence analysis: An improved ordination technique. *Plant Ecol.* 1980; 42:47–58.
78. Ter Braak C.J.F, Prentice IC. A theory of gradient analysis. *Adv Ecol Res.* 1988; 18:271–317.
79. Ter Braak C.J.F. Canonical Correspondence Analysis: A new eigenvector technique for multivariate direct gradient analysis. *Ecology.* 1986; 67:1167–1179.
80. Smol JP, et al. Climate-driven regime shifts in the biological communities of arctic lakes. *Proc Natl Acad Sci USA.* 2005; 102:4397–4402. [PubMed: 15738395]
81. Birks, HJB., Gordon, AD. Numerical methods in Quaternary Pollen Analysis. Academic Press; 1985.
82. Juggins, S. ZONE software, version 1.2. University of Newcastle; 1985.
83. Bennett KD. Determination of the number of zones in a biostratigraphical sequence. *New Phytologist.* 1996; 132:155–170.
84. Ryves DB, Juggins S, Fritz SC, Battarbee RW. Experimental diatom dissolution and the quantification of microfossil preservation in sediments. *Palaeogeogr Palaeoclimatol Palaeoecol.* 2001; 172:99–113.
85. Candy I, et al. The evolution of Palaeolake Flixton and the environmental context of Starr Carr: an oxygen and carbon isotopic record of environmental change for the early Holocene. *Proc Geol Assoc.* 2015; 126:60–71.
86. Domínguez-Rodrigo, M., Barba, R., De la Torre, I., Mora, R. Deconstructing Olduvai: A Taphonomic Study of the Bed I Sites. Domínguez-Rodrigo, M.Barba, R., Egeland, CP., editors. New York: Springer; 2007. p. 101-125.
87. Bunn, HT. Meat-eating and human evolution: Studies on the diet and subsistence patterns of Plio-Pleistocene hominids in East Africa. University of Wisconsin; Madison: 1982. Unpublished PhD thesis
88. Bunn HT, Kroll EM. Systematic butchery by Pilo/Pleistocene Hominids at Olduvai Gorge, Tanzania. *Curr Anthropol.* 1986; 27:431–452.
89. Binford, LR. Faunal remains from Klasies River Mouth. Academic Press; 1984.
90. Andrews P, Cook J. Natural modifications to bones in a temperature setting. *Man.* 1985; 20:675–691.

91. Blumenschine RJ, Selvaggio MM. Percussion marks on bone surfaces as a new diagnostic of hominid behaviour. *Nature*. 1988; 333:763–765.
92. Fisher JW. Bone Surface modifications in zooarchaeology. *J Archaeol Method Theory*. 1995; 2:7–68.
93. Noe-Nygaard N. Man-made trace fossils on bones. *J Hum Evol*. 1989; 4:461–461.
94. Binford, LR. *Bones: Ancient Men and Modern Myths*. Academic Press; 1981.
95. Stiner M, Kuhn S, Weiner S, Bar-Yosef O. Differential burning, recrystallization, and fragmentation of archaeological bone. *J Archaeol Sci*. 1995; 22:223–237.
96. Shipman P, Foster G, Schoeninger M. Bunt bones and teeth: an experimental study of color, morphology, crystal structure and shrinkage. *J Archaeol Sci*. 1984; 11:307–325.
97. Tong HW, Zhang S, Chen F, Li Q. Rongements sélectifs des os par les porcs-épics et autres rongeurs : cas de la grotte Tianyuan, un site avec des restes humains fossiles récemment découvert près de Zhoukoudian (Choukoutien). *Anthropologie*. 2008; 112:353–369.
98. Dart RA. Bone tools and Porcupine gnawing. *Am Anthropol*. 1958; 60:715–724.
99. Behrensmeyer AK. Taphonomic and ecological information from bone weathering. *Paleobiology*. 1978; 4:150–162.
100. Behrensmeyer AK, Gordon K, Yanagi GT. Trampling as a cause of bone surface damage and pseudo-cutmarks. *Nature*. 1986; 319:402–403. [PubMed: 2418361]
101. Villa P, Mahieu E. Breakage pattern of human long bones. *J Hum Evol*. 1991; 21:27–48.
102. Bunn, HT. *Animals and Archaeology, Volume 1*. Clutton-Brock, J., Grigson, C., editors. Vol. 1977. Oxford: BAR International; 1983. p. 143-148.163
103. Scerri EML, Groucutt HS, Jennings RP, Petraglia MD. Unexpected technological heterogeneity in northern Arabia indicates complex Late Pleistocene demography at the gateway to Asia. *J Hum Evol*. 2014; 75:125–142. [PubMed: 25110207]
104. Scerri EML, Gravina B, Blinkhorn J, Delagnes A. Can lithic attribute analyses identify discrete reduction trajectories? A quantitative study using refitted lithic sets. *J Arch Method Theory*. 2016; 23:669–691.
105. Groucutt HS, et al. Late Pleistocene lakeshore settlement in northern Arabia: Middle Palaeolithic technology from Jebel Katefeh, Jubba. *Quatern Int*. 2016; 382:215–236.




Figure 1. Al Wusta location, map of site and stratigraphy.

A: The location of Al Wusta and other key MIS 5 sites in the region 11; B: Al Wusta digital elevation model showing location of AW-1 phalanx, marl beds, lithics and vertebrate fossils, and the locations of the trenches and sections. The inset shows a satellite image of the site; C: Stratigraphic log of Al Wusta showing the sedimentology of the exposed carbonate beds, isotopic values, OSL ages for sand beds and U-series and ESR ages for AW-1 and WU-1601. Sands are shown in yellow: lower massive sands are aeolian (Unit 1), upper laminated sands are waterlain (Unit 3a) and have been locally winnowed to generate a coarse desert pavement (Unit 3b), lacustrine marls are shown (Unit 2) in grey (for full key and description see Supplementary Figures 13 and 14 and Supplementary Information 5).

Section PD40 is shown as it contains the thickest sequence and is most representative of Al Wusta, chronometric age estimates (marked *) from the site are depicted in their relative stratigraphic position, see Supplementary Figure 14 for their absolute positions.




Figure 2. Photographs and micro-CT scans of Al Wusta-1 *Homo sapiens* phalanx.

A: photographs in (left column, top to bottom) distal, palmar and proximal views, and (middle row, left to right) lateral 1, dorsal and lateral 2 views. Micro-CT cross-sections (illustrated at 2x magnification) include B (54% from proximal end) and C (illustrating abnormal bone).




Figure 3. Scatterplot of the first two principal components (PC) scores of the geometric morphometric analysis of the Al Wusta-1 phalanx compared with a sample of primates, including hominins.
 Non-human hominoids: lilac; *Gorilla*: circles, *Pan*: triangles. Cercopithecoids: red; *Colobus*: triangles, *Mandrillus*: squares, *Papio*: circles. Neanderthals: blue diamonds. *H. sapiens*: green; early *H. sapiens*: circles, Holocene *H. sapiens*: squares. Al Wusta-1: black star, circled in red.




Figure 4. Scatterplot of the first two principal component (PC) scores from the geometric morphometric analyses of AW-1 and sample of comparative hominin 2nd, 3rd, and 4th intermediate phalanges.

Wireframes show mean configuration warped to extremes of PC axes in dorsal (left), proximal (middle) and lateral (right) views. Convex hulls added post-hoc to aid visualisation.




Figure 5. Selected Al Wusta lithic artefacts.

A: argillaceous quartzite flake; B: quartz hammerstone; C: ferruginous quartzite Levallois flake; D: chert Levallois flake; E: Quartz recurrent centripetal Levallois core; F: quartzite preferential Levallois core with centripetal preparation and pointed preferential removal.




Figure 6. The chronological and climatic context of Al Wusta.

The Al Wusta lake phase falls chronologically at the end of the time-range of MIS 5 sites from the Mediterranean woodland of the Levant (~130-90 ka) and earlier than the late dispersal(s) (~60-50 ka) as posited in particular by genetic studies. The chronology of these dispersals and occupations correspond with periods of orbitally modulated humid phases in the eastern Mediterranean³⁶ that are important intervals for human dispersals into Eurasia, and are also proposed to correspond with episodes of monsoon driven humidity in the Negev and Arabian desert³⁴. Environmental amelioration of the Saharo-Arabian belt, therefore, appears to be crucial for allowing occupation at key sites that document dispersal out of Africa. A: East Mediterranean speleothem $\delta^{18}\text{O}$ record from Soreq and Pequin Caves³⁶; B: global $\delta^{18}\text{O}$ record³⁷; C: Insolation at 30 degrees north³⁸, showing the temporal position of

key sites relating to dispersal out of Africa^{2,3,11,48}. The chronology for Al Wusta shows the phases defined by the Bayesian model at 2σ .