

RESEARCH ARTICLE

Physiotherapy and related management for childhood obesity: A systematic scoping review

Kim Truong¹, Sandra Park¹, Margarita D. Tsiros², Nikki Milne^{1*}

1 Faculty of Health Sciences and Medicine, Bond Institute of Health and Sport, Bond University, Gold Coast, Queensland, Australia, **2** UniSA Allied Health and Human Performance, Alliance for Research in Exercise, Nutrition and Activity, University of South Australia, Adelaide, South Australia, Australia

* nmilne@bond.edu.au



Abstract

Introduction

Despite targeted efforts globally to address childhood overweight/obesity, it remains poorly understood and challenging to manage. Physiotherapists have the potential to manage children with obesity as they are experts in movement and physical activity. However, their role remains unclear due to a lack of physiotherapy-specific guidelines. This scoping review aims to explore existing literature, critically appraising and synthesising findings to guide physiotherapists in the evidence-based management of childhood overweight/obesity.

Method

A scoping review was conducted, including literature up to May 2020. A review protocol exists on Open Science Framework at <https://osf.io/fap8g/>. Four databases were accessed including PubMed, Embase, CINAHL, Medline via OVID, with grey literature searched through google via "file:pdf". A descriptive synthesis was undertaken to explore the impact of existing interventions and their efficacy.

Results

From the initial capture of 1871 articles, 263 intervention-based articles were included. Interventions included qualitative focused physical activity, quantitative focused physical activity and multicomponent interventions. Various outcome measures were utilised including health-, performance- and behaviour-related outcomes. The general trend for physiotherapy involvement with children who are obese appears to favour: 1) multicomponent interventions, implementing more than one component with environmental modification and parental involvement and 2) quantitative physical activity interventions, focusing on the quantity of bodily movement. These approaches most consistently demonstrated desirable changes across behavioural and health-related outcome measures for multicomponent and quantitative physical activity interventions respectively.

OPEN ACCESS

Citation: Truong K, Park S, Tsiros MD, Milne N (2021) Physiotherapy and related management for childhood obesity: A systematic scoping review. PLoS ONE 16(6): e0252572. <https://doi.org/10.1371/journal.pone.0252572>

Editor: Inmaculada Riquelme, Universitat de les Illes Balears, SPAIN

Received: June 7, 2020

Accepted: May 18, 2021

Published: June 14, 2021

Copyright: © 2021 Truong et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its [Supporting information files](#).

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Conclusion

When managing children with obesity, physiotherapists should consider multicomponent approaches and increasing the quantity of physical activity, given consistent improvements in various obesity-related outcomes. Such approaches are well suited to the scope of physiotherapists and their expertise in physical activity prescription for the management of childhood obesity. Future research should examine the effect of motor skill interventions and consider the role of environmental modification/parental involvement as factors contributing to intervention success.

Introduction

Over 340 million children worldwide are classified as overweight or obese [1]. There are no reports in the empirical literature to demonstrate that any country has been successful with significantly decreasing obesity rates in the last three decades [2].

Obesity is strongly related to both short- and long-term co-morbidities [3]. Common conditions associated with obesity include type 2 diabetes mellitus, hypertension, early onset of puberty, menstrual irregularities, polycystic ovarian syndrome, steatohepatitis, sleep apnoea, asthma, benign intracranial hypertension, musculoskeletal disorders and psychological problems [3]. If obesity is not addressed appropriately during childhood, it can lead to further comorbidities such as a greater risk of health complications in adulthood [4] and increased cardiovascular-related mortality [5–7]. Hence, early interventions for preventing and managing obesity in childhood are important in the prevention of further chronic disease, morbidity or mortality.

Childhood obesity interventions are commonly based on the concept of energy balance [8]. Energy balance is equivalent to energy intake, minus energy expenditure. Based on this theory, if energy intake is greater than energy expenditure, it results in excess adiposity. Currently, two common approaches to obesity management involve either, or a combination of both; decreasing energy intake through nutritional education and healthy eating, or increasing energy expenditure through physical activity [8]. Conversely, Flatt et al. highlight that obesity management is not simplistic but rather multifactorial in nature [9]. Childhood obesity may be influenced by age, gender, genetics, psychological and environmental factors such as school policies, parent's work-related demands and lifestyle [10].

Due to the complexity of obesity, a multicomponent behavioural intervention carried out by a multidisciplinary team is considered best practice and has been shown to be effective [11–15]. Furthermore, Rajjo et al. [16] suggested that physical activity, behavioural, pharmacological, dietary and educational components, with an early intervention approach have a positive impact on weight and healthy lifestyle behaviours. However, the nature of these research-based interventions has limitations regarding the generalisability to wider populations in real-life contexts [16]. The interventions are usually tested in unblinded trials [16]. Additionally, many interventions being used clinically to address child obesity in real world settings are multifaceted and reactive to individual needs. Therefore, it is challenging to replicate these interventions in research in order to acquire evidence to support practice guidelines [16].

Some evidence suggests that childhood obesity may be successfully treated with enhanced fundamental movement skills, motor coordination and physical activity [17]. This field of

research is commonly based on the assumption that children with well-developed fundamental motor skills are more likely to engage in high levels of physical activity than those with poorly developed functional motor skills [17]. Since children and adolescents with obesity have lower coordination, balance, speed, agility and fine and gross motor skills compared to their healthy-weight peers [17], they are often unable to meet the physical activity recommendations and unable to reap the benefits that physical activity offers [18–20]. Physiotherapists play a major role in improving functional motor skills and enhancing physical activity in children with overweight or obesity [21]. However, no previous reviews have explored physiotherapy specific interventions for managing or preventing child obesity. Physiotherapy interventions may be focused on increasing participation in physical activities or improving the quality of movement in physical activity.

In a recently published ‘call-to-action’ for physical therapists, Tsiros and Shultz [22], proposed “Ten Action Points” whereby physiotherapists are encouraged to be aware of physical activity and healthy eating guidelines to set measurable goals around family-level lifestyle behaviours instead of making weight the sole focus of assessment and intervention [22]. Additionally, the National Health and Medical Research Council (NHMRC) recommend that abnormal gait; problems with feet, hips and knees; difficulties with balance and coordination; and hip and knee joint pain should be appropriately managed [14,15]. Physiotherapists have a unique skillset to ensure these aspects are addressed through tailored physical activity programs appropriate to the individual child’s needs (e.g. play-based and family-centred), whilst ensuring injury prevention [20,23].

Despite physiotherapists being identified as health professionals with appropriate skills and knowledge to suitably care for children with obesity, a cross-sectional survey of Australian physiotherapy practice trends and professional needs, found that only half of physiotherapists, who provide care for children, provided services specifically to children with overweight or obesity, with ‘lack of service prioritisation and resources to support their non-acute care’ as the main reasons for not providing services [24]. The findings from this same study demonstrated that just under half of physiotherapists were assessing the motor skills of children who were overweight or obese and this was attributed to not having enough time [24] which may have led to unidentified needs for intervention or not having a physiotherapy-specific evidence-based guideline to support their intervention choices. This suggests that despite their appropriate training and skill set, physiotherapists are currently providing little input into the management of children who are overweight or obese for a variety of reasons. Instead, there is a common trend for other professionals including physical education specialists, exercise physiologists and school nurses, to be implementing interventions that are within the scope of physiotherapists [25] even though physiotherapists have a unique skill set in assessing the underlying reasons for lack of physical activity in children [21].

There are gaps in the literature regarding appropriate and universal physiotherapy protocols for management of childhood obesity. Previous research regarding physiotherapy intervention for managing children with obesity has concluded with recommendations to develop evidence-based guidelines to assist physiotherapists in the development of effective interventions [24]. However, to our knowledge, no prior reviews have examined available literature in the context of physiotherapy practice. Hence, this scoping review aims to (i) explore and critically appraise current evidence regarding physiotherapy and related interventions to manage childhood obesity and; (ii) broadly synthesise the findings of articles regarding interventions to guide physiotherapists in evidence-based management of childhood obesity.

Methods

This scoping review is reported using the PRISMA Extension for Scoping Reviews (S1 Table) [26]. A review protocol exists on Open Science Framework and can be viewed at <https://osf.io/fap8g/>.

Search strategy

The search was initially undertaken on 26th August 2019 and repeated on 23rd May 2020. A combination of search terms was used (S2 Table). Searching of literature occurred in the following five databases: PubMed, Embase, CINAHL, and Medline via OVID. Grey literature was identified with Google searches ending in “file:pdf” and expert referral.

Study selection

After duplicate articles were removed, two authors SP and KT independently conducted title and abstract screening and identified potentially relevant articles for full-text review. A process of consensus for included articles at full text occurred between the two authors during face-to-face meetings. Outstanding disagreements between the two authors were resolved by a third researcher (NM), in collaboration with SP and KT to determine the list of possibly relevant articles to examine at full text.

Studies that appeared to meet the inclusion criteria based on title and abstract screening were retrieved in full text and inclusion/exclusion criteria were applied at full text level. A final consensus between the two authors (SP and KT) was achieved on included articles through face-to-face discussions. Discrepancies were resolved by a third researcher (NM) to achieve a final consensus on included full text articles.

Inclusion/Exclusion criteria

Articles were screened using the below criteria (Table 1).

Critical appraisal tools

Critical appraisal of each included publication was undertaken by two reviewers (SP and KT) independently. A Kappa statistic was calculated to determine the level of agreement in critical

Table 1. Inclusion and exclusion criteria.

| Inclusion | Exclusion |
|--|--|
| <ul style="list-style-type: none"> • Infants <2, children 2–12 years, adolescents 13–18 years who are overweight/obese. • Management/intervention used in article may be physical, behavioural, nutritional or educational in nature (or a combination of all). • Intervention used in article must be delivered by a physiotherapist OR able to be delivered by a physiotherapist according to the scope of physiotherapy (e.g., school-based physical education programs, physical activity programs prescribed by exercise physiologists). • Article published in English language only. • Interventional studies, reviews, protocols, policies, procedures, guidelines, recommendations, position statements or perspectives reporting on obesity management/intervention. | <ul style="list-style-type: none"> • Non-human studies • Pregnancy-related obesity literature (e.g., intervention focused on mother during pregnancy). • Studies where no interventions are provided (e.g., profiling studies, exploring trajectories or correlation/cross-sectional studies, protocol only). • No accompanying full texts made available (e.g., abstract or poster only, unable to source). |

<https://doi.org/10.1371/journal.pone.0252572.t001>

appraisal scoring between the two reviewers. After independently scoring, a process of consensus occurred between the two authors (SP and KT) during face-to-face meetings.

Three appraisal tools were used to assess the quality of included studies/articles, selected according to the relevant study designs. The Mixed Methods Appraisal Tool (MMAT) was used to assess the quality of quantitative and qualitative studies [27]. It has been demonstrated as a tool used to appraise the quality of empirical studies including primary research, based on experiments or observations [28,29]. It appraises quality of five categories: qualitative research, randomised control trials, non-randomised studies, quantitative descriptive studies and mixed-methods studies [27]. The ratings better inform the methodological quality of the included studies to assist with decision making in data synthesis. Criteria were assessed using the following: 'yes' scoring as '1' and 'no' or 'can't tell' scoring as '0', with possible total scores ranging from 0–7. Only studies that achieved a score of five or more out of seven (Critical Appraisal Score (CAS) of $\geq 71\%$) on the MMAT were considered to have strong methodological quality and were included in the descriptive synthesis (see 'Data extraction and synthesis'). This method is consistent with a previous utilisation of the MMAT in a high-quality systematic review [30].

The iCAHE Guideline Quality Checklist was used to assess the quality of guidelines and included grey literature [31]. This tool measures methodological quality of guidelines across six domains: availability, dates, underlying evidence, guideline developers, guideline purpose and users, ease of use. Criteria was assessed using dichotomous responses with 'yes' scoring as '1' and 'no' scoring as '0', with possible total scores ranging from 0–14.

The Joanna Briggs Institute Checklist for Systematic Reviews and Research Syntheses was used to assess the quality of systematic reviews [32]. This tool allows assessment of the methodological quality and the extent to which a study has addressed the possibility of bias in its design, conduct and analysis. Criteria was assessed using the following response options: 'yes' scoring as '1' and 'no' or 'unclear' scoring as '0', with possible total scores ranging from 0–11.

To standardise scoring across multiple critical appraisal tools, after each article was critically appraised with the appropriate tool, a critical appraisal score (CAS) was calculated as a percentage of 'yes' responses over the total possible score, revealing a consistent percentage CAS for all included articles. Articles achieving a CAS of $\geq 71\%$ were considered to have high methodological quality.

Data extraction and synthesis

Data extraction was completed by two authors SP and KT independently using a standardised data extraction form to collect relevant information. This included: first author, year of publication, country of origin, aims/purpose, study population and sample size, methodology (including study design and statistical models used), intervention type, comparator, duration of intervention, outcomes, key findings related to the aims of this scoping review. Discrepancies in data extraction were resolved through a process of consensus using face-to-face meetings after independent data extraction was completed.

All included studies were narratively reported according to study type including guidelines, reviews and clinical trials by authors KT and SP. Clinical trials were explored under presenting key themes highlighting the most predominant form of interventions. The intervention themes were then described within the International Classification of Functioning, Disability and Health—Child and Youth Version (ICF-CY) model [33]. As a classification system recognised by the World Health Organisation (WHO), the ICF-CY uses a universal language for health professionals and acknowledges health and disability from a multidimensional perspective [33]. Studies mapped to the ICF-CY domains were clinical trials of high quality, with a

score of five or more out of seven on the MMAT [30]. Clinical trial interventions were categorised into the taxonomy within the existing ICF-CY model. This included “Body Function and Structure Impairments”, “Activity Limitations”, “Participation Restrictions”, “Environmental Factors”, “Personal Factors” [33]. Multicomponent interventions were replicated amongst applicable categories to demonstrate the overall distribution of intervention modes within the ICF-CY model. The same procedure was repeated to demonstrate the distribution of outcome measures utilised in the included studies.

A descriptive synthesis was conducted to explore major trends regarding intervention types for children with overweight/obesity, considered to be within the scope of physiotherapy practice. Clinical trials were reviewed to identify the most predominant form of intervention and classified in a descriptive synthesis accordingly under “quantitative focused physical activity”, “qualitative focused physical activity” and “multicomponent interventions”. “Quantitative focused physical activity” is defined as the quantity of any bodily movement produced by skeletal muscles, focused on maximal energy expenditure [34]. “Qualitative focused physical activity” is defined as improving quality of movement using planned and structured bodily movement, focused on the development of motor skills [34]. “Multicomponent interventions” have a physical activity intervention with more than one other intervention implemented including diet/nutritional education, healthy lifestyle education and/or addressing environmental factors [11–13]. The outcome measures were reported according to the ICF-CY model including “body structure and function”, “activity limitation”, “personal factors” and “environmental factors”. Outcomes mapped to “body structure and function” were further divided into two categories: health-related physical fitness and performance-related physical fitness measures. For example, health-related physical fitness measures are related to measures of good health including: (i) anthropometry, (ii) cardiovascular/cardiorespiratory fitness, and (iii) blood serum analyses [34]. Performance-related fitness measures are related to measures of enhanced motor proficiency including: (i) coordination, (ii) balance, (iii) speed, (iv) agility, (v) strength, (vi) flexibility, (vii) power, and (viii) posture and gait [34,35]. Outcome measures mapped to “activity limitation” were associated with measuring difficulties an individual may have in the execution of a task or action including physical activity-, sedentary behaviour- and sleep-related outcomes [33]. Outcomes mapped to “personal factors” were behavioural measures [33]. An example of this included readiness for change assessment. Outcomes mapped to “environmental factors” were associated with measuring the physical, social and attitudinal environment in which the individual lives and conducts their daily living including parental engagement and parental satisfaction outcomes [33].

Two stages of coding were undertaken during the descriptive synthesis. Firstly, individual investigations were coded based on whether the effect was positive/desirable (+), negative/undesirable (-) or no effect (0) (i.e., a decrease in waist circumference for children with overweight or obesity is a desirable direction of change, whereas an increase in waist circumference is an undesirable direction of change, and an insignificant change in waist circumference is deemed to have no effect). The second stage of coding was then applied whereby, if at least two thirds (66.6%) of the investigations were reported to have significant results in the same direction of change, then the overall result was considered consistent and the appropriate code was applied [36]. Specifically, if 66.6% or more of the studies included in the descriptive synthesis for the specific intervention mode demonstrated a significant positive or negative change, or no significant change, then summary codes of desirable (D), undesirable (U) or no effect (0) were given respectively. If more than one investigation and less than five investigations were undertaken assessing the impact of an intervention on outcomes for children with obesity or the above criteria was not met, then it was coded as questionable (?).

The total number of studies (n_s) and the total number of participants (n_p) included to determine the summary code which described the impact of the interventions on outcomes for children with obesity, were calculated.

Sensitivity analysis

A post-hoc sensitivity analysis was conducted to determine if the findings of this scoping review would change when we explored the results with all overweight/obesity 'prevention' studies removed from the analysis (i.e., only leaving overweight/obesity 'management' interventions). Management of childhood obesity interventions is specifically inclusive of: (i) a population that is overweight or obese as the intervention group; and (ii) aims to improve measures of health, performance- and behaviour-related outcomes of the targeted population. All included studies not fulfilling these criteria were considered to be preventative studies and were excluded in the sensitivity analysis. The descriptive synthesis method for analysis of data was repeated with the included studies for outcome measures with five or more investigations, as per the original descriptive analysis. When less than five investigations were undertaken, exploring the impact of intervention on a given outcome, they were not included in the sensitivity analysis, as it does not fulfil the criteria for descriptive analysis outlined above.

Results

Study selection

The literature search yielded 1871 titles and abstracts for review. From these, 519 potentially relevant articles were selected for full text review. Two hundred and sixty-three of the full text articles were identified as meeting the inclusion criteria for the present scoping review. This included 14 guidelines, 30 reviews (26 systematic and 4 umbrella) and 219 clinical trials (Fig 1).

Methodological quality assessment

The percentage agreement between reviewers was moderate with 75.50%. Cohen's Kappa analysis confirmed moderate agreement between the two raters, $k = 0.44$ ($p < 0.001$). One hundred percent agreement was achieved on all papers during the consensus process and this was the final CAS reported. One hundred and fifty-eight articles achieved a CAS of $\geq 71\%$ and were considered high quality. One hundred and five articles were graded $< 71\%$. The mean CAS (percentage score) for methodological quality of the included articles was 68.07%.

Guidelines

Fourteen of the 263 articles were guidelines for the management of children with obesity [14,15,21,37–47] as summarised in Fig 2 and S3 Table. The content included in the guidelines covered clinical recommendations and health promotion strategies including the importance of physical activity management.

Reviews

Thirty of the 263 articles were reviews [48–77] as summarised in Fig 2 and S4 Table. Twenty-six were systematic reviews and four were umbrella reviews. Of the 26 systematic reviews included, a total of 556 studies were captured. Thirty-one (14.16%) of the 219 clinical trials included in this scoping review were found to overlap with the 556 studies identified in these systematic reviews. Nine (4.11%) of the 219 clinical trials were included in two systematic reviews [84–92]. Four (1.83%) of the 219 clinical trials were included in three or more




Fig 1. PRISMA flow diagram. Processes for study screening, eligibility and inclusion [26].

<https://doi.org/10.1371/journal.pone.0252572.g001>

systematic reviews [87,88,93,94]. The four umbrella reviews (systematic reviews of systematic reviews) included in this scoping review, captured a total of 86 systematic reviews. Nine (34.62%) of the 26 systematic reviews included in this scoping review were found to overlap with 86 systematic reviews identified in the umbrella reviews. No systematic reviews were duplicated amongst the four umbrella reviews.

14 Guidelines

Published: 2006-2018

Regions: US [38-40], Au [37, 89, 90], Ca [41], Europe [91], UK [92-94], World organisations [21, 46, 47]

Age groups: All targeted children/adolescents and two guidelines also targeted adults [89, 90]

Content/target group: directed towards physiotherapists [21, 37-39, 41, 92], targeted health professionals [21, 38, 39, 41, 89, 90, 92, 94], informative for the general public [37, 46, 91], general health/physical activity guidance [40, 46, 47, 89-91, 93, 94].

30 Reviews (26 SRs, 4 URs)

Published: 2005-2020

Regions: Au [48, 52, 55, 57, 64, 69, 72], Brazil [63, 68, 73, 75], USA [49-51, 58, 66, 67, 70, 71, 74, 76], Belgium [53], Chile [54, 61, 62], Netherlands [56], China [59], Spain [60], Canada [65], Turkey [77].

Age groups: 0-19 years

Interventions: school-based nutrition/physical activity [49, 51-53, 58, 59, 66, 74], exercise only [48, 55, 61, 62], HIIT [54, 60], weight loss [63, 71, 72], FMS [56, 57, 64], physical activity [65, 69, 73], active video games [68, 75], parental involvement [50, 67, 76], preventative school-based interventions [56, 70, 77].

Outcomes: improved anthropometry [49, 54-56, 58, 61, 73], improved physical or sedentary activity [52, 57-59].

Future research needs identified: longer f/u durations needed [53, 68], further exploration of specific components [50, 58, 63, 64, 71, 74], higher quality studies needed [59, 61, 65, 66].

Fig 2. Summary of secondary research. Citations are included in square brackets [21,37-41,46-77,78-83]. Refer to S3 and S4 Tables for detailed information for Guidelines and Reviews respectively. Abbreviations: Au—Australia, Ca—Canada, SRs—systematic reviews, UK—United Kingdom, URs—Umbrella reviews, US—United States.

<https://doi.org/10.1371/journal.pone.0252572.g002>

Clinical trials

Two hundred and nineteen of the 263 included articles were clinical trials [84-94,95-302] as summarised in Table 2 and S5 Table. Clinical trial articles were published between 1970 to 2020 and included from three to 17066 participants aged two to 19 years. The sum of all participants from the 219 included articles was 112,524. Clinical trials included childhood obesity prevention (50%) and management (50%) interventions. Intervention studies included non-randomised (56%) and randomised trials (44%) clinical trials. Clinical trials included in the present scoping review were focused on increasing physical activity (n = 36) [114,119,133,136,146,148,149,153,168,172,188-191,205,207,208,210,214,239,240,242,244,246-248,256,263,271,272,274,276,281,283,286,294]; increasing physical activity and improving diet or nutritional education (n = 19) [93,98,127,129,150,162,164,218,222,229,251,254,258,260,267-269,277,288]; increasing physical activity and offering healthy lifestyle education (n = 13) [84,90,120,160,161,182,204,219,238,241,253,279,293]; enhancing posture and gait (n = 3) [173,213,270]; improving targeted motor skills (n = 7) [107,135,142,209,216,235,249]; increasing aerobic exercise (n = 26) [88,97,101,104,125,140,144,165,166,169,181,183,184,193,197,199,223,224,228,230,236,257,265,285,292,300]; increasing strength exercise (n = 4) [203,250,261,275,292]; increasing strength and aerobic exercise combined (n = 13) [109,123,138,143,158,175,196,202,215,262,264,290,298]; reducing sedentary behaviour (n = 1) [167]; technology-based interventions, including exergaming and mHealth (n = 9) [85,132,145,154,157,195,212,226,234]; multicomponent and multidisciplinary (n = 74) [89,92,94-96,99,100,102,103,105,108,111-113,115-118,121,122,124,128,134,137,139,141,147,151,152,155,156,159,163,170,171,176-180,185-187,192,198,200,201,206,211,216,217,220,221,225,227,231-233,237,243,245,252,255,266,269,273,278,280,282,284,287,289,291,296]; family-based and/or environmental interventions (n = 88) [86,90-92,95,96,98-100,105,108,110-112,114-119,123,127-131,139,141,142,145,149,150,155,156,160-162,170,172,174,176,179,

Table 2. Summary of interventions and outcome measures utilised in the 219 included clinical trials.

| No. of trials & focus | Intervention | Outcome measures |
|--------------------------------------|--|---|
| 36 PA | No./duration of PE classes [146,153,188–191,210,239,244,246,247,256,263,274,276,283], recess/related [119,136,146,148,153,172,207,248,256,281], classroom PA [114,119,189,205,210,240,244,246,256], walking/jogging [114,148,149,168,208,242,286], exercise training [214,239,248,256,271,272,294], adventure education [119], active commuting to/from school [283], family-based and/or environmental interventions [114,119,149,172,188–190,207,247,248,283]. | Anthropometric [114,119,133,136,146,148,153,168,173,188,189,191,205,207,208,210,239,242,244,247,248,263,271,272,274,276,281,294], CV/CR [133,136,168,189,205,207,210,239,242,246,248,263,271,272,274,276,281,286], blood serum analyses [133,189,207,239,248,271,274,281,294], physical performance [189,190,214,263,272], other [114,117,146,149,189,210,214,240,244,246,247,256,263,274,283]. |
| 19 PA + diet/nutrition education | Nutrition education [93,98,127,129,150,218,251,254,260,268], dietary counselling [258,277], dietary changes/restrictions [98,150,222,229,267,277,288], cooking classes [127,162,218], provision of snacks/meals [127,162,164,220,260], family-based and/or environmental interventions [98,127,129,150,218,254,258]. | Anthropometric [93,98,127,129,150,162,164,218,222,229,251,254,260,267,268,277,288], CV/CR [93,127,150,162,164,222,229,251,260,267,288], blood serum analyses [93,129,222,251,258,260,277], physical performance [129,218,229,258], other [98,129,251,254,260,268]. |
| 9 Technology based | Exergaming [132,154,157,212,234], mobile-phone apps [145,195,226], website-based education [85]. Some studies included nutrition and/or healthy lifestyle education [85,145,195,226], family-based interventions [85,145,195]. | Anthropometric [85,132,145,195,226], other [85,132,145,154,157,195,212,226,234]. |
| 1 Sedentary behaviour | "Switch Off Get Active", health education intervention aimed to increase physical activity at the expense of screen time [167]. | Anthropometric, CV/CR, PA/SED, PA self-efficacy [167]. |
| 13 PA & health lifestyle education | Lifestyle education (c-CRTD) e.g., PA/screen time [84,90,120,160,161,182,204,219,238,241,253,279,293], and assistance with academic areas [293]. Family-based and/or environmental interventions [90,160,182,219,241,253]. | Anthropometric [84,90,120,204,219,238,241,253,279,293], CR/CV [120,204,238,293], other [84,160,161,182,204,241,253]. |
| 3 Posture & gait | Movement quality/games [213], lower extremity neuromuscular & core exercises [173], stretching/flexibility/strengthening/agility/aerobic endurance with "locomotion-emphasis" [270]. Diet/nutrition education with parental involvement + posture/gait intervention [270]. | Anthropometric [270], physical performance [173], other outcomes [173,213,270]. |
| 88 Family Based and/or environmental | Individual/group counselling with a multidisciplinary team member (physiotherapist, dietitian, psychologist, paediatrician) [95,96,111,123,139,141,145,174,225,227,232,253,254,258,270,299]; nutrition and/or healthy lifestyle education [92,96,99,100,108,111,112,114,116–118,123,127,128,130,131,139,142,145,150,156,162,170,172,176,179,180,182,192,195,211,219,225,227,231–233,237,241,243,245,252–255,258,260,262,270,278,282,291,296,299,302]; parents supporting behavioural changes [99,115,145,170,187,192,216,225,231,241,243,248,255,262,278,282,301,302]; parents involved in meetings/support groups [108,291]; exercise programs [129,139,160,176,227,258,299,301,302]; sports [100,237,255]; active commuting [283]; motor development [142]; online resources [145,156,172,195,296]; and/or family events/activities [100,105,127,128,221,253]. Environmental interventions include grants/funding [105,108,127,245,301]; professional development e.g., upskilling physical educators [86,90,96,98,100,105,108,112,114,115,117,127,161,172,176,188,189,207,221,248,261,296]; availability of healthy/unhealthy food [91,99,108,117,139,149,190,217,232,233,247,255,262,296]; changes to physical infrastructure [91,99,108,117,139,149,190,217,232,233,247,255,262,296]; changes to school policy [91,96,108,117,127,172,178,182,232,233,259,289,291,296]; provision of social support [115,116,225]; and community physical activity [172]. | Parental engagement and/or satisfaction [108,210,231,266,268]. |
| 26 Aerobic exercise | High/low intensity interval training [88,97,144,193,224,236,257,300]; walking/jogging [165,169,183,197,228,230,295,297], treadmill or cycle ergometer [101,181,184]; aquatic art board [265]; plyometrics [223,236]; group activities [125]; exercise programs [140,166,199,285,292], aerobic exercise + calorie restrictions [101,199], aerobic exercise + nutrition and/or healthy lifestyle education [125,166,181,199,230,257,285,292]. | Anthropometric [88,97,101,104,125,140,144,165,166,169,181,183,197,199,223,224,230,236,257,265,295,297], CV/CR [88,97,104,144,166,169,181,184,193,197,199,224,230,236,257,265,285,292,295], blood serum analyses [101,166,181,183,197,199,230,236,292,295,297], physical performance [140,141,165,181,193,223,224,28,28,285], other [140,165,166,230]. |
| 4 Strength exercise | Exercise programs [261,275,292], exercises using different equipment (e.g. sleeping mats, tennis balls, volleyballs, basketballs, swiss balls, pool floats, elastic bands) [203,250], strength exercises + stretching program [250], strength exercises + motivational interviewing [275]. Some studies also included nutritional and/or healthy lifestyle education [203,261,292]. | Anthropometric [203,275], CV/CR [203,292], blood serum analyses [203,292], physical performance [203,250], other [250,261,275,292]. |
| 13 Strength & aerobic | Resistance band/weight machines [109,143,262], treadmill/elliptical/cycling [109,123,138,143,175,290,298], exercise program/circuit training [123,138,175,196,202,215,262], walking/running [202,215,298], sport [158,262,290,298]. Some studies also included dietary restriction [123,138,143,158,175,196], nutrition/healthy lifestyle education [123,138,158,262], psychological group sessions [123,138,175], family-based and/or environmental interventions [123,196,262]. | Anthropometric [109,123,138,143,175,196,202,215,262,264,290], CV/CR [143,202,264,290], blood serum analyses [109,123,175,196,202,215,262], physical performance [143,203,262], other outcomes [158,202,262]. |
| 7 Motor skills | Motor development [107,142], GM/FM equipment [107], exercise programs [209,216,235,249], tennis-specific training [135], motor skill development with nutrition education/counselling [209,216]. | Anthropometric [142,209,216,249], CV/CR [135,209,235,249], physical performance [107,209,216,249], other [107,216]. |
| 41 Multicomponent/multidisciplinary | Physiotherapy [102,103,206,225,273,280,284], dietitian/nutritionist [95,108,111,124,139,141,152,170,185,198,200,201,206,216,220,232,243,252,269,280,284], psychology [95,111,122,139,141,152,159,163,171,200,201,243,252,280,291], nurse [115,206,291], paediatrician/physician [95,139,220], increasing PE class intensity/no./duration [89,95,96,99,113,118,137,139,163,176–178,206,212,232,237,289,296], exercise program [94,111,112,115–117,122,124,134,141,147,151,152,155,156,163,170,171,176,179,180,185–187,192,198,201,206,211,216,217,220,221,225,227,231–233,237,243,245,252,255,269,273,278,280,282,284,287,291,296], cooking classes [96,139,186,245,266], provision of snacks/meals [94,103,115,155,156,186,237,266], dietary restriction [94,124,141,147,159,163,199,200,220,269,287], and/or behavioural modification [102,103,111,113,115,116,118,122,139,155,170,187,216,221,225,227,231,243,266,282,287,289,291], family-based and/or environmental interventions [92,95,96,99,100,105,108,111,112,115–118,124,128,139,141,155,156,170,176,178–180,187,198,211,216,217,220,221,225,231–233,237,243,245,252,255,266,278,282,289,291,296]. | Anthropometric [89,92,94–96,99,100,102,103,105,111,115–118,122,124,128,134,137,139,141,147,151,152,155,156,159,163,170,171,178–180,185–187,192,198,200,206,211,216,217,220,227,231–233,237,243,245,252,255,266,269,273,278,280,283,284,287,291,296], CV/CR [92,94,95,99,105,116,122,124,134,137,139,152,154,171,180,186,200,211,217,220,225,227,231,233,243,273,287,289], blood serum analyses [89,92,94,113,147,152,155,163,180,186,198,206,211,284,287], physical performance [94,103,151,171,178,216,233,269,296], other [89,92,94,95,99,100,108,113,121,122,128,139,156,176,177,185,186,200,211,216,217,221,225,227,231–233,237,255,266,278,280,291]. |

Note. See S5 Table for detailed information. Examples of 'other' outcomes include: Physical activity/sedentary behaviours, diet/dietary behaviours, knowledge/attitudes/self-efficacy/self-competence/enjoyment relating to various aspects of lifestyle behaviours, quality of life, general health, perceived ratings of exertion, self-esteem, depression, body image/size, menstrual regularity, alignment during gait/posture loading conditions, foot pressure values and/or pain, trunk alignment, metabolic rate, perceived fitness, sleep, cognitive abilities. Abbreviations: CBT—cognitive behaviour therapy, CV/CR—cardiovascular/cardiorespiratory, FM—fine motor, GM—gross motor, PA—physical activity, PA/SED—physical activity and/or sedentary behaviours, PE—physical education, no.—number.

<https://doi.org/10.1371/journal.pone.0252572.t002>

180,182,187–190,192,195,207,211,216,217,219,221,225,227,231–233,237,241,243,245,247, 248,252–255,258–262,266,270,278,282,283,289,291,296,299,301,302]. Studies that were applicable to multiple themes were replicated within appropriate categories and account for the overlap in the total number of clinical trials. Table 2 provides a summary of the interventions and outcome measures utilised in the clinical trials.

Interventions mapped to the International Classification of Functioning, Disability and Health—Child and Youth Version

From the 219 clinical trials included in this scoping review, 138 clinical trials were considered high-quality studies and were consequently mapped on the ICF-CY. “Activity Limitation” interventions were represented by the greatest number of studies ($n_s = 100$) and “Participation Restriction” interventions represented the least number of studies ($n_s = 26$) (Fig 3). Outcome measures in the “Body Function and Structure Impairment” domain of the ICF-CY were represented by the greatest number of studies ($n_s = 134$) and “Participation Restriction” outcome measures were least represented ($n_s = 0$) (Fig 4).

Descriptive synthesis of results

Four of the eight quantitative focused physical activity intervention types resulted in desirable changes in outcome measures; neither of the two qualitative focused physical activity intervention types resulted in desirable changes in outcome measures; and four of the nine multicomponent intervention types resulted in desirable changes based on outcome measures utilised (Table 3; S6 Table).




Fig 3. ICF-CY model with mapped interventions.

<https://doi.org/10.1371/journal.pone.0252572.g003>




Fig 4. ICF-CY model with mapped outcome measures.

<https://doi.org/10.1371/journal.pone.0252572.g004>

Intervention types with desirable changes in anthropometric outcomes included quantitative focused physical activity such as walking/running ($n_s = 5$, $n_p = 1525$), combined aerobic and resistance exercise ($n_s = 4$, $n_p = 345$), physical education and increased physical activity/exercise ($n_s = 12$, $n_p = 3842$), and intensity training ($n_s = 2$, $n_p = 235$). Interventions demonstrating no change based on anthropometric outcomes included multicomponent interventions such as physical activity with environmental changes ($n_s = 6$, $n_p = 8466$).

Interventions with desirable changes in cardiorespiratory/cardiovascular (CRF/CVF) outcomes included quantitative focused physical activity such as physical education and increased physical activity/exercise ($n_s = 9$, $n_p = 1602$); and multicomponent interventions such as physical activity with diet/nutrition education ($n_s = 6$, $n_p = 372$).

Interventions demonstrating desirable changes in blood serum results included quantitative focused physical activity such as walking/running ($n_s = 2$, $n_p = 90$), combined aerobic and resistance exercise ($n_s = 4$, $n_p = 345$), sports-based activities ($n_s = 1$, $n_p = 30$); and multicomponent interventions including physical activity with diet/nutrition education ($n_s = 4$, $n_p = 181$). Interventions demonstrating nil change on blood serum analysis included quantitative focused physical activity such as physical education and increased physical activity/exercise ($n_s = 1$, $n_p = 137$); and multicomponent interventions such as physical activity with environmental changes ($n_s = 3$, $n_p = 1578$).

Interventions resulting in desirable changes on strength included multicomponent interventions such as physical activity with health lifestyle education and diet/nutrition education ($n_s = 5$, $n_p = 336$), physical activity with diet/nutrition education and environmental changes ($n_s = 5$, $n_p = 3032$), and physical activity with healthy lifestyle education and diet/nutrition education and environmental changes ($n_s = 3$, $n_p = 1442$).

Table 3. Descriptive synthesis of results for studies investigating the effects of child overweight and obesity preventative and treatment strategies.

| | Anthropometric | CRF, CVF | Blood serum analysis | Coordination | Balance | Speed | Agility | Strength | Flexibility | Power | Posture + gait | Physical activity assessment | Sedentary behaviour assessment | Sleep quality/duration | Self-esteem, -efficacy and -perception assessment | Food/drink behaviour and consumption |
|--|----------------|----------|----------------------|--------------|---------|-------|---------|----------|-------------|-------|----------------|------------------------------|--------------------------------|------------------------|---|--------------------------------------|
| Quantitative focused physical activity | | | | | | | | | | | | | | | | |
| Walking/running | D | ? | D | ? | ? | | | ? | | | ? | ? | | | | ? |
| Combined aerobic + resistance exercise | D | ? | D | | | | | ? | | | | | | | | ? |
| Physical education + increased physical activity/exercise | D | D | 0 | ? | | ? | | | ? | | | ? | ? | | ? | |
| Sports-based | ? | ? | D | | | | | | | ? | | | | | | |
| Exergaming | ? | | | | | | | | | | | ? | ? | | | |
| Physical activity in classroom (activity breaks) | ? | ? | | ? | | | | | | | | | | | | |
| Plyometrics | ? | ? | | ? | ? | | | ? | ? | | | | | | | |
| Intensity training | D | ? | | | | | | ? | | | | ? | | | | |
| Qualitative focused physical activity | | | | | | | | | | | | | | | | |
| Motor skills | | | | ? | | | | | | | ? | | | | | |
| Alignment during gait | | | | | | | | ? | | | ? | | | | | |
| Multicomponent intervention | | | | | | | | | | | | | | | | |
| Physical activity + healthy lifestyle education | ? | ? | | ? | ? | | | | | | | ? | ? | | ? | ? |
| Healthy lifestyle education + environment | ? | | | | | | | | | | | ? | ? | | ? | |
| Physical activity + diet/nutrition education | ? | D | D | | | | | ? | ? | | | ? | ? | | ? | |
| Physical activity + environment | 0 | ? | 0 | | | | | | | | | ? | ? | | ? | |
| Diet/nutrition education + healthy lifestyle education + environment | ? | ? | ? | | | | | | | | | ? | ? | | ? | ? |
| Physical activity + healthy lifestyle education + diet/nutrition education | ? | ? | ? | ? | ? | | | D | ? | | | ? | ? | ? | ? | ? |
| Physical activity + healthy lifestyle education + environment | ? | | | | | | | | | | | ? | ? | | ? | ? |
| Physical activity + diet/nutrition education + environment | ? | ? | ? | ? | ? | ? | ? | D | ? | ? | ? | ? | ? | | ? | ? |
| Physical activity + healthy lifestyle education + diet/nutrition education + environment | ? | ? | ? | | ? | ? | ? | D | ? | ? | ? | ? | ? | ? | D | ? |

Key: D Desirable, 0 No effect, U Undesirable, ? Questionable, No data.

<https://doi.org/10.1371/journal.pone.0252572.t003>

Multicomponent interventions such as physical activity with healthy lifestyle education, plus diet/nutrition education and environmental changes showed desirable changes with improving quantified physical activity ($n_s = 21$, $n_p = 28565$), reducing sedentary behaviours ($n_s = 10$, $n_p = 6209$) and improving self-esteem, -efficacy and -perception ($n_s = 6$, $n_p = 2037$).

Sensitivity analysis

A sensitivity analysis was undertaken to determine if studies which were focused on prevention (rather than management) of obesity were influencing the results. Once prevention studies were removed, leaving only overweight and obesity management studies, seven intervention modes showed varied results to the initial descriptive synthesis (S7 Table).

Specifically, interventions focused on increasing physical education and increasing physical activity/exercise, and intensity training changed from a desirable to questionable impact on anthropometric outcomes. Further, interventions focused on physical activity with diet/nutrition education changed from questionable to desirable impact on anthropometric outcomes when prevention studies were removed, and interventions focused on enhancing physical activity with environmental modification varied from no effect to a questionable impact. Interventions focused on physical activity with diet/nutrition education and environmental modification varied from questionable effect to no effect.

When obesity prevention studies were removed from the analysis (leaving only overweight and obesity treatment studies), interventions focused on combining physical activity with healthy lifestyle education and diet/nutrition education were shown to have desirable outcomes on CRF/CVF, which was initially questionable.

Interventions focused on increasing physical education and increased physical activity/exercise; and interventions focused on enhancing physical activity with environmental modification changed from no effect to questionable impact on blood serum analyses. Further, interventions focused on physical activity with diet/nutrition education and environmental changes varied from questionable to desirable changes on blood serum analyses.

Interventions focused on physical activity with diet/nutrition, education and environmental modification; and interventions focused on physical activity with healthy lifestyle education, diet/nutrition education and environmental modification varied from a desirable to questionable impact on strength outcomes.

Interventions focused on physical activity with healthy lifestyle education, diet/nutrition education and environmental modification varied from a desirable to questionable impact on sedentary behaviour outcomes.

Interventions focused on physical activity with healthy lifestyle education, diet/nutrition education and environmental modification varied from a desirable to questionable impact on self-esteem, self-efficacy, and self-perception outcomes.

Discussion

The objective of this scoping review was to explore and critically appraise current evidence regarding physiotherapy and related interventions to manage childhood obesity and to broadly synthesise the findings of articles regarding interventions to guide physiotherapists with evidence-based management of childhood obesity. The key finding from our review suggests that quantitative physical activity facilitates desirable changes in health-related outcomes whereas multicomponent interventions facilitate desirable changes in behaviour measures.

From the 263 articles including guidelines, systematic reviews and clinical trials, the interventions most commonly applied were multicomponent interventions targeting physical activity, diet or nutrition education and lifestyle factors, commonly including environmental

modification with parental involvement. The considerable interest in conducting multicomponent interventions may be linked to guidelines by WHO's global strategy on diet, physical activity and health [303]. Due to WHO's recognition of the existing heavy and growing burden of non-communicable diseases like obesity, governments have been recommended to promote applied research, with intervention programs aimed at improving diet and/or physical inactivity [303]. This increase in awareness and targeted funding provided by governments may have contributed to the mass amount of research relating to multicomponent interventions and obesity.

Multicomponent interventions have demonstrated a lack of improvement in anthropometric measures but desirable changes in physical activity; sedentary behaviour; self-esteem, self-efficacy and self-perception measures. To guide the determination of effectiveness or "success" of this intervention, the National Institute for Health and Clinical Excellence guidelines should be considered [44,45]. These guidelines suggest that childhood obesity interventions should not be "weight focused" but instead include behaviour change strategies to increase physical activity levels or decrease inactivity and improve eating behaviours [44,45]. These guidelines suggest that outcome measures focused on improvements in weight (or body composition) alone may not be the most appropriate way to measure the 'success' of an intervention. As demonstrated by this scoping review, multicomponent interventions target the factors and/or behaviours contributing to obesity and have demonstrated improvements in these areas, despite not achieving positive changes in anthropometric measures. This indicates that multicomponent interventions have achieved goals set by guidelines and therefore, are interventions that may be beneficial in the management of childhood obesity.

The most recent Cochrane reviews, inclusive of only long-term (6 to 24 month) randomised controlled trials, explored childhood obesity management and highlighted aspects of diet, physical activity and behavioural intervention components [11–13]. These Cochrane reviews have demonstrated that multidisciplinary or multicomponent interventions are effective for improving obesity-related outcome measures. In contrast to this scoping review, where multicomponent interventions resulted in a lack of improvement in anthropometric measures, the Cochrane reviews found low quality evidence that these interventions reduce BMI and moderate quality evidence of reduced weight in those who are overweight compared to controls [11–13] and that long-term studies led to desirable anthropometric changes [11–13]. Therefore, one may surmise that multicomponent intervention studies induce positive behavioural changes (as demonstrated in this scoping review) and if maintained over a period of 6 to 24 months may demonstrate positive improvements in anthropometric measures.

Most multi-component interventions included in the present scoping review incorporated environmental modification, as it is widely recognised that the physical environments and surrounding infrastructure influences the risk of childhood obesity [304]. However, the environment is complex as it is influenced by socio-economic status, affordability and accessibility to facilities, location of neighbourhood, attractiveness of the environment, geographical distance of home to central business district areas and city centres [305]. For example, pollution, the weight of a child's school bag, footpath condition, lack of sheltered walkways, distance from home to school, and time constraints to extra-curricular activities before or after school, demonstrate the numerous barriers to the simple act of active commuting to school [305]. An environment where children can, under supervision, have the opportunity to engage in activities with some structure, focusing on skill development, whilst allowing children "to be children" and express themselves spontaneously is crucial [306]. This is not as simple as building new playgrounds, which was a common intervention noted in this scoping review. Interventions in this review inclusive of physical activity and environmental modification failed to demonstrate beneficial changes in outcome measures relevant to overweight or obesity. Purposeful

consideration of social factors is likely needed to observe greater strides within the multi-component interventions. By holding greater awareness of environmental factors and its influence on physical activity levels in children, physiotherapists can achieve greater appreciation of such barriers, decrease stigma against obesity perpetuated through beliefs of a lack of perceived effort, to further individualise interventions for greater success [307].

An important part of a child's environment includes the home environment, which commonly involves parental influences. Parents are considered 'agents of change' and have profound influence on various aspects of a child's life [308]. For example, one of the most influential environments for development of eating behaviours and obesity in children is often determined by what is eaten at home [309]. Food exposure or availability, parental control, attitudes and behaviours around food will influence a child's food intake and therefore, may influence the risk of obesity [310]. Furthermore, parental levels of physical activity are directly associated with the physical activity children receive and are exposed to [311]. Particularly in the earlier years of a child's life, the family serves as the dominant source of social support which is linked to good physical, social, mental health and well-being [312]. Equally, poor family function due to poor communication, family conflict and parenting style (i.e., a non-authoritative, unengaged, permissive) have direct links to obesity [312]. The findings from the present scoping review highlight the breadth of multicomponent interventions that included parental components and similarly demonstrate favourable results where parental components were included. Family-based components are crucial in the management of childhood obesity [313].

Despite the benefits associated with family-based interventions, Edmunds conducted a qualitative exploration highlighting how various parental concerns may act as barriers for optimal childhood obesity management [314]. For example, fear that obesity management interventions may further perpetuate negative stereotypes around their child being overweight/obese within a society that values thinness and stigmatises adiposity [314]. Even acknowledging and "realizing that one is overweight is likely to be stressful and psychologically scarring" [315]. Parents may believe that such interventions may inadvertently increase communication to the child that their body size is undesirable and thus affect mental health negatively [315]. Such beliefs may mean that weight gain could go unnoticed or ignored and thus, reduce the likelihood of parents obtaining professional advice [314]. Contrary to these beliefs and concerns, our results demonstrate the opposite effect. Multicomponent interventions observed positive changes in self-esteem, self-efficacy and perception. Lowry et al. in a study investigating weight management programs on self-esteem, reported that through parental involvement, increased social support, targeting self-esteem and related issues, improved locus of control through education and activities that were fun and engaging [316]. Additionally, Lowry et al. reported that self-esteem improvements were related to weight change, implying that the increase in self-esteem was due to the improvement of physical appearance [316]. This was contrary to findings from this scoping review which suggests that multicomponent interventions can improve self-esteem without requiring a change in anthropometric measures.

Multi-component interventions are further supported by previous NHMRC guidelines [14,15]. Part of the multidisciplinary team includes physiotherapists who have a large role and expertise both in physical activity and exercise but also in the understanding of the complex barriers and perceived barriers, that may affect an individual's ability to make positive lifestyle changes [14,15]. Physiotherapy is constantly evolving and involves a holistic approach to the prevention, diagnosis and therapeutic management of movement difficulties or optimisation of function [317]. As demonstrated by this scoping review, quantitative focused physical activity demonstrates a large space available for exploration by physiotherapists in the treatment of childhood obesity. Physiotherapists play a role in the development, prescription,

implementation, supervision and progression of appropriate physical activity, including cardiovascular training as well as resistance exercise, to increase muscle strength, flexibility, and endurance, and maintain weight loss under safe and controlled conditions [42]. Furthermore, physiotherapists have an understanding of physical function and movement and how to address any impairments in physical function that may be limiting movement and activity participation [81]. Participation in sport and being more active in school are major contributors to increased physical activity, energy expenditure and are recommended interventions [44,45]. Physiotherapists could play an integral role in working alongside physical education teachers, in the development of high-quality physical education programs in schools to facilitate maximal improvements in health and well-being of children with overweight or obesity [21].

Physical activity is widely acknowledged to be a critical factor in the effective management of childhood obesity [318]. As reflected by this scoping review, increases in the quantity of physical activity result in improvements in anthropometric, cardiorespiratory or cardiovascular fitness and blood serum analyses outcomes. These outcome measures are also risk factors for cardiovascular disease in later life [319–324]. Children who are overweight have an increased risk of pre-hypertension by 50% and double or triple the odds of hypertension, compared with children who are healthy weight [325]. Children with obesity have been linked to a 12-fold increase in fasting insulin concentration with abnormal levels of triglycerides, total cholesterol, LDL-c and HDL-c compared to children who are of healthy weight [326]. Therefore, by increasing the quantity or amount of physical activity, the improvements in anthropometric, cardiorespiratory or cardiovascular fitness and blood serum analyses outcomes may be associated with a decrease in weight and may lead to a decreased risk of cardiovascular disease.

Several research gaps were identified through this scoping review. For example, there was only one high quality clinical trial article which described motor skill development as an intervention. There is a lack of application of standardised motor skill outcome measures reported in clinical trials. The Bruininks-Oseretsky Test of Motor Proficiency, Second Edition (BOT-2) is an example of a standardised, internationally recognised comprehensive assessment of both gross and fine motor skills for children aged four to 21 years [327]. It objectively assesses manual control, manual coordination, body coordination, strength and agility [327]. Ideally the BOT-2 would support the strength and rigor that many of the observed papers lack for their motor skill outcome measures. However, limitations exist including the costs associated with purchasing the BOT-2 assessment tool and the time required for completion of the tests (often taking 45–60 minutes to administer) [327]. Furthermore, the learning effect of tasks within this outcome measure means that ideally children should not be retested within 6 months; which would therefore require a trial to be greater than 6 months in duration in order to appropriately understand the effects of an intervention on motor performance [327]. These limitations are possible explanations to the existing gap in the literature regarding motor skill interventions for children with overweight or obesity.

Faigenbaum and colleagues discuss a need for experienced paediatric professionals to be providing developmentally appropriate physical activity interventions that recognise the unique physical and psychosocial needs of children [328]. Physiotherapists are ideally suited to the provision of such programs, given that professional entry-level training incorporates paediatric-specific content. However, there is also a perceived need to expand paediatric content delivery in entry-level physiotherapy curricula [328,329], including child obesity-specific content [24]. Furthermore, physiotherapists have identified a need for guidance around obesity management/exercise prescription [24], highlighting the timeliness of the current scoping review. The current scoping review provides evidence of the value and scope of physiotherapy practice for this population, which may serve to advocate for services/resources to increase the engagement of physiotherapists in this important clinical area and to guide evidence-based

practice. Central to physiotherapists, this scoping review has highlighted the importance of physiotherapists utilising health-related measures such as anthropometry to assess the clinical benefits of enhancing physical activity levels, whilst multicomponent interventions may first need to be implemented to induce positive behavioural changes that will support long term health improvements from physical activity related interventions that are fun, engaging and age appropriate. Furthermore, physiotherapists need to be mindful of the social and environmental barriers which may negatively impact their intervention outcomes (e.g., lack of resources or funding to access sports or perpetuating negative beliefs such as a lack of perceived effort from children or parents) when engaging clinically with children who are obese and their families. Emphasis should be focused on family-based components of intervention (i.e., engaging parents in therapy planning and implementation) as these are crucial to enhancing intervention outcomes for children with obesity.

Strengths and limitations

Strengths of this review include the breadth of searches undertaken. Several major databases were searched using a broad inclusion criterion to fully understand the scope of existing literature up to this point. The study selection, critical analysis and extraction processes were performed in duplicate, reducing the risk of reviewer error or bias. Further, a sensitivity analysis was conducted which allowed for the exploration of obesity management with obesity 'prevention' studies removed from the synthesis.

Scoping reviews are intended to be broad, bringing together evidence from a range of study designs, and are particularly useful for a large, complex or heterogenous body of literature [330,331]. While a coding method was applied to create a descriptive synthesis of the results from included trials, we recognise that a large degree of heterogeneity existed between studies. Where an individual included study may have concluded an intervention to be successful overall, the current scoping review synthesised information based on individual study outcome measures which may not have reflected the overall concluding findings of the clinical trials.

Furthermore, due to the inclusion of umbrella reviews, systematic reviews, and clinical trials, it is important to recognise that overlap in primary studies may have occurred. Due to the number of primary studies included more than once in this scoping review, there may have been bias in the interpretation of results. To address this problem, an analysis of the results through a calculated percentage of overlap was created.

Despite the present limitations, the current scoping review provides an increased understanding of the available evidence that may be used as information to guide physiotherapy management of childhood obesity and has highlighted existing gaps in the literature.

Future directions for research

Future reviewers should consider focused systematic reviews and meta-analyses to further the findings collected in this scoping review as part of ongoing improvement of standardised physiotherapy related management guidelines for childhood obesity. In particular, systematic reviews/meta-analyses of physiotherapy-specific interventions stratified by study designs (e.g., RCTs) may add further value. The large number of interventions with a questionable effect on health-, performance- and behaviour- related outcomes highlight areas for further exploration within the literature. Additionally, a multitude of personal and environmental factors may have influenced the observed results in this scoping review to varying degrees. Information from personal and environmental factors are confounding factors and if analysed, may guide direction for future researchers to further our understanding of factors contributing to successful interventions.

Conclusion

Despite the plethora of childhood obesity interventions, findings from this review suggest that increases in the quantity of physical activity facilitates desirable changes in health-related outcomes (i.e., anthropometric measures, cardiorespiratory/cardiovascular measures, and blood serum analyses), whereas multicomponent interventions facilitate desirable changes in behavioural measures (i.e., physical activity and sedentary measures, and self-esteem, -efficacy and -perception measures). Such interventions are ideally suited to the scope and role of physiotherapists, particularly in relation to the prescription and progression of physical activity (either as an isolated intervention, or as part of multicomponent programs). Further research is needed to explore the effectiveness of motor skill interventions for obesity management in addition to the impact of environmental modification and parental involvement as possible moderating factors that may contribute to the success of physiotherapy interventions for children who are overweight or obese.

Supporting information

S1 Table. PRISMA checklist.

(DOC)

S2 Table. Search strategy.

(DOCX)

S3 Table. Clinical guidelines data extraction.

(DOCX)

S4 Table. Reviews data extraction.

(DOCX)

S5 Table. Clinical trials data extraction.

(DOCX)

S6 Table. Descriptive synthesis of studies with strong methodological quality.

(XLSX)

S7 Table. Sensitivity analysis.

(DOCX)

Author Contributions

Conceptualization: Margarita D. Tsiros, Nikki Milne.

Formal analysis: Kim Truong, Sandra Park, Margarita D. Tsiros, Nikki Milne.

Methodology: Kim Truong, Sandra Park, Margarita D. Tsiros, Nikki Milne.

Project administration: Nikki Milne.

Resources: Nikki Milne.

Supervision: Nikki Milne.

Validation: Kim Truong, Sandra Park, Margarita D. Tsiros, Nikki Milne.

Writing – original draft: Kim Truong, Sandra Park.

Writing – review & editing: Kim Truong, Sandra Park, Margarita D. Tsiros, Nikki Milne.

References

1. WHO. Obesity and Overweight Fact Sheet 2018 [<http://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight>].
2. Ng M, Fleming T, Robinson M, Thomson B, Graetz N, Margono C, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *The lancet*. 2014; 384(9945):766–81.
3. Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: public-health crisis, common sense cure. *The lancet*. 2002; 360(9331):473–82. [https://doi.org/10.1016/S0140-6736\(02\)09678-2](https://doi.org/10.1016/S0140-6736(02)09678-2) PMID: 12241736
4. Freedman DS, Khan LK, Dietz WH, Srinivasan SR, Berenson GS. Relationship of childhood obesity to coronary heart disease risk factors in adulthood: the Bogalusa Heart Study. *Pediatrics*. 2001; 108(3):712–8. <https://doi.org/10.1542/peds.108.3.712> PMID: 11533341
5. Reilly JJ, Methven E, McDowell ZC, Hacking B, Alexander D, Stewart L, et al. Health consequences of obesity. *Archives of disease in childhood*. 2003; 88(9):748–52. <https://doi.org/10.1136/adc.88.9.748> PMID: 12937090
6. Lakshman R, Elks CE, Ong KK. Childhood obesity. *Circulation*. 2012; 126(14):1770–9. <https://doi.org/10.1161/CIRCULATIONAHA.111.047738> PMID: 23027812
7. Twig G, Yaniv G, Levine H, Leiba A, Goldberger N, Derazne E, et al. Body-mass index in 2.3 million adolescents and cardiovascular death in adulthood. *New England journal of medicine*. 2016; 374(25):2430–40. <https://doi.org/10.1056/NEJMoa1503840> PMID: 27074389
8. Nga VT, Dung VNT, Chu D-T, Tien NLB, Van Thanh V, Ngoc VTN, et al. School education and childhood obesity: A systemic review. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2019. <https://doi.org/10.1016/j.dsx.2019.07.014> PMID: 31405667
9. Flatt J. Issues and misconceptions about obesity. *Obesity*. 2011; 19(4):676. <https://doi.org/10.1038/oby.2011.7> PMID: 21441937
10. Sahoo K, Sahoo B, Choudhury AK, Sofi NY, Kumar R, Bhadoria AS. Childhood obesity: causes and consequences. *Journal of family medicine and primary care*. 2015; 4(2):187. <https://doi.org/10.4103/2249-4863.154628> PMID: 25949965
11. Al-Khudairy L, Loveman E, Colquitt JL, Mead E, Johnson RE, Fraser H, et al. Diet, physical activity and behavioural interventions for the treatment of overweight or obese adolescents aged 12 to 17 years. *Cochrane database of systematic reviews*. 2017(6). <https://doi.org/10.1002/14651858.CD012691> PMID: 28639320
12. Colquitt JL, Loveman E, O'Malley C, Azevedo LB, Mead E, Al-Khudairy L, et al. Diet, physical activity, and behavioural interventions for the treatment of overweight or obesity in preschool children up to the age of 6 years. *Cochrane Database of Systematic Reviews*. 2016(3). <https://doi.org/10.1002/14651858.CD012105> PMID: 26961576
13. Mead E, Brown T, Rees K, Azevedo LB, Whittaker V, Jones D, et al. Diet, physical activity and behavioural interventions for the treatment of overweight or obese children from the age of 6 to 11 years. *Cochrane Database of Systematic Reviews*. 2017(6). <https://doi.org/10.1002/14651858.CD012651> PMID: 28639319
14. NHMRC. Summary Guide for the Management of Overweight and Obesity in Primary Care 2013 [<https://www.nhmrc.gov.au/about-us/publications/clinical-practice-guidelines-management-overweight-and-obesity#block-views-block-file-attachments-content-block-1>].
15. NHMRC. Clinical practice guidelines for the management of overweight and obesity in adults, adolescents and children in Australia 2013 [<https://www.nhmrc.gov.au/about-us/publications/clinical-practice-guidelines-management-overweight-and-obesity#block-views-block-file-attachments-content-block-1>].
16. Rajjo T, Mohammed K, Alsawas M, Ahmed AT, Farah W, Asi N, et al. Treatment of pediatric obesity: an umbrella systematic review. *The Journal of Clinical Endocrinology & Metabolism*. 2017; 102(3):763–75.
17. Liang J, Matheson B, Kaye W, Boutelle K. Neurocognitive correlates of obesity and obesity-related behaviors in children and adolescents. *International journal of obesity*. 2014; 38(4):494. <https://doi.org/10.1038/ijo.2013.142> PMID: 23913029
18. Lubans DR, Morgan PJ, Cliff DP, Barnett LM, Okely AD. Fundamental movement skills in children and adolescents. *Sports medicine*. 2010; 40(12):1019–35. <https://doi.org/10.2165/11536850-000000000-00000> PMID: 21058749
19. Stodden DF, Goodway JD, Langendorfer SJ, Robertson MA, Rudisill ME, Garcia C, et al. A developmental perspective on the role of motor skill competence in physical activity: An emergent relationship. *Quest*. 2008; 60(2):290–306.

20. Tsiros M, Coates A, Howe P, Grimshaw P, Buckley JD. Obesity: the new childhood disability? *Obesity reviews*. 2011; 12(1):26–36. <https://doi.org/10.1111/j.1467-789X.2009.00706.x> PMID: 20070542
21. WCPT. Promoting physical activity in children, the role of physiotherapists 2018 [<https://www.ercpt.eu/file/164>].
22. Tsiros MD, Shultz SP. Take 10! Action Points for Physical Therapists to Consider When It Comes to Childhood Obesity. *Physical therapy*. 2019; 99(5):490–3. <https://doi.org/10.1093/ptj/pzz016> PMID: 30834434
23. Paulis W, Silva S, Koes B, van Middelkoop M. Overweight and obesity are associated with musculo-skeletal complaints as early as childhood: a systematic review. *Obesity Reviews*. 2014; 15(1):52–67. <https://doi.org/10.1111/obr.12067> PMID: 23941399
24. Milne N, Choy NL, Leong GM, Hughes R, Hing W. Child obesity service provision: a cross-sectional survey of physiotherapy practice trends and professional needs. *Australian journal of primary health*. 2016; 22(2):140–6. <https://doi.org/10.1071/PY14101> PMID: 25586908
25. Bryan C, Broussard L, Bellar D. Effective partnerships: how school nurses and physical education teachers can combat childhood obesity. *NASN School Nurse*. 2013; 28(1):20–3. <https://doi.org/10.1177/1942602X12460891> PMID: 23724595
26. Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Annals of internal medicine*. 2018; 169(7):467–73. <https://doi.org/10.7326/M18-0850> PMID: 30178033
27. Hong QN, Pluye P, Fàbregues S, Bartlett G, Boardman F, Cargo M, et al. Mixed methods appraisal tool (MMAT), version 2018. IC Canadian Intellectual Property Office, Industry Canada. 2018.
28. Porta M. *A dictionary of epidemiology*: Oxford university press; 2014.
29. Abbott A. The causal devolution. *Sociological Methods & Research*. 1998; 27(2):148–81.
30. Jones M, Defever E, Letsinger A, Steele J, Mackintosh KA. A mixed studies systematic review and meta-analysis of school—based interventions to promote physical activity and/or reduce sedentary time in children. *Journal of Sport and Health Science*. 2019. <https://doi.org/10.1016/j.jshs.2019.06.009> PMID: 31921476
31. Grimmer K, Dizon JM, Milanese S, King E, Beaton K, Thorpe O, et al. Efficient clinical evaluation of guideline quality: development and testing of a new tool. *BMC medical research methodology*. 2014; 14(1):63. <https://doi.org/10.1186/1471-2288-14-63> PMID: 24885893
32. Whiting P, Rutjes AW, Reitsma JB, Bossuyt PM, Kleijnen J. The development of QUADAS: a tool for the quality assessment of studies of diagnostic accuracy included in systematic reviews. *BMC medical research methodology*. 2003; 3(1):25. <https://doi.org/10.1186/1471-2288-3-25> PMID: 14606960
33. WHO. International classification of functioning, disability and health: Children & youth version: ICF-CY 2007 [<https://apps.who.int/iris/handle/10665/43737>].
34. Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public health rep*. 1985; 100(2):126–31. PMID: 3920711
35. Corbin CB, Pangrazi RP, Franks BD. Definitions: Health, fitness, and physical activity. President's Council on Physical Fitness and Sports Research Digest. 2000.
36. Van Sluijs EM, McMinn AM, Griffin SJ. Effectiveness of interventions to promote physical activity in children and adolescents: systematic review of controlled trials. *Bmj*. 2007; 335(7622):703. <https://doi.org/10.1136/bmj.39320.843947.BE> PMID: 17884863
37. APA. Health and wellbeing of children and young people 2013 [https://australian.physio/sites/default/files/RESOURCES/2018_HealthWellbeing_ChildrenandYoungPeople.pdf].
38. APPT. Fact sheet: The role and scope of pediatric physical therapy in fitness, wellness, health promotion, and prevention 2012 [<https://pediatricapta.org/includes/fact-sheets/pdfs/12%20Role%20and%20Scope%20in%20Fitness%20Health%20Promo.pdf>].
39. APTA. Clinical recommendations provide guidance for physical therapists treating childhood obesity 2016 [https://www.apta.org/uploadedFiles/APTAorg/Media/Releases/Consumer/2016/APTAPressRelease_PTJChildhoodObesityStudy.pdf].
40. Bagby K, Adams S. Evidence-based practice guideline: increasing physical activity in schools—kindergarten through 8th grade. *J Sch Nurs*. 2007; 23(3):137–43. <https://doi.org/10.1177/10598405070230030301> PMID: 17536917
41. CPA. Physical activity for youth and children 2006 [https://physiotherapy.ca/sites/default/files/positionstatements/physical-activity-for-children-and-youth_en_2.pdf].
42. CSP. Physiotherapy works for obesity 2015 [<https://www.csp.org.uk/publications/physiotherapy-works-obesity>].

43. ECOG. Physical Activity And Play In Children Who Are Obese 2015 [Available from: ebook.ecog-obesity.eu/chapter-energy-expenditure-physical-activity/physical-activity-play-children-obese].
44. NICE. Guidance on the prevention, identification, assessment and management of overweight and obesity in adults and children 2006 [<https://lx.iriss.org.uk/sites/default/files/resources/Obesity.pdf>].
45. NICE. Obesity in children and young people: prevention and lifestyle weight management programmes 2015 [www.nice.org.uk/guidance/qs94].
46. WHO. Taking action on childhood obesity 2017 [<https://www.who.int/end-childhood-obesity/publications/taking-action-childhood-obesity-report/en/>].
47. WPTD. Resources on why physical therapy matters 2015 [https://www.wcpt.org/sites/wcpt.org/files/files/wptday/15/resource-booklet/WPTD2015_Resources_final.pdf].
48. Atlantis E, Barnes EH, Singh MA. Efficacy of exercise for treating overweight in children and adolescents: a systematic review. *Int J Obes (Lond)*. 2006; 30(7):1027–40. <https://doi.org/10.1038/sj.ijo.0803286> PMID: 16534526
49. Brown EC, Buchan DS, Baker JS, Wyatt FB, Bocalini DS, Kilgore L. A Systematised Review of Primary School Whole Class Child Obesity Interventions: Effectiveness, Characteristics, and Strategies. *Biomed Res Int*. 2016; 2016:4902714. <https://doi.org/10.1155/2016/4902714> PMID: 27668254
50. Cislak A, Safron M, Pratt M, Gaspar T, Luszczynska A. Family-related predictors of body weight and weight-related behaviours among children and adolescents: a systematic umbrella review. *Child Care Health Dev*. 2012; 38(3):321–31. <https://doi.org/10.1111/j.1365-2214.2011.01285.x> PMID: 21752064
51. Clark ML, Slemmons M. School programs to reduce the prevalence of obesity in children. *American Journal for Nurse Practitioners*. 2008; 12(10):62–8.
52. Craike M, Wiesner G, Hilland TA, Bengoechea EG. Interventions to improve physical activity among socioeconomically disadvantaged groups: an umbrella review. *Int J Behav Nutr Phys Act*. 2018; 15(1):43. <https://doi.org/10.1186/s12966-018-0676-2> PMID: 29764488
53. De Bourdeaudhuij I, Van Cauwenberghe E, Spittaels H, Oppert JM, Rostami C, Brug J, et al. School-based interventions promoting both physical activity and healthy eating in Europe: a systematic review within the HOPE project. *Obes Rev*. 2011; 12(3):205–16. <https://doi.org/10.1111/j.1467-789X.2009.00711.x> PMID: 20122137
54. Delgado-Floody P, Latorre-Roman P, Jerez-Mayorga D, Caamano-Navarrete F, Garcia-Pinillos F. Feasibility of incorporating high-intensity interval training into physical education programs to improve body composition and cardiorespiratory capacity of overweight and obese children: A systematic review. *J Exerc Sci Fit*. 2019; 17(2):35–40. <https://doi.org/10.1016/j.jesf.2018.11.003> PMID: 30740131
55. Dias KA, Green DJ, Ingul CB, Pavey TG, Coombes JS. Exercise and Vascular Function in Child Obesity: A Meta-Analysis. *Pediatrics*. 2015; 136(3):e648–59. <https://doi.org/10.1542/peds.2015-0616> PMID: 26260721
56. Doak CM, Visscher TL, Renders CM, Seidell JC. The prevention of overweight and obesity in children and adolescents: a review of interventions and programmes. *Obes Rev*. 2006; 7(1):111–36. <https://doi.org/10.1111/j.1467-789X.2006.00234.x> PMID: 16436107
57. Engel AC, Broderick CR, van Doorn N, Hardy LL, Parmenter BJ. Exploring the Relationship Between Fundamental Motor Skill Interventions and Physical Activity Levels in Children: A Systematic Review and Meta-analysis. *Sports Med*. 2018; 48(8):1845–57. <https://doi.org/10.1007/s40279-018-0923-3> PMID: 29687278
58. Errisuriz VL, Golaszewski NM, Born K, Bartholomew JB. Systematic Review of Physical Education-Based Physical Activity Interventions Among Elementary School Children. *J Prim Prev*. 2018; 39(3):303–27. <https://doi.org/10.1007/s10935-018-0507-x> PMID: 29705883
59. Feng L, Wei DM, Lin ST, Maddison R, Ni Mhurchu C, Jiang Y, et al. Systematic review and meta-analysis of school-based obesity interventions in mainland China. *PLoS One*. 2017; 12(9):e0184704. <https://doi.org/10.1371/journal.pone.0184704> PMID: 28910362
60. Garcia-Hermoso A, Cerrillo-Urbina AJ, Herrera-Valenzuela T, Cristi-Montero C, Saavedra JM, Martinez-Vizcaino V. Is high-intensity interval training more effective on improving cardiometabolic risk and aerobic capacity than other forms of exercise in overweight and obese youth? A meta-analysis. *Obes Rev*. 2016; 17(6):531–40. <https://doi.org/10.1111/obr.12395> PMID: 26948135
61. Garcia-Hermoso A, Ramirez-Velez R, Saavedra JM. Exercise, health outcomes, and paediatric obesity: A systematic review of meta-analyses. *J Sci Med Sport*. 2019; 22(1):76–84. <https://doi.org/10.1016/j.jsams.2018.07.006> PMID: 30054135
62. Garcia-Hermoso A, Sanchez-Lopez M, Escalante Y, Saavedra JM, Martinez-Vizcaino V. Exercise-based interventions and C-reactive protein in overweight and obese youths: a meta-analysis of

- randomized controlled trials. *Pediatr Res*. 2016; 79(4):522–7. <https://doi.org/10.1038/pr.2015.274> PMID: 26690715
63. Guerra PH, da Silveira JA, Salvador EP. Physical activity and nutrition education at the school environment aimed at preventing childhood obesity: evidence from systematic reviews. *J Pediatr (Rio J)*. 2016; 92(1):15–23.
 64. Han A, Fu A, Cobley S, Sanders RH. Effectiveness of exercise intervention on improving fundamental movement skills and motor coordination in overweight/obese children and adolescents: A systematic review. *J Sci Med Sport*. 2018; 21(1):89–102. <https://doi.org/10.1016/j.jsams.2017.07.001> PMID: 28728887
 65. Harris KC, Kuramoto LK, Schulzer M, Retallack JE. Effect of school-based physical activity interventions on body mass index in children: a meta-analysis. *Cmaj*. 2009; 180(7):719–26. <https://doi.org/10.1503/cmaj.080966> PMID: 19332753
 66. Lobelo F, Garcia de Quevedo I, Holub CK, Nagle BJ, Arredondo EM, Barquera S, et al. School-based programs aimed at the prevention and treatment of obesity: evidence-based interventions for youth in Latin America. *J Sch Health*. 2013; 83(9):668–77. <https://doi.org/10.1111/josh.12080> PMID: 23879787
 67. Niemeier BS, Hektner JM, Enger KB. Parent participation in weight-related health interventions for children and adolescents: a systematic review and meta-analysis. *Prev Med*. 2012; 55(1):3–13. <https://doi.org/10.1016/j.ypmed.2012.04.021> PMID: 22575353
 68. Oliveira CB, Pinto RZ, Saraiva BTC, Tebar WR, Delfino LD, Franco MR, et al. Effects of active video games on children and adolescents: A systematic review with meta-analysis. *Scand J Med Sci Sports*. 2019. <https://doi.org/10.1111/sms.13539> PMID: 31418915
 69. Salmon J, Booth ML, Phongsavan P, Murphy N, Timperio A. Promoting physical activity participation among children and adolescents. *Epidemiol Rev*. 2007; 29:144–59. <https://doi.org/10.1093/epirev/mxm010> PMID: 17556765
 70. Sharma M. School-based interventions for childhood and adolescent obesity. *Obes Rev*. 2006; 7(3):261–9. <https://doi.org/10.1111/j.1467-789X.2006.00227.x> PMID: 16866974
 71. Stuart WP, Broome ME, Smith BA, Weaver M. An integrative review of interventions for adolescent weight loss. *J Sch Nurs*. 2005; 21(2):77–85. <https://doi.org/10.1177/10598405050210020401> PMID: 15801873
 72. Sun C, Pezic A, Tikellis G, Ponsonby AL, Wake M, Carlin JB, et al. Effects of school-based interventions for direct delivery of physical activity on fitness and cardiometabolic markers in children and adolescents: a systematic review of randomized controlled trials. *Obes Rev*. 2013; 14(10):818–38. <https://doi.org/10.1111/obr.12047> PMID: 23734662
 73. Vasconcellos F, Seabra A, Katzmarzyk PT, Kraemer-Aguiar LG, Bouskela E, Farinatti P. Physical activity in overweight and obese adolescents: systematic review of the effects on physical fitness components and cardiovascular risk factors. *Sports Med*. 2014; 44(8):1139–52. <https://doi.org/10.1007/s40279-014-0193-7> PMID: 24743931
 74. Zenzen W, Kridli S. Integrative review of school-based childhood obesity prevention programs. *J Pediatr Health Care*. 2009; 23(4):242–58. <https://doi.org/10.1016/j.pedhc.2008.04.008> PMID: 19559992
 75. Andrade A, Correia CK, Coimbra DR. The Psychological Effects of Exergames for Children and Adolescents with Obesity: A Systematic Review and Meta-Analysis. *Cyberpsychol Behav Soc Netw*. 2019; 22(11):724–35. <https://doi.org/10.1089/cyber.2019.0341> PMID: 31697604
 76. Morgan EH, Schoonees A, Sriram U, Faure M, Seguin-Fowler RA. Caregiver involvement in interventions for improving children's dietary intake and physical activity behaviors. *Cochrane Database Syst Rev*. 2020; 1(1):Cd012547. <https://doi.org/10.1002/14651858.CD012547.pub2> PMID: 31902132
 77. Yuksel HS, Şahin FN, Maksimovic N, Drid P, Bianco A. School-Based Intervention Programs for Preventing Obesity and Promoting Physical Activity and Fitness: A Systematic Review. *Int J Environ Res Public Health*. 2020; 17(1). <https://doi.org/10.3390/ijerph17010347> PMID: 31947891
 78. NHMRC. Summary Guide for the Management of Overweight and Obesity in Primary Care 2013 2013 [<https://www.nhmrc.gov.au/about-us/publications/clinical-practice-guidelines-management-overweight-and-obesity#block-views-block-file-attachments-content-block-1>].
 79. NHMRC. Clinical practice guidelines for the management of overweight and obesity in adults, adolescents and children in Australia 2013 [<https://www.nhmrc.gov.au/about-us/publications/clinical-practice-guidelines-management-overweight-and-obesity#block-views-block-file-attachments-content-block-1>].
 80. ECOG. Physical activity and play in children who are obese 2015 [<http://ebook.ecog-obesity.eu/chapter-energy-expenditure-physical-activity/physical-activity-play-children-obese>].

81. CSP. Physiotherapy works for obesity 2015 [<https://www.csp.org.uk/publications/physiotherapy-works-obesity>].
82. NICE. Guidance on the prevention, identification, assessment and management of overweight and obesity in adults and children. 2006.
83. NICE. Obesity in children and young people: prevention and lifestyle weight management programmes 2015 [www.nice.org.uk/guidance/qs94].
84. Gortmaker SL, Peterson K, Wiecha J, Sobol AM, Dixit S, Fox MK, et al. Reducing obesity via a school-based interdisciplinary intervention among youth: Planet Health. *Arch Pediatr Adolesc Med*. 1999; 153(4):409–18. <https://doi.org/10.1001/archpedi.153.4.409> PMID: 10201726
85. Jones M, Taylor Lynch K, Kass AE, Burrows A, Williams J, Wilfley DE, et al. Healthy weight regulation and eating disorder prevention in high school students: a universal and targeted Web-based intervention. *J Med Internet Res*. 2014; 16(2):e57. <https://doi.org/10.2196/jmir.2995> PMID: 24583683
86. Kain J, Concha F, Moreno L, Leyton B. School-based obesity prevention intervention in Chilean children: effective in controlling, but not reducing obesity. *J Obes*. 2014; 2014:618293. <https://doi.org/10.1155/2014/618293> PMID: 24872892
87. Kain J, Uauy R, Albala, Vio F, Cerda R, Leyton B. School-based obesity prevention in Chilean primary school children: methodology and evaluation of a controlled study. *Int J Obes Relat Metab Disord*. 2004; 28(4):483–93. <https://doi.org/10.1038/sj.ijo.0802611> PMID: 14993915
88. Lau PW, Wong del P, Ngo JK, Liang Y, Kim CG, Kim HS. Effects of high-intensity intermittent running exercise in overweight children. *Eur J Sport Sci*. 2015; 15(2):182–90. <https://doi.org/10.1080/17461391.2014.933880> PMID: 25012183
89. Li XH, Lin S, Guo H, Huang Y, Wu L, Zhang Z, et al. Effectiveness of a school-based physical activity intervention on obesity in school children: a nonrandomized controlled trial. *BMC Public Health*. 2014; 14:1282. <https://doi.org/10.1186/1471-2458-14-1282> PMID: 25510313
90. Sallis JF, McKenzie TL, Alcaraz JE, Kolody B, Hovell MF, Nader PR. Project SPARK. Effects of physical education on adiposity in children. *Ann N Y Acad Sci*. 1993; 699:127–36. <https://doi.org/10.1111/j.1749-6632.1993.tb18844.x> PMID: 8267303
91. Singh AS, Chin APMJ, Brug J, van Mechelen W. Short-term effects of school-based weight gain prevention among adolescents. *Arch Pediatr Adolesc Med*. 2007; 161(6):565–71. <https://doi.org/10.1001/archpedi.161.6.565> PMID: 17548761
92. Wang JJ, Lau WC, Wang HJ, Ma J. Evaluation of a comprehensive intervention with a behavioural modification strategy for childhood obesity prevention: a nonrandomized cluster controlled trial. *BMC Public Health*. 2015; 15:1206. <https://doi.org/10.1186/s12889-015-2535-2> PMID: 26635229
93. Carrel AL, Clark RR, Peterson SE, Nemeth BA, Sullivan J, Allen DB. Improvement of fitness, body composition, and insulin sensitivity in overweight children in a school-based exercise program: a randomized, controlled study. *Arch Pediatr Adolesc Med*. 2005; 159(10):963–8. <https://doi.org/10.1001/archpedi.159.10.963> PMID: 16203942
94. Donnelly JE, Jacobsen DJ, Whatley JE, Hill JO, Swift LL, Cherrington A, et al. Nutrition and physical activity program to attenuate obesity and promote physical and metabolic fitness in elementary school children. *Obes Res*. 1996; 4(3):229–43. <https://doi.org/10.1002/j.1550-8528.1996.tb00541.x> PMID: 8732957
95. Aires L, Silva G, Martins C, Marques E, Lagoa MJ, Ribeiro JC, et al. Exercise intervention and cardiovascular risk factors in obese children. Comparison between obese youngsters taking part in a physical activity school-based programme with and without individualised diet counselling: the ACORDA project. *Ann Hum Biol*. 2016; 43(3):183–90. <https://doi.org/10.3109/03014460.2015.1059889> PMID: 26193775
96. Alexander AG, Grant WL, Pedrino KJ, Lyons PE. A prospective multifactorial intervention on subpopulations of predominately Hispanic children at high risk for obesity. *Obesity (Silver Spring)*. 2014; 22(1):249–53. <https://doi.org/10.1002/oby.20557> PMID: 23836698
97. Alonso-Fernández D, Fernández-Rodríguez R, Taboada-Iglesias Y, Gutiérrez-Sánchez Á. Impact of a HIIT protocol on body composition and VO2max in adolescents. *Science and Sports*. 2019.
98. An J, DuBose KD, Decker JT, Hatala LE. A school-based mentoring program developing healthy behaviors of adolescents with intellectual and developmental disabilities: A pilot feasibility study. *Disabil Health J*. 2019. <https://doi.org/10.1016/j.dhjo.2019.03.012> PMID: 30987820
99. Angelopoulos PD, Milionis HJ, Grammatikaki E, Moschonis G, Manios Y. Changes in BMI and blood pressure after a school based intervention: the CHILDREN study. *Eur J Public Health*. 2009; 19(3):319–25. <https://doi.org/10.1093/eurpub/ckp004> PMID: 19208697

100. Ariza C, Sanchez-Martinez F, Serral G, Valmayor S, Juarez O, Pasarin MI, et al. The Incidence of Obesity, Assessed as Adiposity, Is Reduced After 1 Year in Primary Schoolchildren by the POIBA Intervention. *J Nutr.* 2019; 149(2):258–69. <https://doi.org/10.1093/jn/nxy259> PMID: 30753540
101. Ashem HN, Abdelsamea GA, Osman DA, Hamada HA, Ayoub HES, Soliman GS. Physical therapy protocol for obese adolescent girls with polycystic ovarian syndrome: A within-subject design. *Annals of Clinical and Analytical Medicine.* 2019; 10(4):496–500.
102. Augustijn M, D'Hondt E, Leemans A, Van Acker L, De Guchteneere A, Lenoir M, et al. Weight loss, behavioral change, and structural neuroplasticity in children with obesity through a multidisciplinary treatment program. *Hum Brain Mapp.* 2019; 40(1):137–50. <https://doi.org/10.1002/hbm.24360> PMID: 30198627
103. Augustijn M, D'Hondt E, Van Acker L, De Guchteneere A, Lenoir M, Caeyenberghs K, et al. Role of Motor Competence and Executive Functioning in Weight Loss: A Study in Children with Obesity. *J Dev Behav Pediatr.* 2018; 39(8):642–51. <https://doi.org/10.1097/DBP.0000000000000589> PMID: 29877989
104. Azad A, Gharakhanlou R, Niknam A, Ghanbari A. Effects of aerobic exercise on lung function in overweight and obese students. *Tanaffos.* 2011; 10(3):24–31. PMID: 25191372
105. Bai Y, Saint-Maurice PF, Welk GJ, Russell DW, Allums-Featherston K, Candelaria N. The Longitudinal Impact of NFL PLAY 60 Programming on Youth Aerobic Capacity and BMI. *Am J Prev Med.* 2017; 52(3):311–23. <https://doi.org/10.1016/j.amepre.2016.10.009> PMID: 27919454
106. Barbeau P, Gutin B, Litaker M, Owens S, Riggs S, Okuyama T. Correlates of individual differences in body-composition changes resulting from physical training in obese children. *Am J Clin Nutr.* 1999; 69(4):705–11. <https://doi.org/10.1093/ajcn/69.4.705> PMID: 10197572
107. Bedard C, Bremer E, Campbell W, Cairney J. A Quasi-Experimental Study of a Movement and Preliteracy Program for 3- and 4-Year-Old Children. *Front Pediatr.* 2017; 5:94. <https://doi.org/10.3389/fped.2017.00094> PMID: 28507981
108. Benjamins MR, Whitman S. A culturally appropriate school wellness initiative: results of a 2-year pilot intervention in 2 Jewish schools. *J Sch Health.* 2010; 80(8):378–86. <https://doi.org/10.1111/j.1746-1561.2010.00517.x> PMID: 20618620
109. Bharath LP, Choi WW, Cho JM, Skobodzinski AA, Wong A, Sweeney TE, et al. Combined resistance and aerobic exercise training reduces insulin resistance and central adiposity in adolescent girls who are obese: randomized clinical trial. *Eur J Appl Physiol.* 2018; 118(8):1653–60. <https://doi.org/10.1007/s00421-018-3898-8> PMID: 29846794
110. Bhave S, Pandit A, Yeravdekar R, Madkaikar V, Chinchwade T, Shaikh N, et al. Effectiveness of a 5-year school-based intervention programme to reduce adiposity and improve fitness and lifestyle in Indian children; the SYM-KEM study. *Arch Dis Child.* 2016; 101(1):33–41. <https://doi.org/10.1136/archdischild-2015-308673> PMID: 26420732
111. Bianchini JA, da Silva DF, Nardo CC, Carolino ID, Hernandez F, Nardo N Jr. Multidisciplinary therapy reduces risk factors for metabolic syndrome in obese adolescents. *Eur J Pediatr.* 2013; 172(2):215–21. <https://doi.org/10.1007/s00431-012-1865-7> PMID: 23097084
112. Bilinska I, Kryst L. Effectiveness of a school-based intervention to reduce the prevalence of overweight and obesity in children aged 7–11 years from Poznan (Poland). *Anthropol Anz.* 2017; 74(2):89–100. <https://doi.org/10.1127/anthranz/2017/0719> PMID: 28492704
113. Branco BHM, Valladares D, de Oliveira FM, Carvalho IZ, Marques DC, Coelho AA, et al. Effects of the Order of Physical Exercises on Body Composition, Physical Fitness, and Cardiometabolic Risk in Adolescents Participating in an Interdisciplinary Program Focusing on the Treatment of Obesity. *Front Physiol.* 2019; 10:1013. <https://doi.org/10.3389/fphys.2019.01013> PMID: 31447700
114. Braun HA, Kay CM, Cheung P, Weiss PS, Gazmararian JA. Impact of an Elementary School-Based Intervention on Physical Activity Time and Aerobic Capacity, Georgia, 2013–2014. *Public Health Rep.* 2017; 132(2_suppl):24s–32s. <https://doi.org/10.1177/0033354917719701> PMID: 29136482
115. Brownell KD, Kaye FS. A school-based behavior modification, nutrition education, and physical activity program for obese children. *Am J Clin Nutr.* 1982; 35(2):277–83. <https://doi.org/10.1093/ajcn/35.2.277> PMID: 7064888
116. Brownell KD, Kelman JH, Stunkard AJ. Treatment of obese children with and without their mothers: changes in weight and blood pressure. *Pediatrics.* 1983; 71(4):515–23. PMID: 6835735
117. Burgi F, Niederer I, Schindler C, Bodenmann P, Marques-Vidal P, Kriemler S, et al. Effect of a lifestyle intervention on adiposity and fitness in socially disadvantaged subgroups of preschoolers: a cluster-randomized trial (Ballabeina). *Prev Med.* 2012; 54(5):335–40. <https://doi.org/10.1016/j.yjmed.2012.02.007> PMID: 22373886

118. Burguera B, Colom A, Pinero E, Yanez A, Caimari M, Tur J, et al. ACTYBOSS: activity, behavioral therapy in young subjects—after-school intervention pilot project on obesity prevention. *Obes Facts*. 2011; 4(5):400–6. <https://doi.org/10.1159/000333436> PMID: 22166761
119. Cadzow RB, Chambers MK, Sandell AM. School-Based Obesity Intervention Associated with Three Year Decrease in Student Weight Status in a Low-Income School District. *J Community Health*. 2015; 40(4):709–13. <https://doi.org/10.1007/s10900-015-9989-0> PMID: 25582637
120. Camhi SM, Phillips J, Young DR. The influence of body mass index on long-term fitness from physical education in adolescent girls. *J Sch Health*. 2011; 81(7):409–16. <https://doi.org/10.1111/j.1746-1561.2011.00609.x> PMID: 21668881
121. Canavera M, Sharma M, Murnan J. Development and pilot testing a social cognitive theory-based intervention to prevent childhood obesity among elementary students in rural Kentucky. *Int Q Community Health Educ*. 2008; 29(1):57–70. <https://doi.org/10.2190/IQ.29.1.e> PMID: 19342357
122. Carlone Baldino Garcia N, Lopes WA, Locatelli JC, Ferraz Simoes C, de Oliveira GH, de Souza Mendes VH, et al. Multidisciplinary obesity treatment program improved health-related quality of life and positively correlated with anthropometric and body composition but not with cardiorespiratory fitness parameters in adolescents. *Qual Life Res*. 2019; 28(7):1803–12. <https://doi.org/10.1007/s11136-019-02141-9> PMID: 30790154
123. Carnier J, de Sanches PL, da Silva PL, de Piano A, Tock L, Campos RM, et al. Obese adolescents with eating disorders: analysis of metabolic and inflammatory states. *Physiol Behav*. 2012; 105(2):175–80. <https://doi.org/10.1016/j.physbeh.2011.08.014> PMID: 21871909
124. Carvalho HM, Milano GE, Lopes WA, Figueiredo AJ, Radominski RB, Leite N. Peak oxygen uptake responses to training in obese adolescents: a multilevel allometric framework to partition the influence of body size and maturity status. *Biomed Res Int*. 2013; 2013:618595. <https://doi.org/10.1155/2013/618595> PMID: 23956992
125. Chehab LG, Pfeffer B, Vargas I, Chen S, Irigoyen M. "Energy Up": a novel approach to the weight management of inner-city teens. *J Adolesc Health*. 2007; 40(5):474–6. <https://doi.org/10.1016/j.jadohealth.2006.12.009> PMID: 17448410
126. Chilton JM. Effect of the Total Girl Wellness Program on Wellness Behaviors in Adolescent Females: University of Texas at Tyler; 2012.
127. Chomitz VR, McGowan RJ, Wendel JM, Williams SA, Cabral HJ, King SE, et al. Healthy Living Cambridge Kids: a community-based participatory effort to promote healthy weight and fitness. *Obesity (Silver Spring)*. 2010; 18 Suppl 1:S45–53. <https://doi.org/10.1038/oby.2009.431> PMID: 20107461
128. Cluss P, Lorigan D, Kinsky S, Nikolajski C, McDermott A, Bhat KB. School-Based Health Promotion Initiative Increases Children's Physical Activity. *American Journal of Health Education*. 2016; 47(6):343–54.
129. Cohen CJ, McMillan CS, Samuelson DR. Long-term effects of a lifestyle modification exercise program on the fitness of sedentary, obese children. *J Sports Med Phys Fitness*. 1991; 31(2):183–8. PMID: 1753725
130. Cohen TR, Hazell TJ, Vanstone CA, Rodd C, Weiler HA. Changes in eating behavior and plasma leptin in children with obesity participating in a family-centered lifestyle intervention. *Appetite*. 2018; 125:81–9. <https://doi.org/10.1016/j.appet.2018.01.017> PMID: 29410008
131. Cohen TR, Hazell TJ, Vanstone CA, Rodd C, Weiler HA. Bone Health is Maintained, While Fat Mass is Reduced in Pre-pubertal Children with Obesity Participating in a 1-Year Family-Centered Lifestyle Intervention. *Calcif Tissue Int*. 2017; 101(6):612–22. <https://doi.org/10.1007/s00223-017-0318-8> PMID: 28866763
132. Coknaz D, Mirzeoglu AD, Atasoy HI, Alkoy S, Coknaz H, Goral K. A digital movement in the world of inactive children: favourable outcomes of playing active video games in a pilot randomized trial. *Eur J Pediatr*. 2019; 178(10):1567–76. <https://doi.org/10.1007/s00431-019-03457-x> PMID: 31471690
133. Cordova A, Villa G, Sureda A, Rodriguez-Marroyo JA, Sanchez-Collado MP. Physical activity and cardiovascular risk factors in Spanish children aged 11–13 years. *Rev Esp Cardiol (Engl Ed)*. 2012; 65(7):620–6. <https://doi.org/10.1016/j.recesp.2012.01.026> PMID: 22633280
134. Crouter SE, Salas C, Wiecha J. Effects of an afterschool community center physical activity program on fitness and body composition in obese youth. *J Sports Sci*. 2017; 35(11):1034–40. <https://doi.org/10.1080/02640414.2016.1209305> PMID: 27433781
135. Crova C, Struzzolino I, Marchetti R, Masci I, Vannozzi G, Forte R, et al. Cognitively challenging physical activity benefits executive function in overweight children. *J Sports Sci*. 2014; 32(3):201–11. <https://doi.org/10.1080/02640414.2013.828849> PMID: 24015968
136. Cvetkovic N, Stojanovic E, Stojiljkovic N, Nikolic D, Scanlan AT, Milanovic Z. Exercise training in overweight and obese children: Recreational football and high-intensity interval training provide similar

- benefits to physical fitness. *Scand J Med Sci Sports*. 2018; 28 Suppl 1:18–32. <https://doi.org/10.1111/sms.13241> PMID: 29979479
137. da Silva DF, Bianchini JA, Lopera CA, Capelato DA, Hintze LJ, Nardo CC, et al. Impact of readiness to change behavior on the effects of a multidisciplinary intervention in obese Brazilian children and adolescents. *Appetite*. 2015; 87:229–35. <https://doi.org/10.1016/j.appet.2014.12.221> PMID: 25558026
 138. da Silveira Campos RM, Landi Masquio DC, Campos Corgosinho F, de Lima Sanches P, de Piano A, Carnier J, et al. Homeostasis Model Assessment-Adiponectin: the role of different types of physical exercise in obese adolescents. *J Sports Med Phys Fitness*. 2017; 57(6):831–8. <https://doi.org/10.23736/S0022-4707.16.06235-6> PMID: 27385541
 139. Dauenhauer B, Keating X, Lambdin D. Effects of a Three-Tiered Intervention Model on Physical Activity and Fitness Levels of Elementary School Children. *J Prim Prev*. 2016; 37(4):313–27. <https://doi.org/10.1007/s10935-016-0430-y> PMID: 27059849
 140. Davis CL, Tomporowski PD, Boyle CA, Waller JL, Miller PH, Naglieri JA, et al. Effects of aerobic exercise on overweight children's cognitive functioning: A randomized controlled trial. *Research Quarterly for Exercise and Sport*. 2007; 78(5):510–9. <https://doi.org/10.1080/02701367.2007.10599450> PMID: 18274222
 141. De Miguel-Etayo P, Moreno LA, Santabarbara J, Bueno G, Martin-Matillas M, Zapatera B, et al. BODY COMPOSITION CHANGES DURING A MULTIDISCIPLINARY TREATMENT PROGRAMME IN OVERWEIGHT ADOLESCENTS: EVASYON STUDY. *Nutr Hosp*. 2015; 32(6):2525–34. <https://doi.org/10.3305/nh.2015.32.6.9663> PMID: 26667699
 142. de Vries AG, Huiting HG, van den Heuvel ER, L'Abée C, Corpeleijn E, Stolk RP. An activity stimulation programme during a child's first year reduces some indicators of adiposity at the age of two-and-a-half. *Acta Paediatr*. 2015; 104(4):414–21. <https://doi.org/10.1111/apa.12880> PMID: 25425024
 143. Deldin A, Kuk JL, Lee S. Influence of Sex on the Changes in Regional Fat and Skeletal Muscle Mass in Response to Exercise Training in Adolescents with Obesity. *Child Obes*. 2019; 15(3):216–22. <https://doi.org/10.1089/chi.2018.0329> PMID: 30694699
 144. Delgado-Floody P, Espinoza-Silva M, García-Pinillos F, Latorre-Román P. Effects of 28 weeks of high-intensity interval training during physical education classes on cardiometabolic risk factors in Chilean schoolchildren: a pilot trial. *European Journal of Pediatrics*. 2018; 177(7):1019–27. <https://doi.org/10.1007/s00431-018-3149-3> PMID: 29680994
 145. Delisle Nystrom C, Sandin S, Henriksson P, Henriksson H, Maddison R, Lof M. A 12-month follow-up of a mobile-based (mHealth) obesity prevention intervention in pre-school children: the MINISTOP randomized controlled trial. *BMC Public Health*. 2018; 18(1):658. <https://doi.org/10.1186/s12889-018-5569-4> PMID: 29793467
 146. DeRenne C, Maeda JK, Chai DX, Ho K, Kaluhiokalani N, Braun KL. Afterschool physical activity program to reduce obesity-related cancer risk: a feasibility study. *J Cancer Educ*. 2008; 23(4):230–4. <https://doi.org/10.1080/08858190802188602> PMID: 19058071
 147. Di Pietro M, Campanaro P, D'Angelo G, Di Ferdinando C, Pomilio M, Verrotti A, et al. Role of camping in the treatment of childhood obesity. *Acta Biomed*. 2004; 75(2):118–21. PMID: 15481701
 148. Eichner JE, Folorunso OA, Moore WE. A Physical Activity Intervention and Changes in Body Mass Index at a Middle School With a Large American Indian Population, Oklahoma, 2004–2009. *Prev Chronic Dis*. 2016; 13:E163. <https://doi.org/10.5888/pcd13.150495> PMID: 27906646
 149. Elder JP, McKenzie TL, Arredondo EM, Crespo NC, Ayala GX. Effects of a multi-pronged intervention on children's activity levels at recess: the Aventuras para Niños study. *Adv Nutr*. 2011; 2(2):171s–6s. <https://doi.org/10.3945/an.111.000380> PMID: 22332049
 150. Epstein LH, Koeske R, Zidansek J, Wing RR. Effects of weight loss on fitness in obese children. *Am J Dis Child*. 1983; 137(7):654–7. <https://doi.org/10.1001/archpedi.1983.02140330038010> PMID: 6858977
 151. Erfle SE, Gamble A. Effects of daily physical education on physical fitness and weight status in middle school adolescents. *J Sch Health*. 2015; 85(1):27–35. <https://doi.org/10.1111/josh.12217> PMID: 25440450
 152. Farah BQ, Ritti-Dias RM, Balagopal PB, Hill JO, Prado WL. Does exercise intensity affect blood pressure and heart rate in obese adolescents? A 6-month multidisciplinary randomized intervention study. *Pediatr Obes*. 2014; 9(2):111–20. <https://doi.org/10.1111/j.2047-6310.2012.00145.x> PMID: 23447453
 153. Farias Edos S, Goncalves EM, Morcillo AM, Guerra-Junior G, Amancio OM. Effects of programmed physical activity on body composition in post-pubertal schoolchildren. *J Pediatr (Rio J)*. 2015; 91(2):122–9. <https://doi.org/10.1016/j.jped.2014.06.004> PMID: 25305637
 154. Fogel VA, Miltenberger RG, Graves R, Koehler S. The effects of exergaming on physical activity among inactive children in a physical education classroom. *J Appl Behav Anal*. 2010; 43(4):591–600. <https://doi.org/10.1901/jaba.2010.43-591> PMID: 21541146

155. Foster GD, Linder B, Baranowski T, Cooper DM, Goldberg L, Harrell JS, et al. A school-based intervention for diabetes risk reduction. *N Engl J Med*. 2010; 363(5):443–53. <https://doi.org/10.1056/NEJMoa1001933> PMID: 20581420
156. Gallotta MC, Iazzoni S, Emerenziani GP, Meucci M, Migliaccio S, Guidetti L, et al. Effects of combined physical education and nutritional programs on schoolchildren's healthy habits. *PeerJ*. 2016; 4:e1880. <https://doi.org/10.7717/peerj.1880> PMID: 27077004
157. Gao Z, Pope Z, Lee JE, Stodden D, Roncesvalles N, Pasco D, et al. Impact of exergaming on young children's school day energy expenditure and moderate-to-vigorous physical activity levels. *J Sport Health Sci*. 2017; 6(1):11–6. <https://doi.org/10.1016/j.jshs.2016.11.008> PMID: 30356552
158. Garcia-Hermoso A, Saavedra JM, Escalante Y, Dominguez AM. The Intention to be Physically Active in Sedentary Obese Children: A Longitudinal Study. *Behav Sci (Basel)*. 2018; 8(1). <https://doi.org/10.3390/bs8010009> PMID: 29324710
159. Gentier I, D'Hondt E, Augustijn M, Tanghe A, De Bourdeaudhuij I, Deforche B, et al. Multidisciplinary residential treatment can improve perceptual-motor function in obese children. *Acta Paediatr*. 2015; 104(6):e263–70. <https://doi.org/10.1111/apa.12899> PMID: 25619529
160. Gesell SB, Scott TA, Barkin SL. Accuracy of perception of body size among overweight Latino preadolescents after a 6-month physical activity skills building intervention. *Clin Pediatr (Phila)*. 2010; 49(4):323–9. <https://doi.org/10.1177/0009922809339386> PMID: 19605865
161. Going S, Thompson J, Cano S, Stewart D, Stone E, Harnack L, et al. The effects of the Pathways Obesity Prevention Program on physical activity in American Indian children. *Prev Med*. 2003; 37(6 Pt 2): S62–9. <https://doi.org/10.1016/j.ypmed.2003.08.005> PMID: 14636810
162. Graf C, Koch B, Bjarnason-Wehrens B, Sreeram N, Brockmeier K, Tokarski W, et al. Who benefits from intervention in, as opposed to screening of, overweight and obese children? *Cardiol Young*. 2006; 16(5):474–80. <https://doi.org/10.1017/S1047951106000667> PMID: 16984699
163. Guerendiain M, Montes R, Lopez-Belmonte G, Martin-Matillas M, Castellote AI, Martin-Bautista E, et al. Changes in plasma fatty acid composition are associated with improvements in obesity and related metabolic disorders: A therapeutic approach to overweight adolescents. *Clin Nutr*. 2018; 37(1):149–56. <https://doi.org/10.1016/j.clnu.2016.11.006> PMID: 27887752
164. Gutin B, Yin Z, Johnson M, Barbeau P. Preliminary findings of the effect of a 3-year after-school physical activity intervention on fitness and body fat: the Medical College of Georgia FitKid Project. *Int J Pediatr Obes*. 2008; 3 Suppl 1:3–9. <https://doi.org/10.1080/17477160801896457> PMID: 18278626
165. Halfon ST, Bronner S. The influence of a physical ability intervention program on improved running time and increased sport motivation among Jerusalem schoolchildren. *Adolescence*. 1988; 23(90):405–16. PMID: 3407501
166. Harrell JS, Gansky SA, McMurray RG, Bangdiwala SI, Frauman AC, Bradley CB. School-based interventions improve heart health in children with multiple cardiovascular disease risk factors. *Pediatrics*. 1998; 102(2 Pt 1):371–80. <https://doi.org/10.1542/peds.102.2.371> PMID: 9685441
167. Harrison M, Burns CF, McGuinness M, Heslin J, Murphy NM. Influence of a health education intervention on physical activity and screen time in primary school children: 'Switch Off—Get Active'. *J Sci Med Sport*. 2006; 9(5):388–94. <https://doi.org/10.1016/j.jsams.2006.06.012> PMID: 16872900
168. Hawthorne A, Shaibi G, Gance-Cleveland B, McFall S. Grand Canyon Trekkers: school-based lunch-time walking program. *J Sch Nurs*. 2011; 27(1):43–50. <https://doi.org/10.1177/1059840510391669> PMID: 21123848
169. Hayashi T, Fujino M, Shindo M, Hiroki T, Arakawa K. Echocardiographic and electrocardiographic measures in obese children after an exercise program. *Int J Obes*. 1987; 11(5):465–72. PMID: 3429110
170. Hills AP, Parker AW. Obesity management via diet and exercise intervention. *Child Care Health Dev*. 1988; 14(6):409–16. <https://doi.org/10.1111/j.1365-2214.1988.tb00592.x> PMID: 3228964
171. Hintze LJ, Cattai GBP, Junior NN. Multidisciplinary program for obesity treatment: Summary of results with adolescents. *Acta Scientiarum—Health Sciences*. 2012; 34(2):137–44.
172. Hollis JL, Sutherland R, Campbell L, Morgan PJ, Lubans DR, Nathan N, et al. Effects of a 'school-based' physical activity intervention on adiposity in adolescents from economically disadvantaged communities: secondary outcomes of the 'Physical Activity 4 Everyone' RCT. *Int J Obes (Lond)*. 2016; 40(10):1486–93. <https://doi.org/10.1038/ijo.2016.107> PMID: 27430652
173. Horsak B, Schwab C, Baca A, Greber-Platzer S, Kreissl A, Nehrer S, et al. Effects of a lower extremity exercise program on gait biomechanics and clinical outcomes in children and adolescents with obesity: A randomized controlled trial. *Gait Posture*. 2019; 70:122–9. <https://doi.org/10.1016/j.gaitpost.2019.02.032> PMID: 30851623

174. Hystad HT, Steinsbekk S, Odegard R, Wichstrom L, Gudbrandsen OA. A randomised study on the effectiveness of therapist-led v. self-help parental intervention for treating childhood obesity. *Br J Nutr*. 2013; 110(6):1143–50. <https://doi.org/10.1017/S0007114513000056> PMID: 23388524
175. Inoue DS, De Mello MT, Foschini D, Lira FS, De Piano Ganen A, Da Silveira Campos RM, et al. Linear and undulating periodized strength plus aerobic training promote similar benefits and lead to improvement of insulin resistance on obese adolescents. *J Diabetes Complications*. 2015; 29(2):258–64. <https://doi.org/10.1016/j.jdiacomp.2014.11.002> PMID: 25441178
176. Irwin CC, Irwin RL, Miller ME, Somes GW, Richey PA. Get Fit with the Grizzlies: a community-school-home initiative to fight childhood obesity. *J Sch Health*. 2010; 80(7):333–9. <https://doi.org/10.1111/j.1746-1561.2010.00510.x> PMID: 20591098
177. Jakubowski TL, Perron T, Farrell A, Kenner C, Hullings C. The Smart Nutrition and Conditioning for Kids (SNACK) Program: An Approach to Increasing Nutrition Knowledge of Second-Grade Students. *MCN Am J Matern Child Nurs*. 2018; 43(5):278–84. <https://doi.org/10.1097/NMC.0000000000000463> PMID: 30113407
178. Jansen W, Borsboom G, Meima A, Zwanenburg EJ, Mackenbach JP, Raat H, et al. Effectiveness of a primary school-based intervention to reduce overweight. *Int J Pediatr Obes*. 2011; 6(2–2):e70–7. <https://doi.org/10.3109/17477166.2011.575151> PMID: 21609245
179. Jiang J, Xia X, Greiner T, Wu G, Lian G, Rosenqvist U. The effects of a 3-year obesity intervention in schoolchildren in Beijing. *Child Care Health Dev*. 2007; 33(5):641–6. <https://doi.org/10.1111/j.1365-2214.2007.00738.x> PMID: 17725789
180. Jones RA, Kelly J, Cliff DP, Batterham M, Okely AD. Acceptability and Potential Efficacy of Single-Sex After-School Activity Programs for Overweight and At-Risk Children: The Wollongong SPORT RCT. *Pediatr Exerc Sci*. 2015; 27(4):535–45. <https://doi.org/10.1123/pes.2015-0116> PMID: 26305240
181. Julian V, Thivel D, Miguet M, Pereira B, Costes F, Coudeyre E, et al. Eccentric cycling is more efficient in reducing fat mass than concentric cycling in adolescents with obesity. *Scand J Med Sci Sports*. 2019; 29(1):4–15. <https://doi.org/10.1111/sms.13301> PMID: 30222208
182. Jurg ME, Kremers SP, Candel MJ, Van der Wal MF, De Meij JS. A controlled trial of a school-based environmental intervention to improve physical activity in Dutch children: JUMP-in, kids in motion. *Health Promot Int*. 2006; 21(4):320–30. <https://doi.org/10.1093/heapro/dal032> PMID: 16963784
183. Karacabey K. The effect of exercise on leptin, insulin, cortisol and lipid profiles in obese children. *J Int Med Res*. 2009; 37(5):1472–8. <https://doi.org/10.1177/147323000903700523> PMID: 19930853
184. Kaufman C, Kelly AS, Kaiser DR, Steinberger J, Dengel DR. Aerobic-exercise training improves ventilatory efficiency in overweight children. *Pediatr Exerc Sci*. 2007; 19(1):82–92. <https://doi.org/10.1123/pes.19.1.82> PMID: 17554160
185. Khammassi M, Miguet M, O'Malley G, Fillon A, Masurier J, Damaso AR, et al. Health-related quality of life and perceived health status of adolescents with obesity are improved by a 10-month multidisciplinary intervention. *Physiol Behav*. 2019; 210:112549. <https://doi.org/10.1016/j.physbeh.2019.05.010> PMID: 31082444
186. King AK, McGill-Meeks K, Beller JP, Burt Solorzano CM. Go Girls!-Dance-Based Fitness to Increase Enjoyment of Exercise in Girls at Risk for PCOS. *Children (Basel)*. 2019; 6(9). <https://doi.org/10.3390/children6090099> PMID: 31500180
187. Kirschenbaum DS, Germann JN, Rich BH. Treatment of morbid obesity in low-income adolescents: effects of parental self-monitoring. *Obes Res*. 2005; 13(9):1527–9. <https://doi.org/10.1038/oby.2005.187> PMID: 16222054
188. Klakk H, Chinapaw M, Heidemann M, Andersen LB, Wedderkopp N. Effect of four additional physical education lessons on body composition in children aged 8–13 years—a prospective study during two school years. *BMC Pediatr*. 2013; 13:170. <https://doi.org/10.1186/1471-2431-13-170> PMID: 24131778
189. Kriemler S, Zahner L, Schindler C, Meyer U, Hartmann T, Hebestreit H, et al. Effect of school based physical activity programme (KISS) on fitness and adiposity in primary schoolchildren: cluster randomised controlled trial. *Bmj*. 2010; 340:c785. <https://doi.org/10.1136/bmj.c785> PMID: 20179126
190. Krombholz H. The impact of a 20-month physical activity intervention in child care centers on motor performance and weight in overweight and healthy-weight preschool children. *Percept Mot Skills*. 2012; 115(3):919–32. <https://doi.org/10.2466/06.10.25.PMS.115.6.919-932> PMID: 23409603
191. Kuhr P, Antunes Lima R, Grontved A, Wedderkopp N, Klakk H. Three times as much physical education reduced the risk of children being overweight or obese after five years. *Acta Paediatr*. 2019. <https://doi.org/10.1111/apa.15005> PMID: 31509297
192. Labayen I, Medrano M, Arenaza L, Maiz E, Osés M, Martínez-Vizcaino V, et al. Effects of Exercise in Addition to a Family-Based Lifestyle Intervention Program on Hepatic Fat in Children With Overweight. *Diabetes Care*. 2019. <https://doi.org/10.2337/dc19-0351> PMID: 31227585

193. Lambrick D, Westrupp N, Kaufmann S, Stoner L, Faulkner J. The effectiveness of a high-intensity games intervention on improving indices of health in young children. *J Sports Sci.* 2016; 34(3):190–8. <https://doi.org/10.1080/02640414.2015.1048521> PMID: 26009003
194. Larsen KT, Huang T, Larsen LR, Olesen LG, Andersen LB, Moller NC. The effect of a multi-component camp-based weight-loss program on children's motor skills and physical fitness: a randomized controlled trial. *BMC Pediatr.* 2016; 16:91. <https://doi.org/10.1186/s12887-016-0627-5> PMID: 27416906
195. Lee RL, Leung C, Chen H, Louie LHT, Brown M, Chen JL, et al. The Impact of a School-Based Weight Management Program Involving Parents via mHealth for Overweight and Obese Children and Adolescents with Intellectual Disability: A Randomized Controlled Trial. *Int J Environ Res Public Health.* 2017; 14(10).
196. Lee S, Libman I, Hughan K, Kuk JL, Jeong JH, Zhang D, et al. Effects of Exercise Modality on Insulin Resistance and Ectopic Fat in Adolescents with Overweight and Obesity: A Randomized Clinical Trial. *J Pediatr.* 2019; 206:91–8.e1. <https://doi.org/10.1016/j.jpeds.2018.10.059> PMID: 30554789
197. Lee SS, Kang S. Effects of regular exercise on obesity and type 2 diabetes mellitus in Korean children: improvements glycemic control and serum adipokines level. *J Phys Ther Sci.* 2015; 27(6):1903–7. <https://doi.org/10.1589/jpts.27.1903> PMID: 26180345
198. Leite N, Carvalho HM, Padez C, Lopes WA, Milano GE, Radominski RB, et al. Age and menarcheal status do not influence metabolic response to aerobic training in overweight girls. *Diabetol Metab Syndr.* 2013; 5(1):7. <https://doi.org/10.1186/1758-5996-5-7> PMID: 23443001
199. Leite N, Milano GE, Cieslak F, Lopes WA, Rodacki A, Radominski RB. Effects of physical exercise and nutritional guidance on metabolic syndrome in obese adolescents. *Brazilian Journal of Physical Therapy/Revista Brasileira de Fisioterapia.* 2009; 13(1):73–81.
200. LoMauro A, Cesareo A, Agosti F, Tringali G, Salvadego D, Grassi B, et al. Effects of a multidisciplinary body weight reduction program on static and dynamic thoraco-abdominal volumes in obese adolescents. *Appl Physiol Nutr Metab.* 2016; 41(6):649–58. <https://doi.org/10.1139/apnm-2015-0269> PMID: 27175804
201. Lopera CA, da Silva DF, Bianchini JA, Locateli JC, Moreira AC, Dada RP, et al. Effect of water- versus land-based exercise training as a component of a multidisciplinary intervention program for overweight and obese adolescents. *Physiol Behav.* 2016; 165:365–73. <https://doi.org/10.1016/j.physbeh.2016.08.019> PMID: 27553575
202. Lopes WA, Leite N, da Silva LR, Brunelli DT, Gaspari AF, Radominski RB, et al. Effects of 12 weeks of combined training without caloric restriction on inflammatory markers in overweight girls. *J Sports Sci.* 2016; 34(20):1902–12. <https://doi.org/10.1080/02640414.2016.1142107> PMID: 26852885
203. Magnani Branco BH, Carvalho IZ, Garcia de Oliveira H, Fanhani AP, Machado Dos Santos MC, Pestillo de Oliveira L, et al. Effects of 2 Types of Resistance Training Models on Obese Adolescents' Body Composition, Cardiometabolic Risk, and Physical Fitness. *J Strength Cond Res.* 2018.
204. Manley D, Cowan P, Graff C, Perlow M, Rice P, Richey P, et al. Self-efficacy, physical activity, and aerobic fitness in middle school children: examination of a pedometer intervention program. *J Pediatr Nurs.* 2014; 29(3):228–37. <https://doi.org/10.1016/j.pedn.2013.10.011> PMID: 24263251
205. Mardones F, Arnaiz P, Soto-Sanchez J, Saavedra J, Dominguez A, Rozowski J, et al. Physical activity in the classroom to prevent childhood obesity: a pilot study in Santiago, Chile. *J Nutr Sci.* 2017; 6:e21. <https://doi.org/10.1017/jns.2017.14> PMID: 28630698
206. Marild S, Gronowitz E, Forsell C, Dahlgren J, Friberg P. A controlled study of lifestyle treatment in primary care for children with obesity. *Pediatr Obes.* 2013; 8(3):207–17. <https://doi.org/10.1111/j.2047-6310.2012.00105.x> PMID: 23172847
207. Martinez Vizcaino V, Salcedo Aguilar F, Franquelo Gutierrez R, Solera Martinez M, Sanchez Lopez M, Serrano Martinez S, et al. Assessment of an after-school physical activity program to prevent obesity among 9- to 10-year-old children: a cluster randomized trial. *Int J Obes (Lond).* 2008; 32(1):12–22. <https://doi.org/10.1038/sj.ijo.0803738> PMID: 17895883
208. Martinez-Lopez EJ, Grao-Cruces A, Moral-Garcia JE, Pantoja-Vallejo A. Intervention for spanish overweight teenagers in physical education lessons. *J Sports Sci Med.* 2012; 11(2):312–21. PMID: 24149205
209. Matvienko O, Ahrabi-Fard I. The effects of a 4-week after-school program on motor skills and fitness of kindergarten and first-grade students. *Am J Health Promot.* 2010; 24(5):299–303. <https://doi.org/10.4278/ajhp.08050146> PMID: 20465142
210. Meyer U, Schindler C, Zahner L, Ernst D, Hebestreit H, van Mechelen W, et al. Long-term effect of a school-based physical activity program (KISS) on fitness and adiposity in children: a cluster-randomized controlled trial. *PLoS One.* 2014; 9(2):e87929. <https://doi.org/10.1371/journal.pone.0087929> PMID: 24498404

211. Militao AG, de Oliveira Karnikowski MG, da Silva FR, Garcez Militao ES, Dos Santos Pereira RM, Grubert Campbell CS. Effects of a recreational physical activity and healthy habits orientation program, using an illustrated diary, on the cardiovascular risk profile of overweight and obese schoolchildren: a pilot study in a public school in Brasília, Federal District, Brazil. *Diabetes Metab Syndr Obes.* 2013; 6:445–51. <https://doi.org/10.2147/DMSO.S52166> PMID: 24348058
212. Miller TA, Vaux-Bjerke A, McDonnell KA, DiPietro L. Can E-Gaming Be Useful for Achieving Recommended Levels of Moderate- to Vigorous-Intensity Physical Activity in Inner-City Children? *Games Health J.* 2013; 2(2):96–102. <https://doi.org/10.1089/g4h.2012.0058> PMID: 26192127
213. Molina-Garcia P, Miranda-Aparicio D, Molina-Molina A, Plaza-Florido A, Migueles JH, Mora-Gonzalez J, et al. Effects of Exercise on Plantar Pressure during Walking in Children with Overweight/Obesity. *Med Sci Sports Exerc.* 2019.
214. Monsalves-Alvarez M, Castro-Sepulveda M, Zapata-Lamana R, Rosales-Soto G, Salazar G. Motor Skills and Nutritional Status Outcomes from a Physical Activity Intervention in Short Breaks on Pre-school Children Conducted by their Educators: A pilot Study. *Nutr Hosp.* 2015; 32(4):1576–81. <https://doi.org/10.3305/nh.2015.32.4.9514> PMID: 26545520
215. Monteiro PA, Chen KY, Lira FS, Saraiva BT, Antunes BM, Campos EZ, et al. Concurrent and aerobic exercise training promote similar benefits in body composition and metabolic profiles in obese adolescents. *Lipids Health Dis.* 2015; 14:153. <https://doi.org/10.1186/s12944-015-0152-9> PMID: 26611872
216. Morano M, Colella D, Rutigliano I, Fiore P, Pettoello-Mantovani M, Campanozzi A. A multi-modal training programme to improve physical activity, physical fitness and perceived physical ability in obese children. *J Sports Sci.* 2014; 32(4):345–53. <https://doi.org/10.1080/02640414.2013.824602> PMID: 23968284
217. Muller I, Schindler C, Adams L, Endes K, Gall S, Gerber M, et al. Effect of a Multidimensional Physical Activity Intervention on Body Mass Index, Skinfolds and Fitness in South African Children: Results from a Cluster-Randomised Controlled Trial. *Int J Environ Res Public Health.* 2019; 16(2). <https://doi.org/10.3390/ijerph16020232> PMID: 30650624
218. Naul R, Schmelt D, Dreiskaemper D, Hoffmann D, l'Hoir M. 'Healthy children in sound communities' (HCSC/gkgk)—a Dutch-German community-based network project to counteract obesity and physical inactivity. *Fam Pract.* 2012; 29 Suppl 1:i110–i6. <https://doi.org/10.1093/fampra/cmr097> PMID: 22399539
219. Nayak BS, Bhat VH. School Based Multicomponent Intervention for Obese Children in Udipi District, South India—A Randomized Controlled Trial. *J Clin Diagn Res.* 2016; 10(12):Sc24–sc8. <https://doi.org/10.7860/JCDR/2016/23766.9116> PMID: 28208968
220. Nemet D, Berger-Shemesh E, Wolach B, Eliakim A. A combined dietary-physical activity intervention affects bone strength in obese children and adolescents. *Int J Sports Med.* 2006; 27(8):666–71. <https://doi.org/10.1055/s-2005-872920> PMID: 16874595
221. Neumark-Sztainer DR, Friend SE, Flattum CF, Hannan PJ, Story MT, Bauer KW, et al. New moves—preventing weight-related problems in adolescent girls a group-randomized study. *Am J Prev Med.* 2010; 39(5):421–32. <https://doi.org/10.1016/j.amepre.2010.07.017> PMID: 20965379
222. Nichols JF, Bigelow DM, Canine KM. Short-term weight loss and exercise training effects on glucose-induced thermogenesis in obese adolescent males during hypocaloric feeding. *Int J Obes.* 1989; 13(5):683–90. PMID: 2583922
223. Nobre GG, de Almeida MB, Nobre IG, Dos Santos FK, Brinco RA, Arruda-Lima TR, et al. Twelve Weeks of Plyometric Training Improves Motor Performance of 7- to 9-Year-Old Boys Who Were Overweight/Obese: A Randomized Controlled Intervention. *J Strength Cond Res.* 2017; 31(8):2091–9. <https://doi.org/10.1519/JSC.0000000000001684> PMID: 27787471
224. Nogueira RC, Weeks BK, Beck BR. An in-school exercise intervention to enhance bone and reduce fat in girls: the CAPO Kids trial. *Bone.* 2014; 68:92–9. <https://doi.org/10.1016/j.bone.2014.08.006> PMID: 25151492
225. Nunez-Gaunaurd A, Kirk-Sanchez N. Health Outcomes of an Extracurricular Family-Based Intervention for Physical Activity in Three Hispanic Male Children: A Case Series. *HPA Resource.* 2011; 11(3): J2–9.
226. Nystrom CD, Sandin S, Henriksson P, Henriksson H, Trolle-Lagerros Y, Larsson C, et al. Mobile-based intervention intended to stop obesity in preschool-aged children: the MINISTOP randomized controlled trial. *Am J Clin Nutr.* 2017; 105(6):1327–35. <https://doi.org/10.3945/ajcn.116.150995> PMID: 28446496
227. Olvera N, Bush JA, Sharma SV, Knox BB, Scherer RL, Butte NF. BOUNCE: a community-based mother-daughter healthy lifestyle intervention for low-income Latino families. *Obesity (Silver Spring).* 2010; 18 Suppl 1:S102–4. <https://doi.org/10.1038/oby.2009.439> PMID: 20107454

228. Paravidino VB, Mediano MF, Hoffman DJ, Sichieri R. Effect of Exercise Intensity on Spontaneous Physical Activity Energy Expenditure in Overweight Boys: A Crossover Study. *PLoS One*. 2016; 11(1): e0147141. <https://doi.org/10.1371/journal.pone.0147141> PMID: 26771742
229. Park KS, Lee MG. Effects of summer school participation and psychosocial outcomes on changes in body composition and physical fitness during summer break. *J Exerc Nutrition Biochem*. 2015; 19(2):81–90. <https://doi.org/10.5717/jenb.2015.15052005> PMID: 26244126
230. Park TG, Hong HR, Lee J, Kang HS. Lifestyle plus exercise intervention improves metabolic syndrome markers without change in adiponectin in obese girls. *Ann Nutr Metab*. 2007; 51(3):197–203. <https://doi.org/10.1159/000104137> PMID: 17587789
231. Peralta LR, Jones RA, Okely AD. Promoting healthy lifestyles among adolescent boys: the Fitness Improvement and Lifestyle Awareness Program RCT. *Prev Med*. 2009; 48(6):537–42. <https://doi.org/10.1016/j.ypmed.2009.04.007> PMID: 19389421
232. Prosper M, Moczulski VL, Qureshi A, Weiss M, Bryars T. Healthy for Life/PE4ME: assessing an intervention targeting childhood obesity. *Californian Journal of Health Promotion*. 2009; 7(S1):1–10.
233. Puder JJ, Marques-Vidal P, Schindler C, Zahner L, Niederer I, Burgi F, et al. Effect of multidimensional lifestyle intervention on fitness and adiposity in predominantly migrant preschool children (Ballabeina): cluster randomised controlled trial. *Bmj*. 2011; 343:d6195. <https://doi.org/10.1136/bmj.d6195> PMID: 21998346
234. Quinn M. Introduction of active video gaming into the middle school curriculum as a school-based childhood obesity intervention. *J Pediatr Health Care*. 2013; 27(1):3–12. <https://doi.org/10.1016/j.pedhc.2011.03.011> PMID: 23237611
235. Racil G, Elmontassar W, Rommene I, Tourny C, Chaouachi A, Coquart JB. Benefits of a regular vs irregular rhythm-based training programme on physical fitness and motor skills in obese girls. *J Endocrinol Invest*. 2017; 40(11):1227–34. <https://doi.org/10.1007/s40618-017-0689-8> PMID: 28528435
236. Racil G, Zouhal H, Elmontassar W, Ben Abderrahmane A, De Sousa MV, Chamari K, et al. Plyometric exercise combined with high-intensity interval training improves metabolic abnormalities in young obese females more so than interval training alone. *Appl Physiol Nutr Metab*. 2016; 41(1):103–9. <https://doi.org/10.1139/apnm-2015-0384> PMID: 26701117
237. Rauber SB, Castro HO, Marinho A, Vicente JB, Ribeiro HL, Monteiro LZ, et al. Effects of a physical activity and nutritional intervention in overweight and obese children through an educational and recreational camp. *Nutr Health*. 2018; 24(3):145–52. <https://doi.org/10.1177/0260106018771519> PMID: 29893164
238. Reed JA, Maslow AL, Long S, Hughey M. Examining the impact of 45 minutes of daily physical education on cognitive ability, fitness performance, and body composition of African American youth. *J Phys Act Health*. 2013; 10(2):185–97. <https://doi.org/10.1123/jpah.10.2.185> PMID: 22820756
239. Regaieg S, Charfi N, Kamoun M, Ghroubi S, Rebai H, Elleuch H, et al. The effects of an exercise training program on body composition and aerobic capacity parameters in Tunisian obese children. *Indian J Endocrinol Metab*. 2013; 17(6):1040–5. <https://doi.org/10.4103/2230-8210.122619> PMID: 24381881
240. Reznik M, Wylie-Rosett J, Kim M, Ozuah PO. A classroom-based physical activity intervention for urban kindergarten and first-grade students: a feasibility study. *Child Obes*. 2015; 11(3):314–24. <https://doi.org/10.1089/chi.2014.0090> PMID: 25747719
241. Rofey DL, Szigethy EM, Noll RB, Dahl RE, Lobst E, Arslanian SA. Cognitive-behavioral therapy for physical and emotional disturbances in adolescents with polycystic ovary syndrome: a pilot study. *J Pediatr Psychol*. 2009; 34(2):156–63. <https://doi.org/10.1093/jpepsy/jsn057> PMID: 18556675
242. Rowland TW, Varzeas MR, Walsh CA. Aerobic responses to walking training in sedentary adolescents. *J Adolesc Health*. 1991; 12(1):30–4. [https://doi.org/10.1016/0197-0070\(91\)90037-m](https://doi.org/10.1016/0197-0070(91)90037-m) PMID: 2007150
243. Ruebel ML, Heelan KA, Bartee T, Foster N. Outcomes of a Family Based Pediatric Obesity Program—Preliminary Results. *Int J Exerc Sci*. 2011; 4(4):217–28. PMID: 27182365
244. Sacchetti R, Cecilian A, Garulli A, Dallolio L, Beltrami P, Leoni E. Effects of a 2-year school-based intervention of enhanced physical education in the primary school. *J Sch Health*. 2013; 83(9):639–46. <https://doi.org/10.1111/josh.12076> PMID: 23879783
245. Sadeghi B, Kaiser LL, Hanbury MM, Tseregounis IE, Shaikh U, Gomez-Camacho R, et al. A three-year multifaceted intervention to prevent obesity in children of Mexican-heritage. *BMC Public Health*. 2019; 19(1):582. <https://doi.org/10.1186/s12889-019-6897-8> PMID: 31096944
246. Saenz KH. The use of heart rate monitors in enabling children to self-regulate physical activity behaviors: University of Central Florida; 2003.

247. Safdie M, Jennings-Aburto N, Levesque L, Janssen I, Campirano-Nunez F, Lopez-Olmedo N, et al. Impact of a school-based intervention program on obesity risk factors in Mexican children. *Salud Publica Mex.* 2013; 55 Suppl 3:374–87. <https://doi.org/10.21149/spm.v55s3.5138> PMID: 24643486
248. Salcedo Aguilar F, Martinez-Vizcaino V, Sanchez Lopez M, Solera Martinez M, Franquelo Gutierrez R, Serrano Martinez S, et al. Impact of an after-school physical activity program on obesity in children. *J Pediatr.* 2010; 157(1):36–42.e3. <https://doi.org/10.1016/j.jpeds.2009.12.046> PMID: 20227726
249. Scheffler C, Ketelhut K, Mohasseb I. Does physical education modify the body composition?—results of a longitudinal study of pre-school children. *Anthropol Anz.* 2007; 65(2):193–201. PMID: 17711151
250. Schwanke NL, Pohl HH, Reuter CP, Borges TS, de Souza S, Burgos MS. Differences in body posture, strength and flexibility in schoolchildren with overweight and obesity: A quasi-experimental study. *Man Ther.* 2016; 22:138–44. <https://doi.org/10.1016/j.math.2015.11.004> PMID: 26683874
251. Seabra A, Katzmarzyk P, Carvalho MJ, Seabra A, Coelho ESM, Abreu S, et al. Effects of 6-month soccer and traditional physical activity programmes on body composition, cardiometabolic risk factors, inflammatory, oxidative stress markers and cardiorespiratory fitness in obese boys. *J Sports Sci.* 2016; 34(19):1822–9. <https://doi.org/10.1080/02640414.2016.1140219> PMID: 26890580
252. Seltzer CC, Mayer J. An effective weight control program in a public school system. *Am J Public Health Nations Health.* 1970; 60(4):679–89. <https://doi.org/10.2105/ajph.60.4.679> PMID: 5461722
253. Serra-Paya N, Ensenyat A, Castro-Vinuales I, Real J, Sinfreu-Bergues X, Zapata A, et al. Effectiveness of a Multi-Component Intervention for Overweight and Obese Children (Nereu Program): A Randomized Controlled Trial. *PLoS One.* 2015; 10(12):e0144502. <https://doi.org/10.1371/journal.pone.0144502> PMID: 26658988
254. Shofan Y, Kedar O, Branski D, Berry E, Wilschanski M. A school-based program of physical activity may prevent obesity. *Eur J Clin Nutr.* 2011; 65(6):768–70. <https://doi.org/10.1038/ejcn.2011.25> PMID: 21427748
255. Siegrist M, Lammel C, Haller B, Christle J, Halle M. Effects of a physical education program on physical activity, fitness, and health in children: the JuvenTUM project. *Scand J Med Sci Sports.* 2013; 23(3):323–30. <https://doi.org/10.1111/j.1600-0838.2011.01387.x> PMID: 22092492
256. Sigmund E, El Ansari W, Sigmundova D. Does school-based physical activity decrease overweight and obesity in children aged 6–9 years? A two-year non-randomized longitudinal intervention study in the Czech Republic. *BMC Public Health.* 2012; 12:570. <https://doi.org/10.1186/1471-2458-12-570> PMID: 22892226
257. Silva HJ, Andersen LB, Lofrano-Prado MC, Barros MV, Freitas IF Jr., Hill J, et al. Improvements on Cardiovascular Diseases Risk Factors in Obese Adolescents: A Randomized Exercise Intervention Study. *J Phys Act Health.* 2015; 12(4):553–60. <https://doi.org/10.1123/jpah.2013-0199> PMID: 24778268
258. Silveira DS, Lemos L, Tassitano RM, Cattuzzo MT, Feitoza AHP, Aires L, et al. Effect of a pilot multi-component intervention on motor performance and metabolic risks in overweight/obese youth. *J Sports Sci.* 2018; 36(20):2317–26. <https://doi.org/10.1080/02640414.2018.1452142> PMID: 29558321
259. Singh AS, Chin APMJ, Brug J, van Mechelen W. Dutch obesity intervention in teenagers: effectiveness of a school-based program on body composition and behavior. *Arch Pediatr Adolesc Med.* 2009; 163(4):309–17. <https://doi.org/10.1001/archpediatrics.2009.2> PMID: 19349559
260. Slawta J, Bentley J, Smith J, Kelly J, Syman-Degler L. Promoting healthy lifestyles in children: a pilot program of be a fit kid. *Health Promot Pract.* 2008; 9(3):305–12. <https://doi.org/10.1177/1524839906289221> PMID: 16803930
261. Smith JJ, Beauchamp MR, Faulkner G, Morgan PJ, Kennedy SG, Lubans DR. Intervention effects and mediators of well-being in a school-based physical activity program for adolescents: The 'Resistance Training for Teens' cluster RCT. *Mental Health and Physical Activity.* 2018; 15:88–94.
262. Smith JJ, Morgan PJ, Plotnikoff RC, Stodden DF, Lubans DR. Mediating effects of resistance training skill competency on health-related fitness and physical activity: the ATLAS cluster randomised controlled trial. *J Sports Sci.* 2016; 34(8):772–9. <https://doi.org/10.1080/02640414.2015.1069383> PMID: 26194449
263. Sollerhed AC, Ejlertsson G. Physical benefits of expanded physical education in primary school: findings from a 3-year intervention study in Sweden. *Scand J Med Sci Sports.* 2008; 18(1):102–7. <https://doi.org/10.1111/j.1600-0838.2007.00636.x> PMID: 17490464
264. Son WM, Sung KD, Bharath LP, Choi KJ, Park SY. Combined exercise training reduces blood pressure, arterial stiffness, and insulin resistance in obese prehypertensive adolescent girls. *Clin Exp Hypertens.* 2017; 39(6):546–52. <https://doi.org/10.1080/10641963.2017.1288742> PMID: 28590143
265. Song JK, Stebbins CL, Kim TK, Kim HB, Kang HJ, Chai JH. Effects of 12 weeks of aerobic exercise on body composition and vascular compliance in obese boys. *J Sports Med Phys Fitness.* 2012; 52(5):522–9. PMID: 22976739

266. Southam MA, Kirkley BG, Murchison A, Berkowitz RI. A summer day camp approach to adolescent weight loss. *Adolescence*. 1984; 19(76):855–68. PMID: [6516934](#)
267. Speaker JG, Schultz C, Grinker JA, Stern JS. Body size estimation and locus of control in obese adolescent boys undergoing weight reduction. *Int J Obes*. 1983; 7(1):73–83. PMID: [6840969](#)
268. Springer NS, Angelocci J. Weight watchers program for trainables. *Mental Retardation*. 1973; 11(5):20–9.
269. Steinberg N, Eliakim A, Pantanowitz M, Kohen-Raz R, Zeev A, Nemet D. The effect of a weight management program on postural balance in obese children. *Eur J Pediatr*. 2013; 172(12):1619–26. <https://doi.org/10.1007/s00431-013-2090-8> PMID: [23881343](#)
270. Steinberg N, Rubinstein M, Nemet D, Ayalon M, Zeev A, Pantanowitz M, et al. Effects of a Program for Improving Biomechanical Characteristics During Walking and Running in Children Who Are Obese. *Pediatr Phys Ther*. 2017; 29(4):330–40. <https://doi.org/10.1097/PEP.0000000000000440> PMID: [28953178](#)
271. Sung KD, Pekas EJ, Scott SD, Son WM, Park SY. The effects of a 12-week jump rope exercise program on abdominal adiposity, vasoactive substances, inflammation, and vascular function in adolescent girls with prehypertension. *Eur J Appl Physiol*. 2019; 119(2):577–85. <https://doi.org/10.1007/s00421-018-4051-4> PMID: [30554386](#)
272. Tan S, Yang C, Wang J. Physical training of 9- to 10-year-old children with obesity to lactate threshold intensity. *Pediatr Exerc Sci*. 2010; 22(3):477–85. <https://doi.org/10.1123/pes.22.3.477> PMID: [20814042](#)
273. Taylor MJ, Mazzone M, Wrotniak BH. Outcome of an exercise and educational intervention for children who are overweight. *Pediatric Physical Therapy*. 2005; 17(3):180–8. <https://doi.org/10.1097/01.pep.0000176576.94482.9a> PMID: [16357672](#)
274. Telford RD, Cunningham RB, Waring P, Telford RM, Olive LS, Abhayaratna WP. Physical education and blood lipid concentrations in children: the LOOK randomized cluster trial. *PLoS One*. 2013; 8(10):e76124. <https://doi.org/10.1371/journal.pone.0076124> PMID: [24204594](#)
275. Ten Hoor GA, Rutten GM, Van Breukelen GJP, Kok G, Ruiter RAC, Meijer K, et al. Strength exercises during physical education classes in secondary schools improve body composition: a cluster randomized controlled trial. *Int J Behav Nutr Phys Act*. 2018; 15(1):92. <https://doi.org/10.1186/s12966-018-0727-8> PMID: [30253776](#)
276. Thivel D, Isacco L, Lazaar N, Aucouturier J, Ratel S, Dore E, et al. Effect of a 6-month school-based physical activity program on body composition and physical fitness in lean and obese schoolchildren. *Eur J Pediatr*. 2011; 170(11):1435–43. <https://doi.org/10.1007/s00431-011-1466-x> PMID: [21475968](#)
277. Togashi K, Masuda H, Iguchi K. Effect of diet and exercise treatment for obese Japanese children on abdominal fat distribution. *Res Sports Med*. 2010; 18(1):62–70. <https://doi.org/10.1080/15438620903423924> PMID: [20391247](#)
278. Tucker JM, Eisenmann JC, Howard K, Guseman EH, Yee KE, DeLaFuente K, et al. FitKids360: design, conduct, and outcomes of a stage 2 pediatric obesity program. *J Obes*. 2014; 2014:370403. <https://doi.org/10.1155/2014/370403> PMID: [25215228](#)
279. Vajda I, Meszaros J, Meszaros Z, Prokai A, Sziva A, Photiou A, et al. Effects of 3 hours a week of physical activity on body fat and cardio-respiratory parameters in obese boys. *Acta Physiol Hung*. 2007; 94(3):191–8. <https://doi.org/10.1556/APhysiol.94.2007.3.4> PMID: [17853771](#)
280. van Middelkoop M, Ligthart KAM, Paulis WD, van Teeffelen J, Kornelisse K, Koes BW. A multidisciplinary intervention programme for overweight and obese children in deprived areas. *Fam Pract*. 2017; 34(6):702–7. <https://doi.org/10.1093/fampra/cmz056> PMID: [28985299](#)
281. Vasconcellos F, Seabra A, Cunha F, Montenegro R, Penha J, Bouskela E, et al. Health markers in obese adolescents improved by a 12-week recreational soccer program: a randomised controlled trial. *J Sports Sci*. 2016; 34(6):564–75. <https://doi.org/10.1080/02640414.2015.1064150> PMID: [26208409](#)
282. Vignolo M, Rossi F, Bardazza G, Pistorio A, Parodi A, Spigno S, et al. Five-year follow-up of a cognitive-behavioural lifestyle multidisciplinary programme for childhood obesity outpatient treatment. *Eur J Clin Nutr*. 2008; 62(9):1047–57. <https://doi.org/10.1038/sj.ejcn.1602819> PMID: [17554247](#)
283. Villa-Gonzalez E, Ruiz JR, Mendoza JA, Chillón P. Effects of a school-based intervention on active commuting to school and health-related fitness. *BMC Public Health*. 2017; 17(1):20. <https://doi.org/10.1186/s12889-016-3934-8> PMID: [28056914](#)
284. Vissers D, De Meulenaere A, Vanroy C, Vanherle K, Van de Sompel A, Truijens S, et al. Effect of a multidisciplinary school-based lifestyle intervention on body weight and metabolic variables in overweight and obese youth. *e-SPEN*. 2008; 3(5):e196–e202.
285. Walther C, Gaede L, Adams V, Gelbrich G, Leichte A, Erbs S, et al. Effect of increased exercise in school children on physical fitness and endothelial progenitor cells: a prospective randomized trial.

- Circulation. 2009; 120(22):2251–9. <https://doi.org/10.1161/CIRCULATIONAHA.109.865808> PMID: 19920000
286. Wanless E, Judge LW, Dieringer ST, Bellar D, Johnson J, Plummer S. Pedometers and aerobic capacity: evaluating an elementary after-school running program. *ScientificWorldJournal*. 2014; 2014:370759. <https://doi.org/10.1155/2014/370759> PMID: 24723803
 287. Webber LS, Osganian SK, Feldman HA, Wu M, McKenzie TL, Nichaman M, et al. Cardiovascular risk factors among children after a 2 1/2-year intervention-The CATCH Study. *Prev Med*. 1996; 25(4):432–41. <https://doi.org/10.1006/pmed.1996.0075> PMID: 8818067
 288. Whipp BJ, Ruff WK. The effect of caloric restriction and physical training on the responses of obese adolescents to graded exercise. *J Sports Med Phys Fitness*. 1971; 11(3):146–53. PMID: 5133370
 289. Willi SM, Hirst K, Jago R, Buse J, Kaufman F, El Ghormli L, et al. Cardiovascular risk factors in multi-ethnic middle school students: the HEALTHY primary prevention trial. *Pediatr Obes*. 2012; 7(3):230–9. <https://doi.org/10.1111/j.2047-6310.2011.00042.x> PMID: 22461375
 290. Wong PC, Chia MY, Tsou IY, Wansaicheong GK, Tan B, Wang JC, et al. Effects of a 12-week exercise training programme on aerobic fitness, body composition, blood lipids and C-reactive protein in adolescents with obesity. *Ann Acad Med Singapore*. 2008; 37(4):286–93. PMID: 18461212
 291. Wright K, Giger JN, Norris K, Suro Z. Impact of a nurse-directed, coordinated school health program to enhance physical activity behaviors and reduce body mass index among minority children: a parallel-group, randomized control trial. *Int J Nurs Stud*. 2013; 50(6):727–37. <https://doi.org/10.1016/j.ijnurstu.2012.09.004> PMID: 23021318
 292. Yetgin MK, Agopyan A, Kucukler FK, Gedikbasi A, Yetgin S, Kayapinar FC, et al. The influence of physical training modalities on basal metabolic rate and leptin on obese adolescent boys. *J Pak Med Assoc*. 2018; 68(6):929–31. PMID: 30325913
 293. Yin Z, Moore JB, Johnson MH, Vernon MM, Gutin B. The impact of a 3-year after-school obesity prevention program in elementary school children. *Child Obes*. 2012; 8(1):60–70. <https://doi.org/10.1089/chi.2011.0085> PMID: 22799482
 294. Zguira MS, Slimani M, Bragazzi NL, Khrouf M, Chaieb F, Saiag B, et al. Effect of an 8-Week Individualized Training Program on Blood Biomarkers, Adipokines and Endothelial Function in Obese Young Adolescents with and without Metabolic Syndrome. *Int J Environ Res Public Health*. 2019; 16(5).
 295. Zhang H, Jiang L, Yang YJ, Ge RK, Zhou M, Hu H, et al. Aerobic exercise improves endothelial function and serum adiponin levels in obese adolescents independent of body weight loss. *Sci Rep*. 2017; 7(1):17717. <https://doi.org/10.1038/s41598-017-18086-3> PMID: 29255252
 296. Zhou Z, Ren H, Yin Z, Wang L, Wang K. A policy-driven multifaceted approach for early childhood physical fitness promotion: impacts on body composition and physical fitness in young Chinese children. *BMC Pediatr*. 2014; 14:118. <https://doi.org/10.1186/1471-2431-14-118> PMID: 24886119
 297. Zorba E, Cengiz T, Karacabey K. Exercise training improves body composition, blood lipid profile and serum insulin levels in obese children. *J Sports Med Phys Fitness*. 2011; 51(4):664–9. PMID: 22212270
 298. Alves ASR, Venâncio TL, Honório SAA, Martins JMC. Multicomponent training with different frequencies on body composition and physical fitness in obese children. *An Acad Bras Cienc*. 2019; 91(4): e20181264. <https://doi.org/10.1590/0001-3765201920181264> PMID: 31778458
 299. Brand C, Martins CML, Lemes VB, Pessoa MLF, Dias AF, Cadore EL, et al. Effects and prevalence of responders after a multicomponent intervention on cardiometabolic risk factors in children and adolescents with overweight/obesity: Action for health study. *J Sports Sci*. 2020; 38(6):682–91. <https://doi.org/10.1080/02640414.2020.1725384> PMID: 32050850
 300. Domaradzki J, Cichy I, Rokita A, Popowczak M. Effects of Tabata Training During Physical Education Classes on Body Composition, Aerobic Capacity, and Anaerobic Performance of Under-, Normal- and Overweight Adolescents. *Int J Environ Res Public Health*. 2020; 17(3).
 301. Gao Z, Lee JE, Zeng N, Pope ZC, Zhang Y, Li X. Home-Based Exergaming on Preschoolers' Energy Expenditure, Cardiovascular Fitness, Body Mass Index and Cognitive Flexibility: A Randomized Controlled Trial. *J Clin Med*. 2019; 8(10).
 302. Sepúlveda AR, Solano S, Blanco M, Lacruz T, Veiga O. Feasibility, acceptability, and effectiveness of a multidisciplinary intervention in childhood obesity from primary care: Nutrition, physical activity, emotional regulation, and family. *Eur Eat Disord Rev*. 2020; 28(2):184–98. <https://doi.org/10.1002/erv.2702> PMID: 31802570
 303. WHO. Global strategy on diet, physical activity and health. 2004.
 304. Volger S, Radler DR, Rothpletz-Puglia P. Early childhood obesity prevention efforts through a life course health development perspective: A scoping review. *PloS one*. 2018; 13(12). <https://doi.org/10.1371/journal.pone.0209787> PMID: 30592757

305. Ziviani J, Kopeshe R, Wadley D. Children walking to school: parent perceptions of environmental and psychosocial influences. *Australian Occupational Therapy Journal*. 2006; 53(1):27–34.
306. Ziviani J, Wadley D, Ward H, Macdonald D, Jenkins D, Rodger S. A place to play: socioeconomic and spatial factors in children's physical activity. *Australian occupational therapy journal*. 2008; 55(1):2–11. <https://doi.org/10.1111/j.1440-1630.2006.00646.x> PMID: 20887428
307. Tomiyama AJ, Carr D, Granberg EM, Major B, Robinson E, Sutin AR, et al. How and why weight stigma drives the obesity 'epidemic' and harms health. *BMC medicine*. 2018; 16(1):123. <https://doi.org/10.1186/s12916-018-1116-5> PMID: 30107800
308. Golan M. Parents as agents of change in childhood obesity—from research to practice. *International journal of pediatric obesity*. 2006; 1(2):66–76. <https://doi.org/10.1080/17477160600644272> PMID: 17907317
309. Rosenkranz RR, Dziewaltowski DA. Model of the home food environment pertaining to childhood obesity. *Nutrition reviews*. 2008; 66(3):123–40. <https://doi.org/10.1111/j.1753-4887.2008.00017.x> PMID: 18289177
310. Tzou IL, Chu N-F. Parental influence on childhood obesity: A review. 2012.
311. Sleddens SF, Gerards SM, Thijs C, De Vries NK, Kremers SP. General parenting, childhood overweight and obesity-inducing behaviors: a review. *International journal of pediatric obesity*. 2011; 6 (sup3):e12–27. <https://doi.org/10.3109/17477166.2011.566339> PMID: 21657834
312. Halliday JA, Palma CL, Mellor D, Green J, Renzaho A. The relationship between family functioning and child and adolescent overweight and obesity: a systematic review. *International journal of obesity*. 2014; 38(4):480–93. <https://doi.org/10.1038/ijo.2013.213> PMID: 24232501
313. Oude Luttikhuis H, Baur L, Jansen H, Shrewsbury VA, O'malley C, Stolk RP, et al. Cochrane review: Interventions for treating obesity in children. *Evidence-based Child Health: A Cochrane Review Journal*. 2009; 4(4):1571–729. <https://doi.org/10.1002/14651858.CD001872.pub2> PMID: 19160202
314. Edmunds L. Parents' perceptions of health professionals' responses when seeking help for their overweight children. *Family practice*. 2005; 22(3):287–92. <https://doi.org/10.1093/fampra/cmh729> PMID: 15772121
315. Robinson E, Sutin AR. Parents' perceptions of their children as overweight and children's weight concerns and weight gain. *Psychological science*. 2017; 28(3):320–9. <https://doi.org/10.1177/0956797616682027> PMID: 28084895
316. Lowry KW, Sallinen BJ, Janicke DM. The effects of weight management programs on self-esteem in pediatric overweight populations. *Journal of Pediatric Psychology*. 2007; 32(10):1179–95. <https://doi.org/10.1093/jpepsy/jsm048> PMID: 17584780
317. APA. Scope of practice: position statement 2009 [https://australian.physio/sites/default/files/RESOURCES/Advocacy_Position_Scope_of_Practice_2009.pdf].
318. Dhar P, Robinson C. Physical activity and childhood obesity. *Applied Economics Letters*. 2016; 23 (8):584–7.
319. Juhola J, Magnussen CG, Viikari JS, Kähönen M, Hutri-Kähönen N, Jula A, et al. Tracking of serum lipid levels, blood pressure, and body mass index from childhood to adulthood: the Cardiovascular Risk in Young Finns Study. *The Journal of pediatrics*. 2011; 159(4):584–90. <https://doi.org/10.1016/j.jpeds.2011.03.021> PMID: 21514597
320. Chen X, Wang Y. Tracking of blood pressure from childhood to adulthood: a systematic review and meta-regression analysis. *Circulation*. 2008; 117(25):3171. <https://doi.org/10.1161/CIRCULATIONAHA.107.730366> PMID: 18559702
321. Freedman D, Patel D, Srinivasan S, Chen W, Tang R, Bond M, et al. The contribution of childhood obesity to adult carotid intima-media thickness: the Bogalusa Heart Study. *International journal of obesity*. 2008; 32(5):749–56. <https://doi.org/10.1038/sj.ijo.0803798> PMID: 18227845
322. Baker JL, Olsen LW, Sørensen TI. Childhood body-mass index and the risk of coronary heart disease in adulthood. *New England journal of medicine*. 2007; 357(23):2329–37. <https://doi.org/10.1056/NEJMoa072515> PMID: 18057335
323. Morrison JA, Glueck CJ, Horn PS, Yeramaneni S, Wang P. Pediatric triglycerides predict cardiovascular disease events in the fourth to fifth decade of life. *Metabolism*. 2009; 58(9):1277–84. <https://doi.org/10.1016/j.metabol.2009.04.009> PMID: 19501856
324. Friedemann C, Heneghan C, Mahtani K, Thompson M, Perera R, Ward AM. Cardiovascular disease risk in healthy children and its association with body mass index: systematic review and meta-analysis. *Bmj*. 2012; 345:e4759. <https://doi.org/10.1136/bmj.e4759> PMID: 23015032
325. Rosner B, Cook N, Portman R, Daniels S, Falkner B. Blood pressure differences by ethnic group among United States children and adolescents. *Hypertension*. 2009; 54(3):502–8. <https://doi.org/10.1161/HYPERTENSIONAHA.109.134049> PMID: 19652080

326. Freedman DS, Dietz WH, Srinivasan SR, Berenson GS. The relation of overweight to cardiovascular risk factors among children and adolescents: the Bogalusa Heart Study. *Pediatrics*. 1999; 103(6):1175–82. <https://doi.org/10.1542/peds.103.6.1175> PMID: 10353925
327. Deitz JC, Kartin D, Kopp K. Review of the Bruininks-Oseretsky test of motor proficiency, (BOT-2). *Physical & occupational therapy in pediatrics*. 2007; 27(4):87–102.
328. Faigenbaum AD, Lloyd RS, Sheehan D, Myer GD. The role of the pediatric exercise specialist in treating exercise deficit disorder in youth. *Strength & Conditioning Journal*. 2013; 35(3):34–41.
329. Mistry K, Yonezawa E, Milne N. Paediatric Physiotherapy curriculum: an audit and survey of Australian entry-level Physiotherapy programs. *BMC medical education*. 2019; 19(1):109. <https://doi.org/10.1186/s12909-019-1540-z> PMID: 30992074
330. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *International journal of social research methodology*. 2005; 8(1):19–32.
331. Peters MD, Godfrey CM, Khalil H, McInerney P, Parker D, Soares CB. Guidance for conducting systematic scoping reviews. *JBI Evidence Implementation*. 2015; 13(3):141–6. <https://doi.org/10.1097/XEB.0000000000000050> PMID: 26134548