

Antioxidants inhibit low density lipoprotein oxidation less at lysosomal pH: A possible explanation as to why the clinical trials of antioxidants might have failed

Feroz Ahmad*, David S. Leake

Institute of Cardiovascular and Metabolic Research, University of Reading, Reading, United Kingdom

ARTICLE INFO

Keywords:

Low density lipoproteins
Macrophages
Lipid peroxidation
Probulcol
Cysteamine
N,N-diphenyl-1,4-phenylenediamine

ABSTRACT

Oxidised low density lipoprotein (LDL) was considered to be important in the pathogenesis of atherosclerosis, but the large clinical trials of antioxidants, including the first one using probucol (the PQRST Trial), failed to show benefit and have cast doubt on the importance of oxidised LDL. We have shown previously that LDL oxidation can be catalysed by iron in the lysosomes of macrophages. The aim of this study was therefore to investigate the effectiveness of antioxidants in preventing LDL oxidation at lysosomal pH and also establish the possible mechanism of oxidation. Probulcol did not effectively inhibit the oxidation of LDL at lysosomal pH, as measured by conjugated dienes or oxidised cholesteryl esters or tryptophan residues in isolated LDL or by ceroid formation in the lysosomes of macrophage-like cells, in marked contrast to its highly effective inhibition of LDL oxidation at pH 7.4. LDL oxidation at lysosomal pH was inhibited very effectively for long periods by *N,N*-diphenyl-1,4-phenylenediamine, which is more hydrophobic than probucol and has been shown by others to inhibit atherosclerosis in rabbits, and by cysteamine, which is a hydrophilic antioxidant that accumulates in lysosomes. Iron-induced LDL oxidation might be due to the formation of the superoxide radical, which protonates at lysosomal pH to form the much more reactive, hydrophobic hydroperoxyl radical, which can enter LDL and reach its core. Probulcol resides mainly in the surface monolayer of LDL and would not effectively scavenge hydroperoxyl radicals in the core of LDL. This might explain why probucol failed to protect against atherosclerosis in various clinical trials. The oxidised LDL hypothesis of atherosclerosis now needs to be re-evaluated using different and more effective antioxidants that protect against the lysosomal oxidation of LDL.

1. Introduction

Oxidation of low density lipoprotein (LDL) was widely believed to be a critical step in the pathogenesis of atherosclerosis, the underlying cause of myocardial infarctions and thrombotic strokes (Steinberg and Witztum, 2010). The oxidised LDL hypothesis proposes that cells in the arterial wall oxidise LDL in the extracellular space and then take it up rapidly, leading to the formation of the foam cells characteristic of atherosclerosis (Steinberg and Witztum, 2010). Two key problems with this conventional view are that the oxidation is inhibited by low concentrations of interstitial fluid (or serum) (Dabbagh and Frei, 1995), and that large clinical trials have shown no protection by antioxidants against cardiovascular disease (Steinberg and Witztum, 2002). To address the shortcomings in the conventional LDL oxidation theory, we proposed that nonoxidatively modified LDL is rapidly endocytosed by macrophages and then oxidised within lysosomes (Wen and Leake, 2007). We showed that 7 days after taking up mechanically-aggregated

(vortexed) LDL, mouse J774 cells (a macrophage-like cell line) and human monocyte-derived macrophages (HMDM) generated ceroid (an advanced lipid oxidation product present in human atherosclerotic lesions) in their lysosomes (Wen and Leake, 2007). Furthermore, there was an increased production of oxysterols (e.g. 7-ketocholesterol) in the J774 cells when incubated with nonoxidized acetylated LDL which is rapidly endocytosed by macrophages, and then incubated in the absence of extracellular lipoproteins. Chloroquine, a weak base that concentrates in lysosomes and increases their pH, inhibited the oxidation of LDL, suggesting that the oxidation takes place at acidic pH. It was further shown that the oxidation in lysosomes is mediated by iron which is highly effective in oxidising LDL at pH 4.5, the approximate pH of lysosomes, but very poor at doing so at pH 7.4 in a simple buffer (Satchell and Leake, 2012; Wen and Leake, 2007).

Iron is a transition metal capable of redox cycling between the oxidation states of Fe^{2+} and Fe^{3+} . Whether iron in lysosomes exists in ferric or ferrous state has been a matter of discussion, (Collins et al.,

* Corresponding author at: School of Biological Sciences, Hopkins Building, University of Reading, Whiteknights, Reading, Berkshire, RG6 6UB, United Kingdom.
E-mail address: f.ahmad@reading.ac.uk (F. Ahmad).

1991) and both might be present (Meguro et al., 2005), but because of the reducing environment within lysosomes the ferrous oxidation state of iron is likely to be favoured (Terman and Kurz, 2013). Various theories have been proposed for the mechanism of LDL oxidation by iron. Some suggest that iron may be able to catalyse the oxidation of LDL by the production of hydroxyl radicals from hydrogen peroxide (Halliwell and Gutteridge, 1986), while others believe that superoxide ion is required for the initiation of LDL oxidation (Lynch and Frei, 1993). It is also believed that iron may be able to initiate the oxidation of LDL by reacting with pre-existing lipid hydroperoxides (LOOH) in LDL particles (Esterbauer et al., 1990).

These mechanisms are based on reactions occurring at physiological pH (Lynch and Frei, 1993), at which iron is practically insoluble and precipitates unless bound to various molecules, and might not be indicative of the reactions that occur at lysosomal pH. We show here that antioxidants that have been used to test the oxidised LDL hypothesis do not inhibit the oxidation of LDL effectively at lysosomal pH and hence might not be a true test for this hypothesis. We propose another antioxidant, cysteamine, which accumulates in lysosomes and inhibits the oxidation of LDL effectively at acidic pH, which might provide a better test for the oxidised LDL hypothesis.

2. Materials and methods

Chemicals and reagents used in this study were purchased from Sigma-Aldrich, Dorset, UK, or Fisher Scientific Ltd, Loughborough, UK, unless otherwise stated. Solutions were prepared using ultrapure water generated from a Barnstead Nanopure system. Organic solvents were HPLC or molecular biology grade.

2.1. LDL isolation

Blood was taken from healthy volunteers after overnight fasting using EDTA (final concentration 3 mmol/L) as the anticoagulant. LDL (1.019–1.063 g/mL) was isolated from the plasma by sequential density ultracentrifugation at 4 °C, as described previously (Wilkins and Leake, 1994). LDL was stored in the presence of 100 µmol/L EDTA in the dark at 4 °C and used within 1 month.

2.2. Aggregation of LDL with sphingomyelinase

LDL was aggregated by incubating native LDL (2 mg protein/mL) with sphingomyelinase (*Bacillus cereus*, catalogue number S9396-25UN, Sigma) at 10 mU/mL (Walters and Wrenn, 2010) until the attenuation (absorbance plus light scattering) at 680 nm increased from about 0.0017–0.027. Sphingomyelinase-aggregated LDL (SMase-LDL) was dialysed against a phosphate buffer containing EDTA and sterilised by membrane filtration (0.45 µm).

2.3. Measurement of conjugated dienes

LDL (50 µg LDL protein/ml) was oxidised with freshly dissolved FeSO₄ (5 µmol/L) or FeCl₃ (5 µmol/L) at 37 °C in a NaCl/sodium acetate buffer (NaCl 150 mmol/L, sodium acetate 10 mmol/L; pH 4.5) in capped quartz cuvettes and conjugated dienes were monitored in the presence or absence of antioxidants using a method based on that of Esterbauer et al. (Esterbauer et al., 1989). The change in attenuation at 234 nm was measured at 37 °C against reference cuvettes containing all the components except LDL. Measurements were taken at one minute intervals in a Lambda-2 6-cell or a Lambda Bio 40 8-cell spectrophotometer with UV Winlab software. The attenuation at time zero was subtracted from all values, and the time taken to reach an attenuation of 0.1 was used as a reference since this corresponds to the rapid oxidation phase (Satchell and Leake, 2012).

2.4. Loss of LDL-tryptophan fluorescence measurement

The decrease in lipoprotein bound tryptophan fluorescence was measured on a Cary Eclipse fluorescence spectrophotometer using the time-drive method at an emission wavelength of 331 nm, with excitation set at 282 nm (Giessauf et al., 1995). The emission and excitation slits were set at 10 nm to obtain optimal fluorescence output. LDL (50 µg LDL protein/ml) was oxidised by freshly dissolved FeSO₄ (5 µmol/L) at 37 °C in the NaCl/sodium acetate buffer, pH 4.5 in capped quartz cuvettes and the tryptophan fluorescence was measured every 10 min.

2.5. Monitoring of Fe²⁺ levels using bathophenanthrolinedisulfonic acid

The ferrous iron chelator bathophenanthrolinedisulfonic acid (BP) was used to measure ferrous iron concentrations. BP forms a red color complex with Fe (II) with a molar absorption coefficient of 22,140 L mol⁻¹ cm⁻¹ at 535 nm (Pieroni et al., 2001). LDL (50 µg protein/ml) was oxidised with freshly prepared FeSO₄ (5 µmol/L) in the NaCl/sodium acetate buffer in polypropylene tubes in a water bath at 37 °C. Samples of 1 mL were taken at different time points into new tubes, 30 µL of BP (10 mmol/L) was added to them and absorbance was measured immediately at 535 nm with a spectrophotometer.

2.6. HPLC analysis

Lipids were extracted from oxidised LDL for HPLC analysis using methanol and hexane (Kritharides et al., 1993; Satchell and Leake, 2012). The upper hexane layer was collected and evaporated at ambient temperature in a SpeedVac Concentrator System (ThermoFisher), the residue was dissolved in the relevant mobile phase before injection into the HPLC. Lipid species were separated by reverse phase HPLC in a Waters C18 column (250 mm × 4.6 mm, 5 µm particle size, 5 µm guard column) with a Agilent 1100 HPLC system. Cholesteryl esters were detected at 210 nm using an acetonitrile/2-propanol/water mobile phase (44/54/2, by volume) and a flow rate of 1.2 mL/min. Cholesteryl linoleate hydroperoxide and 7-ketocholesterol were detected at 234 nm using an acetonitrile/2-propanol/water mobile phase (50/48.8/1.2, by volume) and a flow rate of 1 mL/min. The identities of the peaks were confirmed by mass spectrometry (data not shown), and the lipids were quantified using commercially available standards.

2.7. Measurement of lipid hydroperoxides

Lipid hydroperoxides (LOOH) were measured using a method adapted from that described by el-Saadani et al. (1989) Standard concentrations of H₂O₂ were made up using pure water. Samples and standards (250 µL) were added to assay tubes in triplicate and 1 mL of color reagent (0.2 mol/L potassium phosphate, 0.12 mol/L potassium iodide, 0.15 mmol/L sodium azide, polyethyleneglycol mono[*p*-(1,1',3,3'-tetramethyl-butyl)-phenyl]ether (2 g/L), alkylbenzyltrimethylammonium chloride (0.1 g/L), ammonium molybdate (10 µmol/L); pH 6.0–6.2) was added to each tube. After leaving the tubes in the dark for 1 h, the absorbance at 365 nm was measured. The levels of lipid hydroperoxides were also confirmed by HPLC method as described above (Fig. S1).

2.8. Cell culture

THP-1 cells were purchased from the European Collection of Cell Cultures (Salisbury, UK), and cultured in RPMI 1640 containing L-glutamine to which was added penicillin (50 IU/mL), streptomycin (50 µg/mL) and amphotericin B (0.95 µg/mL). THP-1 cells were differentiated into macrophages on 18 × 18 mm glass coverslips in 6-well tissue culture plates using 25 nM phorbol 12-myristate 13-acetate (25 nmol/L) for 72 h. The cells were then washed three times with PBS




Fig. 1. Oxidation of LDL catalysed by FeSO₄ and FeCl₃ at pH 4.5. LDL (50 µg protein/ml) in NaCl/sodium acetate buffer (pH 4.5) was incubated with either FeSO₄ or FeCl₃ (5 µmol/L) at 37 °C in quartz cuvettes. LDL oxidation was monitored by measuring the change in attenuance at 234 nm against appropriate reference cuvettes (A). Time taken to reach an attenuance increase of 0.1 is shown (B). Tryptophan fluorescence was measured every 10 min at an excitation wavelength of 282 nm and emission wavelength 331 nm in a spectrofluorometer (C). The decrease in fluorescence after 150 min is shown (D). Decrease of Fe²⁺ concentration during LDL oxidation by FeSO₄ (E). These are representative examples of at least three independent experiments (A and C). * indicates $p < 0.05$; paired t -test (B and D).

lipid peroxidation by scavenging LOO• (Takahashi et al., 1989). DPPD has polar surface area of 24.1 Å² (PubChem) which is much lower than that of probucol which has a polar surface area of 91.1 Å² (PubChem) and hence DPPD would be expected to concentrate more in the hydrophobic core of LDL than does probucol. DPPD completely inhibited

cholesteryl linoleate hydroperoxide (CLOOH) formation by iron at pH 4.5, whereas probucol had no inhibitory effect (Fig. 4A). DPPD completely protected against the loss of cholesteryl linoleate and cholesteryl arachidonate, whereas probucol had no effect (Fig. 4B and C) (a representative HPLC chromatogram is shown in Fig. S3). Furthermore,




Fig. 2. Effect of HPODE on LDL oxidation catalysed by $5 \mu\text{M}$ FeSO_4 at pH 4.5. Native LDL and HPODE-rich native LDL (30 or 60 nmol/mg LDL protein) were incubated with $5 \mu\text{M}$ FeSO_4 at pH 4.5 in NaCl/sodium acetate buffer (pH 4.5) at 37°C in quartz cuvettes. Addition of lipid hydroperoxide did not increase the rate of LDL oxidation measured at 234 nm (A). There was no significant difference in the times the samples took to reach an attenuation of 0.1 (B). Addition of HPODE had no effect on the loss of LDL-tryptophan fluorescence during LDL oxidation (C). The decrease in tryptophan fluorescence at 150 min in the absence or presence of HPODE is shown (D). These are representative examples of at least three independent experiments. (***) indicates $p < 0.001$ compared to the control without iron.

DPPD prevented conjugated diene formation during oxidation with iron (Fig. 4D). Probucol has limited access to the core so it is unable to prevent the oxidation of the cholesteryl esters, whereas DPPD gains access to the core of LDL and is able to completely inhibit its oxidation.

4.6. Lysosomal ceroid formation

To confirm the above findings of probucol's failure as an antioxidant for the initial oxidation of LDL at lysosomal pH, we measured ceroid in the lysosomes of cultured macrophages. Ceroid is a polymerised product of lipid and protein oxidation found within foam cells in human atherosclerotic lesions (Mitchinson et al., 1985). As mentioned above, we have previously shown that mechanically aggregated LDL is rapidly taken up by human macrophages and oxidised in lysosomes generating ceroid (Wen and Leake, 2007). We have also shown that LDL aggregated by sphingomyelinase is internalised by macrophages and oxidised in lysosomes (Wen et al., 2014). Cysteamine (2-aminoethanethiol) (structure shown in Fig. 6E) is an antioxidant, which concentrates in lysosomes by several orders of magnitude, and is used in the treatment of a rare lysosomal storage disorder called cystinosis (Brodin-Sartorius et al., 2012). THP-1 macrophages were treated with native LDL or SMase-LDL (200 μg protein/ml) or without LDL for 24 h and were washed and incubated for 7 days in lipoprotein-deficient medium with or without the antioxidants, probucol (10 $\mu\text{mol/L}$) and cysteamine (10 $\mu\text{mol/L}$). The cells were then stained for ceroid after other lipids had been removed by organic solvents. Ceroid was clearly visible in the form of Oil Red O stained, irregularly shaped granules in

cells treated with SMase-LDL (Fig. 5C) (Mitchinson, 1982). Little ceroid was present in cells that had been incubated without LDL (Fig. 5A), but was increased in cells incubated with native LDL ($p < 0.05$) (Fig. 5B). There was no significant decrease in the ceroid development in cells incubated with probucol (Fig. 5D), but cells which were treated with cysteamine (Fig. 5E) showed a significant reduction in ceroid formation. These results show that lysosomal oxidation of LDL is not prevented by probucol but can be inhibited by the lysosomotropic drug cysteamine.

4.7. Effect of cysteamine on LDL oxidation catalysed by iron at lysosomal pH

As cysteamine inhibited ceroid formation, we investigated the effect of cysteamine on LDL oxidation catalysed by iron at pH 4.5. Cysteamine showed a concentration-dependent inhibition of LDL oxidation catalysed by iron with 25 $\mu\text{mol/L}$ preventing oxidation for about 8 h and 250 $\mu\text{mol/L}$ for over 30 h (Fig. 6A and B). The lower concentration was chosen to conduct further experiments. Cysteamine (25 $\mu\text{mol/L}$) concentration decreased the oxidation of tryptophan residues of apoB-100 by iron for about first 8 h (Fig. 6C and D). Furthermore, cysteamine completely prevented the loss of cholesteryl arachidonate (Fig. 7A) and cholesteryl linoleate (Fig. 7B) for 6 h and also prevented the formation of cholesteryl linoleate hydroperoxide (Fig. 7C) and 7-ketocholesterol (Fig. 7D) for 6 h. These results indicate that cysteamine is able to prevent the oxidation of the hydrophobic cholesteryl ester core of LDL and of nonesterified cholesterol, which is located mainly in the surface




Fig. 3. Effect of probucol on LDL oxidation catalysed by iron at pH 4.5. LDL (50 µg protein/ml) was incubated with FeSO₄ in the presence or absence of 2 µM probucol at pH 4.5 in NaCl/sodium acetate buffer at 37 °C in quartz cuvettes; Oxidation was monitored by measuring the change in attenuation at 234 nm against appropriate reference cuvettes (A). The time required for the attenuation to reach 0.1 was not significantly increased by probucol (B). Probucol was also unable to effectively protect the initial loss of LDL-tryptophan fluorescence during incubation with iron at pH 4.5 (C). After 80 min of LDL oxidation the fluorescence intensity decrease was not significantly different (*P* > 0.05) from that in the presence of probucol (D). Structure of probucol (E). These are representative examples of at least three independent experiments. Mean ± SEM, paired *t*-test.

monolayer of LDL (Lund-Katz and Phillips, 1986) as well as the oxidation of tryptophan residues in apoB-100.

5. Discussion

We previously showed that nonoxidatively modified LDL is endocytosed by macrophages and oxidised by iron in their lysosomes

(Wen and Leake, 2007). Iron levels are increased by an order of magnitude in human and animal atherosclerotic lesions (Sullivan, 2009). Iron is highly effective in oxidising LDL at pH 4.5, the approximate pH of lysosomes, but very poor at doing so at pH 7.4 in a simple buffer (Wen and Leake, 2007). Animal studies have shown a positive correlation between iron-deposition within plaques and the severity of atherosclerosis (Lee et al., 1999; Yuan et al., 1996). Low molecular






Fig. 4. HPLC analysis of LDL oxidation catalysed by FeSO₄ at pH 4.5. LDL (50 µg protein/mL) in NaCl/sodium acetate buffer at pH 4.5 was incubated with 5 µM FeSO₄ at 37 °C in the presence or absence of probucol (2 µmol/L) or DPPD (2 µmol/L) and samples were taken every 30 min to be analysed by reverse-phase HPLC for cholesterol linoleate hydroperoxide (CLOOH) (A) at 234 nm and cholesteryl linoleate (B) and cholesteryl arachidonate (C) at 210 nm. Measure of conjugated diene formation in presence of DPPD (D). Structure of DPPD (E). The statistics shown are two way ANOVA followed by Bonferroni post-test ($p < 0.0001$) compared to the control. Graphs represent the mean \pm the SEM of at least three independent experiments.

weight, probably redox-active iron has been shown to be present in lysosomes of macrophage foam cells in human atherosclerotic lesions, probably because of the degradation of ferritin, the autophagy of iron-containing organelles or the phagocytosis of erythrocytes (Kurz et al., 2008). Further to this, L-ferritin levels are increased in coronary arteries from patients with coronary artery disease, indicating that iron accumulates in atherosclerotic plaques (You et al., 2003). The epidemiology of iron status and coronary heart disease is inconsistent (Sempos, 2002). This might be because measuring iron in plasma is not a good measure of iron inside macrophages. For instance, hepcidin is increased by inflammation and decreases iron in plasma but increases it in macrophages (Sullivan, 2009).

It has been proposed that transition metals may be able to initiate the oxidation of LDL by reacting with pre-existing lipid hydroperoxides in the LDL particles (Frei and Gaziano, 1993) and ferrous iron is believed to react faster with lipid hydroperoxides than does ferric iron

(Esterbauer et al., 1992).



The above reactions have been proposed on the basis of rate constants for the analogous reactions of Fe²⁺ and Fe³⁺ with H₂O₂ and not the actual lipid hydroperoxides (Minotti and Aust, 1992). In addition, the rate of these reactions is dependent on pH and ferric iron is insoluble at pH 7.4 and precipitates (Tachiev et al., 2000).

The current study shows that the initial addition of a lipid hydroperoxide (13-HPODE) does not increase the rate of LDL oxidation with iron at pH 4.5. In fact, addition of a higher amount of 13-HPODE delayed the later stage of oxidation of LDL, maybe because it was converting Fe²⁺ to Fe³⁺, which is less effective in oxidising LDL (Fig. 1A). It has previously been shown that the rate of oxidation of LDL by copper




Fig. 5. Inhibition of ceroid formation in THP-1 macrophages. THP-1 macrophage-like cells were incubated with native or SMase-LDL (200 μg protein/mL) for 24 h. They were then incubated for a further 7 days in the absence of lipoproteins in RPMI with 10% (v/v) LPDS in the presence or absence of probucol or cysteamine. A shows cells incubated without LDL or antioxidants, B shows cells incubated with native LDL only, C shows cells incubated with SMase-LDL but without antioxidants, D shows cells incubated with SMase-LDL and then probucol (10 μmol/L) and E shows cells incubated with SMase-LDL and then cysteamine (10 μmol/L). The ceroid levels in the cells were quantified using ImageJ as mean integrated density of at least 100 cells and expressed as percent decrease of cells treated with only SMase-LDL (F). Mean ± SEM of 3 independent experiments. * $p < 0.05$, ** $p < 0.01$, compared to SMase-LDL, ANOVA and Dunnett's test.

at pH 7.4 is increased considerably by HPODE (Patterson et al., 2003). Therefore, the present data suggest that the initial oxidation of LDL induced by iron at lysosomal pH is independent of the presence of lipid hydroperoxides.

The concentration of Fe^{2+} decreased rapidly to low levels during the first 60 min of LDL oxidation at pH 4.5 (Fig. 1E), which corresponds almost exactly to the rapid phase of LDL oxidation (Fig. 1A). Our previous work has also shown that adding higher concentrations of Fe^{2+}

shortened the lag phase and increased the rates of the rapid, slow, and aggregation phases of LDL oxidation, whereas adding higher concentrations of Fe^{3+} had little effect (Satchell and Leake, 2012). The formation of conjugated dienes (Fig. 1A) and decrease in tryptophan fluorescence (Fig. 1C) both occurred faster with Fe^{2+} than with Fe^{3+} , again suggesting the important role of ferrous compared to ferric iron.

Probucol efficiently inhibits LDL oxidation by cells or copper at physiological pH (Parthasarathy et al., 1986), by scavenging lipid




Fig. 6. Effect of cysteamine on LDL oxidation catalysed by iron at pH 4.5. LDL (50 μg protein/ml) was oxidised in the presence or absence of cysteamine by FeSO_4 (5 $\mu\text{mol/L}$) at pH 4.5 in NaCl/sodium acetate buffer at 37 $^{\circ}\text{C}$. Oxidation was monitored by measuring the change in attenuation at 234 nm against appropriate reference cuvettes (A). The time required for the attenuation to reach 0.1 was significantly increased by cysteamine (25 $\mu\text{mol/L}$) ($p < 0.001$) (B) mean \pm SEM, paired t -test, $***p < 0.001$. Cysteamine (25 $\mu\text{mol/L}$) prevented the initial loss of LDL-tryptophan fluorescence on oxidation with iron (C). After 150 min of LDL oxidation with iron, the fluorescence intensity decrease was significantly less ($p < 0.001$) in the presence of cysteamine (D), mean \pm SEM, paired t -test. (E) shows the structure of cysteamine.

radicals ($\text{L}\cdot$, $\text{LO}\cdot$, $\text{LOO}\cdot$) and hence inhibiting the lipid peroxidation chain reaction. The initial oxidation of LDL catalysed by iron at pH 4.5, however, was not effectively inhibited by probucol measured by either conjugated dienes (Fig. 3A, B) or loss of tryptophan fluorescence (Fig. 3C, D). In support of this, the oxidation of the cholesteryl esters in the core of LDL was not inhibited by probucol, as measured in terms of the formation of cholesteryl linoleate hydroperoxide or loss of cholesteryl linoleate or arachidonate (Fig. 5). In contrast, cholesteryl ester oxidation was inhibited entirely by the very hydrophobic antioxidant DPPD. Probucol inhibited the oxidation of LDL after about 100 min and this might be due to an antioxidant effect on the phospholipid monolayer of LDL. This suggests that the oxidation of LDL catalysed by iron at pH 4.5 starts in the hydrophobic core (containing cholesteryl esters and triacylglycerols), where DPPD would be expected to accumulate, rather than in the surface monolayer (containing mainly phospholipids and nonesterified cholesterol) where probucol mainly resides (Bard et al., 1994),

Tryptophan loss is one of the most significant oxidative changes in oxidised proteins due to the high susceptibility of this amino acid to reactive oxygen species (Ronsein et al., 2011). ApoB-100 of LDL contains 37 tryptophan residues and these have been proposed to be responsible for the initiation of LDL oxidation by copper ions (Giessauf et al., 1995). We found that there was loss of LDL-tryptophan fluorescence during iron-catalysed oxidation at pH 4.5, which was unaffected by the addition of hydroperoxides and was not prevented by probucol. Although, DPPD greatly prevented the formation of conjugated dienes (Fig. 5D), we could not investigate the effect of DPPD on the loss of tryptophan fluorescence because DPPD absorbs at 331 nm (Linschitz et al., 1967) and quenched the fluorescence of tryptophan (Fig. S2). It was also noted that the loss of LDL-tryptophan fluorescence occurs faster when LDL is oxidised with Fe^{2+} than Fe^{3+} (Fig. 1C, D). Lynch and Frei (1993) suggested that reduction of Fe^{3+} to Fe^{2+} in the presence of an exogenous reductant such as superoxide ($\text{O}_2^{\cdot-}$) is essential for iron-catalysed LDL oxidation at pH 7.4.






Fig. 7. HPLC analysis of LDL oxidation catalysed by FeSO_4 with or without cysteamine. LDL (50 μg protein/ml) was incubated with FeSO_4 (5 $\mu\text{mol/L}$) in the presence or absence of cysteamine at pH 4.5 in NaCl/sodium acetate buffer at 37 $^\circ\text{C}$. Samples were analysed by reverse-phase HPLC for cholesteryl linoleate (A), cholesterol arachidonate (B) and cholesteryl linoleate hydroperoxide (C) and 7-ketocholesterol (D). The statistics shown are two way ANOVA followed by Bonferroni post-test (* $p < 0.05$, *** $p < 0.001$) compared to the control. Graphs represent the mean \pm the SEM of at least three independent experiments.



The concentration of Fe^{2+} falls rapidly when LDL is oxidised at pH 4.5 (Fig. 1E) and is converted presumably to Fe^{3+} generating superoxide radicals (Morgan and Lahav, 2007).



The superoxide radical undergoes protonation at pH 4.5 to form its conjugate acid, hydroperoxyl radical, which is highly reactive (De Grey, 2002).






At pH 7.4, only about 0.25% of superoxide would be present as hydroperoxyl radical, whereas at pH 4.5 about 67% of it would be present as the hydroperoxyl radical. Hydroperoxyl radical is a much more potent oxidant than superoxide anion and capable of abstracting a hydrogen atom from a polyunsaturated fatty acid (De Grey, 2002) as well as tryptophan (Dubinina et al., 2002) and oxidising LDL (Bedwell et al., 1989). Superoxide anions are negatively charged and hydrophilic and might not be able to enter the core of LDL or the hydrocarbon chain region of the surface phospholipid monolayer. Hydroperoxyl radicals, as well as being more reactive than superoxide anions, are not charged and would be able to diffuse from the aqueous phase into the core of the LDL particles and abstract hydrogen atoms from the polyunsaturated fatty acyl groups of cholesteryl esters and triacylglycerols (Bedwell et al., 1989).




Lipid radicals and tryptophan radicals would react with oxygen to

form their respective peroxy radicals. The lipid peroxy radicals would then lead to the lipid peroxidation chain reaction.



Antioxidants containing thiol groups are potent nucleophiles and ready to interact with electrophilic groups of reactive oxygen species (ROS) (Güngör et al., 2011). Thiols, like cysteamine are capable of scavenging the superoxide radical in both its anionic ($\text{O}_2^{\cdot -}$) and protonated forms (HO_2^{\cdot}) (Cardey et al., 2007).



Ceroid (lipofuscin) is a final product of lipid oxidation that consists of insoluble polymerized lipid and protein complexes and is found within foam cells in atherosclerotic lesions (Mitchinson, 1982). Cysteamine inhibited lysosomal ceroid formation in human macrophage-like cells possibly by scavenging the hydroperoxyl radical in the lysosomes, whereas probucol had little effect (Fig. 5). Cysteamine was able to completely prevent lipid peroxidation for long periods of time, as measured in terms of conjugated dienes, cholesteryl linoleate hydroperoxide and 7-ketocholesterol formation and loss of cholesteryl linoleate and arachidonate, as well as oxidation of the tryptophan residues of apoB-100 of LDL (Figs. 6 and 7).

Based on our data, we conclude that the initial oxidation of LDL catalysed by iron at lysosomal pH is mediated by hydroperoxyl radicals and is not dependent on pre-existing lipid hydroperoxide levels.




Fig. 8. Oxidation of LDL by hydroperoxyl radicals at lysosomal pH. Iron in the lysosomes of macrophages leads to production of superoxide, which becomes protonated to the hydroperoxyl radical (HO_2^\bullet), which attacks lipids (LH) in the both phospholipid monolayer and hydrophobic core of LDL and tryptophan residues in apoB-100. Probuco in the monolayer is unable to reach the hydrophobic core and thus cannot inhibit oxidation of cholesteryl esters, but does inhibit oxidation of the phospholipid monolayer. DPPD is more hydrophobic than probuco and inhibits oxidation in both the monolayer and core. Cysteamine scavenges hydroperoxyl or superoxide radicals in the aqueous phase and is able to inhibit the oxidation of LDL in both the core and monolayer.

Furthermore the oxidation of LDL in the lysosomes of macrophages is prevented by the lysosomotropic drug cysteamine, which scavenges hydroperoxyl and superoxide radicals in the aqueous phase (Fig. 8).

Probuco was initially introduced as a cholesterol-lowering drug in early 1970's and later it was found to attenuate atherosclerotic lesion development in most animal models (Carew et al., 1987). However, the findings of The Probuco Quantitative Regression Swedish Trial (PQRST) in 1995 showed that the treatment of hypercholesterolaemic patients with probuco does not decrease the volume of femoral artery atherosclerotic lesions (Walldius et al., 1994). Two other clinical studies, The Aggressive Reduction of Inflammation Stop Events (ARISE) (Tardif et al., 2018) and Probuco Observational Study Illuminating Therapeutic Impact on Vascular Events (POSITIVE) (Yamashita et al., 2008), were conducted to assess different aspects of the treatment with probuco or its analog succinobuco, but both failed to demonstrate any significant benefits for the primary endpoints, although there were some beneficial effects. Our finding that probuco gives no protection against oxidation in the cholesteryl ester core of LDL at lysosomal pH offers an explanation about probuco's low efficacy in the clinical trials, if LDL oxidation takes place mainly in the lysosomes of macrophages. We have previously shown that enrichment of LDL with α -tocopherol has some pro-oxidant effects in the presence of iron at lysosomal pH (Satchell and Leake, 2012). There are other possibilities to explain the low efficacy of these antioxidants in the clinical trials, however, such as low penetration into atherosclerotic lesions, an inappropriate dose or that the treatment was started too late.

DPPD inhibited LDL oxidation very effectively in both the core and monolayer. It is of interest that it protects considerably against atherosclerosis in cholesterol-fed rabbits (Sparrow et al., 1992), but it cannot be used in humans because it is mutagenic. Water-soluble antioxidants that accumulate in lysosomes, such as cysteamine, that can scavenge hydroperoxyl radicals in the acidic aqueous phase of lysosomes have potential to treat atherosclerosis in humans.

In summary, the most effective antioxidants to test the oxidised LDL hypothesis of atherosclerosis might be lysosomotropic water-soluble antioxidants or highly hydrophobic antioxidants, which can protect both the core and monolayer of LDL from oxidation, rather than moderately hydrophobic antioxidants, such as probuco, which can only protect the monolayer but not the core of LDL from oxidation.

Conflict of interest

The authors declare no competing financial interests.

Sources of funding

This work was supported by Felix trust and British Heart Foundation (grant no PG/15/98/31864).

Author contributions

Both authors contributed equally to this work.

Acknowledgements

The authors are grateful to Dr Kim Jackson and Rada Mihaylova for taking blood from our volunteers.

References

- Bard, J.M., Urien, S., Fruchart, J.C., Tillement, J.P., 1994. Location of probuco in lipoproteins inferred from compositional analysis of lipoprotein particles. An in-vitro study. *J. Pharm. Pharmacol.* 46, 797–800.
- Bedwell, S., Dean, R.T., Jessup, W., 1989. The action of defined oxygen-centred free radicals on human low-density lipoprotein. *Biochem. J.* 262, 707–712.
- Brodin-Sartorius, A., Tete, M.J., Niaudet, P., Antignac, C., Guest, G., Ottolenghi, C., Charbit, M., Moysé, D., Legendre, C., Lesavre, P., Cochat, P., Servais, A., 2012. Cysteamine therapy delays the progression of nephropathic cystinosis in late adolescents and adults. *Kidney Int.* 81, 179–189.
- Cardey, B., Foley, S., Enescu, M., 2007. Mechanism of thiol oxidation by the superoxide radical. *J. Phys. Chem. A* 111, 13046–13052.
- Carew, T.E., Schwenke, D.C., Steinberg, D., 1987. Antiatherogenic effect of probuco unrelated to its hypocholesterolemic effect: evidence that antioxidants in vivo can selectively inhibit low density lipoprotein degradation in macrophage-rich fatty streaks and slow the progression of atherosclerosis in the Watanabe heritable hyperlipidemic rabbit. *Proc. Natl. Acad. Sci. U. S. A.* 84, 7725–7729.
- Collins, D.S., Unanue, E.R., Harding, C.V., 1991. Reduction of disulfide bonds within lysosomes is a key step in antigen processing. *J. Immunol.* 147, 4054–4059.
- Dabbagh, A.J., Frei, B., 1995. Human suction blister interstitial fluid prevents metal ion-dependent oxidation of low density lipoprotein by macrophages and in cell-free systems. *J. Clin. Invest.* 96, 1958–1966.
- De Grey, A.D., 2002. HO_2^\bullet : the forgotten radical. *DNA Cell Biol.* 21, 251–257.
- Dubinina, E.E., Gavrovskaya, S.V., Kuzmich, E.V., Leonova, N.V., Morozova, M.G., Kovrugina, S.V., Smirnova, T.A., 2002. Oxidative modification of proteins: oxidation of tryptophan and production of dihydroxytryptophan in purified proteins using fenton's system. *Biochemistry (Moscow)* 67, 343–350.
- el-Saadani, M., Esterbauer, H., el-Sayed, M., Goher, M., Nassar, A.Y., Jurgens, G., 1989. A spectrophotometric assay for lipid peroxides in serum lipoproteins using a commercially available reagent. *J. Lipid Res.* 30, 627–630.
- Esterbauer, H., Striegl, G., Puhl, H., Rotheneder, M., 1989. Continuous monitoring of in vitro oxidation of human low density lipoprotein. *Free Radic. Res. Commun.* 6, 67–75.
- Esterbauer, H., Dieber-Rotheneder, M., Waeg, G., Striegl, G., Jurgens, G., 1990. Biochemical, structural, and functional properties of oxidized low-density lipoprotein. *Chem. Res. Toxicol.* 3, 77–92.
- Esterbauer, H., Gebicki, J., Puhl, H., Jurgens, G., 1992. The role of lipid peroxidation and antioxidants in oxidative modification of LDL. *Free Radic. Biol. Med.* 13, 341–390.
- Frei, B., Gaziano, J.M., 1993. Content of antioxidants, preformed lipid hydroperoxides, and cholesterol as predictors of the susceptibility of human LDL to metal ion-

- dependent and – independent oxidation. *J. Lipid Res.* 34, 2135–2145.
- Güngör, N., Özyürek, M., Güçlü, K., Çekiç, S.D., Apak, R., 2011. Comparative evaluation of antioxidant capacities of thiol-based antioxidants measured by different in vitro methods. *Talanta* 83, 1650–1658.
- Giessauf, A., Steiner, E., Esterbauer, H., 1995. Early destruction of tryptophan residues of apolipoprotein B is a vitamin E-independent process during copper-mediated oxidation of LDL. *Biochim. Biophys. Acta* 1256, 221–232.
- Giessauf, A., van Wickern, B., Simat, T., Steinhart, H., Esterbauer, H., 1996. Formation of N-formylkynurenine suggests the involvement of apolipoprotein B-100 centered tryptophan radicals in the initiation of LDL lipid peroxidation. *FEBS Lett.* 389, 136–140.
- Halliwell, B., Gutteridge, J.M., 1986. Oxygen free radicals and iron in relation to biology and medicine: some problems and concepts. *Arch. Biochem. Biophys.* 246, 501–514.
- Jackson, R.L., Barnhart, R.L., Mao, S.J., 1991. Probucol and its mechanisms for reducing atherosclerosis. *Adv. Exp. Med. Biol.* 285, 367–372.
- Kritharides, L., Jessup, W., Gifford, J., Dean, R.T., 1993. A method for defining the stages of low-density lipoprotein oxidation by the separation of cholesterol- and cholesteryl ester-oxidation products using HPLC. *Anal. Biochem.* 213, 79–89.
- Kurz, T., Terman, A., Gustafsson, B., Brunk, U.T., 2008. Lysosomes in iron metabolism, ageing and apoptosis. *Histochem. Cell Biol.* 129, 389–406.
- Lee, T.S., Lee, F.Y., Pang, J.H., Chau, L.Y., 1999. Erythrophagocytosis and iron deposition in atherosclerotic lesions. *Chin. J. Physiol.* 42, 17–23.
- Linschitz, H., Ottolenghi, M., Bensasson, R., 1967. One-electron oxidation of triplet diphenyl-p-phenylenediamine by the diimine. *J. Am. Chem. Soc.* 89, 4592–4599.
- Lund-Katz, S., Phillips, M.C., 1986. Packing of cholesterol molecules in human low-density lipoprotein. *Biochemistry* 25, 1562–1568.
- Lynch, S.M., Frei, B., 1993. Mechanisms of copper- and iron-dependent oxidative modification of human low density lipoprotein. *J. Lipid Res.* 34, 1745–1753.
- Meguro, R., Asano, Y., Odagiri, S., Li, C., Iwatsuki, H., Shoumura, K., 2005. The presence of ferric and ferrous iron in the nonheme iron store of resident macrophages in different tissues and organs: histochemical demonstrations by the perfusion-Perls and –Turnbull methods in the rat. *Arch. Histol. Cytol.* 68, 171–183.
- Minotti, G., Aust, S.D., 1992. Redox cycling of iron and lipid peroxidation. *Lipids* 27, 219–226.
- Mitchinson, M.J., Hothersall, D.C., Brooks, P.N., De Burbure, C.Y., 1985. The distribution of ceroid in human atherosclerosis. *J. Pathol.* 145, 177–183.
- Mitchinson, M.J., 1982. Insoluble lipids in human atherosclerotic plaques. *Atherosclerosis* 45, 11–15.
- Morgan, B., Lahav, O., 2007. The effect of pH on the kinetics of spontaneous Fe(II) oxidation by O₂ in aqueous solution—basic principles and a simple heuristic description. *Chemosphere* 68, 2080–2084.
- Parthasarathy, S., Young, S.G., Witztum, J.L., Pittman, R.C., Steinberg, D., 1986. Probucol inhibits oxidative modification of low density lipoprotein. *J. Clin. Invest.* 77, 641–644.
- Patterson, R.A., Horsley, E.T., Leake, D.S., 2003. Prooxidant and antioxidant properties of human serum ultrafiltrates toward LDL: important role of uric acid. *J. Lipid Res.* 44, 512–521.
- Pieroni, L., Khalil, L., Charlotte, F., Poinard, T., Piton, A., Hainque, B., Imbert-Bismut, F., 2001. Comparison of bathophenanthroline sulfonate and ferene as chromogens in colorimetric measurement of low hepatic iron concentration. *Clin. Chem.* 47, 2059–2061.
- PubChem, National Center for Biotechnology Information, PubChem Compound Database.
- Ronsein, G.E., de Oliveira, M.C., de Medeiros, M.H., Di Mascio, P., 2011. Mechanism of dioxindolylalanine formation by singlet molecular oxygen-mediated oxidation of tryptophan residues. *Photochem. Photobiol. Sci.* 10, 1727–1730.
- Satchell, L., Leake, D.S., 2012. Oxidation of low-density lipoprotein by iron at lysosomal pH: implications for atherosclerosis. *Biochemistry* 51, 3767–3775.
- Sempos, C.T., 2002. Do body iron stores increase the risk of developing coronary heart disease? *Am. J. Clin. Nutr.* 76, 501–503.
- Sparrow, C.P., Doebber, T.W., Olszewski, J., Wu, M.S., Ventre, J., Stevens, K.A., Chao, Y.S., 1992. Low density lipoprotein is protected from oxidation and the progression of atherosclerosis is slowed in cholesterol-fed rabbits by the antioxidant N,N'-diphenylphenylenediamine. *J. Clin. Invest.* 89, 1885–1891.
- Steinberg, D., Witztum, J.L., 2002. Is the oxidative modification hypothesis relevant to human atherosclerosis? Do the antioxidant trials conducted to date refute the hypothesis? *Circulation* 105, 2107–2111.
- Steinberg, D., Witztum, J.L., 2010. Oxidized low-density lipoprotein and atherosclerosis. *Arterioscler. Thromb. Vasc. Biol.* 30, 2311–2316.
- Sullivan, J.L., 2009. Iron in arterial plaque: modifiable risk factor for atherosclerosis. *Biochim. Biophys. Acta* 1790, 718–723.
- Tachiev, G., Roth, J.A., Bowers, A.R., 2000. Kinetics of hydrogen peroxide decomposition with complexed and free iron catalysts. *Int. J. Chem. Kinet.* 32, 24–35.
- Takahashi, M., Tsuchiya, J., Niki, E., 1989. Oxidation of lipids. XVI. Inhibition of auto-oxidation of methyl linoleate by diarylamines. *Bull. Chem. Soc. Jpn.* 62, 1880–1884.
- Tardif, J.-C., McMurray, J.J.V., Klug, E., Small, R., Schumi, J., Choi, J., Cooper, J., Scott, R., Lewis, E.F., L'Allier, P.L., Pfeffer, M.A., 2018. Effects of succinobucol (AGI-1067) after an acute coronary syndrome: a randomised, double-blind, placebo-controlled trial. *Lancet* 371, 1761–1768.
- Terman, A., Kurz, T., 2013. Lysosomal iron, iron chelation, and cell death. *Antioxid. Redox Signal.* 18, 888–898.
- Walldius, G., Erikson, U., Olsson, A.G., Bergstrand, L., Hadell, K., Johansson, J., Kaijser, L., Lassvik, C., Molgaard, J., Nilsson, S., et al., 1994. The effect of probucol on femoral atherosclerosis: the probucol quantitative regression swedish trial (PQRST). *Am. J. Cardiol.* 74, 875–883.
- Walters, M.J., Wrenn, S.P., 2010. Size-selective uptake of colloidal low density lipoprotein aggregates by cultured white blood cells. *J. Colloid Interface Sci.* 350, 494–501.
- Wen, Y., Leake, D.S., 2007. Low density lipoprotein undergoes oxidation within lysosomes in cells. *Circ. Res.* 100, 1337–1343.
- Wen, Y., Satchell, L., Gibson, T.M., Weinberg, P.D., Leake, D.S., 2014. Low density lipoprotein aggregated by sphingomyelinase is internalised by macrophages and oxidised in lysosomes. *Atherosclerosis* 232, e5–e6.
- Wilkins, G.M., Leake, D.S., 1994. The effect of inhibitors of free radical generating-enzymes on low-density lipoprotein oxidation by macrophages. *Biochim. Biophys. Acta* 1211, 69–78.
- Yamashita, S., Hbujo, H., Arai, H., Harada-Shiba, M., Matsui, S., Fukushima, M., Saito, Y., Kita, T., Matsuzawa, Y., 2008. Long-term probucol treatment prevents secondary cardiovascular events: a cohort study of patients with heterozygous familial hypercholesterolemia in Japan. *J. Atheroscler. Thromb.* 15, 292–303.
- You, S.A., Archacki, S.R., Angheloiu, G., Moravec, C.S., Rao, S., Kinter, M., Topol, E.J., Wang, Q., 2003. Proteomic approach to coronary atherosclerosis shows ferritin light chain as a significant marker: evidence consistent with iron hypothesis in atherosclerosis. *Physiol. Genomics* 13, 25–30.
- Yuan, X.M., Anders, W.L., Olsson, A.G., Brunk, U.T., 1996. Iron in human atheroma and LDL oxidation by macrophages following erythrophagocytosis. *Atherosclerosis* 124, 61–73.