

SCIENTIFIC REPORTS



OPEN

Chemokine CXCL1 mediated neutrophil recruitment: Role of glycosaminoglycan interactions

Kirti V. Sawant¹, Krishna Mohan Poluri¹, Amit K. Dutta¹, Krishna Mohan Sepuru¹, Anna Troshkina¹, Roberto P. Garofalo^{2,3} & Krishna Rajarathnam^{1,2,4}

Received: 15 March 2016

Accepted: 22 August 2016

Published: 14 September 2016

The chemokine CXCL1/MGSA plays a pivotal role in the host immune response by recruiting and activating neutrophils for microbial killing at the tissue site. CXCL1 exists reversibly as monomers and dimers, and mediates its function by binding glycosaminoglycans (GAG) and CXCR2 receptor. We recently showed that both monomers and dimers are potent CXCR2 agonists, the dimer is the high-affinity GAG ligand, lysine and arginine residues located in two non-overlapping domains mediate GAG interactions, and there is extensive overlap between GAG and receptor-binding domains. To understand how these structural properties influence *in vivo* function, we characterized peritoneal neutrophil recruitment of a trapped monomer and trapped dimer and a panel of WT lysine/arginine to alanine mutants. Monomers and dimers were active, but WT was more active indicating synergistic interactions promote recruitment. Mutants from both domains showed reduced GAG heparin binding affinities and reduced neutrophil recruitment, providing compelling evidence that both GAG-binding domains mediate *in vivo* trafficking. Further, mutant of a residue that is involved in both GAG binding and receptor signaling showed the highest reduction in recruitment. We conclude that GAG interactions and receptor activity of CXCL1 monomers and dimers are fine-tuned to regulate neutrophil trafficking for successful resolution of tissue injury.

Circulating neutrophils are rapidly recruited in response to microbial infections and form the first line in host defense. Impaired recruitment and/or impaired activation could result in runaway infection leading to sepsis, whereas uncontrolled recruitment and/or sustained activation could result in collateral tissue damage and disease^{1–4}. Clinical studies and animal models have shown that the chemokine CXCL1 plays dual roles in the host immune response by recruiting and activating neutrophils to combat infection. First, it acts as a directional cue in routing peripheral neutrophils to the site of infection. Second, it activates the release of proteases and reactive oxygen species (ROS) for microbial killing in the tissue^{5–9}.

CXCL1 (also known as MGSA and Gro- α in humans and KC in mice) mediates function by signaling via CXCR2 on neutrophils and by binding to glycosaminoglycans (GAG) on endothelial and epithelial cells and the extracellular matrix (ECM). GAGs such as heparan sulfate are a family of linear polysaccharides covalently attached to core proteins called proteoglycans, and are present on both the luminal and abluminal sides of the endothelium and epithelium and as non-covalent complexes with proteins such as collagen and laminin in the ECM^{10,11}. In principle, GAG interactions can result in soluble chemotactic and tissue-bound haptotactic gradients, and the nature of these gradients can vary in the vasculature, across the endothelium, and across the epithelium^{12–14}. ECM GAGs can also be cleaved by proteases during the recruitment process, and soluble GAG-bound chemokine gradients have been shown to regulate neutrophil migration¹⁵. GAG interactions can also regulate recruitment by increasing chemokines' lifetime by preventing its proteolysis and from being washed away by blood flow in the vasculature¹⁶. All of these processes collectively control the flux, duration, and kinetics of circulating neutrophil egress to the tissue for successful resolution of inflammation.

CXCL1 levels are negligible under basal conditions but increase significantly during active infection. CXCL1 exists as monomers and dimers (dimer dissociation constant $\sim 10 \mu\text{M}$), and therefore one or both forms could

¹Department of Biochemistry and Molecular Biology, The University of Texas Medical Branch, Galveston, TX, USA.

²Department of Microbiology and Immunology, The University of Texas Medical Branch, Galveston, TX, USA.

³Department of Pediatrics, The University of Texas Medical Branch, Galveston, TX, USA. ⁴Sealy Center for Structural Biology and Molecular Biophysics, The University of Texas Medical Branch, Galveston, TX, USA. Correspondence and requests for materials should be addressed to K.R. (email: krrajara@utmb.edu)




Figure 1. Neutrophil recruitment profiles of CXCL1 wild type, monomer and dimer at different doses. BALB/c mice were injected intraperitoneally with different doses (0.1, 1 and 10 µg) of the CXCL1 variants. Peritoneal cells were harvested 6 h post-injection and total cells were counted along with differential counting to measure neutrophil levels. The results are expressed as means ± standard error from two independent experiments using 4 animals/group. The levels of neutrophils for both monomer and dimer are compared to WT at each dose and the differences between the variants were compared using ANOVA followed by Tukey's post hoc analysis. ** $p < 0.01$, *** $p < 0.001$.

mediate trafficking. We recently showed that both monomers and dimers are agonists for the CXCR2 receptor and that the dimer binds GAG with higher affinity^{17,18}. We also showed that basic residues located in two distinct domains on opposite faces of the protein orchestrate GAG interactions, and some of the GAG-binding residues are also located in the receptor-binding domain indicating that GAG-bound chemokine cannot activate the receptor¹⁸.

In this study, we characterized how CXCL1's structural properties influence *in vivo* function, by characterizing the peritoneal neutrophil recruitment of the WT (that can exist as monomers and dimers), a trapped non-associating monomer, a trapped non-dissociating dimer, and alanine (Ala) mutants of lysines and arginines from both GAG-binding domains on the WT background. On the basis of our current and previous studies on the related chemokine CXCL8, we propose that GAG interactions and receptor activity of the monomer and dimer collectively determine *in vivo* gradients in a chemokine-specific manner to direct neutrophils to the target site.

Results

Recruitment profiles of CXCL1 variants. We characterized the peritoneal neutrophil recruitment of WT CXCL1 and of the trapped monomer and dimer at different doses with PBS as the control (Fig. 1). The trapped monomer cannot dimerize and the trapped dimer cannot dissociate at any concentration or condition. Accordingly, the recruitment can be unambiguously attributed to the monomer or dimer. We have shown previously that the structures of the trapped variants are essentially similar to the native chemokine^{17,19}.

We observed that the recruitment profiles vary, and in particular, the monomer and dimer were minimally active at low doses, and the WT was more active than the monomer and dimer at all doses. CXCL1-mediated trafficking involves GAG interactions and CXCR2 activation. We have previously shown that both monomer and dimer are potent agonists for receptor activation and that the dimer is the high affinity GAG ligand^{17,20}. Therefore, we attribute the low recruitment activity of the dimer at lower doses to it predominantly existing in the GAG-bound form, and of the monomer to sub-optimal gradients due to its low GAG affinity. The higher activity of the WT, which is especially evident at the 1 µg data, must be due to its ability to exist as monomers and dimers, which allow optimal gradients for efficient recruitment to the tissue.

We also measured protein levels in the peritoneum as an indicator of tissue damage. Robust recruitment and/or premature neutrophil activation could damage the integrity of the epithelium resulting in the leakage of plasma proteins into the peritoneum^{21,22}. We observed that the protein levels of the WT and monomer and dimer variants at all doses were similar, and also no different from the PBS control (Fig. 2). In particular, that robust recruitment observed at the 10 µg dose, which is comparable to that seen in disease models, does not cause any tissue damage is quite striking. These observations suggest that the process of chemokine-mediated recruitment itself is highly regulated, and tissue damage observed during infection can be attributed to premature or uncontrolled neutrophil activation.

Design of CXCL1 GAG mutants. We recently showed that the GAG-binding residues are located in two non-overlapping domains and that the chemokine-GAG interface is extensive¹⁸ (Fig. 3). Residues Lys21 from the N-loop, Lys45 from the 40 s turn, and Lys60, Lys61, and Lys65 from the C-helix from both monomers of the dimer, constitute one domain that we define as the α-domain. Residues Arg8 from the N-terminus, Lys29 from the 1st β-strand, Lys48 from the 40 s turn, and Lys49 from the 3rd β-strand from both monomers of the dimer, constitute the other domain that we define as the β-domain. In principle, role of α- and β-domains and of the individual residues in each domain could be the same or different. For instance, one of the domains and/or specific residues from each domain could play a more prominent role and differentially regulate function. In order to




Figure 2. Protein levels in peritoneal fluid. Peritoneal fluid supernatants from mice were estimated using the BCA assay kit (Pierce). Each data set is representative of two experiments using 4 animals/group. Protein levels for each variant were compared to the PBS control using ANOVA followed by Tukey's post hoc analysis ($p < 0.05$). (ns - not significant).




Figure 3. GAG binding domains in CXCL1. (A) Ribbon representation of the CXCL1 monomer. α -domain and β -domain GAG-binding residues are highlighted in red and blue, respectively. (B) Surface representation of the CXCL1 dimer showing the GAG-binding residues of the α -domain and β -domain are in red and blue, respectively.

explicitly address the potential roles for GAG interactions, we individually mutated these basic residues to alanine (Ala) and characterized their GAG binding affinities and neutrophil recruitment activities.

GAG binding affinities of CXCL1 mutants. The binding affinities of the CXCL1 mutants were measured for GAG heparin using surface plasmon resonance (SPR). Heparin is commonly used for structural and biophysical studies for the reason it is commercially available, and has been shown to be useful for determining relative affinities and for identifying GAG-binding residues. The most common GAG on cell surfaces is heparan sulfate that has a modular structure of sulfated (NS) and acetylated (NA) domains whereas heparin is more uniformly sulfated. It has been shown that two NS domains separated by an NA domain in heparan sulfate binds chemokine CXCL8 dimer, and also that native chemokines dimerize on binding to cell surface GAGs^{23,24}.

Binding affinities (K_d) were calculated from surface plasmon resonance (SPR) measurements²⁵. Binding profiles for the WT and K29A mutant and a summary of the binding affinities are shown (Fig. 4). The binding affinities for the WT and dimer were similar $\sim 2 \pm 1 \mu\text{M}$ suggesting that the WT binds heparin as a dimer. The binding constants were calculated using the steady state model. Data could be fitted for all mutants with low χ^2 values (between 0.2 and 2.5), and the calculated binding constants show that the mutants bind heparin with 2 to 3 fold lower affinity (Fig. 4D). We also fitted the SPR data to Langmuir model, where the binding constant was calculated from the kinetic on and off constants. These data show that the mutants bind with 3 to 8-fold lower affinity, with χ^2 values between 2.7 and 8.8. The calculated and experimental fits for the WT and K29A mutant using the Langmuir model are shown in panels A and B, respectively.

The reported binding constants are apparent binding constants and have contributions from both monomer-dimer equilibrium and binding affinities of the monomer and dimer. Binding affinities of the monomer and dimer will vary, and we have shown previously using NMR spectroscopy that the dimer is the high-affinity ligand¹⁸. Mutations do not influence the monomer-dimer equilibrium and contribution from the monomer to overall binding will be less due to its lower binding affinity. Therefore, any differences in binding affinity essentially arise from reduced binding of the mutant dimer due to loss of H-bonding/electrostatic interaction mediated




Figure 4. Binding affinities of CXCL1 GAG mutants. SPR sensorgrams for (A) CXCL1 WT and (B) K29A mutant passed over immobilized heparin on a BIAcore SA chip. The curves represent the maximum response signal (RU) minus the reference signal (no GAG). Kinetic fits for the raw data are shown in red lines. (C) Steady state K_d analysis. Maximum response units were plotted for each concentration of CXCL1 WT and K29A mutant. (D) Relative binding affinities of the α -domain and β -domain GAG mutants compared to the WT.




Figure 5. Receptor activity of GAG mutants. Receptor activity of the CXCL1 mutants was characterized by measuring calcium release in CXCR2 transfected HL60 cells using FLIPR assay. 10 nM of CXCL1 mutants were added to the cells in quadruplicate and raw peak fluorescence was obtained as relative fluorescence units (RFU). Data shown are mean \pm SEM of the four measurements, and the graph is a representative of three independent experiments. The calcium release activity for each mutant was compared with the WT using ANOVA followed by Tukey's post hoc analysis. *** $p < 0.001$.

by that basic residue. The *in vivo* recruitment activity of the mutants is lower and correlates with the reduced heparin binding affinities.

Receptor activity of CXCL1 GAG mutants. CXCR2 receptor activation can be characterized by downstream functions like calcium release^{17,26}. We therefore characterized calcium release as a read-out for receptor function (Fig. 5). Our data show all the mutants, except R8A, are as active as the WT indicating that the altered recruitment is essentially due to altered GAG interactions. The R8A mutant showed no activity and was also highly impaired in the recruitment studies.




Figure 6. Neutrophil recruitment of CXCL1 GAG mutants in the peritoneum. Mice were injected intraperitoneally with 10 μ g of each CXCL1 mutant and cells were harvested at 6 h post-injection. Total cells were counted and a differential cell count was done. The results are expressed as means \pm standard error from two independent experiments using 4 animals/group. The neutrophil levels for the mutants are compared to the WT using ANOVA followed by Tukey's post hoc analysis. ** $p < 0.01$, *** $p < 0.001$.

Peritoneal recruitment of CXCL1 mutants. Peritoneal neutrophil recruitment of the GAG mutants was characterized at the 10 μ g dose because the recruitment levels seen at this dose are comparable to those seen in disease models^{2,8}. All mutants were less active providing unambiguous evidence that both α -domain and β -domain GAG interactions mediate *in vivo* recruitment (Fig. 6). Highly impaired recruitment of N-terminal Arg8 mutant must be due to its dual role of binding to GAG and activating the CXCR2 receptor. Previous studies have established that chemokine N-terminal ELR residues are crucial for receptor activation²⁷. β -domain N-terminal Arg8 is part of the highly conserved ELR triad, and the Arg8 mutant was also highly impaired in alveolar recruitment²⁸.

Discussion

Chemokines were first identified as critical regulators of immunity about thirty years ago, and it has now become increasingly clear that their function is intimately coupled to not only their receptor activation, but also to their dimerization/oligomerization and GAG binding properties^{18,25,29–33}. However, the molecular basis of how these properties orchestrate *in vivo* function is not well understood. CXCL1 belongs to a subset of seven chemokines that are characterized by the highly conserved N-terminal ELR triad and are agonists for the CXCR2 receptor. In this study, we show that both CXCL1 monomer and dimer are competent in mediating *in vivo* neutrophil recruitment, but the WT is more active. We attribute this to WT's ability to reversibly exist as monomers and dimers. Previous studies using zebra fish CXCL8 mutants that fail to bind proteoglycans *in situ* have shown GAG interactions not only mediate directional guidance but also restrict neutrophil motility suggesting that these distinct functions are most likely mediated via differential activities of the monomers and dimers³⁴. We also show that residues located at two distinct domains mediate *in vivo* GAG interactions, and that interactions from both domains are important as mutating even one of the GAG-binding residues impacts recruitment.

Comparison of the recruitment properties of CXCL1 and related chemokine CXCL8 show similarities and differences³². For both chemokines, WT is more active than the monomer or dimer, but the recruitment profiles vary for the WT and monomer but not for the dimer. For both chemokines, the dimer binds GAG with higher affinity but their GAG interactions are quite different. CXCL8 binds only residues of the α -domain, as well as some peripheral residues in the same face of the protein. Conversely, in addition to binding to the α -domain, CXCL1 binds another domain (defined as the β -domain) on the opposite face of the protein^{18,33}.

Migration of circulating neutrophils to the injury site is a multi-step process, and involves initial arrest in the vasculature, and subsequent migration across the endothelium, extracellular matrix, and epithelium. Chemokines mediate recruitment at every step of the way by forming gradients that involve GAG interactions^{12–14}. Structural studies have shown that the N-loop and N-terminal residues of CXCL1 in the GAG-bound form are not accessible for receptor binding, indicating that only soluble and not GAG-bound gradients mediate neutrophil trafficking¹⁸. Structural studies of CXCL8 in the GAG-bound form have also shown that the N-loop residues are occluded and not accessible for receptor binding suggesting soluble chemotactic gradients mediate trafficking³³. However gradient characteristics such as steepness at different locations along the migration route most likely vary between the two chemokines, which could explain the differences in their neutrophil trafficking profiles. We propose that crosstalk between GAG and receptor interactions of monomer and dimer dictate gradients at different locations, and that differences in gradients determine chemokine-specific differences in neutrophil trafficking.

Our previous studies have shown that the structural basis underlying GAG interactions of CXCL1 and CXCL8 are distinctly different. Comparison of the ELR chemokine sequences reveal similarities and sequence-specific differences in the distribution of the lysine and arginine residues (Fig. 7). For instance, a 'BB' motif with one residue from the 40 s turn and the other from the 3rd β -strand, are conserved in CXCL1, CXCL3, and CXCL7, whereas other members including CXCL8 have only one of the two basic residues. On the other hand, N-terminal arginine is absolutely conserved, which is involved in GAG interactions in CXCL1 but not in CXCL8. These observations suggest that presence or absence of a few basic residues in the context of structure could elicit strikingly different GAG interactions. Our studies on CXCL5 have already shown that its GAG interactions differ from those of CXCL1 and CXCL8. Chemokine receptor activation triggers G-protein and β -arrestin signaling pathways, and both pathways play a role in the migration process. However, related chemokines can have similar potencies

fixed, stained with Wright-Giemsa stain, and a differential leukocyte count was performed. Total leukocytes were counted using a hemocytometer after staining with Turk's solution.

Statistical analysis. All data are represented as mean \pm SEM. The differences between experimental groups were analyzed using one way analysis of variance (ANOVA) followed by Tukey's post hoc test. Analysis was done using Graph Pad Prism 5 (GraphPad Software). $P < 0.05$ was considered to be statistically significant.

References

- Adams, J. M. *et al.* Early trauma polymorphonuclear neutrophil responses to chemokines are associated with development of sepsis, pneumonia, and organ failure. *J. Trauma* **51**, 452–456, 456–457 (2001).
- Craciun, F. L., Schuller, E. R. & Remick, D. G. Early enhanced local neutrophil recruitment in peritonitis-induced sepsis improves bacterial clearance and survival. *J. Immunol.* **185**, 6930–6938 (2010).
- Pinheiro da Silva, F. & Soriano, F. G. Neutrophils recruitment during sepsis: Critical points and crossroads. *Front Biosci* **14**, 4464–4476 (2009).
- Kolaczowska, E. & Kubes, P. Neutrophil recruitment and function in health and inflammation. *Nat. Rev. Immunol.* **13**, 159–175 (2013).
- Cummings, C. J. *et al.* Expression and function of the chemokine receptors CXCR1 and CXCR2 in sepsis. *J. Immunol.* **162**, 2341–2346 (1999).
- Ritzman, A. M. *et al.* The chemokine receptor CXCR2 ligand KC (CXCL1) mediates neutrophil recruitment and is critical for development of experimental Lyme arthritis and carditis. *Infect. Immun.* **78**, 4593–4600 (2010).
- Jin, L., Batra, S., Douda, D. N., Palaniyar, N. & Jeyaseelan, S. CXCL1 contributes to host defense in polymicrobial sepsis via modulating T cell and neutrophil functions. *J. Immunol.* **193**, 3549–3558 (2014).
- De Filippo, K. *et al.* Mast cell and macrophage chemokines CXCL1/CXCL2 control the early stage of neutrophil recruitment during tissue inflammation. *Blood* **121**, 4930–4937 (2013).
- Shea-Donohue, T. *et al.* Mice deficient in the CXCR2 ligand, CXCL1 (KC/GRO- α), exhibit increased susceptibility to dextran sodium sulfate (DSS)-induced colitis. *Innate Immun* **14**, 117–124 (2008).
- Parish, C. R. The role of heparan sulphate in inflammation. *Nat. Rev. Immunol.* **6**, 633–643 (2006).
- Sarrazin, S., Lamanna, W. C. & Esko, J. D. Heparan sulfate proteoglycans. *Cold Spring Harb. Perspect. Biol.* **3** (2011).
- Frevert, C. W. *et al.* Binding of interleukin-8 to heparan sulfate and chondroitin sulfate in lung tissue. *Am. J. Respir. Cell Mol. Biol.* **28**, 464–472 (2003).
- Rot, A. Chemokine patterning by glycosaminoglycans and interceptors. *Front Biosci (Landmark Ed)* **15**, 645–660 (2010).
- Tanino, Y. *et al.* Kinetics of Chemokine-Glycosaminoglycan Interactions Control Neutrophil Migration into the Airspaces of the Lungs. *J. Immunol.* **184**, 2677–2685 (2010).
- Li, Q. L., Park, P. W., Wilson, C. L. & Parks, W. C. Matrilysin shedding of syndecan-1 regulates chemokine mobilization and transepithelial efflux of neutrophils in acute lung injury. *Cell* **111**, 635–646 (2002).
- Wolf, M., Albrecht, S. & Marki, C. Proteolytic processing of chemokines: implications in physiological and pathological conditions. *Int. J. Biochem. Cell Biol.* **40**, 1185–1198 (2008).
- Ravindran, A., Sawant, K. V., Sarmiento, J., Navarro, J. & Rajarathnam, K. Chemokine CXCL1 Dimer Is a Potent Agonist for the CXCR2 Receptor. *J. Biol. Chem.* **288**, 12244–12252 (2013).
- Sepuru, K. M. & Rajarathnam, K. CXCL1/MGSA Is a Novel Glycosaminoglycan (GAG)-binding Chemokine: Structural Evidence for Two Distinct Non-Overlapping Binding Domains. *J. Biol. Chem.* **291**, 4247–4255 (2016).
- Rajarathnam, K., Clark-Lewis, I. & Sykes, B. D. 1H NMR solution structure of an active monomeric interleukin-8. *Biochemistry* **34**, 12983–12990 (1995).
- Rajarathnam, K. *et al.* Neutrophil-activating peptide-2 and melanoma growth-stimulatory activity are functional as monomers for neutrophil activation. *J. Biol. Chem.* **272**, 1725–1729 (1997).
- Benjamin, C. F., Canetti, C., Cunha, F. Q., Kunkel, S. L. & Peters-Golden, M. Opposing and hierarchical roles of leukotrienes in local innate immune versus vascular responses in a model of sepsis. *J. Immunol.* **174**, 1616–1620 (2005).
- Holter, J. E., Weiland, J. E., Pacht, E. R., Gadek, J. E. & Davis, W. B. Protein permeability in the adult respiratory distress syndrome. Loss of size selectivity of the alveolar epithelium. *J. Clin. Invest.* **78**, 1513–1522 (1986).
- Hoogewerf, A. J. *et al.* Glycosaminoglycans mediate cell surface oligomerization of chemokines. *Biochemistry* **36**, 13570–13578 (1997).
- Spillmann, D., Witt, D. & Lindahl, U. Defining the interleukin-8-binding domain of heparan sulfate. *J. Biol. Chem.* **273**, 15487–15493 (1998).
- Dyer, D. P., Salanga, C. L., Volkman, B. F., Kawamura, T. & Handel, T. M. The dependence of chemokine-glycosaminoglycan interactions on chemokine oligomerization. *Glycobiology* **26**, 312–326 (2016).
- Nasser, M. W. *et al.* Differential activation and regulation of CXCR1 and CXCR2 by CXCL8 monomer and dimer. *J. Immunol.* **183**, 3425–3432 (2009).
- Rajagopalan, L. & Rajarathnam, K. Structural basis of chemokine receptor function—a model for binding affinity and ligand selectivity. *Biosci. Rep.* **26**, 325–339 (2006).
- Sawant, K. V. *et al.* Chemokine CXCL1-Mediated Neutrophil Trafficking in the Lung: Role of CXCR2 Activation. *J. Innate Immun.* **7**, 647–658 (2015).
- Das, S. T. *et al.* Monomeric and dimeric CXCL8 are both essential for *in vivo* neutrophil recruitment. *PLoS ONE* **5**, e11754 (2010).
- Proudfoot, A. E. *et al.* Glycosaminoglycan binding and oligomerization are essential for the *in vivo* activity of certain chemokines. *Proc. Natl. Acad. Sci. USA* **100**, 1885–1890 (2003).
- Kuschert, G. S. *et al.* Glycosaminoglycans interact selectively with chemokines and modulate receptor binding and cellular responses. *Biochemistry* **38**, 12959–12968 (1999).
- Gangavarapu, P. *et al.* The monomer-dimer equilibrium and glycosaminoglycan interactions of chemokine CXCL8 regulate tissue-specific neutrophil recruitment. *J. Leukoc. Biol.* **91**, 259–265 (2012).
- Joseph, P. R., Mosier, P. D., Desai, U. R. & Rajarathnam, K. Solution NMR characterization of chemokine CXCL8/IL-8 monomer and dimer binding to glycosaminoglycans: structural plasticity mediates differential binding interactions. *Biochem. J.* **472**, 121–133 (2015).
- Sarris, M. *et al.* Inflammatory chemokines direct and restrict leukocyte migration within live tissues as Glycan-Bound gradients. *Curr. Biol* **22**, 2375–2382 (2012).
- Stillie, R., Farooq, S. M., Gordon, J. R. & Stadnyk, A. W. The functional significance behind expressing two IL-8 receptor types on PMN. *J. Leukoc. Biol.* **86**, 529–543 (2009).

Acknowledgements

We thank Ms. Renling Xu for assistance with protein expression, Dr. Jiqing Sai of Vanderbilt University for the HL60 cells, and Dr. Heather Lander for editorial assistance. This work was supported by a grant from National Institutes of Health (P01 HL107152) to K.R.

Author Contributions

K.V.S., K.M.P., K.M.S., R.P.G. and K.R. designed the experiments. K.V.S., K.M.P., A.K.D., K.M.S. and A.T. performed the experiments and analyzed the data. K.V.S. and K.R. wrote the paper.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Sawant, K. V. *et al.* Chemokine CXCL1 mediated neutrophil recruitment: Role of glycosaminoglycan interactions. *Sci. Rep.* **6**, 33123; doi: 10.1038/srep33123 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016