

# Reactive Oxygen Species Production and Mitochondrial Dysfunction Contribute to Quercetin Induced Death in *Leishmania amazonensis*

Fernanda Fonseca-Silva, Job D. F. Inacio, Marilene M. Canto-Cavalheiro, Elmo Eduardo Almeida-Amaral\*

Laboratório de Bioquímica de Tripanosomatídeos, Instituto Oswaldo Cruz (IOC), Fundação Oswaldo Cruz (FIOCRUZ), Rio de Janeiro, Rio de Janeiro, Brazil

## Abstract

**Background:** Leishmaniasis, a parasitic disease caused by protozoa of the genus *Leishmania*, affects more than 12 million people worldwide. Quercetin has generated considerable interest as a pharmaceutical compound with a wide range of therapeutic activities. One such activity is exhibited against the bloodstream parasite *Trypanosoma brucei* and amastigotes of *Leishmania donovani*. However, the mechanism of protozoan action of quercetin has not been studied.

**Methodology/Principal Findings:** In the present study, we report here the mechanism for the antileishmanial activity of quercetin against *Leishmania amazonensis* promastigotes. Quercetin inhibited *L. amazonensis* promastigote growth in a dose- and time- dependent manner beginning at 48 hours of treatment and with maximum growth inhibition observed at 96 hours. The IC<sub>50</sub> for quercetin at 48 hours was 31.4 μM. Quercetin increased ROS generation in a dose-dependent manner after 48 hours of treatment. The antioxidant GSH and NAC each significantly reduced quercetin-induced cell death. In addition, quercetin caused mitochondrial dysfunction due to collapse of mitochondrial membrane potential.

**Conclusions/Significance:** The effects of several drugs that interfere directly with mitochondrial physiology in parasites such as *Leishmania* have been described. The unique mitochondrial features of *Leishmania* make this organelle an ideal drug target while minimizing toxicity. Quercetin has been described as a pro-oxidant, generating ROS which are responsible for cell death in some cancer cells. Mitochondrial membrane potential loss can be brought about by ROS added directly *in vitro* or induced by chemical agents. Taken together, our results demonstrate that quercetin eventually exerts its antileishmanial effect on *L. amazonensis* promastigotes due to the generation of ROS and disrupted parasite mitochondrial function.

**Citation:** Fonseca-Silva F, Inacio JDF, Canto-Cavalheiro MM, Almeida-Amaral EE (2011) Reactive Oxygen Species Production and Mitochondrial Dysfunction Contribute to Quercetin Induced Death in *Leishmania amazonensis*. PLoS ONE 6(2): e14666. doi:10.1371/journal.pone.0014666

**Editor:** Photini Sinnis, New York University, United States of America

**Received:** April 20, 2010; **Accepted:** January 13, 2011; **Published:** February 8, 2011

**Copyright:** © 2011 Fonseca-Silva et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was supported by grants from Fundacao Carlos Chagas Filho de Amparo À Pesquisa do Estado do Rio de Janeiro (FAPERJ) and Fundacao Oswaldo Cruz (FIOCRUZ). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: elmo@ioc.fiocruz.br

## Introduction

Recently, the effects of several drugs that interfere directly with mitochondrial physiology in parasites such as *Leishmania* have been described [1,2]. The unique mitochondrial features of *Leishmania* make this organelle an ideal drug target while minimizing toxicity. *Leishmania* has a single large mitochondrion which is distributed in branches under the subpellicular microtubules and a specialized region rich in DNA called the kinetoplast [3].

Leishmaniasis, a parasitic disease caused by protozoa of the genus *Leishmania*, affects more than 12 million people worldwide [4]. Treatment of leishmaniasis is based on pentavalent antimonials, drugs developed more than 50 years ago that are toxic and prone to drug resistance [5]. Several drug screens of natural compounds have been successful in discovering novel compounds for treating some parasitic diseases [6]. Extracts obtained from plants, as well as pure compounds including some kinds of flavonoids, have been reported to possess significant antiprotozoan activities [7–9].

Quercetin is the most common polyphenolic flavonoids present in plants such as onions, ginkgo biloba and tea and can be absorbed

by humans. Quercetin has a wide range of reported biological effects including antioxidant, anti-hypertension, anti-inflammatory, antimicrobial and antiprotozoan activities [10,11], although, the precise molecular mechanism of action of quercetin has not yet been demonstrated. Quercetin can induce the production of superoxide anion, hydrogen peroxide, and other reactive oxygen species (ROS) [12–15].

ROS are generated in cells infected by pathogens to combat infection. ROS can also be generated in response to some drugs, and the same principle works for certain antiprotozoan drugs in killing parasites in an infected cell. This property of a drug, to generate ROS to cause the destruction of cellular macromolecular components, is important because this action can be modulated to derive maximal effect.

The aim of the present work is to study the antileishmanial activity of quercetin and describe the mechanism of quercetin against promastigotes of *Leishmania amazonensis*. Our results demonstrate that the effect of the quercetin is associated with ROS production leading to mitochondrial dysfunction, ultimately causing parasite death.

## Materials and Methods

### 1. Reagents

H<sub>2</sub>DCFDA (2',7'-dichlorodihydrofluorescein diacetate) was obtained from Invitrogen Molecular Probes (Leiden, The Netherlands). Others reagents were purchased from Merck (São Paulo, Brazil) or Sigma-Aldrich (St Louis, MO). Deionized distilled water was obtained using a MilliQ system of resins (Millipore Corp., Bedford, MA) and was used in the preparation of all solutions.

### 2. Parasites

Promastigotes of *L. amazonensis* (MHOM/BR/LTB0016 strain) were grown at 26°C in Schneider's *Drosophila* medium pH 7.2 supplemented with 10% (v/v) heat-inactivated fetal calf serum. The number of parasites was determined by direct counting with a Neubauer chamber.

### 3. Cell proliferation

Promastigotes of *L. amazonensis* were harvested, washed twice and seed into fresh medium in the absence or in the presence of different concentration of quercetin (3 µM–96 µM) for 24 to 96 at 26°C. The cell density was estimated in a Neubauer chamber and the growth curve was initiated with 1.0×10<sup>6</sup> cells/ml. The cell proliferation was verified by the counting of the cell number in a Neubauer chamber.

### 4. Determination of mitochondrial membrane potential ( $\Delta\Psi_m$ )

**4.1. Flow cytometry studies.** Promastigotes of *L. amazonensis* (1×10<sup>6</sup> cells/ml) were treated for 48 hours with or without 24 µM or 96 µM quercetin and then incubated with 10 µg/ml rhodamine 123 for 20 minutes. Samples were kept on ice until analysis. Data acquisition and analysis were performed using a FACSCalibur flow cytometer (Becton Dickinson, Franklin Lakes, USA) equipped with the Cell Quest software (Joseph Trotter, Scripps Research Institute, La Jolla, USA). A total of 10,000 events were acquired in the region previously established as corresponding to the parasites. Alterations in the fluorescence for Rh123 were quantified using an index of variation (IV) obtained by the equation (MT–MC)/MC, where MT is the median of fluorescence for treated parasites and MC is the median of control parasites. Negative IV values correspond to depolarization of the mitochondrial membrane [16,17].

**4.2. JC-1.** The cationic JC-1 was used as a probe to determine the mitochondrial membrane potential ( $\Delta\Psi_m$ ) as described [9]. Promastigotes (1×10<sup>6</sup> cells/ml) were cultured for 48 hours in the absence or presence of 24 µM or 96 µM quercetin. Cells were harvested, re-suspended in Hank's Balanced Salt Solution (HBSS) and the cell number was counted in a Neubauer chamber. Promastigotes (2×10<sup>6</sup> cells/ml) were incubated with JC-1 (10 µg/ml) for 10 minutes at 37°C. After washing twice with HBSS, fluorescence was measured spectrofluorometrically at both 530 nm and 590 nm using an excitation wavelength of 480 nm. The ratio of values obtained at 590 nm and 530 nm was plotted as the relative  $\Delta\Psi_m$ .

### 5. Alamar Blue assay

Promastigotes of *L. amazonensis* (1×10<sup>6</sup> cells/ml) were treated for 48 hours with or without different concentration of quercetin (3 µM–96 µM). Cells were harvested, re-suspended in Hank's Balanced Salt Solution (HBSS) and the cell number was counted in a Neubauer chamber. Promastigotes (5×10<sup>6</sup> cells/ml) were incubated with Alamar Blue (10% v/v) for 6 hours at 26°C. The

absorbance was measured at 570 nm with a spectrophotometer. *L. amazonensis* cells lysed by addition of 0.1% Triton X-100 were used as positive control.

### 6. Measurement of reactive oxygen species (ROS) levels

Intracellular ROS levels were measured in treated and untreated cells. Promastigotes (1×10<sup>6</sup> cells/ml) were cultured for 48 hours in the absence or presence of 24 µM or 96 µM quercetin. Promastigotes were then harvested, re-suspended in Phosphate Buffered Saline (PBS) and the cell number was counted in a Neubauer chamber. Promastigotes (2×10<sup>6</sup> cells/ml) were incubated with H<sub>2</sub>DCFDA (20 µM) for 20 minutes at 37°C. Fluorescence was measured spectrofluorometrically at 530 nm using an excitation wavelength of 507 nm. For all measurements, basal fluorescence was subtracted. Positive control was obtained by addition of 20 units/ml glucose oxidase+60 mM glucose for 20 minutes.

### 7. Statistical analysis


All experiments were performed in triplicate. The data were analyzed statistically using Student's *t*-test and two-way analysis of variance (ANOVA). In all cases, the level of significance was set at  $p < 0.05$  or  $p < 0.01$  as indicated. The data are expressed as means ± standard errors.

## Results and Discussion


Quercetin, an abundant naturally occurring flavonoid, has generated considerable interest as a pharmaceutical compound with a wide range of therapeutic activities. One such activity is exhibited against the bloodstream parasite *Trypanosoma brucei* and amastigotes of *Leishmania donovani* [11,18]. To determine the effect of quercetin on the growth of *L. amazonensis*, we incubated the parasites with different concentrations of quercetin (3 µM–96 µM) for 24 to 96 hours. Incubation with quercetin inhibited *L. amazonensis* promastigote growth in a time- and dose-dependent manner (Fig. 1). This inhibitory effect was equal to 66% after 48 hours with 96 µM. The IC<sub>50</sub> for quercetin at 48 hours was 31.4 µM. Total growth arrest was observed after incubating *L. amazonensis* promastigotes with 96 µM quercetin at 96 hours. This result demonstrates the antileishmanial activity of quercetin against *L. amazonensis*.

Mitochondria are essential cellular organelles that play a central role in energy metabolism and are critical for the survival of any cell. During oxidative phosphorylation, electrons are moved thorough the mitochondrial respiratory chain, and a proton gradient is established across the inner mitochondrial membrane as the energy source for ATP production. The maintenance of mitochondrial membrane potential ( $\Delta\Psi_m$ ) is vital for this metabolic process as well as for cell survival [3,19]. Studies have shown that variations in  $\Delta\Psi_m$  induced by drugs are associated with cell survival in *Trypanosoma cruzi* [9,16], *Leishmania donovani* [19] and *L. amazonensis* [20].

To determine if quercetin induces an alteration in the  $\Delta\Psi_m$  of *L. amazonensis* promastigotes, we measured the  $\Delta\Psi_m$  by using the fluorescent probe rhodamine 123, which accumulates within energized mitochondria, in conjunction with flow cytometry analysis. Figure 2 shows that *L. amazonensis* treated with 24 µM or 96 µM quercetin for 48 hours decreased rhodamine 123 fluorescence and thus altered  $\Delta\Psi_m$ . The quantification of this decrease was evaluated by unit IV. *L. amazonensis* treated with 24 µM or 96 µM led to IVs of  $-0.20 \pm 0.018$  and  $-0.95 \pm 0.014$  respectively. Similarly, a decrease in rhodamine 123 fluorescence and IV values ( $-0.29 \pm 0.026$ ) was also observed following



**Figure 1. Growth of *L. amazonensis* cultivated in Schneider's *Drosophila* medium at 26°C in the absence or in the presence of Quercetin.** *L. amazonensis* was cultivated in Schneider's *Drosophila* medium at 26°C as described in Materials and Methods for 96 h, in the absence (●) or presence of 3 μM (○), 12 μM (▼), 24 μM (△) 72 μM (■) or 96 μM (□) quercetin. The number of parasites was determined by direct counting with a Neubauer chamber. Values shown are the mean ± standard error of three different experiments. In the control curve (absence of quercetin), the same volume of vehicle (DMSO 0.2%) was added to the growth medium. No differences were observed on lengths of flagella and cell body of promastigote in the presence of quercetin when compared with parasite growth in the absence of quercetin. doi:10.1371/journal.pone.0014666.g001



**Figure 2. Flow cytometry analysis of *Leishmania amazonensis* treated with quercetin.** *Leishmania amazonensis* was cultivated in Schneider's *Drosophila* medium at 26°C as for 48 h, in the absence or in the presence of 24 μM and 96 μM quercetin. Promastigotes were subsequently labeled with rhodamine 123 (10 μg/ml). Data acquisition and analysis were performed using a FACSCalibur flow cytometer (equipped with the Cell Quest software). A total of 10,000 events were acquired in the regions previously established as those corresponding to each form of *L. amazonensis*. Positive control was obtained by addition of FCCP (20 μM) for 20 minutes. In the control (absence of quercetin) the same volume of vehicle (DMSO 0.2%) was added to the growth medium. doi:10.1371/journal.pone.0014666.g002

treatment with a mitochondrial uncoupling agent, carbonyl cyanide *p*-trifluoromethoxyphenylhydrazone (FCCP) at 20  $\mu\text{M}$  [21,22].

To confirm that quercetin induces  $\Delta\Psi_m$  collapse, we used JC-1, which is a cationic mitochondrial vital dye. This dye is lipophilic and becomes concentrated in the mitochondria in proportion to the membrane potential such that more dye accumulates in mitochondria with greater  $\Delta\Psi_m$ . Spectrofluorometric data presented in Fig. 3 showed a marked decrease in relative fluorescence intensity ( $\Delta\Psi_m$  values) in a dose-dependent manner, indicating depolarization of the membrane potential in cells following treatment with 24  $\mu\text{M}$  or 96  $\mu\text{M}$  of quercetin, with  $\Delta\Psi_m$  reductions of 42.61% and 77.62%, respectively. Furthermore, cells incubated with 20  $\mu\text{M}$  FCCP had a similar decrease in the relative fluorescence intensity values (79.82% of reduction). Taken together, these data suggest that quercetin inhibit *L. amazonensis* growth by affecting parasite mitochondrial function.

The reduced fluorescence of rhodamine 123 and JC-1 was not due to measuring membrane potential in the remaining live cells, because the percentage of viable cells was similar in treated and untreated parasites (Fig. 4).


In mammalian cells, increased cellular ROS production has been suggested to be responsible for the depolarization of  $\Delta\Psi_m$  and subsequent cell death [23,24]. To investigate if the mitochondrial dysfunction observed in *L. amazonensis* promastigote treated with quercetin is promoted by ROS production, we measured ROS levels using the cell-permeable dye  $\text{H}_2\text{DCFDA}$  [25–27]. The levels of ROS in quercetin-treated cells were 2.4-fold and 4.4-fold higher for 24  $\mu\text{M}$  and 96  $\mu\text{M}$ , respectively, compared to the level of ROS in control cells throughout the experiment (Fig. 5). Since glucose oxidase catalyses the oxidation of D-glucose generating  $\text{H}_2\text{O}_2$ , it was employed as a positive control. Glucose/glucose oxidase led to an increase in ROS levels

compared to control (2.6-folds, compared to the level of ROS in the control).

Reduced glutathione (GSH) is an important molecule for protecting kinetoplasts from ROS or toxic compounds, and intracellular GSH levels can be increased by the antioxidant compound N-Acetyl-L-cysteine (NAC) [28,29]. In *L. amazonensis*, glutathione is a component of trypanothione, a major antioxidant of this parasite [30]. Thus, we tested if pre-incubation of *L. amazonensis* promastigotes with GSH or NAC (300  $\mu\text{M}$ ) could prevent the inhibitory effect of quercetin and found that these intracellular antioxidants protected *L. amazonensis* from inhibition by quercetin (Fig. 6A) and reduced the levels of ROS in quercetin-treated cells (Fig. 6B). Alternatively, oxidized glutathione (GSSG) did not reduce the levels of ROS and did not protect the growth inhibition promoted by quercetin. Taken together, these results indicate that inhibition of growth and disruption of  $\Delta\Psi_m$  promoted by quercetin in *L. amazonensis* is mediated by ROS production which might alter the cellular redox status.

Mitochondria are responsible for respiration and oxidative phosphorylation in eukaryotes, including trypanosomes, and they provide ATP through respiratory-coupled oxidative phosphorylation [31,32]. The mitochondria of *Leishmania* exhibit a unique structure and are functionally distinct from mammalian mitochondria, making this organelle an exceptional chemotherapeutic target. Here, we demonstrate for the first time that quercetin exerts its antileishmanial effect on *L. amazonensis* promastigotes by generating ROS and affecting parasite mitochondrial function.


To elucidate the mechanism of cell death induced by quercetin, we chose to measure the mitochondrial membrane potential because previous studies have shown that the single mitochondrion of the kinetoplastid parasite is a good indicator of cellular dysfunction [16,19,33]. *L. amazonensis* promastigotes were incubated with different concentrations of quercetin for 48 hours, the




**Figure 3. Effect of treatment of quercetin on the mitochondrial membrane potential of *Leishmania amazonensis* promastigote.**

*Leishmania amazonensis* was cultivated in Schneider's *Drosophila* medium at 26°C as for 48 h, in the absence or presence of 24  $\mu\text{M}$  or 96  $\mu\text{M}$  quercetin. Promastigotes were labeled with potentiometric probe JC-1 (10  $\mu\text{g}/\text{ml}$ ). Positive control was obtained by addition of FCCP (20  $\mu\text{M}$ ) for 20 minutes. In the control (absence of quercetin), the same volume of vehicle (DMSO 0.2%) was added to the growth medium. Dose-dependent changes in relative  $\Delta\Psi_m$  values are expressed as the ratio of the fluorescence measurement carried out at 590 nm (for J-aggregate) versus 530 nm (for J-monomer). Data are expressed as means  $\pm$  standard errors of three different experiments. \* indicates significant difference relative to the control group ( $p < 0.05$ ); \*\* indicates significant difference relative to the control group ( $p < 0.01$ ); # indicates significant difference relative to the 24  $\mu\text{M}$  quercetin treated group ( $p < 0.05$ ).


doi:10.1371/journal.pone.0014666.g003



**Figure 4. Comparison of the *L. amazonensis* untreated and treated with different concentration of quercetin.** *Leishmania amazonensis* was cultivated in Schneider's *Drosophila* medium at 26°C as for 48 h in the absence or presence of different concentration of quercetin (3 μM–96 μM). Cellular viability was measured using Alamar Blue assay as described under Material and Methods. Values shown are the mean±standard error of three different experiments. In the control (absence of quercetin), the same volume of vehicle (DMSO 0.2%) was added to the growth medium. Positive control (disrupted cells) was obtained by addition of 0.1% of Triton X-100 \* indicates significant difference relative to the control group ( $p < 0.05$ ). T—0.1% Triton X-100. doi:10.1371/journal.pone.0014666.g004



**Figure 5. Quercetin induced ROS formation.** *Leishmania amazonensis* was cultivated in Schneider's *Drosophila* medium at 26°C as for 48 h in the absence or presence of 24 μM or 96 μM quercetin. Generation of ROS was measured using fluorescent dye H<sub>2</sub>DCFDA as described under Materials and Methods. Data are expressed as fold increase in ROS production relative to control. Values shown are the mean±standard error of three different experiments. Positive control was obtained by addition of 20 units/ml glucose oxidase+60 mM glucose for 20 minutes. \* indicates significant difference relative to the control group ( $p < 0.05$ ); \*\* indicates significant difference relative to the control group ( $p < 0.01$ ); # indicates significant difference relative to the 24 μM quercetin treated group ( $p < 0.05$ ). G/GO—Glucose+Glucose oxidase. doi:10.1371/journal.pone.0014666.g005



**Figure 6. Effect of N-Acetyl-L-cysteine, Reduced Glutathione and Oxidized Glutathione on quercetin-induced cell death (A) and ROS formation (B).** *L. amazonensis* was cultivated in Schneider's *Drosophila* medium at 26°C as described in Materials and Methods for 48 h in the presence of N-Acetyl-L-cysteine, reduced glutathione and oxidized glutathione in the absence or in the presence of quercetin. N-Acetyl-L-cysteine, reduced glutathione and oxidized glutathione were solubilized in PBS and quercetin was solubilized in DMSO. N-Acetyl-L-cysteine, reduced glutathione and oxidized glutathione were added to the culture to a final concentration of 300  $\mu$ M, and quercetin was added to the culture to a final concentration of 96  $\mu$ M. Values shown are the mean  $\pm$  standard error of three different experiments. In the control (absence of quercetin), the same volume of vehicle (DMSO 0.2%) was added to the growth medium. Generation of ROS was measured using fluorescent dye H<sub>2</sub>DCFDA as described under Materials and Methods. Data are expressed as fold increase in ROS production relative to control. Values shown are the mean  $\pm$  standard error of three different experiments. Positive control for ROS generation was obtained by addition of 20 units/ml glucose oxidase+60 mM glucose for 20 minutes. CTRL - control; NAC - N-Acetyl-L-cysteine; GSH - Reduced Glutathione; GSSG - Oxidized Glutathione, Q—Quercetin and G/GO—Glucose+Glucose oxidase. \*\* indicates significant difference relative to the control group ( $p < 0.01$ ); \* indicates significant difference relative to the control group ( $p < 0.05$ ).

doi:10.1371/journal.pone.0014666.g006

shortest time that demonstrated inhibition of cellular growth (66% of inhibition). The concentrations of quercetin used in these experiments were based on the IC<sub>50</sub> for 48 hours (31.4  $\mu$ M). A decrease in rhodamine 123 fluorescence intensity and the JC-1 fluorescence ratio when *L. amazonensis* was treated with different concentrations of quercetin suggests increased proton permeability across the inner mitochondrial membrane with increasing quercetin. This could decrease ATP synthesis and result in parasite death. Collapse of  $\Delta\Psi_m$  was observed in *Leishmania* spp. [20,25–27] and *T. cruzi* [16,17,34] after treatment with different drugs.

It has been demonstrated that quercetin inhibited the growth of *Leishmania donovani* promastigotes, as well as, amastigote in infected macrophages *in vitro*, induced topoisomerase II mediated linearization of kDNA, causing cell cycle arrest, leading to apoptosis [35]. However, quercetin has been described as a pro-oxidant, generating ROS which are responsible for cell death in some cancer cells [12,36].  $\Delta\Psi_m$  loss can be brought about by ROS added directly *in vitro* or induced by chemical agents [37,38].

## References

- Rodrigues JC, de Souza W (2008) Ultrastructural alterations in organelles of parasitic protozoa induced by different classes of metabolic inhibitors. *Curr Pharm Des* 14: 925–938.
- Sen N, Majumder HK (2008) Mitochondrion of protozoan parasite emerges as potent therapeutic target: exciting drugs are on the horizon. *Curr Pharm Des* 14: 839–846.
- de Souza W, Attias M, Rodrigues JC (2009) Particularities of mitochondrial structure in parasitic protists (Apicomplexa and Kinetoplastida). *Int J Biochem Cell Biol* 41: 2069–2080.
- Desjeux P (2004) Leishmaniasis: current situation and new perspectives. *Comp Immunol Microbiol Infect Dis* 27: 305–318.
- Croft SL, Coombs GH (2003) Leishmaniasis—current chemotherapy and recent advances in the search for novel drugs. *Trends Parasitol* 19: 502–508.
- Kayser O, Kiderlen AF, Croft SL (2003) Natural products as antiparasitic drugs. *Parasitol Res* 90 Suppl 2: S55–62.
- Muzitano MF, Tinoco LW, Guette C, Kaiser CR, Rossi-Bergmann B, et al. (2006) The antileishmanial activity assessment of unusual flavonoids from *Kalanchoe pinnata*. *Phytochemistry* 67: 2071–2077.
- Guida MC, Esteva MI, Camino A, Flavia MM, Torres HN, et al. (2007) Trypanosoma cruzi: *in vitro* and *in vivo* antiproliferative effects of epigallocatechin gallate (EGCg). *Exp Parasitol* 117: 188–194.
- Mukherjee P, Majee SB, Ghosh S, Hazra B (2009) Apoptosis-like death in *Leishmania donovani* promastigotes induced by diospyrin and its ethanolamine derivative. *Int J Antimicrob Agents* 34: 596–601.
- Bischoff SC (2008) Quercetin: potentials in the prevention and therapy of disease. *Curr Opin Clin Nutr Metab Care* 11: 733–740.

11. Mamani-Matsuda M, Rambert J, Malvy D, Lejoly-Boisseau H, Daulouede S, et al. (2004) Quercetin induces apoptosis of *Trypanosoma brucei gambiense* and decreases the proinflammatory response of human macrophages. *Antimicrob Agents Chemother* 48: 924–929.
12. Lapidot T, Walker MD, Kanner J (2002) Antioxidant and prooxidant effects of phenolics on pancreatic beta-cells in vitro. *J Agric Food Chem* 50: 7220–7225.
13. De Marchi U, Biasutto L, Garbisa S, Toninello A, Zoratti M (2009) Quercetin can act either as an inhibitor or an inducer of the mitochondrial permeability transition pore: A demonstration of the ambivalent redox character of polyphenols. *Biochim Biophys Acta* 1787: 1425–1432.
14. Cao G, Sofic E, Prior RL (1997) Antioxidant and prooxidant behavior of flavonoids: structure-activity relationships. *Free Radic Biol Med* 22: 749–760.
15. Miura YH, Tomita I, Watanabe T, Hirayama T, Fukui S (1998) Active oxygen generation by flavonoids. *Biol Pharm Bull* 21: 93–96.
16. Menna-Barreto RF, Goncalves RL, Costa EM, Silva RS, Pinto AV, et al. (2009) The effects on *Trypanosoma cruzi* of novel synthetic naphthoquinones are mediated by mitochondrial dysfunction. *Free Radic Biol Med* 47: 644–653.
17. Menna-Barreto RF, Correa JR, Pinto AV, Soares MJ, de Castro SL (2007) Mitochondrial disruption and DNA fragmentation in *Trypanosoma cruzi* induced by naphthoimidazoles synthesized from beta-lapachone. *Parasitol Res* 101: 895–905.
18. Tasdemir D, Kaiser M, Brun R, Yardley V, Schmidt TJ, et al. (2006) Antitrypanosomal and antileishmanial activities of flavonoids and their analogues: in vitro, in vivo, structure-activity relationship, and quantitative structure-activity relationship studies. *Antimicrob Agents Chemother* 50: 1352–1364.
19. Mehta A, Shaha C (2006) Mechanism of metalloinduced death in *Leishmania* spp.: role of iron, reactive oxygen species, Ca<sup>2+</sup>, and glutathione. *Free Radic Biol Med* 40: 1857–1868.
20. Rodrigues JC, Bernardes CF, Visbal G, Urbina JA, Vercesi AE, et al. (2007) Sterol methenyl transferase inhibitors alter the ultrastructure and function of the *Leishmania amazonensis* mitochondrion leading to potent growth inhibition. *Protist* 158: 447–456.
21. Gunter TE, Pfeiffer DR (1990) Mechanisms by which mitochondria transport calcium. *Am J Physiol* 258: C755–786.
22. Boveris A, Chance B (1973) The mitochondrial generation of hydrogen peroxide. General properties and effect of hyperbaric oxygen. *Biochem J* 134: 707–716.
23. Banki K, Hutter E, Gonchoroff NJ, Perl A (1999) Elevation of mitochondrial transmembrane potential and reactive oxygen intermediate levels are early events and occur independently from activation of caspases in Fas signaling. *J Immunol* 162: 1466–1479.
24. Johnson TM, Yu ZX, Ferrans VJ, Lowenstein RA, Finkel T (1996) Reactive oxygen species are downstream mediators of p53-dependent apoptosis. *Proc Natl Acad Sci U S A* 93: 11848–11852.
25. Roy A, Ganguly A, Bosedasgupta S, Das BB, Pal C, et al. (2008) Mitochondria dependent ROS-mediated programmed cell death (PCD) induced by 3,3'-Diindolylmethane (DIM) through Inhibition of FoF1-ATP synthase in unicellular protozoan parasite *Leishmania donovani*. *Mol Pharmacol*.
26. Sen N, Banerjee B, Das BB, Ganguly A, Sen T, et al. (2007) Apoptosis is induced in leishmanial cells by a novel protein kinase inhibitor withaferin A and is facilitated by apoptotic topoisomerase I-DNA complex. *Cell Death Differ* 14: 358–367.
27. Sen N, Das BB, Ganguly A, Mukherjee T, Tripathi G, et al. (2004) Camptothecin induced mitochondrial dysfunction leading to programmed cell death in unicellular hemoflagellate *Leishmania donovani*. *Cell Death Differ* 11: 924–936.
28. Sudhandiran G, Shaha C (2003) Antimonial-induced increase in intracellular Ca<sup>2+</sup> through non-selective cation channels in the host and the parasite is responsible for apoptosis of intracellular *Leishmania donovani* amastigotes. *J Biol Chem* 278: 25120–25132.
29. Leite MS, Thomaz R, Oliveira JH, Oliveira PL, Meyer-Fernandes JR (2009) *Trypanosoma brucei brucei*: effects of ferrous iron and heme on ecto-nucleoside triphosphate diphosphohydrolase activity. *Exp Parasitol* 121: 137–143.
30. Castro-Pinto DB, Lima EL, Cunha AS, Genestra M, De Leo RM, et al. (2007) *Leishmania amazonensis* trypanothione reductase: evaluation of the effect of glutathione analogs on parasite growth, infectivity and enzyme activity. *J Enzyme Inhib Med Chem* 22: 71–75.
31. Zilberstein D, Dwyer DM (1988) Identification of a surface membrane proton-translocating ATPase in promastigotes of the parasitic protozoan *Leishmania donovani*. *Biochem J* 256: 13–21.
32. Affranchino JL, De Tarlovsky MN, Stoppani AO (1985) Respiratory control in mitochondria from *Trypanosoma cruzi*. *Mol Biochem Parasitol* 16: 289–298.
33. Luque-Ortega JR, Rivero-Lezcano OM, Croft SL, Rivas L (2001) In vivo monitoring of intracellular ATP levels in *Leishmania donovani* promastigotes as a rapid method to screen drugs targeting bioenergetic metabolism. *Antimicrob Agents Chemother* 45: 1121–1125.
34. Santa-Rita RM, Lira R, Barbosa HS, Urbina JA, de Castro SL (2005) Anti-proliferative synergy of lysophospholipid analogues and ketoconazole against *Trypanosoma cruzi* (Kinetoplastida: Trypanosomatidae): cellular and ultrastructural analysis. *J Antimicrob Chemother* 55: 780–784.
35. Mitra B, Saha A, Chowdhury AR, Pal C, Mandal S, et al. (2000) Luteolin, an abundant dietary component is a potent anti-leishmanial agent that acts by inducing topoisomerase II-mediated kinetoplast DNA cleavage leading to apoptosis. *Mol Med* 6: 527–541.
36. Halliwell B (2008) Are polyphenols antioxidants or pro-oxidants? What do we learn from cell culture and in vivo studies? *Arch Biochem Biophys* 476: 107–112.
37. Satoh T, Enokido Y, Aoshima H, Uchiyama Y, Hatanaka H (1997) Changes in mitochondrial membrane potential during oxidative stress-induced apoptosis in PC12 cells. *J Neurosci Res* 50: 413–420.
38. Dvorakova K, Waltmire CN, Payne CM, Tome ME, Briehl MM, et al. (2001) Induction of mitochondrial changes in myeloma cells by imexon. *Blood* 97: 3544–3551.
39. Mukherjee SB, Das M, Sudhandiran G, Shaha C (2002) Increase in cytosolic Ca<sup>2+</sup> levels through the activation of non-selective cation channels induced by oxidative stress causes mitochondrial depolarization leading to apoptosis-like death in *Leishmania donovani* promastigotes. *J Biol Chem* 277: 24717–24727.