

Cardiovascular RNA markers and artificial intelligence may improve COVID-19 outcome: a position paper from the EU-CardioRNA COST Action CA17129

Lina Badimon ¹, Emma L. Robinson ^{2,3}, Amela Jusic ⁴, Irina Carpusca ⁴, Leon J. deWindt⁵, Costanza Emanuelli ⁶, Péter Ferdinandy ^{7,8}, Wei Gu ⁹, Mariann Gyöngyösi¹⁰, Matthias Hackl ¹¹, Kanita Karaduzovic-Hadziabdic ¹², Mitja Lustrek ¹³, Fabio Martelli ¹⁴, Eric Nham ¹⁵, Ines Potočnjak¹⁶, Venkata Satagopam ⁹, Reinhard Schneider ⁹, Thomas Thum ^{17,18}, and Yvan Devaux ^{4,*}; on behalf of EU-CardioRNA COST Action CA17129

¹Cardiovascular Science Program-ICCC, IR-Hospital de la Santa Creu i Santa Pau, Ciber CV, Autonomous University of Barcelona, Barcelona, Spain; ²Department of Cardiology, School for Cardiovascular Diseases, Faculty of Health, Medicine and Life Sciences, Maastricht University, Maastricht, the Netherlands; ³Division of Cardiology, Department of Medicine, University of Colorado Anschutz Medical Campus, Aurora, CO, USA; ⁴Cardiovascular Research Unit, Department of Population Health, Luxembourg Institute of Health, 1A-B rue Edison, L-1445 Strassen, Luxembourg; ⁵Department of Molecular Genetics, Faculty of Science and Engineering, Faculty of Health, Medicine and Life Sciences, Maastricht University, Maastricht, the Netherlands; ⁶National Heart & Lung Institute, Faculty of Medicine, Imperial College London, London, UK; ⁷Cardiometabolic Research Group and MTA-SE System Pharmacology Research Group, Department of Pharmacology and Pharmacotherapy, Semmelweis University, Budapest, Hungary; ⁸Pharmahungary Group, Szeged, Hungary; ⁹Luxembourg Center for Systems Biomedicine, University of Luxembourg, Esch sur Alzette, Luxembourg; ¹⁰Department of Cardiology, Medical University of Vienna, Vienna, Austria; ¹¹TAmiRNA GmbH, Vienna, Austria; ¹²Faculty of Engineering and Natural Sciences, International University of Sarajevo, Sarajevo, Bosnia and Herzegovina; ¹³Department of Intelligent Systems, Jozef Stefan Institute, Ljubljana, Slovenia; ¹⁴Molecular Cardiology Laboratory, IRCCS Policlinico San Donato, San Donato Milanese, Milan 20097, Italy; ¹⁵University of Zagreb School of Medicine, Zagreb, Croatia; ¹⁶Institute for Clinical Medical Research and Education, University Hospital Centre Sisters of Charity, Zagreb, Croatia; ¹⁷Institute of Molecular and Translational Therapeutic Strategies (IMTTS), Fraunhofer Institute for Toxicology and Experimental Medicine, Hannover, Germany; and ¹⁸REBIRTH Center for Translational Regenerative Medicine, Hannover Medical School, Hannover, Germany

Received 16 September 2020; editorial decision 7 March 2021; accepted 8 April 2021; online publish-ahead-of-print 11 April 2021

Abstract

The coronavirus disease 2019 (COVID-19) pandemic has been as unprecedented as unexpected, affecting more than 105 million people worldwide as of 8 February 2020 and causing more than 2.3 million deaths according to the World Health Organization (WHO). Not only affecting the lungs but also provoking acute respiratory distress, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is able to infect multiple cell types including cardiac and vascular cells. Hence a significant proportion of infected patients develop cardiac events, such as arrhythmias and heart failure. Patients with cardiovascular comorbidities are at highest risk of cardiac death. To face the pandemic and limit its burden, health authorities have launched several fast-track calls for research projects aiming to develop rapid strategies to combat the disease, as well as longer-term projects to prepare for the future. Biomarkers have the possibility to aid in clinical decision-making and tailoring healthcare in order to improve patient quality of life. The biomarker potential of circulating RNAs has been recognized in several disease conditions, including cardiovascular disease. RNA biomarkers may be useful in the current COVID-19 situation. The discovery, validation, and marketing of novel biomarkers, including RNA biomarkers, require multi-centre studies by large and interdisciplinary collaborative networks, involving both the academia and the industry. Here, members of the EU-CardioRNA COST Action CA17129 summarize the current knowledge about the strain that COVID-19 places on

*Corresponding author. Tel: +352 26 970 300; fax: +352 26 970 396, E-mail: yvan.deviaux@lih.lu

© The Author(s) 2021. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

the cardiovascular system and discuss how RNA biomarkers can aid to limit this burden. They present the benefits and challenges of the discovery of novel RNA biomarkers, the need for networking efforts, and the added value of artificial intelligence to achieve reliable advances.

Keywords

Biomarkers • Artificial intelligence • RNAs • Genomics

1. SARS-CoV-2 in 2020

The effect of the coronavirus disease 2019 (COVID-19) pandemic on the cardiovascular system is alarming. More research focusing on the collateral damage associated with COVID-19 infection is needed. COVID-19 causes pneumonia with multi-organ disease. Infection can be asymptomatic or may cause a wide spectrum of symptoms, from mild upper respiratory tract infection to life-threatening sepsis with generalized endothelial damage, inflammation, and thrombosis. COVID-19 first emerged in December 2019 in Wuhan, China, and as of 8 February 2020 has affected people in more than 200 countries, with more than 105 million identified cases and with over 2.3 million confirmed deaths (WHO Coronavirus Disease Dashboard). It is clear that one of the causes for the significant differences in the severity of symptoms and mortality may derive from patient susceptibility to infection. Moreover, a significant proportion of COVID-19 survivors suffer cardiovascular damage. As such, there is a clinical need for novel biomarkers which would aid in the identification of patients at risk of suffering a severe form of the disease or that may identify those patients prone to develop collateral damage in the vascular, cardiac, and cerebrovascular systems that may jeopardize their future well-being. We need to investigate and innovate to detain the next pandemic wave of COVID-related cardiovascular disease.

To face the pandemic and limit its medical, social, and economic burden, health authorities have launched several fast-track calls for research projects aiming to develop rapid strategies to combat the disease, as well as longer-term projects to learn and draw lessons from the current pandemic and prepare for the future.¹

A myriad of potential biomarkers of COVID-19, for both diagnostic and prognostic purposes, have been highlighted in an extremely high number of published articles within the few months following the beginning of the pandemic. Although it is difficult to identify from all these reports the most relevant biomarkers with serious translational potential, artificial intelligence approaches could constitute a key component of such endeavours. Cardiovascular and blood RNA markers, coupled with artificial intelligence methods, represent a still poorly explored yet rich reservoir of novel biomarkers with some potential to aid in personalizing healthcare of COVID-19 patients. Recent single-cell RNA sequencing experiments support this assumption.²

2. Epidemiology of SARS-CoV-2 and cardiovascular disease

SARS-CoV-2 infection affects mostly the ageing population with pre-existing cardiovascular diseases, such as coronary artery diseases, heart failure or respiratory failure of any origin. Moreover, individuals with pre-existing risk factors for cardiovascular disease or with co-morbidities affecting the cardiovascular system are at high risk for worse clinical outcome during the infection.³ Frequent involvement of cardiovascular comorbidities is detected in patients with SARS-CoV-2 infection and up to 33% of hospitalized patients with a COVID-19-positive test have

cardiac injury⁴ evidenced by elevated cardiac troponin I and troponin T levels. These patients are prone to develop acute heart failure and have a high (up to 44.4% reported) burden of arrhythmias.^{5–8} Patients with SARS-CoV-2 infection and acute cardiac injury have a substantially higher rate of in-hospital mortality (up to 71.2%), as compared with the mortality of patients with SARS-CoV-2 infection and no evidence of cardiac injury.^{3,7} Patients with pre-existing heart failure and SARS-CoV-2 infection have a two-fold higher risk of 30-day mortality as compared to patients without pre-existing heart failure and SARS-CoV-2 infection, independently of the category of heart failure (reduced, mid-range, or preserved ejection fraction).⁹ Multi-organ failure due to hypoxia caused by respiratory failure, acute kidney injury, electrolyte disturbances, systemic inflammation, and cytokine storm contribute to the cardiac injury in patients with SARS-CoV-2. The cytokine storm seems to contribute to a large extent to cardiac and vascular events. However, there are reports asking for a more concise definition of the cytokine storm and its real impact in the pathogenesis of the infection.^{10–12} Altered coagulation may lead to thrombotic complications including microthrombosis, microvascular damage, and generalized thromboembolic disorder. Recent empirical drugs against COVID-19, such as chloroquine, antiviral or anti-rheumatic drugs, monoclonal antibodies, or antibiotics may also aggravate cardiovascular symptoms by prolonging QT interval leading to arrhythmias, or resulting in drug-induced cardiomyopathies or cardiotoxicity.⁴ Since SARS-CoV-2 has a strong affinity for the angiotensin-converting enzyme 2 (ACE2) cell receptor, it was plausible to assume that antihypertensive treatment with ACE inhibitors or angiotensin receptor blockers (ARBs) might aggravate the disease. To date, however, no clinical evidence can confirm this assumption; thus, ACE inhibitor and ARB treatments continue to be administered to SARS-CoV-2-positive patients.^{13,14} The lockdown regulations and subsequent closure of outpatient clinics have led to major re-organizational efforts of the management of patients with cardiovascular disease. A paradoxical decrease of documented acute myocardial infarction has also been observed, which could be attributed to the lack of preventive control of patients with chest pain, and the self-quarantining of the patients fearing from the risk of nosocomial infection.¹⁵

3. Pathophysiology of SARS-CoV-2 infection phases and effects on the heart

SARS-CoV-2, a member of the family of coronaviruses, is an enveloped, positive-sense, single-stranded RNA virus that is able to infect various host species.¹⁶ Amongst the viral-encoded proteins, the SARS-CoV-2 spike (S) transmembrane glycoprotein protrudes from the viral surface and is essential for target cell binding and infection. ACE2 has been identified as the SARS-CoV-2 receptor,^{17–20} and ACE2 is highly expressed in the lung, heart, ileum, kidney, and bladder.²¹ The majority of adaptive immune cells that invade the infected lung tissue consist of T cells, since a




Figure 1 Disease mechanisms of SARS-CoV-2 infection.

proportional decrease in circulating T cells has been observed in COVID-19 patients. IL-8 and IL-6, recognized chemo-attractants for T cells and neutrophils, are produced by SARS-CoV-2-compromised lung epithelial cells (Figure 1A).²² As neutrophils function in adaptive immunity but can also provoke further damage to the lung, these cells are regarded as double-edged swords in the context of COVID-19.²³ Circulating monocytes are attracted from the circulation by granulocyte macrophage colony-stimulating factor that is produced by local T cells in infected tissue. In addition, elevated CD14⁺CD16⁺ inflammatory monocytes producing high levels of IL-6 are found in COVID-19 patients, suggesting that also monocytes actively contribute to the systemic inflammatory response. Finally, thrombosis and pulmonary embolism are commonly observed in severely ill COVID-19 patients (Figure 1B), likely indicating the presence of significant endothelial injury and microvascular permeability, which may further exacerbate viral invasion.

The symptoms of COVID-19 patients are heterogeneous, ranging from minimal symptoms to significant hypoxia with acute respiratory

distress, shock, coagulation dysfunction, and multi-organ involvement, including acute kidney injury, encephalopathy, myocardial injury, and heart failure. Indeed, epidemiological, clinical, and biological evidence shows a clear cardiac involvement in COVID-19 patients, due to direct myocardial infection and injury and/or to indirect mechanisms, linked to the underlying pathophysiology of the disease.²⁴

In keeping with a direct effect on heart function of SARS-CoV-2 (Figure 1C), its receptor ACE2 is expressed by cardiomyocytes, fibroblasts, endothelial cells, pericytes, macrophages, and the epicardial fat.²¹ Moreover, ACE2 levels are increased in failing hearts, and its high expression in arterial vascular cells of fibrotic lungs may facilitate the bloodstream spreading of SARS-CoV-2.²⁵ Cardiomyocytes derived from human-induced pluripotent stem cells can be infected efficiently by SARS-CoV-2.^{26,27} The SARS-CoV-2 genome has been identified in endomyocardial biopsies of patients with suspected myocarditis.²⁸ However, whilst cardiomyocyte damage was present, no viral particles were detected in cardiomyocytes and endothelium, suggesting that the

particles were due to infected macrophage migration. Thus, direct myocardial infection may not be the main mechanism of myocardial damage explaining the frequently observed troponin increases. The release of inflammatory cytokines (Figure 1C), a hallmark of severe COVID-19, can also lead to a form of myocarditis resembling Takotsubo syndrome.²⁹ Moreover, the pro-thrombotic state of COVID-19 patients, associated to D-dimers increase, may lead to microvascular dysfunction, coronary thrombosis or embolism (Figure 1B).³⁰ Along with the pro-coagulant profile of patients with COVID-19,³¹ other forms of stress may facilitate cardiomyopathy occurrence, such as hypoxemia caused by respiratory dysfunction, endothelial dysfunction leading to small arterial obliteration,²⁸ and the increased metabolic demands (Figure 1D).

4. Treatments: what is available, what is needed

4.1 Remdesivir

Remdesivir is the first medicinal product for human use for the treatment of COVID-19 which was granted a conditional marketing authorization of the European Parliament and of the Council.³² It is a nucleotide analogue with a broad-spectrum antiviral activity. The European Medicines Agency, specifically the Committee for Medicinal Products for Human Use, has granted a conditional marketing authorization to Veklury (*remdesivir*) for the treatment of COVID-19 in adults and adolescents with pneumonia who require supplemental oxygen (O₂).³³ The recommendation of *remdesivir* is mainly based on the results of the Adaptive COVID-19 Treatment Trial (ACTT)-1 sponsored by the US National Institute of Allergy and Infectious Diseases, and supporting data from other studies on *remdesivir*.^{33,34} According to the ACTT-1 study, patients in the *remdesivir* group had a shorter time to recovery than patients in the placebo group (median 10 vs. 15 days).³⁵ Kaplan–Meier estimates of mortality at Day 29 were 11.4% in the *remdesivir* group and 15.2% in the placebo group (hazard ratio 0.73; 95% CI 0.52–1.03).³⁵ The Food and Drug Administration (FDA) issued an emergency use authorization.³⁶ The use of *remdesivir* has shown shortening of recovery time in severe patients with O₂ saturation ≤ 94%, and cases requiring supplemental O₂, mechanical ventilation, or extracorporeal membrane oxygenation.^{37,38} It is recommended to start the treatment on Day 1 with 200-mg infusion, followed by 100-mg infusion daily for at least 4 days and maximum 9 days.³³ According to the WHO SOLIDARITY trial (results in preprint), death rate ratios for *remdesivir* are RR = 0.95 (95% CI 0.81–1.11, P = 0.50).³⁹ Comparative results from other studies are shown in Table 1. Overall, *remdesivir*, while improving time to recovery in patients with mild symptoms in ACTT1 trial, fails to improve mortality.

4.2 Dexamethasone

According to the RECOVERY trial results, in the *dexamethasone* group, the incidence of death was lower than in the usual care group amongst patients receiving invasive mechanical ventilation (29.3% vs. 41.4%) and amongst those receiving O₂ without invasive mechanical ventilation (23.3% vs. 26.2%) but not amongst those who were not receiving respiratory support at randomization (17.8% vs. 14.0%).⁴⁰ Based on these results, 6 mg of *dexamethasone* is recommended once daily for up to 10 days in COVID-19 patients on mechanical ventilation or who require supplemental O₂ but who are not on mechanical ventilation.^{38,41}

4.3 Chloroquine or hydroxychloroquine, lopinavir–ritonavir

Although *chloroquine* or *hydroxychloroquine* was one of the medications which appeared to show great potential at the beginning of COVID-19 pandemic, their use has been stopped due to lack of efficacy. Numerous companies donated these medications for treating COVID-19 patients; however, the FDA revoked the emergency use authorization for this drug. Furthermore, the combined use of *hydroxychloroquine* and *azithromycin* is not recommended because of the potential adverse reactions. *Lopinavir/ritonavir* also did not demonstrate benefit in patients with COVID-19. As reported in Table 1, the interim WHO SOLIDARITY trial results indicate that *remdesivir*, *hydroxychloroquine*, *lopinavir*, and *interferon* treatments had little or no effect on hospitalized COVID-19 patients, as indicated by overall mortality, initiation of ventilation, and duration of hospital stay.³⁹

4.4 Immunomodulatory medications

Several medications used in modulating the immune response, such as interleukin-1 (*anakinra*) or interleukin-6 (*sarilumab*, *siltuximab*, *tocilizumab*) inhibitors, are being used off-label and are being investigated. These medications have been proposed to suppress the cytokine storm.⁴²

4.5 Convalescent plasma

The convalescent plasma containing antibodies against SARS-CoV-2 virus collected from recovered COVID-19 patients is also being widely investigated. A randomized clinical trial with convalescent plasma therapy did not show any statistically significant improvement in clinical status or death rate.⁴³ However, this trial provided valuable information on the potential benefits of convalescent plasma, which may be useful in combination with antiviral drugs. According to some preliminary research, early administration of high-dose intravenous immunoglobulin therapy may improve the prognosis of critically ill patients.⁴⁴ On 23 August 2020, FDA issued an emergency use authorization for convalescent plasma for the treatment of COVID-19 in hospitalized patients.⁴⁵

5. Markers of disease evolution: what is available, what is needed

As the world faces the COVID-19 pandemic, markers enabling to predict the development of severe symptoms after SARS-CoV-2 infection are highly needed. Presence of cardiovascular risk factors (particularly arterial hypertension, diabetes mellitus, and aging) and previous cardiovascular diseases reportedly expose to an unfavourable progression of COVID-19.⁴⁶ As such, they can already provide an initial and rudimentary model to risk stratify patients.

Mortality rate after COVID-19 is associated with elevation in the ‘classic’ cardiac damage biomarkers, such as troponin T (TnT) and/or BNP/NT-proBNP.^{3,47} In line with that, COVID-19 patients who do not have significantly increased TnT levels show a lower mortality compared to patients without cardiovascular disease.^{5,48} This suggests that TnT and BNP/NT-proBNP concentration should be closely followed in patients with COVID-19 both for diagnostic (cardiac involvement) and for prognostic purposes. Elevations of D-Dimers have also been associated with poor outcome.⁴⁹ The addition of other biomarkers such as the inflammatory cytokine IL6 and lymphocyte count will be also helpful to determine the individual risk of a patient.

Table 1 Comparison of 28-day mortality of patients with SARS-CoV-2 treated with remdesivir, dexamethasone, hydroxychloroquine, lopinavir, and interferon with/without O₂ from the SOLIDARITY,³⁹ ACTT-1,³⁵ and RECOVERY⁴⁰ trials

Drug	28-day mortality	No O ₂	Low/hi-O ₂	Ventilation
Remdesivir* (N = 2743)	301/2743 (12.5%)	11/661 (2.0%)	192/1828 (12.2%)	98/254 (43.0%)
Control (N = 2708)	303/2708 (12.7%)	13/664 (2.1%)	219/1811 (13.8%)	71/233 (37.8%)
Remdesivir** (N = 541)	59/541 (10.9%)	3/75 (4.1%)	28/327 (8.6%)	28/131 (21.9%)
Placebo (N = 521)	77/521 (14.8%)	3/63 (4.8%)	45/301 (15.0%)	29/154 (19.3%)
Dexamethasone*** (N = 2104)	482/2104 (22.9%)	89/501 (17.8%)	298/1279 (23.3%)	95/324 (29.3%)
Usual care (N = 4321)	1110/4321 (25.7%)	145/1034 (14.0%)	682/2604 (26.2%)	283/683 (41.4%)
Hydroxychloroquine**** Control	104/947 (10.2%) 84/906 (8.9%)	69/862 (7.4%) 57/824 (6.6%)		35/85 (39.2%) 27/82 (32.3%)
Lopinavir***** Control	148/1399 (9.7%) 146/1372 (10.3%)	113/1287 (8.1%) 111/1258 (8.7%)		35/112 (28.1%) 35/114 (28.7%)
Interferon-β1a **** Control	243/2050 (12.9%) 216/2050 (11.0%)	188/1911 (10.9%) 176/1920 (9.5%)		55/139 (42.4%) 40/130 (33.8%)

Remdesivir*—SOLIDARITY trial. Day 0: 200 mg; Day 1–9: 100 mg i.v.;

Remdesivir**—ACTT. Day 1: 200 mg; Day 2–10: 100 mg compared to placebo;

Dexamethasone***—RECOVERY 6 mg oral/i.v. for up to 10 days;

Hydroxychloroquine****—SOLIDARITY trial. Hydroxychloroquine sulphate a 200 mg tbl at Hour 0, four tablets; Hour 6, four tablets; Hour 12, begin two tablets twice daily for 10 days;

Lopinavir*****—SOLIDARITY trial. Lopinavir a 200 mg+ ritonavir 50 mg 2x 2 tablets for 14 days;

Interferon****—SOLIDARITY trial. Three doses over six days of 44 μg subcutaneous Interferon-β1a.

Omics-based approaches recently discovered interesting metabolites in plasma of patients with COVID-19. Using both targeted and untargeted tandem mass spectrometry to profile the plasma lipidome and metabolome of COVID-19 patients with various degrees of severity and healthy controls, a panel of 10 plasma metabolites was found to distinguish COVID-19 patients from healthy controls with an area under the receiver-operating characteristic curve (AUC) of 0.975.⁵⁰

Biomarkers that might be useful in indicating progression from mild-to-severe multi-organ complication in COVID-19 patients are summarized in *Tables 2 and 3*, which have been inspired in part by two important review articles and meta-analyses.^{51,52} A myriad of recent publications have reported associations between classical and emerging biomarkers and COVID-19 prognosis. Yet, only meta-analyses enrolling more than 200 patients are included in *Tables 2 and 3*. Amongst inflammatory and cardiac injury markers, decreased number of white blood cells, lymphopenia, and thrombocytopenia and increased CRP, D-dimers, procalcitonin (PCT), lactate dehydrogenase (LDH), aspartate aminotransferase (AST), alanine aminotransferase (ALT), IL-6, cardiac troponin, and CK-MB are associated with poor outcomes of COVID-19 patients, indicating their potential to aid in risk stratification and prediction of severe and fatal outcomes (*Tables 2 and 3*).

The role of the cardiovascular expression/activity of the putative SARS-CoV-2 receptor ACE2 as well as of the use of renin-angiotensin-aldosterone system (RAAS) inhibitors in SARS-CoV-2 susceptibility and COVID-19 disease severity has been a matter of debate.^{68–71} However, the clear recommendation is to continue the administering of RAAS inhibitors or blockers in SARS-CoV-2-positive patients with underlying

cardiovascular disease. Elevated angiotensin II levels have been found to correlate with lung injury and viral load, suggesting that administration of angiotensin 1-7 and angiotensin 1-9 may help in restoration of normal functioning of renin-angiotensin system by antagonizing the effect of abnormally increased angiotensin II.⁷²

Circulating RNAs represent a rich source of biomarkers with clinical utility due to their biological relevance, dynamic regulation in response to onset and progression of disease, tissue-specificity, and accessibility for non-invasive analysis using biofluids (liquid biopsies). Especially for diseases with diverse symptoms and complications such as COVID-19, RNA biomarkers could provide important decision support. RNAs have shown some potential as cardiovascular disease biomarkers and may help in predicting unexpected cardiovascular events in COVID-19 patients. Although some clinical trials on miRNAs in COVID-19 have been started or are even completed (nine trials registered in clinicaltrials.gov database as of November 2020), none of them have been specifically designed to identify (mi)RNA predictors of cardiovascular outcome of COVID-19 patients. *Table 4* gathers the currently available studies reporting regulations of non-coding RNAs (ncRNAs) in patients infected with SARS-CoV-2. Predicted messenger RNA targets as well as their proposed role in COVID-19 are also included in this table.

Given the disproportionate impact of COVID-19 in ethnic minorities, it is essential to clarify if biomarkers are of use in such populations and if so how they could be ad-hoc adapted. Not only cardiac but also endothelial biomarkers deserve attention.⁷³ Gender-medicine considerations for COVID-19 cardiovascular risk stratification are also of paramount importance. Women appear to be better protected, as men display

Table 2 Laboratory markers associated with poor outcomes after SARS-CoV-2 infection

Sample size	WBC	Lymphocytes	Platelets	D-dimer	CRP	PCT	IL-6	AST	ALT	Ref.
	↑ ↓ OR [95% CI]	↑ ↓ OR [95% CI]	↑ ↓ OR [95% CI]	↑ ↓ OR [95% CI]	↑ ↓ OR [51% CI]	↑ ↓ OR [95% CI]	↑ ↓ OR [95% CI]	↑ ↓ OR [95% CI]	↑ ↓ OR [95% CI]	
3962	-	-	-	-	↑ -	↑ -	↑ -	-	-	52
10 491	-	↓ 3.33 [2.51-4.41]	↓ 2.36 [1.64-3.40]	↑ 3.39 [2.66-4.33]	↑ 4.37 [3.37-5.68]	↑ 6.33 [4.24-9.45]	-	↑ 2.75 [2.30-3.29]	↑ 1.7 [1.32-2.20]	51
1955	↓ -	↓ -	↓ -	↑ -	↑ -	↑ -	-	↑ -	↑ -	54
4662	↓ -	↓ 4.5 [3.3-6.0]	↓ -	↑ -	↑ 3.00 [2.1-4.4]	↑ -	↑ 53.1% [36.0/70.0%]	↑ -	↑ -	55
-	↓ 0.93 [0.46-1.86]	↓ 1.66 [1.26-2.20]	↓ 0.88 [0.26-2.95]	↑ 1.50 [0.89-2.56]	↑ 1.41 [1.17-1.70]	↑ 2.94 [2.09-4.15]	-	↑ 2.27 [1.76-2.94]	↑ 1.60 [1.34-1.90]	56
6320	↓ 1.75 [1.21-2.54]	↓ 0.30 [0.19-0.47]	↓ 0.56 [0.42-0.74]	↑ 3.97 [2.62-6.02]	↑ 6.36 [3.22-12.5]	↑ 4.76 [2.48-9.14]	↑ 2.10 [1.02-4.32]	-	-	57
-	↓ -	↓ -	↓ -	↑ -	↑ -	↑ -	↑ -	↑ -	↑ -	58
91 621	↓ -	↓ -	↓ -	↑ -	↑ -	↑ -	↑ -	↑ -	↑ -	59
3027	↓ 0.30 [0.17-0.51]	-	-	↑ 43.24 [9.92-188.49]	-	↑ 43.24 [9.92-188.49]	-	↑ 4.00 [2.46-6.52]	-	60
51 225	↓ 2.75 [2.02-3.9]	↓ -0.6 (-2.55-1.38)	↓ -36.06 (-49.24; -22.77)	↑ 3.22 [2.84-3.61]	↑ 68.31 [53.11-83.50]	↑ 0.52 [0.42-0.62]	↑ 43.64 [30.92-56.35]	↑ 17.41 [13.99-20.83]	↑ 2.18 [0.09-4.28]	61
4631	-	-	-	-	-	-	↑ RR 0.54 [0.27-0.81]	-	-	62
5626	-	-	-	↑ 1.4 [-2.04-(-0.77)]	↑ 64.03 [-68.88-(-59.19)]	-	-	-	-	63

The hyphen means not studied. Poor outcomes include in-hospital admission, intensive care unit admission, oxygen saturation <90%, severe disease, utilization of invasive mechanical ventilation, and mortality. Adapted from two references.^{51,52}

ALT, alanine aminotransferase; AST, aspartate aminotransferase; CRP, C-reactive protein; IL-6, interleukin 6; MA, meta-analysis; OR, odds ratio; PCT, procalcitonin; WBC, white blood cells; ↑, increased; ↓, decreased.

Table 3 Cardiac injury biomarkers associated with poor outcomes in COVID-19 patients

Sample size	LDH		CK		Creatinine		Troponin I		CK-MB		Ref.
	↑↓	OR [95% CI] ^c	↑↓	OR [95% CI] ^c	↑↓	OR [95% CI] ^c	↑↓	SMD [95%CI]	↑↓	SMD [95%CI]	
–	↑	6.7 [2.4–18.9]	–	–	–	–	↑	0.71 [0.42; 1.00]	↑	0.68 [0.48; 0.87]	56
6320	↑	2.03 [1.42–2.90]	↑	–	↑	–	–	–	–	–	51
491	–	–	–	–	–	–	↑	–	–	–	58
91 621	↑	–	↑	–	↑	–	↑	16% [11–22]	–	–	59
3027	↑	–	↑	–	↑	–	↑	43.24 [9.92–188.49]	–	–	60
51 225	↑	8.86 [2.72–28.89]	–	–	↑	5.30 [2.19–12.83]	↑	0.02 [0.02; 0.02]	–	–	61
4631	↑	180.26 [131.02–229.51]	–	–	↑	21.72 [16.72–26.71]	↑	0.74 [0.19–1.30]	–	–	62
5626	↑	RR 2.20 [1.55–31.12]	↑	RR 1.89 [1.50–2.61]	–	–	↑	-1.55 [-2.23; -0.88]	↑	-4.75 [13.31; 3.82]	63
341	–	–	–	–	–	–	↑	25.6 [6.8–44.5]	–	–	64
3118	–	–	–	–	–	–	↑	21.15 [10.19–43.94]	–	–	65
4189	–	–	–	–	–	–	↑	0.53 [0.30–0.75]	↑	0.62 [0.28–0.97]	66
982	–	–	–	–	–	–	↑	HR 2.48 [1.50–4.11]	–	–	67

Poor outcomes include in-hospital admission, intensive care unit admission, oxygen saturation <90%, severe disease, utilization of invasive mechanical ventilation, and mortality. The hyphen means not studied. Adapted from two references.^{51,52}

CK-MB, creatinine kinase-MB; HR, hazard ratio; LDH, lactate dehydrogenase; OR, odds ratio; RR, risk ratio; SMD, standardized mean difference; ↑, increased; ↓, decreased.

higher mortality rates (ranging from 60 to 75%).⁷⁸ Should this be due to a protective effect of oestrogens, perimenopausal, and postmenopausal women without hormonal replacement therapy could be considered at higher risk of cardiovascular death following COVID-19. Preclinical evidence suggests that sex may influence the expression of the ACE2 receptor.⁷⁸ Hence, the examination of sex differences should be an integral part of COVID-19-directed research projects. This is especially crucial as sex-specific RNA biomarkers may help in tailoring future healthcare.

6. Networking and coordination efforts for multinational, multi-centre studies on cardiovascular RNA markers

For a global pandemic of this kind, worldwide efforts are needed to understand the infectious agent, to develop diagnostic tools, treatments, and also to monitor the well-being of those infected with SARS-CoV-2 in the following years. For robust development of biomarkers or treatments, their effectivity must be validated in numerous cohorts, internationally in different demographics and on a large scale.

Addressing the increasing challenges posed by communicable diseases thus calls for multidisciplinary and multi-centre international cooperation to link available data, tools, and expertise, which will otherwise only be suboptimally exploited at regional or national levels. A truly integrated approach coordinating and facilitating the access to and sharing of biological resources, data, advanced technological facilities, and expertise, within a common research roadmap, is needed to exploit the full potential of the various resources. As COVID-19 incidence and clinical outcomes have been shown to be greatly influenced by many biological and environmental factors, the need to integrate data across the various settings worldwide is critical to increase the precision of analyses and to deliver meaningful results.

Through the EU-CardioRNA COST Action,⁷⁹ in April 2020, a call was placed to assemble a taskforce of clinicians and translational scientists working with COVID-19 patients to join forces in an international effort. This was communicated internally within the Action network as well as externally on the Action website and professional (social) media (<https://cardioma.eu/news/cost-actions-unite-efforts-in-the-fight-against-covid-19/>).⁸⁰ In total, 38 institutions and 22 countries responded to the call. Members of the taskforce have access to COVID-19 patient clinical data, blood samples, and other biospecimen and/or expertise in analysis of biomarkers in liquid biopsies.

With the clear elevated risk of COVID-19 in aged individuals and patients with cardiovascular disease, the task force aims to monitor (re-) hospitalization rates, mortality rates from cardiovascular disease in those with exposure to SARS-CoV-2, as well as to identify RNA biomarkers reflecting cardiovascular health.⁸¹ We encourage medical professionals in the taskforce to perform functional follow-up of COVID-19 infection including echocardiography or cardiovascular magnetic resonance imaging, where possible. Using the numerous and extensive patient databases across multiple centres involved, epidemiological analyses and clinical statistics will be performed to identify differential risk of COVID-19 infection and response as well as cardiovascular effects according to comorbidities and patient characteristics. This is especially important to identify individuals most at risk in the event of long-term presence of the SARS-CoV-2 in the population, as well as to further understand the mechanism of infection and morbidity.

Regulatory RNAs are emerging as stable reliable circulating molecular indicators of cardiovascular health.⁸² With expertise in regulatory RNA biomarkers detectable in peripheral blood samples, the EU-CardioRNA taskforce will analyse these RNAs and overlay with clinical information in search of biomarkers for cardiovascular outcomes of COVID-19 infection. Such a study and the translation of the results into clinical application would not be feasible without the integration of complementary expertise and resources from the various actors (cardiology and infectious disease, biomarkers, RNA, cardiovascular research, clinical studies, biobanking, artificial intelligence, data management, biostatistics, bioinformatics, technology transfer, etc.).

Table 4 Potential ncRNA biomarkers of COVID-19


ncRNA	Sample size	Type of sample	Regulation	Number of predicted target genes	Experimentally validated target genes in any disease	Experimentally validated target genes in COVID-19 patient samples	Proposed role in COVID-19	Reference
miR-16-2-3p	14	Blood	↑	71	FGFR2, PDPK1	—	—	73
miR-6501-5p				88	—	—	—	—
miR-618				11	MTDH, TLR-4, ATP6V1E1, HAT1, MCTS1, TGF- β 2	TLR-4, HAT1, TGF- β 2	TLR4 regulates inflammation; HAT1: mitochondrial function, cellular senescence, and telomere attrition; TGF- β 2 induces expression of furin in HBE cells	—
miR-183-5p			↓	220	PTEN, PIK3CA	PTEN	Regulator of SARS-CoV-2 ACE2-TMPRSS2-Furin-DPP4 axis	—
miR-627-5p				25	CDK6, SOX-2, LINC00958, lnc-UCA1	—	—	—
miR-144-3p				80	PTEN, APP, FoxO1	FOXO1 PTEN	FOXO1 regulates cell death downstream of several signalling pathways including CDK1, PKB/AKT1, and STK4/MST1	—
lncRNA DANCR	563	Lung tissue and blood	↓	—	miR-496/mTOR axis; miR-335-5p/miR-1972 and ROCK1 axis	mTOR	PTEN signalling is increased after SARS-CoV-2 infection	74
lncRNA NEAT1			↑	—	miR-129-5p/CLK7 axis;	RUNX3, SPI1	Regulator of Akt/mTOR/HIF-1 signalling pathway	—
miR-21-5p			↑	139	TGFBI, MAPK1	DANCR, NEAT1	RUNX3 regulates DANCR and is related to inflammatory reaction in the lung; SPI1 controls DANCR expression in the brain and in epithelial cells	—
miR-22-3p				162	WRNIP1, HMGB1	HMGB1	DANCR and NEAT1 can block inflammation via interacting with other ncRNAs, sponging miRNAs, or affecting TFs (e.g. STAT3)	—
miR-335-5p				63	Rb1, CARE, SGK3	—	Exogenous HMGB1 induces the expression of SARS-CoV-2 entry receptor ACE2	—
miR-19a-3p				241	UBAP2L, PSG10P, IL1RAP	—	—	—
miR-1207-5p	18	Lung tissue	↑	147	CSF1	CSF1	Enhances macrophage recruitment and activation and its over	75

Continued

Table 4 Continued

ncRNA	Sample size	Type of sample	Regulation	Number of predicted target genes	Experimentally validated target genes in any disease	Experimentally validated target genes in COVID-19 patient samples	Proposed role in COVID-19	Reference
miR-21-5p	Discovery: 33 Validation: 65	Serum	↑	41	RASGRP1, BCL2, SMARCA4, SPRY2, DUSP10, TIMP3, SOX5, MTAP, RECK, PIAS3, TGFB2, PTEN, E2F1, LRRFIP1, TPM1, NFIB, APAF1, BTG2, PDCD4, RHOB, ANP32A, SERPINB5, BMPR2, DAXX, TP63, MSH2, MSH6, ISCU, EIF4A2, ANKRD46, CDK2AP1, PPARA, FASLG, SMAD7, SERPINI1, DDAH1, HPGD, MYD88, IRAK1, VHL, GDF5, IL12A, CASC2, DNIM1L	TIMP3 PTEN	expression may contribute to acute inflammation SARS-CoV-2 reduces TIMP3 mRNA expression in alveolar epithelial cells, that likely promotes greater ADAM17 activity in COVID-19 patients. PTEN signalling is increased after SARS-CoV-2 infection	76
miR-155-5p			↑	70	MEIS1, TAB2, MECP2, SOCS1, MLH1, INPP5D, SMAD5, HIVEP2, ZNF652, BACH1, APC, SMAD1, SDCBP, MYO10, CLDN1, CEBPB, RHOA, AGTR1, RNF123, TP53INP1, IKBKE, KDM3A, SPI1, FOXO3, RUNX2, JUN, ETS1, CYR61, SMAD2, MYB, SKI, CKAP5, SOX6, CSF1R, FADD, NOS3, MYLK, PSIP1, ANXA2, HBP1, NFKB1, E2F2, PIK3R1, MMP16, MYC, SEL1L, DOCK1, RAD51, MXI1	TAB2 SOCS1 TP53INP1 FADD	TAB2 is associated with vascular inflammation SOCS1 is a key checkpoint regulator of the immune system TP53INP1 induced cell death by an autophagy- and caspase-dependent mechanism The FADD/caspase-8 axis regulates TNF- α and IFN- γ co-treatment-induced inflammatory cell death independent of intrinsic apoptosis in macrophages	
miR-208a-3p			↑	3	CDKN1A, MED13, ETS1	–	Down-regulated upon SARS-CoV-2 infection, associated with cellular signalling	
miR-499-5p			↑	43	FOXO4, PDCD4, ETS1	FOXO4		

Predicted miRNA-target interactions were performed using miRWalk 3.0, miRDB 6.0, and miRTarBase 8.0 databases. Experimentally validated target genes in any disease (mostly cancer) were obtained from miRTarBase 8.0. Experimentally validated target genes in COVID 19 and their proposed roles were obtained through literature search. The authors apologize for the many references that could not be added to this table due to space restrictions.
 †: up-regulated; ↓: down-regulated. CSF1, colony stimulating factor 1; DANCR, anti-differentiation lncRNA; FADD, Fas associated with death domain; FOXO4, forkhead box O4; FOXO1, forkhead box O1; HAT1, Histone acetyltransferase 1; HMGB1, high-mobility group protein 1; lncRNA, long non-coding RNA; mTOR, mechanistic target of rapamycin kinase; ncRNAs, non-coding RNAs; NEAT1, nuclear paraspeckle assembly transcript 1; PTEN, phosphatase and tensin homolog; RUNX3, RUNX family transcription factor 3; SOCS1, suppressor of cytokine signalling 1; SPI1, Spi-1 proto-oncogene; TAB2, TGF-beta activated kinase 1 (MAP3K7) binding protein 2; TGF- β 2, transforming growth factor beta 2; TIMP3, TIMP metalloproteinase inhibitor 3; TLR-4, toll-like receptor 4; TP53INP1, tumour protein p53 inducible nuclear protein 1.



7. Technical challenges and requirements in the RNA-study

The quantitative analysis of RNAs in biological samples faces several technical challenges that must be overcome in order to generate robust and reproducible results. Specifically, the analysis of circulating RNAs is complicated by a variety of pre-analytical settings that impact the analysis as well as the analytical challenge to deal with very low RNA concentrations.

To date, whole blood, serum, and plasma are the most widely explored liquid matrices for circulating RNA analysis. Analysis of whole blood can be biased by red blood cells and platelets, which are a rich source of small RNAs despite being anucleate.⁸³ Thus, protocols for specific depletion of certain types of RNAs have been developed for whole blood that improve sensitivity for other types of RNAs.⁸⁴ Serum and plasma as the liquid components of blood can behave quite differently due to the release of RNAs during platelet activation and blood coagulation after which serum is collected.⁸⁵ Therefore, results for RNA biomarker analysis are oftentimes not comparable between serum and plasma.⁸⁶ In addition, contamination of serum or plasma with cellular RNA derived from red blood cells due to haemolysis,^{87,88} or platelets due to variable pre-analytical processing,⁸⁹ can confound the analysis and lead to false-positive or false-negative results.⁹⁰

Currently, only few studies have attempted to address sources of bias for other types of liquid biopsies. For example, in case of urine, it is known that donor-dependent differences in volume based on hydration

status result in highly variable RNA concentrations that require normalization prior to analysis using for example urinary creatinine levels.⁹¹

In biofluids, RNAs are associated with two main types of RNA carriers, which facilitate transport and protect their RNA cargo from degradation: protein complexes and extracellular vesicles (EVs). At least in terms of small RNAs, it is known that the majority of extracellular RNAs in plasma or conditioned media is associated with protein complexes.^{92,93} This means that total RNA isolation and analysis from these matrices mainly reflects the protein-associated RNA fraction, and that the separate analysis of RNAs that are selectively released via EVs can reveal different results.⁹⁴ It is important to note that RNA analysis in EVs is anything but trivial and requires careful optimization of EV isolation and characterization and reporting according to the MISEV standard developed by the International Society of Extracellular Vesicles.⁹⁵

The analysis of RNA integrity and abundance obtained by RNA isolation is hampered by low concentrations. Thus, either highly sensitive methods using RNA specific dyes should be used and internal process controls such as spike-in oligonucleotides (spike-ins) can be useful to monitor RNA recovery and analytical variability and to normalize RNA expression data in biofluids in the absence of robust endogenous RNA references.

Analytical methods for circulating RNA quantification must also be highly sensitive to cope with low concentrations. Reverse-transcription quantitative PCR (RT-qPCR) is a gold-standard technology for this purpose. However, low throughput and high cost for using RT-qPCR in genome-wide RNA biomarker discovery have restricted its use to targeted




Figure 3 Biomarker identification and COVID-19 risk prediction workflow.

analyses for biomarker validation. This limitation resulted in the uptake of next-generation sequencing (NGS) for untargeted RNA biomarker discovery. Since early on it was observed that the abundance and stability of small RNAs in biofluids was surprisingly high, small RNA sequencing was rapidly adopted for biomarker identification in liquid biopsies.⁹⁶

The challenges for using small RNA NGS for circulating RNA analysis are as follows: (i) the extended PCR pre-amplification that is needed to obtain sufficient input material but is potentially resulting in PCR duplicates, (ii) adapter-ligation bias leading to over- and under-representation of certain RNAs in the library, and (iii) the relative quantification that restricts the main use to cross-sectional comparisons between selected groups. To overcome these challenges, unique molecular indices can be included in the adapter sequences to identify and remove PCR duplicates prior to data analysis.⁹⁷ Secondly, sophisticated adapter design such as randomized ends or single ligation protocols has been shown to reduce the ligation bias and reduce adapter dimers.^{98,99} Finally, the addition of spike-in calibrators with randomized ends and optimized concentration ranges can be used to normalize small RNA NGS data and achieve absolute quantification that is less sensitive towards changes in the (small) RNA composition of a sample.¹⁰⁰

Recently, also the application of total RNA sequencing for RNA biomarkers discovery in liquid biopsies has advanced to explore the full spectrum of RNAs. A stranded total RNA sequencing kit appeared to be sufficiently robust, accurate, and precise to quantify thousands of genes in platelet-rich and platelet-free plasma, urine, and conditioned medium as well as EVs isolated from these matrices.¹⁰¹ EVs from platelet-free plasma showed a large percentage (>80%) of short reads that were too short to be aligned. This was not observed for total RNA from platelet-free plasma and platelet-rich plasma, and total RNA as well as EV-RNA from urine and conditioned medium. This might suggest that RNA released from cells via EVs into the blood stream might be fragmented endogenously. In terms of gene biotypes, protein-coding genes made up the majority (>70%) of reads for all matrices except platelet-rich plasma, followed by pseudogenes, long noncoding RNAs (lncRNAs), and miscellaneous RNAs.¹⁰¹

Overall, the planning of ideal RNA biomarker study should in the first step consciously decide which biological matrix and RNA carrier are most relevant and practical, secondly, implement standardized protocols for sample collection and sample quality control at the study sites, and thirdly, take advantage of a well-characterized, fit-for-purpose validated, NGS protocol for genome-wide total RNA and small RNA quantification in low RNA input samples.

8. Data handling and integration

Data infrastructure that curates, integrates, and analyses clinical and experimental data from several COVID-19 cohorts is pivotal to make harmonized data available to research network members in order to unravel cardiovascular RNA markers of SARS-CoV-2 infection. Systematic collection and application of standards play an important role in managing and handling cohort data and its metadata efficiently. They facilitate findable, accessible, interoperable and reusable (FAIR) use of the data, which provides a solid foundation for systematically discovering, retrieving, understanding, integrating, disseminating, exchanging, reusing the data, and reproducing research results and outcome.

8.1 Making data findable, including provisions for metadata

In order to make the data discoverable, the following rules should be ensured:

- Data sets need to be assigned a unique identifier within the project. The data management team ensures that the identifier is globally unique.
- Accompanying metadata such as the study protocol, experimental parameters etc. should be provided. This will make it possible for members of research networks to fully grasp the experimental set-up and data content.
- Project should follow standards defined for the different data domains, for example the clinical data interchange standards




Figure 4 Architecture of an ANN. Input layer contains the collected patient phenotype (demographic and clinical) and molecular data (RNA biomarkers), followed by two input layers, and an output layer which in this case predicts mortality, but could also predict MACE or other clinical outcomes.

consortium (CDISC) standards for collecting clinical and non-clinical data, or the minimum information about a microarray experiment (MIAME) for microarray experiments. An extensive list of recommended standards is defined in the eTRIKS—Standards Starter Pack Standards Guideline.¹⁰²

- The metadata of datasets should be collected by using the templates developed by ongoing efforts such as IMI-FAIRplus, COVID-19 Research Data Alliance working groups. The metadata should be published in a searchable registry or data-catalogue (e.g. IMI FAIRplus catalogue¹⁰³) to enable findability of datasets.

8.2 Making data accessible

Data should be made available to broader audience in accordance with the access model that will be defined by participant-informed consent and ethics/institutional review board approvals. This should include descriptions and data formats and in compliance with legal obligations, in particular the General Data Protection Regulation (GDPR). Data security is of paramount importance for protection of personally identifiable information.

8.3 Making data interoperable

Harmonization of data and metadata by applying standard ontologies, controlled terminologies, and state-of-the-art data models is pivotal for interoperability of the data that will facilitate cross-study analysis. Clinical and phenotype data should be standardized by using state-of-the-art standards such as the CDISC standards: Study Data Tabulation Model (SDTM), Clinical Data Acquisition Standards Harmonization (CDASH), and Analysis Data Model (ADaM). All clinical datasets from various cohorts should be mapped to International Severe Acute Respiratory and emerging Infection Consortium (ISARIC) COVID-19 eCRF.¹⁰⁴ In addition, application of controlled terminologies and ontologies described

in the eTRIKS Standards Starter Pack including ICD-11, MedDRA, WTO ATC codes, Human Phenotype Ontology (HPO), LOINC etc., is important to standardize and harmonize contents/values of clinical variables.

Omics or molecular data and associated metadata from both single-cell and bulk samples should be standardized using investigation, study and assay (ISA) framework¹⁰⁵ and corresponding minimum information guidelines (MIGs) such as MIAME format for transcriptome data, MIGS-MIMS (minimum information about a genome/metagenome sequence), and MIINSEQE (minimum information about a high-throughput NucLeo Pharmatide sequencing experiment) for both genome and transcriptome data. Study data tabulation model implementation guide (SDTMIG) pharmacogenomics/genetics (PGx) standards are useful to represent the genetic biomarkers including genetic variation, genotyping, and RNA expression data. To represent molecular entities within omics data, it is important to use identifiers from standard databases such as ENSEMBL gene, NCBI gene, ENSEMBL transcript for messenger RNA, UniProt for proteins, NCDB dbSNP for SNPs, GO for gene ontology, and KEGG (Kyoto Encyclopedia of Genes and Genomes)/Ingenuity for pathways. Molecular data, for example, the transcriptome of the biosamples and each transcript, are mapped to ENSEMBL transcript identifier (one of the stable and persistent identifiers). The corresponding genes and proteins are mapped to ENSEMBL genes, NCBI genes, UniProt identifiers and involved biological processes, cellular components, molecular functions using GO and KEGG identifiers. These stable identifiers provide cross-references to other biological databases and thus facilitate the interoperability of the molecular (-OMICS) data.

In addition to applying the state-of-the-art standards for clinical and omics data, application of standards in data management to guarantee data security and data privacy in compliance with GDPR and ethical guidelines is necessary. Given the sensitive nature of human data, the data and computing environment must be access-controlled and in/

output data flows should be encrypted, site restricted, and equipped with two-factor authentication wherever needed.

8.4 Increase data reuse (through clarifying licences)

The long-term sustainability for the database, analysis portal, and related outputs (results, tools, software modules, and algorithms) should be planned in advance. For archiving, preservation, and long-term usage of the data and software tools/algorithms, research network partners should have the capacity to provide long-term sustainability of translational research data through GDPR compliant hosting and tools. The process should follow well-defined access criteria and data protection needs. We recommend to prepare a sustainability plan for defining the rules to fulfil the legal processes (including addressing the issue of institutional data access committee responsibility), governance, and the economic viability of the database.

8.5 Data integration

A robust and secure data management and analysis platform, for example through a software portal and database, is important for the collection and integration of harmonized clinical, healthcare (electronic health records) data and pre-processed omics (molecular) data, imaging data, and real-world sensor/mobile data, biobank sample data, and metadata from various COVID-19 projects (Figure 2).

Such a data portal should also provide secure, easy, and robust interface for the input and integration of new data from ongoing recruitment of cohort studies. Analytical tools from existing initiatives/packages such as I2B2,¹⁰⁶ tranSMART,¹⁰⁷ SmartR,¹⁰⁸ European Genome-phenome Archive (EGA),¹⁰⁹ and eTRIKS platform¹¹⁰ are very useful to perform integrated data analysis and hypothesis generation. In order to store, process, and analyse imaging data, for example chest X-ray images from COVID-19 patients, a dedicated open-source imaging informatics solution such as XNAT¹¹¹ should be integrated into the platform instead of only storing the images in a file system. Such a portal will enable researchers to perform cross-study comparisons, slice and dice the cohorts based on certain clinical features, and run built-in workflows from the graphical user interface. An application programming interface to enable batch/programmatic interaction with the portal will provide structured and harmonized data to bioinformaticians, statisticians, and data scientists working with large amounts of data.

9. Data analysis, biostatistics, and artificial intelligence

After the data on RNA and clinical data are collected, secured, pre-processed, and integrated, most informative biomarkers to predict major adverse cardiovascular events (MACEs) and mortality of COVID-19 patients shall be identified. This identification can rely on biostatistical and machine-learning (ML) methods. Afterwards, ML should be utilized to build a classifier to predict MACE and mortality based on these biomarkers. For this approach to be used, RNA expression data accompanied by demographic and clinical data of patients are required, as well as information on MACE and mortality. To our knowledge, such data is not yet available—or has not yet been compiled from different patient cohorts—in a sufficient number of patients, allowing for application of ML methods. The dataset—once available—needs to be properly

organized for analysis. It should be split into training, validation, and test datasets. The training dataset is intended for biomarker discovery and model training, the validation dataset allows model selection and hyperparameter optimization, and the test dataset is for final testing. If the available dataset does not contain data from a large number of patients, k-fold cross-validation may be used to split the data. If the distribution of classes (with vs. without MACE, or dead vs. alive) is imbalanced, resampling of the training data (either undersampling the majority class or oversampling the minority class) may be appropriate.^{112,113}

9.1 Biomarker identification

The most basic approach to identify predictive RNAs is differential expression analysis: RNAs that are significantly over- or under-expressed in patients who experienced a MACE or died, compared to those who did not, are potential biomarkers. Various statistical methods can be used for this.¹¹⁴ However, this approach is simplistic, mainly in that it does not take into account interactions between the RNAs, so it can only serve as the first step. Two more sophisticated approaches can be explored: Bayesian variable selection (BVS) and feature selection.

Bayesian variable selection is a state-of-the-art statistical approach for selecting informative predictors such as RNA biomarkers.¹¹⁵ One first picks a class of models, such as linear or logistic regression models, to predict the end-point of interest (e.g. MACE or mortality) based on the predictors (RNA quantities). The goal is to select from this class of models those able to accurately predict end-points. To do so, prior probability distributions of their parameters need to be set first. The most appropriate strategies to do this are subject of ongoing research, but one of accepted automatic methods can certainly be used. We believe, though, that information on RNA's biological function from the NONCODE database,¹¹⁶ or overlap with genomic loci related to cardiovascular disease, could yield more informative biomarkers. Based on the models' prior probabilities and the collected data, one computes their posterior probability using the Bayes rule, where good models are the ones with a high posterior probability. Since the space of models is too large to search exhaustively, Monte-Carlo sampling is used, which can relatively quickly identify accurate models.

Feature selection is an approach that selects informative features (RNA biomarkers) to be used to train ML models that predict the end-point of interest (MACE or mortality).¹¹⁷ There are three main groups of feature-selection methods. Filter methods consider each feature in isolation and are similar to differential expression analysis, so they are rarely the best option. Embedded methods are a part of some ML algorithms. Their quality depends on the quality of the algorithm they are derived from, but they can take into account some interactions between features. Wrapper methods are the most complex ones and are conceptually similar to BVS. They search the space of feature combinations, and evaluate each combination by training a model on it and checking the model's accuracy. Since the space of feature combinations is again too large to search exhaustively, various types of greedy search are typically used. The main advantage of simple approaches, such as differential expression analysis or filter feature selection, is the clear justification for the selection of each biomarker. The disadvantage is that they can provide redundant biomarkers or fail to identify RNAs having biomarker potential only when combined with others. The advantage of BVS and more advanced feature selection is that they provide sets of biomarkers that perform well in combination. The disadvantages are that they are somewhat opaque and computationally expensive. Wrapper methods

appear to be the most flexible and potentially most powerful methods to identify predictive biomarkers.

From the two approaches, we recommend Bayesian variable selection and feature selection, either the one that results in better risk-prediction models on the validation dataset, or the combination of both can be used. They can be combined in sequence (one making the first selection and the other refining it) or in parallel (by using the intersection or union of the biomarkers selected by the two approaches). The best approach depends on the dataset and the outcome to predict, and needs to be determined experimentally.

9.2 Cardiovascular/COVID-19 risk prediction

After identifying the most informative RNA biomarkers, these—together with phenotype (demographic and clinical) data—are fed into ML algorithms to build risk-prediction models.

Figure 3 depicts the workflow of biomarker identification and COVID-19 risk prediction. Details of data collection and data management are depicted in Figure 2. The workflow starts with the data collection and management. This is followed by biomarker identification using the training dataset (cf. section Biomarker identification) and machine-learning model development with the validation dataset. Note that even though biomarker identification can be done independently of phenotype and clinical data, such data are often included in the prediction models. This enables one to analyse their capacity to predict MACE and mortality alongside with the RNA biomarkers. Finally, the prediction model is thoroughly evaluated using the test dataset.

The most common ML algorithms that have been successfully applied to problems that use omics and clinical data include artificial neural networks (ANNs), support vector machines (SVMs), and ensemble methods such as random forest.

ANNs have been designed to mimic human neural architecture. ANNs are able to effectively capture complex non-linear relationships in the data and are thus suitable for complex RNA data combined with clinical data. However, they are often computationally demanding, and compared to other algorithms they have many parameters that require tuning in order to optimize the prediction accuracy. Deep learning models are ANNs with multiple hidden layers. Many different deep neural network architectures exist.¹¹⁸ Figure 4 depicts an example of an ANN, with input data using the phenotype (demographic and clinical) and molecular data (RNA biomarkers). ANN consists of interconnected neurons, arranged in input layer, one or more hidden layers and an output layer. During training of ANN, the model learns from the examples provided in the training set.

SVM is another ML algorithm that is able to capture data non-linearity. SVM applies a kernel to map data into multidimensional space. A SVM model is a hyperplane that splits the classes in this multidimensional space in a way that minimizes the prediction error during data classification. The selection of the kernel function is crucial to the algorithm's performance. Compared to ANNs, SVMs tend to be more resistant to overfitting (better handle noise in the training data) and require less memory.

Ensemble methods are a popular approach that has been successfully applied to high-dimensional biomedical datasets with small sample size. The idea behind ensemble methods is to combine several base classifiers that will produce better classification results than a single classifier. One of the most successful ensemble methods is random forest. Random forest uses a set of decision trees that form a forest. In order to avoid

overfitting, each decision tree in the forest uses a random subset of samples from the training set, and a random subset of features. Classification is then performed based on the majority vote of the trees. For example, the FEELnc tool uses random forest for annotation of lncRNAs and achieves an AUC of 0.97.¹¹⁹ The lncLocator tool uses an ensemble of support vector machine and random forest classifiers to predict lncRNA subcellular localization.¹²⁰

Considering the recent success of deep learning, we believe this method to be worth investigating. Due to the many parameters in deep ANNs, it typically requires more data than other methods, hence it may become unsuitable for datasets limited in size. In this case, ensemble methods are likely to provide better results. To evaluate the performance of the classification methods, various measures such as classification accuracy, sensitivity, specificity and precision can be used.¹²¹ The AUC is a particularly suitable performance indicator, since it evaluates the performance of models over all possible trade-offs between type 1 and 2 errors.

10. Translational aspects: development of diagnostics and therapy for COVID-19

The rapid spread, high mortality in some geographical areas, and the yet largely unknown long-term consequences of COVID-19 including cardiovascular pathologies all highlighted the need to develop effective diagnostic and prognostic biomarkers and therapeutics against SARS-CoV-2. Basic research and development of novel biomarkers and therapeutics run in parallel on the basis of broad collaboration between key players of the biomedical field including industrial and academic partners, national governments, and regulatory agencies as well as investors. Ongoing repositioning of existing drugs as well as development of novel drugs, vaccines, and a variety of medical devices for prevention and treatment of COVID-19 have been at the front line of very recent research activities.

Development of RNA diagnostics and therapeutics, especially small non-coding RNA compounds, attracted the attention of the pharmaceutical industry in the past few years that has been further accelerated by the rapid outbreak of COVID-19. Indeed, development of RNA molecules for diagnosis, prognosis, and treatment of SARS-CoV-2 and other RNA viruses has been recently proposed.^{122,123} Currently, there are nine ongoing or completed clinical trials when searching for miRNA and COVID-19 in the clinicaltrials.gov platform, showing the rapidly increasing activity of translational research in this field. Moreover, extracellular vesicles—as important players in the life cycle of RNA viruses as well as cargo particles for non-coding RNAs—may provide opportunities for more sensitive diagnosis and targeted therapies for SARS-CoV-2.^{124,125} Although there are currently no examples of molecular diagnostic assays based on cardiovascular RNA biomarkers of COVID-19 and utilizing digital PCR as a means to quantify circulating RNA transcripts, we believe that this technology holds great promise and may rapidly be applied to COVID-19 tests.

Another aspect of the COVID-19 pandemic is the need for cardioprotective strategies to prevent the long-term cardiovascular consequences of the disease. Yet, despite intensive efforts, the development of cardioprotective therapies has been unsuccessful in the last three decades.¹²⁶ Small non-coding RNA fingerprints of COVID-19 itself and the different comorbidities and their co-medications that affect the infection may

provide a useful tool to develop diagnostic and prognostic markers and to discover novel drug targets to prevent and treat COVID-19 and its cardiovascular consequences.¹²⁷ Understanding the molecular interactions between SARS-CoV-2 and its host as well as the influence of cardiovascular risk factors, comorbidities, and medications on clinical outcomes may significantly speed up the lengthy process of development of diagnostics and therapeutics not only against COVID-19 but also other diseases.^{127–129}

11. Conclusion and perspectives

COVID-19 has brought about an unexpected and unprecedented historical period, worldwide. Despite the tremendous efforts and reactivity of all stakeholders from the broad healthcare sector—clinicians, healthcare staff, researchers, funding bodies, and regulatory authorities—the burden of COVID-19 is enormous, medically, socially, and economically.

The research field has been very reactive, and multiple networks of experts and task forces have been formed to tackle the challenge of finding drugs and biomarkers of COVID-19. Building an effective coordination of large interdisciplinary networks involved in multi-centre studies is key for success of biomarker projects. RNA biomarkers combined with artificial intelligence-based strategies will certainly help in building algorithms to aid in clinical decision making and personalization of healthcare through risk stratification of patients. Efficient academia–industry partnerships are essential to rapid marketing and clinical use of novel disease biomarkers. Novel tools based on systems biomedicine concepts and artificial intelligence methods are needed to speed up the translational process and clinical application.

Whilst it is obvious that the cardiovascular burden associated with SARS-CoV-2 infection is alarming and deserves great attention during healthcare of COVID-19 patients, it is also important to keep in mind that more than a third of hospitalized COVID-19 patients present psychological distress and neurological manifestations such as headache, ischemic stroke, seizures, and other diverse encephalopathies.¹³⁰ SARS-CoV-2 has been detected in the brain and cerebrospinal fluid,¹³¹ and is associated with encephalitis. Various neurological sequelae have been associated with the Spanish influenza pandemic and other coronaviruses.¹³² Therefore, a deeper knowledge of the host–pathogen interactions involving regulatory RNAs⁸² in the brain–heart axis¹³³ may provide novel avenues for discovery of biomarkers and therapeutic pathways to improve healthcare and prepare for future pandemics.

Acknowledgements

This article is based upon work from EU-CardioRNA COST Action CA17129 (www.cardiorna.eu) supported by COST (European Cooperation in Science and Technology), and from the COVIRNA project which received funding from the European Union's Horizon 2020 Research and Innovation Programme under the grant agreement n°101016072. The open access fee of this article is covered by EU-CardioRNA COST Action CA17129, funded by COST.

Conflict of interest: L.B. is founder and shareholder of Glycardial Diagnosis SL; received a research grant from AstraZeneca; hold advisory board work for Sanofi, Bayer and AstraZeneca; and received speaker fees from Lilly, MSD-Boehringer and AstraZeneca (all outside of this work). L.D.W. is founder and shareholder of Mirabilis Therapeutics BV. P.F. is the founder and CEO of Pharmahungary Group, a group of R&D companies. M.H. is employed by TAmiRNA GmbH and company

shareholder. M.H. holds patents related to diagnostic and therapeutic applications of microRNAs. F.M. acted as consultant for Amicus Therapeutics on a topic not related to the manuscript. T.T. is founder and shareholder of Cardior Pharmaceuticals GmbH. T.T. received support (including speaker fees) and/or holds advisory board activities for Boehringer Ingelheim, Sanofi-Aventis, Takeda, Amicus Therapeutics, Novo Nordisk (all outside the field of this manuscript). Y.D. holds patents related to diagnostic and therapeutic applications of RNAs.

Funding

L.B. is funded by the EU Horizon 2020 project COVIRNA (Grant Agreement # 101016072), the Spanish Ministry of Economy and Competitiveness of Science [PNS2016-76819-R]; the Carlos III Institute of Health [CIBERCV CB16/11/00411 and Red Terapia Celular TerCel RD16/0011/0018] cofounded by FEDER; and the Fundación Investigación Cardiovascular-Fundación Jesus Serra.E.L.R. is funded by the CardioVascular Onderzoek Nederland (CVON) EARLY-HFPEF-2015 consortium (Dutch Heart Foundation). A.J. is funded by a Horizon 2020 Marie Skłodowska-Curie Action (H2020-MSCA-IF-EF-ST 2019 Grant: 893435). I.C. is funded by the EU Horizon 2020 project COVIRNA (Grant Agreement 101016072). L.D.W. acknowledges support from the Netherlands CardioVascular Research Initiative: the Dutch Heart Foundation, Dutch Federation of University Medical Centers, ZonMW, and the Royal Netherlands Academy of Sciences (CVON2017-ARENA PRIME). L.D.W. was further supported by ERC Consolidator Grant 311549 CALMIRS, a VICI award 918-156-47 from NWO and a Marie Skłodowska-Curie grant agreement No. 813716. P.F. is funded by the National Research, Development and Innovation Office of Hungary (Research Excellence Program—TKP, National Heart Program NVKP 16-1-2016-0017, Investment into the Future—COVID-19 project at Semmelweis University); the Higher Education Institutional Excellence Program of the Ministry of Human Capacities in Hungary, within the framework of the Therapeutic Development thematic program of the Semmelweis University; and EU Horizon 2020 projects COVIRNA (Grant #101016072) and CRYSTAL3 (MSCA-RISE Project #101007931). M.G. is funded by the EU Horizon 2020 grants SCIENCE (643478), CRESpace (732170), and ReGenHeart (731532). M.H. is funded by Eurostars Grant No. 871562 and FFG Early Stage No. 874078. M. L. is funded by the EU Horizon 2020 project COVIRNA (Grant Agreement 101016072) and the Slovenian Research Agency (research core funding No. P2-0209). F.M. is funded by the Italian Ministry of Health, (Ricerca Corrente and 5 × 1000), the Telethon Foundation (# GGP19035A), AFM-Telethon grant (# 23054), and EU Horizon 2020 projects COVIRNA (Grant #101016072). T.T. is funded by the EU Horizon 2020 project Cardioregenix. Y.D. is funded by the EU Horizon 2020 project COVIRNA (Grant Agreement # 101016072), the National Research Fund (grants # C14/BM/8225223, C17/BM/11613033, and COVID-19/2020-1/14719577/miRCOVID.), the Ministry of Higher Education and Research, and the Heart Foundation—Daniel Wagner of Luxembourg.

References

- <https://ec.europa.eu/info/funding-tenders/opportunities/portal/screen/covid-19>.
- Zhang J-Y, Wang X-M, Xing X, Xu Z, Zhang C, Song J-W, Fan X, Xia P, Fu J-L, Wang S-Y, Xu R-N, Dai X-P, Shi L, Huang L, Jiang T-J, Shi M, Zhang Y, Zumla A, Maeurer M, Bai F, Wang F-S. Single-cell landscape of immunological responses in patients with COVID-19. *Nat Immunol* 2020;**21**:1107–1118.
- Shi S, Qin M, Shen B, Cai Y, Liu T, Yang F, Gong W, Liu X, Liang J, Zhao Q, Huang H, Yang B, Huang C. Association of cardiac injury with mortality in hospitalized patients with COVID-19 in Wuhan, China. *JAMA Cardiol* 2020;**5**:802.
- Naksuk N, Lazar S, Peerapattit TB. Cardiac safety of off-label COVID-19 drug therapy: a review and proposed monitoring protocol. *Eur Heart J Acute Cardiovasc Care* 2020;**9**:215–221.
- Guo T, Fan Y, Chen M, Wu X, Zhang L, He T, Wang H, Wan J, Wang X, Lu Z. Cardiovascular implications of fatal outcomes of patients with coronavirus disease 2019 (COVID-19). *JAMA Cardiol* 2020;**5**:811.

6. Judson GL, Kelemen BW, Njoroge JN, Mahadevan VS. Cardiovascular implications and therapeutic considerations in COVID-19 infection. *Cardiol Ther* 2020;**9**:293–305.
7. Si D, Du B, Ni L, Yang B, Sun H, Jiang N, Liu G, Massé S, Jin L, Nanthakumar J, Bhaskaran A, Yang P, Nanthakumar K. Death, discharge and arrhythmias among patients with COVID-19 and cardiac injury. *CMAJ* 2020;**192**:E791–E798.
8. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020.
9. Alvarez-Garcia J, Lee S, Gupta A, Cagliostro M, Joshi AA, Rivas-Lasarte M, Contreras J, Mitter SS, LaRocca G, Tlachi P, Brunjes D, Glicksberg BS, Levin MA, Nadkarni G, Fayad Z, Fuster V, Mancini D, Lala A. Prognostic impact of prior heart failure in patients hospitalized with COVID-19. *J Am Coll Cardiol* 2020;**76**:2334–2348.
10. Gao Y, Wang C, Kang K, Peng Y, Luo Y, Liu H, et al. Cytokine storm may not be the chief culprit for the deterioration of COVID-19. *Viral Immunol* 2020.
11. Mudd PA, Crawford JC, Turner JS, Souquette A, Reynolds D, Bender D, et al. Distinct inflammatory profiles distinguish COVID-19 from influenza with limited contributions from cytokine storm. *Sci Adv* 2020.
12. Sinha P, Matthay MA, Calfee CS. Is a “cytokine storm” relevant to COVID-19? *JAMA Intern Med* 2020;**180**:1152–1154.
13. Mancia G, Rea F, Ludergnani M, Apolone G, Corrao G. Renin-angiotensin-aldosterone system blockers and the risk of Covid-19. *N Engl J Med* 2020;**382**:2431–2440.
14. Reynolds HR, Adhikari S, Pulgarin C, Troxel AB, Iturrate E, Johnson SB, Hausvater A, Newman JD, Berger JS, Bangalore S, Katz SD, Fishman GI, Kunichoff D, Chen Y, Ogedegbe G, Hochman JS. Renin-angiotensin-aldosterone system inhibitors and risk of Covid-19. *N Engl J Med* 2020;**382**:2441–2448.
15. Metzler B, Siotrzonek P, Binder RK, Bauer A, Reinstadler SJ. Decline of acute coronary syndrome admissions in Austria since the outbreak of COVID-19: the pandemic response causes cardiac collateral damage. *Eur Heart J* 2020;**41**:1852–1853.
16. Channappanavar R, Zhao J, Perlman S. T cell-mediated immune response to respiratory coronaviruses. *Immunol Res* 2014;**59**:118–128.
17. Li W, Moore MJ, Vasilieva N, Sui J, Wong SK, Berne MA, Somasundaram M, Sullivan JL, Luzuriaga K, Greenough TC, Choe H, Farzan M. Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. *Nature* 2003;**426**:450–454.
18. Chen Y, Guo Y, Pan Y, Zhao ZJ. Structure analysis of the receptor binding of nCoV. *Biochem Biophys Res Commun* 2019;**2020**.
19. Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Veesler D. Structure, function, and antigenicity of the SARS-CoV-2 spike glycoprotein. *Cell* 2020;**181**:281–292.
20. Letko M, Marzi A, Munster V. Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nat Microbiol* 2020;**5**:562–569.
21. Zou X, Chen K, Zou J, Han P, Hao J, Han Z. Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Front Med* 2020;**14**:185–192.
22. Yoshikawa T, Hill T, Li K, Peters CJ, Tseng CT. Severe acute respiratory syndrome (SARS) coronavirus-induced lung epithelial cytokines exacerbate SARS pathogenesis by modulating intrinsic functions of monocyte-derived macrophages and dendritic cells. *J Virol* 2009;**83**:3039–3048.
23. Liu S, Su X, Pan P, Zhang L, Hu Y, Tan H, Wu D, Liu B, Li H, Li H, Li Y, Dai M, Li Y, Hu C, Tsung A. Neutrophil extracellular traps are indirectly triggered by lipopolysaccharide and contribute to acute lung injury. *Sci Rep* 2016;**6**:37252.
24. Everaert BR, Muylle J, Bartholomeus T. T. Emerging cardiological issues during the COVID-19 pandemic. *Eur J Clin Invest* 2020;**e13270**.
25. Guo J, Wei X, Li Q, Li L, Yang Z, Shi Y, Qin Y, Zhang X, Wang X, Zhi X, Meng D. Single-cell RNA analysis on ACE2 expression provides insights into SARS-CoV-2 potential entry into the bloodstream and heart injury. *J Cell Physiol* 2020;**235**:9884–9894.
26. Yang L, Han Y, Nilsson-Payant BE, Gupta V, Wang P, Duan X, Tang X, Zhu J, Zhao Z, Jaffré F, Zhang T, Kim TW, Harschnitz O, Redmond D, Houghton S, Liu C, Naji A, Ciceri G, Guttkonda S, Bram Y, Nguyen D-HT, Cioffi M, Chandar V, Hoagland DA, Huang Y, Xiang J, Wang H, Lyden D, Borczuk A, Chen HJ, Studer L, Pan FC, Ho DD, tenOever BR, Evans T, Schwartz RE, Chen S. A human pluripotent stem cell-based platform to study SARS-CoV-2 tropism and model virus infection in human cells and organoids. *Cell Stem Cell* 2020;**27**:125–136.
27. Bojkova D, Wagner JUG, Shumliakivska M, Aslan GS, Saleem U, Hansen A, et al. SARS-CoV-2 infects and induces cytotoxic effects in human cardiomyocytes. *Cardiovasc Res* 2020.
28. Escher F, Pietsch H, Aleshcheva G, Bock T, Baumeier C, Elsaesser A, et al. Detection of viral SARS-CoV-2 genomes and histopathological changes in endomyocardial biopsies. *ESC Heart Fail* 2020.
29. Meyer P, Degrauwe S, Van Delden C, Ghadri JR, Templin C. Typical takotsubo syndrome triggered by SARS-CoV-2 infection. *Eur Heart J* 2020;**41**:1860.
30. Bangalore S, Sharma A, Slotwiner A, Yatskar L, Harari R, Shah B, Ibrahim H, Friedman GH, Thompson C, Alviar CL, Chadow HL, Fishman GI, Reynolds HR, Keller N, Hochman JS. ST-segment elevation in patients with Covid-19 - a case series. *N Engl J Med* 2020;**382**:2478–2480.
31. Ranucci M, Ballotta A, Di Dedda U, Bayshnikova E, Dei Poli M, Resta M, Falco M, Albano G, Menicanti L. The procoagulant pattern of patients with COVID-19 acute respiratory distress syndrome. *J Thromb Haemost* 2020;**18**:1747–1751.
32. Commission Implementing Decision of 3.7.2020 granting a conditional marketing authorisation under Regulation (EC) No 726/2004 of the European Parliament and of the Council for “Veklury - remdesivir”, a medicinal product for human use. https://ec.europa.eu/health/documents/community-register/2020/20200703148664/dec_148664_en.pdf (22 July 2020, date last accessed).
33. First COVID-19 treatment recommended for EU authorisation. <https://www.ema.europa.eu/en/news/first-covid-19-treatment-recommended-eu-authorisation> (30 June 2020, date last accessed).
34. NIH Clinical Trial Shows Remdesivir Accelerates Recovery from Advanced COVID-19. <https://www.niaid.nih.gov/news-events/nih-clinical-trial-shows-remdesivir-accelerates-recovery-advanced-covid-19> (2 July 2020, date last accessed).
35. Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, Hohmann E, Chu HY, Luetkemeyer A, Kline S, Lopez de Castilla D, Finberg RW, Dierberg K, Tapson V, Hsieh L, Patterson TF, Paredes R, Sweeney DA, Short WR, Touloumi G, Lye DC, Ohmagari N, Oh M-D, Ruiz-Palacios GM, Benfield T, Fätkenheuer G, Kortepeter MG, Atmar RL, Creech CB, Lundgren J, Babiker AG, Pett S, Neaton JD, Remdesivir TH, Bonnett T, Green M, Makowski M, Osinusi A, Nayak S, Lane HC. Remdesivir for the treatment of Covid-19 - final report. *N Engl J Med* 2020;**383**:1813–1826.
36. Coronavirus (COVID-19) Update: FDA Issues Emergency Use Authorization for Potential COVID-19 Treatment. <https://www.fda.gov/news-events/press-announcements/coronavirus-covid-19-update-fda-issues-emergency-use-authorization-potential-covid-19-treatment> (30 June 2020, date last accessed).
37. Fact Sheet for Health Care Providers Emergency Use Authorization (EUA) of Remdesivir (Gs-5734™).
38. COVID-19 Treatment Guidelines Panel. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines. National Institutes of Health. <https://www.covid19treatmentguidelines.nih.gov/> (30 June 2020, date last accessed).
39. Peto Q, Abdool Karim MA, AMHenao-Restrepo, C Hernández García, M-P Kieny, R Malekzadeh, S Murthy, M-P Preziosi, S Reddy, M Roses Periago Vasee Sathiyamoorthy, J-AR, SwaminathanS; WHO Solidarity Trial Consortium HP. Repurposed antiviral drugs for COVID-19 – interim WHO SOLIDARITY trial results. *medRxiv Preprint* 2020. doi:10.1101/2020.10.15.20209817. This version posted October 15, 2020. This article is a preprint and has not been peer-reviewed.
40. Group RC, Horby P, Lim WS, Emberson JR, Mafham M, Bell JL, et al. Dexamethasone in hospitalized patients with Covid-19 - preliminary report. *N Engl J Med* 2020.
41. Randomised Evaluation of COVID-19 Therapy (RECOVERY). Low-cost dexamethasone reduces death by up to one third in hospitalised patients with severe respiratory complications of COVID-19. 2020. <https://www.recoverytrial.net/news/low-cost-dexamethasone-reduces-death-by-up-to-one-third-in-hospitalised-patients-with-severe-respiratory-complications-of-covid-19> (30 June 2020, date last accessed).
42. Mehta P, McAuley DF, Brown M, Sanchez E, Tattersall RS, Manson JJ. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet* 2020;**395**:1033–1034.
43. Li L, Zhang W, Hu Y, Tong X, Zheng S, Yang J, et al. Effect of convalescent plasma therapy on time to clinical improvement in patients with severe and life-threatening COVID-19: a randomized clinical trial. *JAMA* 2020.
44. Shao Z, Feng Y, Zhong L, Xie Q, Lei M, Liu Z, et al. Clinical efficacy of intravenous immunoglobulin therapy in critical patients with COVID-19: a multicenter retrospective cohort study. *medRxiv* 2020. 2020.2004.2011.20061739.
45. FDA Issues Emergency Use Authorization for Convalescent Plasma as Potential Promising COVID-19 Treatment, Another Achievement in Administration’s Fight Against Pandemic. <https://www.fda.gov/news-events/press-announcements/fda-issues-emergency-use-authorization-convalescent-plasma-potential-promising-covid-19-treatment> (1 September 2020, date last accessed).
46. Driggin E, Madhavan MV, Bikdeli B, Chuich T, Laracy J, Biondi-Zoccai G, Brown TS, Der Nigoghossian C, Zidar DA, Haythe J, Brodie D, Beckman JA, Kirtane AJ, Stone GW, Krumholz HM, Parikh SA. Cardiovascular considerations for patients, health care workers, and health systems during the COVID-19 pandemic. *J Am Coll Cardiol* 2020;**75**:2352–2371.
47. Mueller C, McDonald K, de Boer RA, Maisel A, Cleland JGF, Kozhuharov N, Coats AJS, Metra M, Mebazaa A, Ruschitzka F, Lainscak M, Filippatos G, Seferovic PM, Meijers WC, Bayes-Genis A, Mueller T, Richards M, Januzzi JL; on behalf of the Heart Failure Association of the European Society of Cardiology. Heart failure association of the European society of cardiology practical guidance on the use of natriuretic peptide concentrations. *Eur J Heart Fail* 2019;**21**:715–731.
48. Bonow RO, Fonarow GC, O’Gara PT, Yancy CV. Association of coronavirus disease 2019 (COVID-19) with myocardial injury and mortality. *JAMA Cardiol* 2020;**5**:751.
49. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, Qiu Y, Wang J, Liu Y, Wei Y, Xia J, Yu T, Zhang X, Zhang L. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* 2020;**395**:507–513.

50. Song JW, Lam SM, Fan X, Cao WJ, Wang SY, Tian H. Omics-driven systems interrogation of metabolic dysregulation in COVID-19 pathogenesis. *Cell Metab* 2020.
51. Malik P, Patel U, Mehta D, Patel N, Kelkar R, Akrmah M, et al. Biomarkers and outcomes of COVID-19 hospitalisations: systematic review and meta-analysis. *BMJ Evid Based Med* 2020.
52. Sandoval Y, Januzzi JL, Jaffe AS. Cardiac troponin for assessment of myocardial injury in COVID-19. *J Am Coll Cardiol* 2020;**76**:1244–1258.
53. Zeng F, Huang Y, Guo Y, Yin M, Chen X, Xiao L, Deng G. Association of inflammatory markers with the severity of COVID-19: a meta-analysis. *Int J Infect Dis* 2020;**96**:467–474.
54. Li L-Q, Huang T, Wang Y-Q, Wang Z-P, Liang Y, Huang T-B, Zhang H-y, Sun W, Wang Y. COVID-19 patients' clinical characteristics, discharge rate, and fatality rate of meta-analysis. *J Med Virol* 2020;**92**:577–583.
55. Zhang ZL, Hou YL, Li DT, Li FZ. Laboratory findings of COVID-19: a systematic review and meta-analysis. *Scand J Clin Lab Invest* 2020;**80**:441–447.
56. Danwang C, Endomba FT, Nkeck JR, Wouna DLA, Robert A, Noubiap JJ. A meta-analysis of potential biomarkers associated with severity of coronavirus disease 2019 (COVID-19). *Biomark Res* 2020;**8**.
57. Elshazli RM, Toraih EA, Elgaml A, El-Mowafy M, El-Mesery M, Amin MN, Hussein MH, Killackey MT, Fawzy MS, Kandil E. Diagnostic and prognostic value of hematological and immunological markers in COVID-19 infection: a meta-analysis of 6320 patients. *PLoS One* 2020;**15**:e0238160.
58. Velavan TP, Meyer CG. Mild versus severe COVID-19: laboratory markers. *Int J Infect Dis* 2020;**95**:304–307.
59. Borges do Nascimento IJ, von Groote TC, O'Mathúna DP, Abdulazeem HM, Henderson C, Jayarajah U, Weerasekera I, Poklepovic Pericic T, Klapproth HEG, Puljak L, Cacic N, Zakarija-Grkovic I, Guimarães SMM, Atallah AN, Bragazzi NL, Marcolino MS, Marusic A, Jeronic A; On behalf of the International Task Force Network of Coronavirus Disease 2019 (InterNetCOVID-19). Clinical, laboratory and radiological characteristics and outcomes of novel coronavirus (SARS-CoV-2) infection in humans: a systematic review and series of meta-analyses. *PLoS One* 2020;**15**:e0239235.
60. Zheng Z, Peng F, Xu B, Zhao J, Liu H, Peng J, Li Q, Jiang C, Zhou Y, Liu S, Ye C, Zhang P, Xing Y, Guo H, Tang W. Risk factors of critical & mortal COVID-19 cases: a systematic literature review and meta-analysis. *J Infect* 2020;**81**:e16–e25.
61. Mesas AE, Caverro-Redondo I, Álvarez-Bueno C, Sarriá Cabrera MA, Maffei de Andrade S, Sequí-Dominguez I, Martínez-Vizcaíno V. Predictors of in-hospital COVID-19 mortality: a comprehensive systematic review and meta-analysis exploring differences by age, sex and health conditions. *PLoS One* 2020;**15**:e0241742.
62. Li X, Pan X, Li Y, An N, Xing Y, Yang F, Tian L, Sun J, Gao Y, Shang H, Xing Y. Cardiac injury associated with severe disease or ICU admission and death in hospitalized patients with COVID-19: a meta-analysis and systematic review. *Crit Care* 2020;**24**.
63. Walker C, Deb S, Ling H, Wang Z. Assessing the elevation of cardiac biomarkers and the severity of COVID-19 infection: a meta-analysis. *J Pharm Pharm Sci* 2020;**23**:396–405.
64. Lippi G, Lavie CJ, Sanchis-Gomar F. Cardiac troponin I in patients with coronavirus disease 2019 (COVID-19): evidence from a meta-analysis. *Prog Cardiovasc Dis* 2020;**63**:390–391.
65. Li X, Guan B, Su T, Liu W, Chen M, Bin Waleed K, Guan X, Gary T, Zhu Z. Impact of cardiovascular disease and cardiac injury on in-hospital mortality in patients with COVID-19: a systematic review and meta-analysis. *Heart* 2020;**106**:1142–1147.
66. Li J-W, Han T-W, Woodward M, Anderson CS, Zhou H, Chen Y-D, Neal B. The impact of 2019 novel coronavirus on heart injury: a systematic review and meta-analysis. *Prog Cardiovasc Dis* 2020;**63**:518–524.
67. Vrsalovic M, Vrsalovic PA. Cardiac troponins predict mortality in patients with COVID-19: a meta-analysis of adjusted risk estimates. *J Infect* 2020;**81**:e99–e100.
68. Cao Y, Li L, Feng Z, Wan S, Huang P, Sun X, Wen F, Huang X, Ning G, Wang W. Comparative genetic analysis of the novel coronavirus (2019-nCoV/SARS-CoV-2) receptor ACE2 in different populations. *Cell Discov* 2020;**6**:11.
69. Groß S, Jahn C, Cushman S, Bär C, Thum T. SARS-CoV-2 receptor ACE2-dependent implications on the cardiovascular system: from basic science to clinical implications. *J Mol Cell Cardiol* 2020;**144**:47–53.
70. Mehta N, Kalra A, Nowacki AS, Anjewierden S, Han Z, Bhat P, Carmona-Rubio AE, Jacob M, Procop GW, Harrington S, Milinovich A, Svensson LG, Jehi L, Young JB, Chung MK. Association of use of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers with testing positive for coronavirus disease 2019 (COVID-19). *JAMA Cardiol* 2020;**5**:1020.
71. Thum T. SARS-CoV-2 receptor ACE2 expression in the human heart: cause of a post-pandemic wave of heart failure? *Eur Heart J* 2020;**41**:1807–1809.
72. Muslim S, Nasrin N, Alotaibi FO, Prasad G, Singh SK, Alam I. Treatment options available for COVID-19 and an analysis on possible role of combination of rhACE2, angiotensin (1-7) and angiotensin (1-9) as effective therapeutic measure. *SN Compr Clin Med* 2020;1–6.
73. Li C, Hu X, Li L, Li JH. Differential microRNA expression in the peripheral blood from human patients with COVID-19. *J Clin Lab Anal* 2020;**34**:e23590.
74. Meydan C, Madrer N, Soreq H. The near death of COVID-19: NEAT1, DANCR, and co-modulated cholinergic RNAs link to inflammation. *Front Immunol* 2020;**11**.
75. Bertolazzi G, Cipollina C, Benos PV, Tumminello M, Coronello C. miR-1207-5p can contribute to dysregulation of inflammatory response in COVID-19 via targeting SARS-CoV-2 RNA. *Front Cell Infect Microbiol* 2020;**10**.
76. Garg A, Seeliger B, Derda AA, Xiao K, Gietz A, Scherf K. Circulating cardiovascular microRNAs in critically ill COVID-19 patients Short title: microRNA signatures in COVID-19. *Eur J Heart Fail* 2021.
77. Evans PC, Ed Rainger G, Mason JC, Guzik TJ, Osto E, Stamatakis Z, et al. Endothelial dysfunction in COVID-19: a position paper of the ESC working group for atherosclerosis and vascular biology, and the ESC council of basic cardiovascular science. *Cardiovasc Res* 2020.
78. Bienvenu LA, Noonan J, Wang X, Peter K. Higher mortality of COVID-19 in males: sex differences in immune response and cardiovascular comorbidities. *Cardiovasc Res* 2020;**116**:2197–2206.
79. Badimon L, Devaux Y. Transcriptomics research to improve cardiovascular health-care the EU-CardioRNA COST Action CA17129. *Eur Heart J* 2020.
80. Emanuel C, Badimon L, Martelli F, Potočnjak I, Carpusca I, Robinson EL, Devaux Y. Call to action for the cardiovascular side of COVID-19: a call for cooperative action from the EU-CardioRNA COST Action. *Eur Heart J* 2020;**41**:1796–1797.
81. Guzik TJ, Mohiddin SA, Dimarco A, Patel V, Savvatis K, Marelli-Berg FM, et al. COVID-19 and the cardiovascular system: implications for risk assessment, diagnosis, and treatment options. *Cardiovasc Res* 2020.
82. Gomes C, P D C, Schroen B, Kuster GM, Robinson EL, Ford K, Squire IB, Heymans S, Martelli F, Emanuel C, Devaux Y; on behalf of the EU-CardioRNA COST Action (CA17129). Regulatory RNAs in heart failure. *Circulation* 2020;**141**:313–328.
83. Landry P, Plante I, Ouellet DL, Perron MP, Rousseau G, Provost P. Existence of a microRNA pathway in anucleate platelets. *Nat Struct Mol Biol* 2009;**16**:961–966.
84. Juzenas S, Lindqvist CM, Ito G, Dolshanskaya Y, Halfvarson J, Franke A, Hemmrich-Stanisak G. Depletion of erythropoietic miR-486-5p and miR-451a improves detectability of rare microRNAs in peripheral blood-derived small RNA sequencing libraries. *NAR Genomics Bioinf* 2020;**2**.
85. Krammer TL, Mayr M, Hackl M. microRNAs as promising biomarkers of platelet activity in antiplatelet therapy monitoring. *Int J Mol Sci* 2020;**21**.
86. Wang K, Yuan Y, Cho J-H, McClarty S, Baxter D, Galas DJ. Comparing the microRNA spectrum between serum and plasma. *PLoS One* 2012;**7**:e41561.
87. Blondal T, Jensby Nielsen S, Baker A, Andreasen D, Mouritzen P, Wrang Teilm M, Dahlsveen IK. Assessing sample and miRNA profile quality in serum and plasma or other biofluids. *Methods* 2013;**59**:S1–S6.
88. Kirschner MB, Kao SC, Edelman JJ, Armstrong NJ, Valley MP, van Zandwijk N, Reid G. Haemolysis during sample preparation alters microRNA content of plasma. *PLoS One* 2011;**6**:e24145.
89. Cheng HH, Yi HS, Kim Y, Kroh EM, Chien JW, Eaton KD, Goodman MT, Tait JF, Tewari M, Pritchard CC. Plasma processing conditions substantially influence circulating microRNA biomarker levels. *PLoS One* 2013;**8**:e64795.
90. Mitchell AJ, Gray WD, Hayek SS, Ko Y-A, Thomas S, Rooney K, Awad M, Roback JD, Quyyumi A, Searles CD. Platelets confound the measurement of extracellular miRNA in archived plasma. *Sci Rep* 2016;**6**:32651.
91. Pavkovic M, Riefke B, Ellinger-Ziegelbauer H. Urinary microRNA profiling for identification of biomarkers after cisplatin-induced kidney injury. *Toxicology* 2014;**324**:147–157.
92. Arroyo JD, Chevillet JR, Kroh EM, Ruf IK, Pritchard CC, Gibson DF, Mitchell PS, Bennett CF, Pogosova-Agadjanyan EL, Stirewalt DL, Tait JF, Tewari M. Argonaute2 complexes carry a population of circulating microRNAs independent of vesicles in human plasma. *Proc Natl Acad Sci USA* 2011;**108**:5003–5008.
93. Priglinger E, Strohmeier K, Weigl M, Lindner C, Auer D, Gimona M, Barsch M, Jacak J, Redl H, Grillari J, Sandhofer M, Hackl M, Wolbank S. SVF-derived extracellular vesicles carry characteristic miRNAs in lipedema. *Sci Rep* 2020;**10**.
94. Dickman CTD, Lawson J, Jabalee J, MacLellan SA, LePard NE, Bennenwith KL, Garnis C. Selective extracellular vesicle exclusion of miR-142-3p by oral cancer cells promotes both internal and extracellular malignant phenotypes. *Oncotarget* 2017;**8**:15252–15266.
95. Théry C, Witwer KW, Aikawa E, Alcaraz MJ, Anderson JC, Andriantsitohaina R, Antoniou A, Arab T, Archer F, Atkin-Smith GK, Ayre DC, Bach J-M, Bachurski D, Baharvand H, Balaj L, Baldacchino S, Bauer NN, Baxter AA, Bebawy M, Beckham C, Bedina Zavec A, Benmoussa A, Berardi AC, Bergese P, Bielska E, Blenkiron C, Bobis-Wozowicz S, Boilard E, Boireau V, Bongiovanni A, Borrás FE, Bosch S, Boulanger CM, Breakfield X, Breglio AM, Brennan MÁ, Brigstock DR, Brisson A, Broekman ML, Bromberg JF, Bryl-Górecka P, Buch S, Buck AH, Burger D, Busatto S, Buschmann D, Bussolati B, Buzás EI, Byrd JB, Camussi G, Carter DR, Caruso S, Chamley LW, Chang Y-T, Chen C, Chen S, Cheng L, Chin AR, Clayton A, Clerici SP, Cocks A, Cocucci E, Coffey RJ, Cordeiro-da-Silva A, Couch Y, Coumans FA, Coyle B, Crescitelli R, Criado MF, D'Souza-Schorey C, Das S, Datta Chaudhuri A, de Candia P, De Santana EF, De Wever O, del Portillo HA, Demaret T, Deville S, Devitt A, Dhondt B, Di Vizio D, Dieterich LC, Dolo V, Dominguez Rubio AP, Dominici M, Dourado MR, Driedonks TA, Duarte FV, Duncan HM, Eichenberger RM, Ekström K, EL Andaloussi S, Etie-Caille C, Erdbrügger U, Falcón-Pérez JM, Fatima F, Fish JE, Flores-Bellver M, Försonits A, Frelet-Barrand A, Fricke F, Fuhrmann G, Gabrielsson S, Gámez-Valero A, Gardiner C, Gärtner K, Gaudin R, Gho YS, Giesel B, Gilbert C, Girona M, Giusti I, Goberdhan DC, Görgens A, Gorski SM, Greening DW, Gross JC, Gualerzi A, Gupta GN, Gustafson D,

- Handberg A, Haraszti RA, Harrison P, Hegyesi H, Hendrix A, Hill AF, Hochberg FH, Hoffmann KF, Holder B, Holthofer H, Hosseinkhani B, Hu G, Huang Y, Huber V, Hunt S, Ibrahim AG-E, Ikezu T, Inal JM, Isin M, Ivanova A, Jackson HK, Jacobsen S, Jay SM, Jayachandran M, Jenster G, Jiang L, Johnson SM, Jones JC, Jong A, Jovanovic-Talisman T, Jung S, Kalluri R, Kano S-I, Kaur S, Kawamura Y, Keller ET, Khamari D, Khomyakova E, Khvorova A, Kierulff P, Kim KP, Kislinger T, Klingeborn M, Klinke DJ, Kornek M, Kosanovic MM, Kovács ÁF, Krämer-Albers E-M, Krasemann S, Krause M, Kurochkin IV, Kusuma GD, Kuypers S, Laitinen S, Langevin SM, Languino LR, Lannigan J, Lässer C, Laurent LC, Lavieou G, Lázaro-Ibáñez E, Le Lay S, Lee M-S, Lee YXF, Lemos DS, Lenassi M, Leszczynska A, Li IT, Liao K, Libregts SF, Ligeti E, Lim R, Lim SK, Liné A, Linnemannstons K, Llorente A, Lombard CA, Lorenovic MJ, Lörcincz ÁM, Lötvall J, Lovett J, Lowry MC, Loyer X, Lu Q, Lukomska B, Lunavat TR, Maas SL, Malhi H, Marcilla A, Mariani J, Mariscal J, Martens-Uzunova ES, Martin-Jaular L, Martinez MC, Martins VR, Mathieu M, Mathivanan S, Maugeri M, McGinnis LK, McVey MJ, Meckes DG, Meehan KL, Mertens I, Minciacci VR, Möller A, Møller Jørgensen M, Morales-Kastresana A, Morhayim J, Mullier F, Muraca M, Musante L, Mussack V, Muth DC, Myburgh KH, Najrana T, Nawaz M, Nazarencenko I, Nejsum P, Neri C, Neri T, Nieuwland R, Nimrichter L, Nolan JP, Nolte-t Hoen EN, Noren Hooten N, O'Driscoll L, O'Grady T, O'Loughlin A, Ochiya T, Olivier M, Ortiz A, Ortiz LA, Osteikoetxea X, Østergaard O, Ostrowski M, Park J, Pegtel DM, Peinado H, Perut F, Pfaffl MW, Phinney DG, Pieters BC, Pink RC, Pisetky DS, Pogge von Strandmann E, Polakovicova I, Poon IK, Powell BH, Prada I, Pulliam L, Quesenberry P, Radeghieri A, Raffai RL, Raimondo S, Rak J, Ramirez MI, Raposo G, Rayyan MS, Regeve-Rudzik N, Ricklefs FL, Robbins PD, Roberts DD, Rodrigues SC, Rohde E, Rome S, Rouschop KM, Rugheiti A, Russell AE, Saá P, Sahoo S, Salas-Huenoleo E, Sánchez C, Saugstad JA, Saul MJ, Schiffelers RM, Schneider R, Schøyen TH, Scott A, Shahaj E, Sharma S, Shatnyeva O, Shekari F, Shelke GV, Shetty AK, Shiba K, Siljander PR-M, Silva AM, Skowronek A, Snyder OL, Soares RP, Sódar BW, Soekmadji C, Sotillo J, Stahl PD, Stoorvogel W, Stott SL, Strasser EF, Swift S, Tahara H, Tewari M, Timms K, Tiwari S, Tixeira R, Tkach M, Toh WS, Tomasini R, Torrecillas AC, Tosar JP, Toxavidis V, Urbanelli L, Vader P, van Balkom BW, van der Grein SG, Van Deun J, van Herwijnen MJ, Van Keuren-Jensen K, van Niel G, van Royen ME, van Wijnen AJ, Vasconcelos MH, Vechetti IJ, Veit TD, Vella LJ, Velot É, Verweij FJ, Vestad B, Viñas JL, Visnovitz T, Vukman KV, Wahlgren J, Watson DC, Wauben MHM, Weaver A, Webber JP, Weber V, Wehman AM, Weiss DJ, Welsh JA, Wendt S, Wheelock AM, Wiener Z, Witte L, Wolfram J, Xagorari A, Xander P, Xu J, Yan X, Yáñez-Mó M, Yin H, Yuana Y, Zappulli V, Zarubova J, Žekas V, Zhang J-y, Zhao Z, Zheng L, Zheutlin AR, Zickler AM, Zimmermann P, Zivkovic AM, Zocco D, Zuba-Surma EK. Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *J Extracell Vesicles* 2018;**7**:1535750.
96. Buschmann D, Haberberger A, Kirchner B, Spornraft M, Riedmaier I, Schelling G, Pfaffl MW. Toward reliable biomarker signatures in the age of liquid biopsies - how to standardize the small RNA-Seq workflow. *Nucleic Acids Res* 2016;**44**:5995–6018.
97. Fu Y, Wu PH, Beane T, Zamore PD, Weng Z. Elimination of PCR duplicates in RNA-seq and small RNA-seq using unique molecular identifiers. *BMC Genomics* 2018;**19**.
98. Barberán-Soler S, Vo JM, Hogans RE, Dallas A, Johnston BH, Kazakov SA. Decreasing miRNA sequencing bias using a single adapter and circularization approach. *Genome Biol* 2018;**19**.
99. Wong RKY, MacMahon M, Woodside JV, Simpson DA. A comparison of RNA extraction and sequencing protocols for detection of small RNAs in plasma. *BMC Genomics* 2019;**20**:446.
100. Lutzmayer S, Enugutti B, Nodine MD. Novel small RNA spike-in oligonucleotides enable absolute normalization of small RNA-Seq data. *Sci Rep* 2017;**7**:5913.
101. Everaert C, Helmoortel H, Decock A, Hulstaert E, Van Paemel R, Verniers K, Nuytens J, Anckaert J, Nijs N, Tulkens J, Dhondt B, Hendrix A, Mestdagh P, Vandesompele J. Performance assessment of total RNA sequencing of human biofluids and extracellular vesicles. *Sci Rep* 2019;**9**:17574.
102. Bussery J, Denis LA, Guillon B, Liu P, Marchetti G, Rahal G. eTRIKS platform: conception and operation of a highly scalable cloud-based platform for translational research and applications development. *Comput Biol Med* 2018;**95**:99–106.
103. <https://datacatalog.elixir-luxembourg.org/> (20 April 2021, date last accessed).
104. <https://isarc.tghn.org/covid-19-clinical-research-resources> (20 April 2021, date last accessed).
105. Sansone S-A, Rocca-Serra P, Field D, Maguire E, Taylor C, Hofmann O, Fang H, Neumann S, Tong W, Amaral-Zettler L, Begley K, Booth T, Bougueleret L, Burns G, Chapman B, Clark T, Coleman L-A, Copeland J, Das S, de Daruvar A, de Matos P, Dix I, Edmunds S, Evelo CT, Forster MJ, Gaudet P, Gilbert J, Goble C, Griffin JL, Jacob D, Kleinjans J, Harland L, Haug K, Hermjakob H, Sui SJH, Laederach A, Liang S, Marshall S, McGrath A, Merrill E, Reilly D, Roux M, Shamu CE, Shang CA, Steinbeck C, Trefethen A, Williams-Jones B, Wolstencroft K, Xenarios I, Hide W. Toward interoperable bioscience data. *Nat Genet* 2012;**44**:121–126.
106. Murphy SN, Weber G, Mendis M, Gainer V, Chueh HC, Churchill S, Kohane I. Serving the enterprise and beyond with informatics for integrating biology and the bedside (i2b2). *J Am Med Assoc* 2010;**303**:124–130.
107. Athey BD, Braxenthaler M, Haas M, Guo Y. transSMART: an open source and community-driven informatics and data sharing platform for clinical and translational research. *AMIA Summits Transl Sci Proc* 2013;**2013**:6–8.
108. Herzinger S, Gu W, Satagopam V, Eifes S, Rege K, Barbosa-Silva A, Schneider R, eTRIKS Consortium. SmartR: an open-source platform for interactive visual analytics for translational research data. *Bioinformatics* 2017;**33**:2229–2231.
109. <https://ega-archive.org/> (20 April 2021, date last accessed).
110. <https://public.etriks.org/> (20 April 2021, date last accessed).
111. www.xnat.org (20 April 2021, date last accessed).
112. Chawla NV, Lazarevic A, Hall LO, Bowyer KW. *SMOTEBoost: Improving Prediction of the Minority Class in Boosting*. Berlin, Heidelberg: Springer Berlin Heidelberg; 2003. p107–119.
113. Chen JJ, Tsai CA, Young JF, Kodell RL. Classification ensembles for unbalanced class sizes in predictive toxicology. *SAR QSAR Environ Res* 2005;**16**:517–529.
114. Van den Berge K, Hembach KM, Soneson C, Tiberi S, Clement L, Love MI, et al. RNA sequencing data: hitchhiker's guide to expression analysis. *Ann Rev Biomed Data Sci* 2019;**2**:139–173.
115. Bottolo L, Chadeau-Hyam M, Hastie DJ, Zeller T, Liqueur B, Newcombe P, Yengo L, Wild PS, Schillert A, Ziegler A, Nielsen SF, Butterworth AS, Ho WK, Castagné R, Munzel T, Tregouet D, Falchi M, Cambien F, Nordestgaard BG, Fumeron F, Tybjaerg-Hansen A, Froguel P, Danesh J, Petretto E, Blankenberg S, Tiret L, Richardson S. GUESS-ing polygenic associations with multiple phenotypes using a GPU-based evolutionary stochastic search algorithm. *PLoS Genet* 2013;**9**:e1003657.
116. Liu C, Bai B, Skogerbo G, Cai L, Deng W, Zhang Y, et al. NONCODE: an integrated knowledge database of non-coding RNAs. *Nucleic Acids Res* 2004;**33**:D112–D115.
117. Wang L, Wang Y, Chang Q. Feature selection methods for big data bioinformatics: A survey from the search perspective. *Methods* 2016;**111**:21–31.
118. LeCun Y, Bengio Y, Hinton G. Deep learning. *Nature* 2015;**521**:436–444.
119. Wucher V, Legeai F, Hedan B, Rizk G, Lagoutte L, Leeb T, et al. FEELnc: a tool for long non-coding RNA annotation and its application to the dog transcriptome. *Nucleic Acids Res* 2017;**45**:e57.
120. Cao Z, Pan X, Yang Y, Huang Y, Shen H-B. The IncLocator: a subcellular localization predictor for long non-coding RNAs based on a stacked ensemble classifier. *Bioinformatics* 2018;**34**:2185–2194.
121. Jiao Y, Du P. Performance measures in evaluating machine learning based bioinformatics predictors for classifications. *Quant Biol* 2016;**4**:320–330.
122. Chen L, Zhong L. Genomics functional analysis and drug screening of SARS-CoV-2. *Genes Dis* 2020.
123. Gutierrez A, de Azeredo Lima CH, Miranda RL, Gadelha MR. What is the potential function of microRNAs as biomarkers and therapeutic targets in COVID-19? *Infect Genet Evol* 2020;**85**:104417.
124. Hassanpour M, Rezaie J, Nouri M, Panahi Y. The role of extracellular vesicles in COVID-19 virus infection. *Infect Genet Evol* 2020;**85**:104422.
125. Sluijter JPG, Davidson SM, Boulanger CM, Buzás EI, de Kleijn DPV, Engel FB, Giricz Z, Hausenloy DJ, Kishore R, Lecour S, Leor J, Madonna R, Perrino C, Prunier F, Sahoo S, Schiffelers RM, Schulz R, Van Laake LW, Ytrehus K, Ferdinandy P. Extracellular vesicles in diagnostics and therapy of the ischaemic heart: position Paper from the Working Group on Cellular Biology of the Heart of the European Society of Cardiology. *Cardiovasc Res* 2018;**114**:19–34.
126. Hausenloy DJ, Garcia-Dorado D, Bøtker HE, Davidson SM, Downey J, Engel FB, Jennings R, Lecour S, Leor J, Madonna R, Ovize M, Perrino C, Prunier F, Schulz R, Sluijter JPG, Van Laake LW, Vinten-Johansen J, Yellon DM, Ytrehus K, Heusch G, Ferdinandy P. Novel targets and future strategies for acute cardioprotection: position paper of the European Society of Cardiology Working Group on Cellular Biology of the Heart. *Cardiovasc Res* 2017;**113**:564–585.
127. Perrino C, Barabási A-L, Condorelli G, Davidson SM, De Windt L, Dimmeler S, Engel FB, Hausenloy DJ, Hill JA, Van Laake LW, Lecour S, Leor J, Madonna R, Mayr M, Prunier F, Sluijter JPG, Schulz R, Thum T, Ytrehus K, Ferdinandy P. Epigenomic and transcriptomic approaches in the post-genomic era: path to novel targets for diagnosis and therapy of the ischaemic heart? Position Paper of the European Society of Cardiology Working Group on Cellular Biology of the Heart. *Cardiovasc Res* 2017;**113**:725–736.
128. Ferdinandy P, Hausenloy DJ, Heusch G, Baxter GF, Schulz R. Interaction of risk factors, comorbidities, and comedications with ischemia/reperfusion injury and cardioprotection by preconditioning, postconditioning, and remote conditioning. *Pharmacol Rev* 2014;**66**:1142–1174.
129. Parini P, Altucci L, Balligand JL, Baumbach J, Ferdinandy P, Filetti S, et al. The network medicine imperative and the need for an international network medicine consortium. *Am J Med* 2020.
130. Mao L, Wang MD, Chen SH, He QW, Chang J, Hong CD, et al. Neurological manifestations of hospitalized patients with COVID-19 in Wuhan, China: a retrospective case series study. *MedRxiv* 2020. 2020.2002.2022.20026500.
131. Xiang P, Xu XM, Gao LL, Wang HZ, Xiong HF, Li RH. First case of 2019 novel coronavirus disease with encephalitis. *ChinaXiv* 200015 2020:T202003.
132. Wu Y, Xu X, Chen Z, Duan J, Hashimoto K, Yang L, Liu C, Yang C. Nervous system involvement after infection with COVID-19 and other coronaviruses. *Brain Behav Immun* 2020;**87**:18–22.
133. Beis D, Zerr I, Martelli F, Doehner W, Devaux Y. RNAs in brain and heart diseases. *Int J Mol Sci* 2020;**21**:3717.