



Prenatal Exposure to Per- and Polyfluoroalkyl Substances, Maternal Thyroid Dysfunction, and Child Autism Spectrum Disorder

Hyeong-Moo Shin¹, Jiwon Oh², Rebecca J. Schmidt^{2,3}, Elizabeth N. Pearce⁴

¹Department of Environmental Science, Baylor University, Waco, TX; ²Department of Public Health Sciences, University of California, Davis, CA; ³UC Davis MIND (Medical Investigations of Neurodevelopmental Disorders) Institute, Sacramento, CA; ⁴Section of Endocrinology, Diabetes, and Nutrition, Department of Medicine, Boston University School of Medicine, Boston, MA, USA

Autism spectrum disorder (ASD), with its high economic and societal costs, is a growing public health concern whose prevalence has risen steadily over the last two decades. Although actual increased incidence versus improved diagnosis remains controversial, the increased prevalence of ASD suggests non-inherited factors as likely contributors. There is increasing epidemiologic evidence that abnormal maternal thyroid function during pregnancy is associated with increased risk of child ASD and other neurodevelopmental disorders. Prenatal exposure to endocrine-disrupting chemicals such as per- and polyfluoroalkyl substances (PFAS) is known to disrupt thyroid function and can affect early brain development; thus, thyroid dysfunction is hypothesized to mediate this relationship. The concept of a potential pathway from prenatal PFAS exposure through thyroid dysfunction to ASD etiology is not new; however, the extant literature on this topic is scant. The aim of this review is to evaluate and summarize reports with regard to potential mechanisms in this pathway.

Keywords: Antibodies; Autism spectrum disorder; Endocrine disrupting chemicals; Environmental exposure; Pregnancy; Thyroid

INTRODUCTION

Autism spectrum disorder (ASD) is a complex neurodevelopmental condition characterized by limited interests, repetitive behaviors, and impaired social interaction and communication [1]. ASD is a growing public health concern in part due to its high economic and societal costs, especially in developed countries [2]. Annual costs (direct medical, direct non-medical, and productivity combined) of ASD in 2025 are projected to reach

nearly one-half trillion dollars in the United States [3]. The prevalence of ASD has risen steadily in the last two decades [4]; in the United States in 2018, one of every 44 children (3 to 8 years old) was estimated to have ASD [5]. Although actual increased incidence versus improved diagnosis remains controversial [6-8], the rapid rise in ASD prevalence suggests that environmental factors may contribute to ASD etiology [9,10]. In the last decade, environmental research linking modifiable factors to ASD has proliferated, with replication or meta-analysis

Received: 4 October 2022, **Revised:** 16 October 2022,
Accepted: 25 October 2022

Corresponding author: Hyeong-Moo Shin
Department of Environmental Science, Baylor University, One Bear Place
#97266, Waco, TX 76798, USA
Tel: +1-254-710-7627, **Fax:** +1-254-710-3409
E-mail: hyeongmoo_shin@baylor.edu

Copyright © 2022 Korean Endocrine Society

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

covering pesticides [11,12], air pollution [13,14], maternal fever during pregnancy [15,16], periconceptional nutrition [17-19], maternal diabetes or obesity [20,21], preeclampsia [22], and interpregnancy interval [23-25].

Thyroid hormones (THs) are essential for brain development and influence brain function throughout life [26,27]. Animal studies have shown that THs regulate crucial processes of brain development in mammals, including proliferation, migration, and differentiation of neuronal cells [28-30]. There is epidemiologic evidence that abnormal maternal thyroid function during pregnancy is associated with increased risk of child ASD and other neurodevelopmental disorders [31]. In addition, the prevalence of thyroid peroxidase antibody (TPO-Ab), a marker for thyroid autoimmunity, was reported to be higher in families with autism probands than in comparison subjects [32].

Simultaneous with a growing understanding of the importance of maternal thyroid homeostasis for fetal brain development, chemical production volumes have increased 300-fold since the 1970s [33], leading to widespread human exposure to compounds known as endocrine-disrupting chemicals (EDCs) [34-36]. EDCs are defined as exogenous chemicals that interfere with hormone actions, resulting in increased risk of adverse health effects [37]. A wide range of EDCs disrupt thyroid homeostasis in laboratory animal studies [38]. Hundreds of synthetic chemicals interfere with the production, transport, and metabolism of THs [39]. Studies have shown that a broad range of EDCs can bind to TH receptors, may produce complex effects on TH signaling [40-42], and either alone or in combination, act at many levels in the thyroid system [43].

Among a large number of EDCs, this review focuses on per- and polyfluoroalkyl substances (PFAS), a class of synthetic chemicals widely used in consumer (e.g., cookware, dental floss) and industrial (e.g., lining of gas pipes, surfactant) applications [44]. Recently PFAS have received significant public attention due to increasing evidence of their widespread environmental contamination and adverse health effects. As PFAS-containing products are widely used in daily life, many common PFAS compounds have been detected in the blood of most of the United States general population [45]. PFAS have also been detected in cord blood [46,47] and in amniotic fluid [48,49]. Importantly, both animal studies [50-52] and epidemiologic studies [53-55] have shown that prenatal exposure to PFAS disrupts thyroid function and immune systems, which can alter early brain development (Fig. 1). Moreover, there is epidemiologic evidence that PFAS exposure is associated with child neurodevelopmental disorders such as attention-deficit/hyperactivity disorder [56,57], indicating that PFAS may adversely affect child brain development.

The aim of this review is to assess evidence for a potential pathway from prenatal PFAS exposure through abnormal thyroid function to ASD etiology. Hypothesizing that maternal thyroid dysfunction mediates a relationship between prenatal PFAS exposure and child ASD, we have focused on potential mechanisms related to thyroid dysfunction. This review also discusses antibody-mediated immune dysregulation that may cause thyroid dysfunction [58,59]. Building on the current report, subsequent research may help set the stage in support of prenatal thyroid treatment and strategies to prevent or reduce PFAS exposure.




Fig. 1. Per- and polyfluoroalkyl substances (PFAS) are known to disrupt immune systems and the hypothalamus-pituitary-thyroid axis, either alone or in combination. Thyroid peroxidase antibody (TPO-Ab) and thyroglobulin antibody (Tg-Ab), which are common in autoimmune thyroid disorders, may cause thyroid dysfunction. Primary and secondary pathways discussed in this review are represented in blue and magenta, respectively. TSH, thyroid stimulating hormone; T3, triiodothyronine; T4, thyroxine.

THYROID FUNCTION

Thyroid function is assessed with thyroid stimulating hormone (TSH) and free thyroxine (FT4). When TSH is high and FT4 is low or within a normal range, hypothyroidism is diagnosed. According to the American Thyroid Association guidelines, the population-based trimester-specific normal reference range for serum TSH should be used when assessing thyroid function during pregnancy [60]. However, reference ranges which are assay-, laboratory-, cohort-, and population-specific are preferred when available. Some studies have defined maternal abnormal thyroid function based on hospital diagnosis codes for hyperthyroidism or hypothyroidism or on prescriptions for THs or anti-thyroid drugs [61,62].

POTENTIAL MECHANISMS

Abnormal thyroid function and ASD

Epidemiologic studies have suggested that maternal gestational

hypothyroidism, hyperthyroidism, and hypothyroxinemia were associated with increased risk of ASD in children (Table 1) [61-66]. In addition, low FT4 levels in cord blood were associated with increased ASD risk [67,68]. Overt maternal hypothyroidism is associated with impaired offspring cognition, which reflects that placental transfer of maternal THs to the fetus is essential for the regulation of fetal brain development [69,70]. Severe iodine deficiency (in which inadequate substrate for TH synthesis causes both maternal and fetal TH levels to be low) may cause cretinism, a syndrome of profoundly impaired growth and neurodevelopment. Maternal thyroid dysfunction during pregnancy is also known to be associated with adverse maternal and fetal outcomes such as preterm delivery, preeclampsia, and low birth weight [71-73], which are known risk factors for ASD [22,74-77].

Antibody-mediated immune dysregulation, ASD, and thyroid dysfunction

There is substantial evidence that autoimmunity and immune

Table 1. Associations between Prenatal Maternal Thyroid Function or Thyroid Antibodies and Autistic Outcomes in Children

Study	Population (countries)	Sample size, <i>n</i>	ASD diagnosis method	Thyroid or antibody measurement	Key thyroid dysfunctions or antibody positivity ^a	Estimate (95% CI) ^b
Roman et al. (2013) [64]	Population-based cohort (The Netherlands)	4,309	PDP-CBCL SRS	TSH FT4 TPO-Ab	Severe hypothyroxinemia	OR, 3.89 (1.83–8.20)
Andersen et al. (2014) [61]	Population-based cohort (Denmark)	857,014	ICD-10	Not measured ^c	Hyperthyroidism Hypothyroidism	HR, 1.34 (1.14–1.59)
Andersen et al. (2018) [66]	Population-based cohort (Denmark)	7,624	ICD-10	TSH FT4	Hypothyroidism Overt hyperthyroidism	HR, 1.75 (1.12–2.73) HR, 2.18 (1.08–4.39)
Getahun et al. (2018) [63]	Retrospective cohort (USA)	397,201	DSM-IV	TSH FT4	Hypothyroidism	HR, 1.31 (1.13–1.53)
Levie et al. (2018) [65]	Population-based cohort (Spain, the Netherlands, the United Kingdom)	9,036	CAST PDP-CBCL SCDC	TSH FT4 TPO-Ab	FT4 <5th percentile FT4 >95th percentile FT4 <2.5th percentile FT4 >97.5th percentile	OR, 1.5 (1.0–2.3) OR, 1.2 (0.7–2.1) OR, 1.3 (0.7–2.5) OR, 1.9 (1.0–3.4)
Rotem et al. (2020) [62]	Population-based cohort (Israel)	437,222	ICD-9	TSH FT4	Hypothyroidism Hyperthyroidism Other thyroid conditions	OR, 1.28 (1.11–1.49) OR, 1.39 (0.88–2.18) OR, 1.22 (1.05–1.42)
Brown et al. (2015) [32]	Nested case-control study (Finland)	960	ICD-10	TPO-Ab	TPO-Ab+	OR, 1.78 (1.16–2.75)

ASD, autism spectrum disorder; CI, confidence interval; PDP-CBCL, the Pervasive Developmental Problems Subscale of the Child Behavior Checklist for Toddlers; SRS, social responsiveness scale; TSH, thyroid stimulating hormone; FT4, free thyroxine; TPO-Ab, thyroid peroxidase antibody; OR, odds ratio; ICD-9 or 10, the ninth or tenth revision of the International Classification of Diseases; HR, hazard ratio; DSM-IV, Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition; CAST, Childhood Autism Spectrum Test; SCDC, Social Communication Disorder Checklist; TPO-Ab+, positive to TPO-Ab.

^aEach study defined thyroid dysfunction using various levels of TSH and/or FT4; ^bAdjusted for various confounders and covariates in each study; ^cHyperthyroidism and hypothyroidism were defined by the various combinations of first hospital diagnosis of hyperthyroidism or hypothyroidism, number of anti-thyroid medications, and number of prescriptions of thyroid hormones.

system dysfunction likely play a role in the development of ASD [78]. A body of epidemiologic evidence has shown that autoimmune disorders are significantly more frequent in families of autism probands than in those of comparison subjects [79-81]. More mothers of children with ASD had ASD-specific autoantibodies to proteins in the developing brain, compared with mothers of typically developing children [82-87]. In addition, a higher prevalence of maternal ASD-specific autoantibodies during pregnancy is associated with increased risk of child ASD and other neurodevelopmental disorders [88-90].

In a case-control Finnish study, maternal TPO-Ab positivity during pregnancy was associated with increased risk of child ASD (odds ratio, 1.78; 95% confidence interval, 1.16 to 2.75) (Table 1) [32], implying that there is a potential role of thyroid autoimmunity in ASD etiology, although in that analysis maternal FT4 and TSH levels were not independently associated with ASD. It has been demonstrated that women with TPO-Ab positivity have a blunting of the typical thyroidal response to human chorionic gonadotropin in early gestation [91,92], resulting in lower serum FT4 levels and higher serum TSH levels. Thus, thyroid autoimmunity (high TPO-Ab and/or thyroglobulin antibody [Tg-Ab]) may be a secondary intermediate outcome (1) between PFAS exposure and ASD or (2) between PFAS exposure and thyroid dysfunction (Fig. 1).

Prenatal PFAS exposure and ASD

To our knowledge, seven epidemiologic studies to date have examined associations between maternal PFAS exposure and child ASD [49,93-98]. Although results differed, three studies showed that higher prenatal exposure to perfluorohexane sulfonate, perfluorononanoate, perfluorooctanoate (PFOA), or perfluorooctane sulfonate was associated with increased risk of child ASD (Table 2) [96-98]. Potential reasons for inconsistent results among these studies include differences in timing of exposure measures in pregnancy, characteristics of study populations, methods of identification or confirmation of ASD cases, and genetic factors. In addition, because PFAS were moderately correlated with each other and one PFAS may confound another, consideration of a single compound in the model may explain, at least in part, the inconsistent findings. The number of ASD cases is relatively small in three prospective birth cohorts [49, 93,97], potentially resulting in inadequate power to detect associations.

Prenatal PFAS exposure, thyroid dysfunction, and immune dysregulation

Many epidemiologic studies have shown that higher prenatal PFAS concentrations in maternal blood are associated with altered TH levels in maternal blood or cord blood (Table 3) [99-

Table 2. Associations between Prenatal Maternal PFAS Exposure and Autistic Outcomes in Children

Study	Population (country)	Sample size, n ^a	ASD diagnosis method	Blood sample type	PFAS with notable findings	Estimate (95% CI) ^b
Braun et al. (2014) [93]	Prospective birth cohort (USA)	175	SRS	Prenatal maternal serum	PFOS	β , -2.0 (-4.4 to 0.4)
Liew et al. (2015) [94]	Nested case-control study (Denmark)	770	ICD-10	Prenatal maternal plasma	PFOA PFOS	RR, 0.98 (0.82-1.16) RR, 0.87 (0.74-1.02)
Lyll et al. (2018) [95]	Nested case-control study (USA)	986	DSM-IV	Prenatal maternal serum	PFOA PFOS	OR, 0.92 (0.74-1.15) OR, 0.92 (0.73-1.17)
Long et al. (2019) [49]	Retrospective cohort (Denmark)	210	ICD-8 ICD-10	Amniotic fluid	PFOS	OR, 0.41 (0.17-0.97)
Shin et al. (2020) [96]	Case-control study (USA)	453	ADI-R ADOS-G	Postnatal maternal serum ^c	PFHxS PFOS	OR, 1.46 (0.98-2.18) OR, 1.03 (0.99-1.08)
Oh et al. (2021) [97]	Prospective birth cohort (USA)	173	ADOS	Prenatal maternal serum	PFOA PFNA	RR, 1.31 (1.04-1.65) RR, 1.79 (1.13-2.85)
Skogheim et al. (2021) [98]	Prospective birth cohort (Norway)	1,380	ICD-10	Prenatal maternal plasma	PFOA	OR, 1.71 (1.20-2.45)

PFAS, per- and polyfluoroalkyl substances; ASD, autism spectrum disorder; CI, confidence interval; SRS, social responsiveness scale; PFOS, perfluorooctane sulfonate; ICD-8 or 10, the eighth or tenth revision of the International Classification of Diseases; PFOA, perfluorooctanoate; RR, relative risk; DSM-IV, Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition; OR, odds ratio; ADI-R, Autism Diagnostic Interview-Revised; ADOS-G, Autism Diagnostic Observation Schedules-Generic; PFHxS, perfluorohexane sulfonate; PFNA, perfluorononanoate.

^aNumber of mother-child pairs in which the child has SRS or a final diagnosis of ASD; ^bAdjusted for various confounders and covariates in each study; ^cReconstructed maternal PFAS serum concentrations at the time of pregnancy using a simple pharmacokinetic model and maternal blood samples collected when the child was 2 to 5 years old in a case-control study.

110]. Although study results are not entirely consistent, increases or decreases in THs indicate that prenatal PFAS exposure may disrupt maternal or neonatal thyroid homeostasis. In addition, three studies have reported that higher exposure to a mixture of PFAS was associated with increased or decreased THs [107,109,110], implying that PFAS can disrupt thyroid either alone or in combination. Some PFAS levels were significantly higher in infants with congenital hypothyroidism compared with healthy infants [111]. Two studies reported the relationship between prenatal PFAS exposure and thyroid autoimmunity; PFOA was inversely associated with TPO-Ab [106], whereas

perfluorododecanoic acid was positively associated with TPO-Ab [110].

Studies have shown that PFAS exposure alone was not associated with TH levels among those with normal TPO-Ab but was associated with increases and decreases in THs among those with high TPO-Ab levels or low iodine concentrations. In a prospective birth cohort study, higher prenatal PFAS levels were associated with increased TSH levels only among pregnant women with high TPO-Ab (≥ 9 IU/mL) [112]. Another prospective birth cohort showed that higher prenatal PFOA levels were associated with lower prevalence of TPO-Ab in mater-

Table 3. Associations between PFAS Exposure and Thyroid Dysfunction in Pregnant Women or Neonates

Study	Population (country)	Sample size, <i>n</i> ^a	Blood sample type	Thyroid or antibody measurement	Relationships between PFAS and thyroid ^b
Kim et al. (2011) [99]	Prospective birth cohort (South Korea)	44 mothers 43 infants	Prenatal maternal plasma Cord serum	TT3, TT4, TSH	No relationship ^c
de Cock et al. (2014) [100]	Prospective birth cohort (The Netherlands)	83	Cord plasma	TT4	PFOA (↑) → TT4 (↑)
Wang et al. (2014) [101]	Prospective birth cohort (Taiwan)	285 mothers 116 neonates	Prenatal maternal plasma Cord serum	TT3, TT4 FT4, TSH	PFNA (↑), PFUnDA (↑), PFDoA (↑) → FT4 (↓), TT4 (↓)
Berg et al. (2015) [102]	Prospective birth cohort (Norway)	515	Prenatal maternal serum	TT3, TT4 FT4, TSH	PFOS (↑) → TSH (↑)
Shah-Kulkarni et al. (2016) [103]	Retrospective birth cohort (South Korea)	279	Cord serum	TT3, TT4, TSH	PFPeA (↑) → TT4 (↑)
Berg et al. (2017) [104]	Prospective birth cohort (Norway)	391	Prenatal maternal serum	TT3, TT4 FT4, TSH, TPO-Ab	PFOS (↑) → TSH (↑)
Preston et al. (2018) [105]	Prospective birth cohort (USA)	732 mothers 480 neonates	Prenatal maternal serum Cord serum	TT4, FT4I T3U, TSH	PFOA (↑), PFHxS (↑), MeFOSAA (↑) → FT4I (↓)
Itoh et al. (2019) [106]	Prospective birth cohort (Japan)	701	Prenatal maternal serum Cord serum	FT3, FT4, TSH, TPO-Ab, Tg-Ab	PFOS (↑) → TSH (↑) PFOA (↑) → TPO-Ab (↓)
Lebeaux et al. (2020) [108]	Prospective birth cohort (USA)	468	Prenatal maternal serum Cord serum	FT3, TT3, TT4 FT4, TSH	PFOA (↑), PFOS (↑), PFHxS (↑) → FT4 (↓) TPO-Ab (↑) → FT4 (↓)
Preston et al. (2020) [107]	Prospective birth cohort (USA)	726 mothers 465 neonates	Prenatal maternal plasma Cord serum	TT4, FT4I T3U, TSH	PFAS mixture (↑) → FT4I (↓)
Liang et al. (2020) [109]	Prospective birth cohort (China)	300	Cord plasma	FT3, TT4 FT4, TSH	PFAS mixture (↑) → FT3 (↑)
Guo et al. (2021) [110]	Prospective birth cohort (China)	490	Cord serum	FT3, TT3, TT4 FT4, TSH, TPO-Ab, Tg-Ab	PFHpA (↑), PFNA (↑) → TSH (↓) PFOA (↑), PFOS (↑), PFNA (↑), PFUnDA (↑) → TT4 (↑) PFOS (↑), PFUnDA (↑), PFDoA (↑) → FT4 (↑) PFDoA (↑) → TT3 (↑), TPO-Ab (↑) PFAS mixture (↑) → TT4 (↑), FT4 (↑)

PFAS, per- and polyfluoroalkyl substances; TT3, total triiodothyronine; TT4, total thyroxine; TSH, thyroid stimulating hormone; PFOA, perfluorooctanoate; FT4, free thyroxine; PFNA, perfluorononanoate; PFUnDA, perfluoroundecanoic acid; PFDoA, perfluorododecanoic acid; PFOS, perfluorooctane sulfonate; PFPeA, perfluoro-n-pentanoic acid; TPO-Ab, thyroid peroxidase antibody; FT4I, free thyroxine index; PFHxS, perfluorohexane sulfonate; MeFOSAA, 2-(N-methyl-perfluorooctane sulfonamido) acetate; FT3, free triiodothyronine; Tg-Ab, thyroglobulin antibody; T3U, triiodothyronine resin uptake; PFHpA, perfluoroheptanoic acid.

^aNumber of mother-child pairs, unless otherwise noted; ^bAdjusted for various confounders and covariates in each study; ^cResults were not statistically significant even after adjusting for major covariates.

nal blood and that PFAS-induced thyroid disruption and susceptibility may vary by the presence of two maternal TPO-Ab and Tg-Ab [106]. In a subset of United States adults, PFAS exposure was more likely to be associated with thyroid disruption in individuals with both TPO-Ab positivity and a urinary iodine concentration (UIC) $<100 \mu\text{g/L}$ than in individuals with TPO-Ab positivity or low UIC alone, or TPO-Ab negative individuals with $\text{UIC} \geq 100 \mu\text{g/L}$ [113].

CONCLUSIONS

To date, no studies have examined a potential pathway from prenatal PFAS exposure through thyroid dysfunction and/or thyroid autoimmunity to ASD etiology within a well-characterized ASD population. Iodine deficiency is associated with increased risk of hypothyroidism [114] and known to cause brain damage [65,115]. However, most studies included in Tables 1, 3 have failed to measure important biomarkers that might affect maternal thyroid function, such as iodine status or thyroid antibodies. Thus, this review highlights that more rigorous studies are needed to yield robust and generalizable information about this potential pathway. Moreover, the evidence on mechanisms of this pathway summarized in this review suggests that thyroid dysfunction could mediate a relationship between prenatal PFAS exposure and child ASD, and this potential mediation effect could help explain significant findings from only three of the seven studies on an association between PFAS exposure and child ASD [96-98]. Therefore, future studies need to carefully disentangle the relationships among all potential mechanisms through mediation analysis [116-118] to help explain the underlying mechanism of any relationship between PFAS exposure and child ASD.

CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

ACKNOWLEDGMENTS

This work was supported by the grant from the National Institute of Environmental Health Sciences (R21-ES033389).

ORCID

Hyeong-Moo Shin <https://orcid.org/0000-0002-7001-2555>

REFERENCES

1. American Psychiatric Association. Diagnostic and statistical manual of mental disorders: DSM-5. Arlington: American Psychiatric Association; 2013.
2. Rogge N, Janssen J. The economic costs of autism spectrum disorder: a literature review. *J Autism Dev Disord* 2019;49:2873-900.
3. Leigh JP, Du J. Brief report: forecasting the economic burden of autism in 2015 and 2025 in the United States. *J Autism Dev Disord* 2015;45:4135-9.
4. Myers SM, Voigt RG, Colligan RC, Weaver AL, Storlie CB, Stoeckel RE, et al. Autism spectrum disorder: incidence and time trends over two decades in a population-based birth cohort. *J Autism Dev Disord* 2019;49:1455-74.
5. Maenner MJ, Shaw KA, Bakian AV, Bilder DA, Durkin MS, Esler A, et al. Prevalence and characteristics of autism spectrum disorder among children aged 8 years: autism and developmental disabilities monitoring network, 11 sites, United States, 2018. *MMWR Surveill Summ* 2021;70:1-16.
6. Grinker RR, Leventhal BL. Estimating the incidence of autism. *Epidemiology* 2009;20:622-3.
7. Hertz-Picciotto I, Delwiche L. The rise in autism and the role of age at diagnosis. *Epidemiology* 2009;20:84-90.
8. King M, Bearman P. Diagnostic change and the increased prevalence of autism. *Int J Epidemiol* 2009;38:1224-34.
9. Hallmayer J, Cleveland S, Torres A, Phillips J, Cohen B, Torigoe T, et al. Genetic heritability and shared environmental factors among twin pairs with autism. *Arch Gen Psychiatry* 2011;68:1095-102.
10. Sandin S, Lichtenstein P, Kuja-Halkola R, Larsson H, Hultman CM, Reichenberg A. The familial risk of autism. *JAMA* 2014;311:1770-7.
11. Roberts EM, English PB, Grether JK, Windham GC, Somberg L, Wolff C. Maternal residence near agricultural pesticide applications and autism spectrum disorders among children in the California central valley. *Environ Health Perspect* 2007;115:1482-9.
12. Shelton JF, Geraghty EM, Tancredi DJ, Delwiche LD, Schmidt RJ, Ritz B, et al. Neurodevelopmental disorders and prenatal residential proximity to agricultural pesticides: the CHARGE study. *Environ Health Perspect* 2014;122:1103-9.
13. Volk HE, Hertz-Picciotto I, Delwiche L, Lurmann F, McConnell R. Residential proximity to freeways and autism in the CHARGE study. *Environ Health Perspect* 2011;119:

- 873-7.
14. Volk HE, Lurmann F, Penfold B, Hertz-Picciotto I, McConnell R. Traffic-related air pollution, particulate matter, and autism. *JAMA Psychiatry* 2013;70:71-7.
 15. Brucato M, Ladd-Acosta C, Li M, Caruso D, Hong X, Kaczaniuk J, et al. Prenatal exposure to fever is associated with autism spectrum disorder in the Boston Birth Cohort. *Autism Res* 2017;10:1878-90.
 16. Croen LA, Qian Y, Ashwood P, Zerbo O, Schendel D, Pinto-Martin J, et al. Infection and fever in pregnancy and autism spectrum disorders: findings from the study to explore early development. *Autism Res* 2019;12:1551-61.
 17. Schmidt RJ. Maternal folic acid supplements associated with reduced autism risk in the child. *Evid Based Med* 2013;18:e53.
 18. Schmidt RJ, Hansen RL, Hartiala J, Allayee H, Schmidt LC, Tancredi DJ, et al. Prenatal vitamins, one-carbon metabolism gene variants, and risk for autism. *Epidemiology* 2011;22:476-85.
 19. Schmidt RJ, Tancredi DJ, Ozonoff S, Hansen RL, Hartiala J, Allayee H, et al. Maternal periconceptional folic acid intake and risk of autism spectrum disorders and developmental delay in the CHARGE (Childhood Autism Risks from Genetics and Environment) case-control study. *Am J Clin Nutr* 2012;96:80-9.
 20. Krakowiak P, Walker CK, Bremer AA, Baker AS, Ozonoff S, Hansen RL, et al. Maternal metabolic conditions and risk for autism and other neurodevelopmental disorders. *Pediatrics* 2012;129:e1121-8.
 21. Li M, Fallin MD, Riley A, Landa R, Walker SO, Silverstein M, et al. The association of maternal obesity and diabetes with autism and other developmental disabilities. *Pediatrics* 2016;137:e20152206.
 22. Mann JR, McDermott S, Bao H, Hardin J, Gregg A. Pre-eclampsia, birth weight, and autism spectrum disorders. *J Autism Dev Disord* 2010;40:548-54.
 23. Cheslack-Postava K, Liu K, Bearman PS. Closely spaced pregnancies are associated with increased odds of autism in California sibling births. *Pediatrics* 2011;127:246-53.
 24. Gunnes N, Suren P, Bresnahan M, Hornig M, Lie KK, Lipkin WI, et al. Interpregnancy interval and risk of autistic disorder. *Epidemiology* 2013;24:906-12.
 25. Zerbo O, Yoshida C, Gunderson EP, Dorward K, Croen LA. Interpregnancy interval and risk of autism spectrum disorders. *Pediatrics* 2015;136:651-7.
 26. Feingold KR, Anawalt B, Boyce A, Chrousos G, de Herder WW, Dhatariya K, et al. *Endotext*. South Dartmouth: MD-Text.com Inc.; 2000. Chapter, Thyroid hormones in brain development and function [cited 2022 Oct 27]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK285549>.
 27. Rovet JF. The role of thyroid hormones for brain development and cognitive function. *Endocr Dev* 2014;26:26-43.
 28. Bernal J. Thyroid hormones and brain development. *Vitam Horm* 2005;71:95-122.
 29. Lavado-Autric R, Auso E, Garcia-Velasco JV, Arufe Mdel C, Escobar del Rey F, Berbel P, et al. Early maternal hypothyroxinemia alters histogenesis and cerebral cortex cytoarchitecture of the progeny. *J Clin Invest* 2003;111:1073-82.
 30. Stepien BK, Huttner WB. Transport, metabolism, and function of thyroid hormones in the developing mammalian brain. *Front Endocrinol (Lausanne)* 2019;10:209.
 31. Ge GM, Leung MT, Man KK, Leung WC, Ip P, Li GH, et al. Maternal thyroid dysfunction during pregnancy and the risk of adverse outcomes in the offspring: a systematic review and meta-analysis. *J Clin Endocrinol Metab* 2020;105:dga555.
 32. Brown AS, Surcel HM, Hinkka-Yli-Salomaki S, Cheslack-Postava K, Bao Y, Sourander A. Maternal thyroid autoantibody and elevated risk of autism in a national birth cohort. *Prog Neuropsychopharmacol Biol Psychiatry* 2015;57:86-92.
 33. UN Environment Programme. *Global chemicals outlook: towards sound management of chemicals*. Nairobi: UNEP; 2012. p. 44.
 34. Shin HM, Moschet C, Young TM, Bennett DH. Measured concentrations of consumer product chemicals in California house dust: implications for sources, exposure, and toxicity potential. *Indoor Air* 2020;30:60-75.
 35. Diamanti-Kandarakis E, Bourguignon JP, Giudice LC, Hauser R, Prins GS, Soto AM. Endocrine-disrupting chemicals: an endocrine society scientific statement. *Endocr Rev* 2009;30:293-342.
 36. Mughal BB, Fini JB, Demeneix BA. Thyroid-disrupting chemicals and brain development: an update. *Endocr Connect* 2018;7:R160-86.
 37. La Merrill MA, Vandenberg LN, Smith MT, Goodson W, Browne P, Patisaul HB, et al. Consensus on the key characteristics of endocrine-disrupting chemicals as a basis for hazard identification. *Nat Rev Endocrinol* 2020;16:45-57.
 38. Brucker-Davis F. Effects of environmental synthetic chemicals on thyroid function. *Thyroid* 1998;8:827-56.
 39. Howdeshell KL. A model of the development of the brain

- as a construct of the thyroid system. *Environ Health Perspect* 2002;110 Suppl 3:337-48.
40. Talsness CE. Overview of toxicological aspects of polybrominated diphenyl ethers: a flame-retardant additive in several consumer products. *Environ Res* 2008;108:158-67.
 41. Moriyama K, Tagami T, Akamizu T, Usui T, Saijo M, Kanamoto N, et al. Thyroid hormone action is disrupted by bisphenol A as an antagonist. *J Clin Endocrinol Metab* 2002;87:5185-90.
 42. Fritsche E, Cline JE, Nguyen NH, Scanlan TS, Abel J. Polychlorinated biphenyls disturb differentiation of normal human neural progenitor cells: clue for involvement of thyroid hormone receptors. *Environ Health Perspect* 2005;113:871-6.
 43. Boas M, Feldt-Rasmussen U, Skakkebaek NE, Main KM. Environmental chemicals and thyroid function. *Eur J Endocrinol* 2006;154:599-611.
 44. Prevedouros K, Cousins IT, Buck RC, Korzeniowski SH. Sources, fate and transport of perfluorocarboxylates. *Environ Sci Technol* 2006;40:32-44.
 45. Centers for Disease Control and Prevention. Fourth national report on human exposure to environmental chemicals. Atlanta: CDC; 2015 [cited 2022 Oct 28]. Available from: https://www.cdc.gov/biomonitoring/pdf/fourthreport_updatedtables_feb2015.pdf.
 46. Fromme H, Mosch C, Morovitz M, Alba-Alejandre I, Boehmer S, Kiranoglu M, et al. Pre- and postnatal exposure to perfluorinated compounds (PFCs). *Environ Sci Technol* 2010;44:7123-9.
 47. Monroy R, Morrison K, Teo K, Atkinson S, Kubwabo C, Stewart B, et al. Serum levels of perfluoroalkyl compounds in human maternal and umbilical cord blood samples. *Environ Res* 2008;108:56-62.
 48. Jensen MS, Norgaard-Pedersen B, Toft G, Hougaard DM, Bonde JP, Cohen A, et al. Phthalates and perfluorooctanesulfonic acid in human amniotic fluid: temporal trends and timing of amniocentesis in pregnancy. *Environ Health Perspect* 2012;120:897-903.
 49. Long M, Ghisari M, Kjeldsen L, Wielsoe M, Norgaard-Pedersen B, Mortensen EL, et al. Autism spectrum disorders, endocrine disrupting compounds, and heavy metals in amniotic fluid: a case-control study. *Mol Autism* 2019;10:1.
 50. Boas M, Feldt-Rasmussen U, Main KM. Thyroid effects of endocrine disrupting chemicals. *Mol Cell Endocrinol* 2012;355:240-8.
 51. Torres L, Redko A, Limper C, Imbiakha B, Chang S, August A. Effect of perfluorooctanesulfonic acid (PFOS) on immune cell development and function in mice. *Immunol Lett* 2021;233:31-41.
 52. Guillette TC, McCord J, Guillette M, Polera ME, Rachels KT, Morgeson C, et al. Elevated levels of per- and polyfluoroalkyl substances in cape fear river striped bass (*Morone saxatilis*) are associated with biomarkers of altered immune and liver function. *Environ Int* 2020;136:105358.
 53. Chang ET, Adami HO, Boffetta P, Wedner HJ, Mandel JS. A critical review of perfluorooctanoate and perfluorooctanesulfonate exposure and immunological health conditions in humans. *Crit Rev Toxicol* 2016;46:279-331.
 54. Osuna CE, Grandjean P, Weihe P, El-Fawal HA. Autoantibodies associated with prenatal and childhood exposure to environmental chemicals in Faroese children. *Toxicol Sci* 2014;142:158-66.
 55. Coperchini F, Awwad O, Rotondi M, Santini F, Imbriani M, Chiovato L. Thyroid disruption by perfluorooctane sulfonate (PFOS) and perfluorooctanoate (PFOA). *J Endocrinol Invest* 2017;40:105-21.
 56. Hoffman K, Webster TF, Weisskopf MG, Weinberg J, Vieira VM. Exposure to polyfluoroalkyl chemicals and attention deficit/hyperactivity disorder in U.S. children 12-15 years of age. *Environ Health Perspect* 2010;118:1762-7.
 57. Stein CR, Savitz DA. Serum perfluorinated compound concentration and attention deficit/hyperactivity disorder in children 5-18 years of age. *Environ Health Perspect* 2011;119:1466-71.
 58. Ushijima J, Furukawa S, Sameshima H. The presence of thyroid peroxidase antibody is associated with lower placental weight in maternal thyroid dysfunction. *Tohoku J Exp Med* 2019;249:231-6.
 59. Iddah MA, Macharia BN. Autoimmune thyroid disorders. *ISRN Endocrinol* 2013;2013:509764.
 60. Alexander EK, Pearce EN, Brent GA, Brown RS, Chen H, Dosiou C, et al. 2017 Guidelines of the American Thyroid Association for the diagnosis and management of thyroid disease during pregnancy and the postpartum. *Thyroid* 2017;27:315-89.
 61. Andersen SL, Laurberg P, Wu CS, Olsen J. Attention deficit hyperactivity disorder and autism spectrum disorder in children born to mothers with thyroid dysfunction: a Danish nationwide cohort study. *BJOG* 2014;121:1365-74.
 62. Rotem RS, Chodick G, Shalev V, Davidovitch M, Koren G, Hauser R, et al. Maternal thyroid disorders and risk of autism spectrum disorder in progeny. *Epidemiology* 2020;31:

- 409-17.
63. Getahun D, Jacobsen SJ, Fassett MJ, Wing DA, Xiang AH, Chiu VY, et al. Association between maternal hypothyroidism and autism spectrum disorders in children. *Pediatr Res* 2018;83:580-8.
 64. Roman GC, Ghassabian A, Bongers-Schokking JJ, Jaddoe VW, Hofman A, de Rijke YB, et al. Association of gestational maternal hypothyroxinemia and increased autism risk. *Ann Neurol* 2013;74:733-42.
 65. Levie D, Korevaar TI, Bath SC, Dalmau-Bueno A, Murcia M, Espada M, et al. Thyroid function in early pregnancy, child IQ, and autistic traits: a meta-analysis of individual participant data. *J Clin Endocrinol Metab* 2018;103:2967-79.
 66. Andersen SL, Andersen S, Vestergaard P, Olsen J. Maternal thyroid function in early pregnancy and child neurodevelopmental disorders: a Danish nationwide case-cohort study. *Thyroid* 2018;28:537-46.
 67. Hoshiko S, Grether JK, Windham GC, Smith D, Fessel K. Are thyroid hormone concentrations at birth associated with subsequent autism diagnosis? *Autism Res* 2011;4:456-63.
 68. Lyall K, Anderson M, Kharrazi M, Windham GC. Neonatal thyroid hormone levels in association with autism spectrum disorder. *Autism Res* 2017;10:585-92.
 69. Korevaar TI, Tiemeier H, Peeters RP. Clinical associations of maternal thyroid function with foetal brain development: epidemiological interpretation and overview of available evidence. *Clin Endocrinol (Oxf)* 2018;89:129-38.
 70. Pearce EN, Lazarus JH, Moreno-Reyes R, Zimmermann MB. Consequences of iodine deficiency and excess in pregnant women: an overview of current knowns and unknowns. *Am J Clin Nutr* 2016;104 Suppl 3:918S-23S.
 71. Tingi E, Syed AA, Kyriacou A, Mastorakos G, Kyriacou A. Benign thyroid disease in pregnancy: a state of the art review. *J Clin Transl Endocrinol* 2016;6:37-49.
 72. Consortium on Thyroid and Pregnancy-Study Group on Preterm Birth, Korevaar TI, Derakhshan A, Taylor PN, Meima M, Chen L, et al. Association of thyroid function test abnormalities and thyroid autoimmunity with preterm birth: a systematic review and meta-analysis. *JAMA* 2019;322:632-41.
 73. Derakhshan A, Peeters RP, Taylor PN, Bliddal S, Carty DM, Meems M, et al. Association of maternal thyroid function with birthweight: a systematic review and individual-participant data meta-analysis. *Lancet Diabetes Endocrinol* 2020;8:501-10.
 74. Harel-Gadassi A, Friedlander E, Yaari M, Bar-Oz B, Even-tov-Friedman S, Mankuta D, et al. Risk for ASD in preterm infants: a three-year follow-up study. *Autism Res Treat* 2018;2018:8316212.
 75. Buchmayer S, Johansson S, Johansson A, Hultman CM, Sparen P, Cnattingius S. Can association between preterm birth and autism be explained by maternal or neonatal morbidity? *Pediatrics* 2009;124:e817-25.
 76. Burstyn I, Sithole F, Zwaigenbaum L. Autism spectrum disorders, maternal characteristics and obstetric complications among singletons born in Alberta, Canada. *Chronic Dis Can* 2010;30:125-34.
 77. Lampi KM, Lehtonen L, Tran PL, Suominen A, Lehti V, Banerjee PN, et al. Risk of autism spectrum disorders in low birth weight and small for gestational age infants. *J Pediatr* 2012;161:830-6.
 78. Hughes HK, Mills Ko E, Rose D, Ashwood P. Immune dysfunction and autoimmunity as pathological mechanisms in autism spectrum disorders. *Front Cell Neurosci* 2018;12:405.
 79. Comi AM, Zimmerman AW, Frye VH, Law PA, Peeden JN. Familial clustering of autoimmune disorders and evaluation of medical risk factors in autism. *J Child Neurol* 1999;14:388-94.
 80. Spann MN, Timonen-Soivio L, Suominen A, Cheslack-Postava K, McKeague IW, Sourander A, et al. Proband and familial autoimmune diseases are associated with proband diagnosis of autism spectrum disorders. *J Am Acad Child Adolesc Psychiatry* 2019;58:496-505.
 81. Chen SW, Zhong XS, Jiang LN, Zheng XY, Xiong YQ, Ma SJ, et al. Maternal autoimmune diseases and the risk of autism spectrum disorders in offspring: a systematic review and meta-analysis. *Behav Brain Res* 2016;296:61-9.
 82. Braunschweig D, Krakowiak P, Duncanson P, Boyce R, Hansen RL, Ashwood P, et al. Autism-specific maternal autoantibodies recognize critical proteins in developing brain. *Transl Psychiatry* 2013;3:e277.
 83. Braunschweig D, Ashwood P, Krakowiak P, Hertz-Picciotto I, Hansen R, Croen LA, et al. Autism: maternally derived antibodies specific for fetal brain proteins. *Neurotoxicology* 2008;29:226-31.
 84. Braunschweig D, Duncanson P, Boyce R, Hansen R, Ashwood P, Pessah IN, et al. Behavioral correlates of maternal antibody status among children with autism. *J Autism Dev Disord* 2012;42:1435-45.
 85. Rossi CC, Fuentes J, Van de Water J, Amaral DG. Brief report: antibodies reacting to brain tissue in Basque Spanish

- children with autism spectrum disorder and their mothers. *J Autism Dev Disord* 2014;44:459-65.
86. Zimmerman AW, Connors SL, Matteson KJ, Lee LC, Singer HS, Castaneda JA, et al. Maternal antibrain antibodies in autism. *Brain Behav Immun* 2007;21:351-7.
87. Singer HS, Morris CM, Gause CD, Gillin PK, Crawford S, Zimmerman AW. Antibodies against fetal brain in sera of mothers with autistic children. *J Neuroimmunol* 2008;194:165-72.
88. Croen LA, Braunschweig D, Haapanen L, Yoshida CK, Fireman B, Grether JK, et al. Maternal mid-pregnancy autoantibodies to fetal brain protein: the early markers for autism study. *Biol Psychiatry* 2008;64:583-8.
89. Heuer L, Ashwood P, Schauer J, Goines P, Krakowiak P, Hertz-Picciotto I, et al. Reduced levels of immunoglobulin in children with autism correlates with behavioral symptoms. *Autism Res* 2008;1:275-83.
90. Grether JK, Ashwood P, Van de Water J, Yolken RH, Anderson MC, Torres AR, et al. Prenatal and newborn immunoglobulin levels from mother-child pairs and risk of autism spectrum disorders. *Front Neurosci* 2016;10:218.
91. Korevaar TI, Steegers EA, Pop VJ, Broeren MA, Chaker L, de Rijke YB, et al. Thyroid autoimmunity impairs the thyroidal response to human chorionic gonadotropin: two population-based prospective cohort studies. *J Clin Endocrinol Metab* 2017;102:69-77.
92. Osinga JA, Derakhshan A, Palomaki GE, Ashoor G, Manisto T, Maraka S, et al. TSH and FT4 reference intervals in pregnancy: a systematic review and individual participant data meta-analysis. *J Clin Endocrinol Metab* 2022;107:2925-33.
93. Braun JM, Kalkbrenner AE, Just AC, Yolton K, Calafat AM, Sjodin A, et al. Gestational exposure to endocrine-disrupting chemicals and reciprocal social, repetitive, and stereotypic behaviors in 4- and 5-year-old children: the HOME study. *Environ Health Perspect* 2014;122:513-20.
94. Liew Z, Ritz B, von Ehrenstein OS, Bech BH, Nohr EA, Fei C, et al. Attention deficit/hyperactivity disorder and childhood autism in association with prenatal exposure to perfluoroalkyl substances: a nested case-control study in the Danish national birth cohort. *Environ Health Perspect* 2015;123:367-73.
95. Lyall K, Yau VM, Hansen R, Kharrazi M, Yoshida CK, Calafat AM, et al. Prenatal maternal serum concentrations of per- and polyfluoroalkyl substances in association with autism spectrum disorder and intellectual disability. *Environ Health Perspect* 2018;126:017001.
96. Shin HM, Bennett DH, Calafat AM, Tancredi D, Hertz-Picciotto I. Modeled prenatal exposure to per- and polyfluoroalkyl substances in association with child autism spectrum disorder: a case-control study. *Environ Res* 2020;186:109514.
97. Oh J, Bennett DH, Calafat AM, Tancredi D, Roa DL, Schmidt RJ, et al. Prenatal exposure to per- and polyfluoroalkyl substances in association with autism spectrum disorder in the MARBLES study. *Environ Int* 2021;147:106328.
98. Skogheim TS, Weyde KV, Aase H, Engel SM, Suren P, Oie MG, et al. Prenatal exposure to per- and polyfluoroalkyl substances (PFAS) and associations with attention-deficit/hyperactivity disorder and autism spectrum disorder in children. *Environ Res* 2021;202:111692.
99. Kim S, Choi K, Ji K, Seo J, Kho Y, Park J, et al. Trans-placental transfer of thirteen perfluorinated compounds and relations with fetal thyroid hormones. *Environ Sci Technol* 2011;45:7465-72.
100. de Cock M, de Boer MR, Lamoree M, Legler J, van de Bor M. Prenatal exposure to endocrine disrupting chemicals in relation to thyroid hormone levels in infants: a Dutch prospective cohort study. *Environ Health* 2014;13:106.
101. Wang Y, Rogan WJ, Chen PC, Lien GW, Chen HY, Tseng YC, et al. Association between maternal serum perfluoroalkyl substances during pregnancy and maternal and cord thyroid hormones: Taiwan maternal and infant cohort study. *Environ Health Perspect* 2014;122:529-34.
102. Berg V, Nost TH, Hansen S, Elverland A, Veyhe AS, Jorde R, et al. Assessing the relationship between perfluoroalkyl substances, thyroid hormones and binding proteins in pregnant women; a longitudinal mixed effects approach. *Environ Int* 2015;77:63-9.
103. Shah-Kulkarni S, Kim BM, Hong YC, Kim HS, Kwon EJ, Park H, et al. Prenatal exposure to perfluorinated compounds affects thyroid hormone levels in newborn girls. *Environ Int* 2016;94:607-13.
104. Berg V, Nost TH, Pettersen RD, Hansen S, Veyhe AS, Jorde R, et al. Persistent organic pollutants and the association with maternal and infant thyroid homeostasis: a multipollutant assessment. *Environ Health Perspect* 2017;125:127-33.
105. Preston EV, Webster TF, Oken E, Claus Henn B, McClean MD, Rifas-Shiman SL, et al. Maternal plasma per- and polyfluoroalkyl substance concentrations in early pregnancy and maternal and neonatal thyroid function in a prospec-

- tive birth cohort: Project Viva (USA). *Environ Health Perspect* 2018;126:027013.
106. Itoh S, Araki A, Miyashita C, Yamazaki K, Goudarzi H, Minatoya M, et al. Association between perfluoroalkyl substance exposure and thyroid hormone/thyroid antibody levels in maternal and cord blood: the Hokkaido study. *Environ Int* 2019;133(Pt A):105139.
107. Preston EV, Webster TF, Claus Henn B, McClean MD, Gennings C, Oken E, et al. Prenatal exposure to per- and polyfluoroalkyl substances and maternal and neonatal thyroid function in the project viva cohort: a mixtures approach. *Environ Int* 2020;139:105728.
108. Lebeaux RM, Doherty BT, Gallagher LG, Zoeller RT, Hoofnagle AN, Calafat AM, et al. Maternal serum perfluoroalkyl substance mixtures and thyroid hormone concentrations in maternal and cord sera: the HOME study. *Environ Res* 2020;185:109395.
109. Liang H, Wang Z, Miao M, Tian Y, Zhou Y, Wen S, et al. Prenatal exposure to perfluoroalkyl substances and thyroid hormone concentrations in cord plasma in a Chinese birth cohort. *Environ Health* 2020;19:127.
110. Guo J, Zhang J, Wang Z, Zhang L, Qi X, Zhang Y, et al. Umbilical cord serum perfluoroalkyl substance mixtures in relation to thyroid function of newborns: findings from sheyang mini birth cohort study. *Chemosphere* 2021;273:129664.
111. Kim DH, Kim UJ, Kim HY, Choi SD, Oh JE. Perfluoroalkyl substances in serum from South Korean infants with congenital hypothyroidism and healthy infants: its relationship with thyroid hormones. *Environ Res* 2016;147:399-404.
112. Webster GM, Venners SA, Mattman A, Martin JW. Associations between perfluoroalkyl acids (PFASs) and maternal thyroid hormones in early pregnancy: a population-based cohort study. *Environ Res* 2014;133:338-47.
113. Webster GM, Rauch SA, Marie NS, Mattman A, Lanphear BP, Venners SA. Cross-sectional associations of serum perfluoroalkyl acids and thyroid hormones in U.S. adults: variation according to TPOAB and iodine status (NHANES 2007-2008). *Environ Health Perspect* 2016;124:935-42.
114. de Escobar GM, Obregon MJ, del Rey FE. Iodine deficiency and brain development in the first half of pregnancy. *Public Health Nutr* 2007;10:1554-70.
115. Bath SC, Steer CD, Golding J, Emmett P, Rayman MP. Effect of inadequate iodine status in UK pregnant women on cognitive outcomes in their children: results from the Avon Longitudinal Study of Parents and Children (ALSPAC). *Lancet* 2013;382:331-7.
116. Valeri L, Vanderweele TJ. Mediation analysis allowing for exposure-mediator interactions and causal interpretation: theoretical assumptions and implementation with SAS and SPSS macros. *Psychol Methods* 2013;18:137-50.
117. Shrout PE, Bolger N. Mediation in experimental and non-experimental studies: new procedures and recommendations. *Psychol Methods* 2002;7:422-45.
118. Hayes AF. Introduction to mediation, moderation, and conditional process analysis: a regression-based approach. New York: The Guilford Press; 2013.