

Effect of *Chlorella vulgaris* intake on cadmium detoxification in rats fed cadmium*

You Jin Kim, Sanghee Kwon and Mi Kyung Kim[§]

Department of Nutritional Science and Food Management, Ewha Womans University, 11-1 Daehyeon-dong, Seodaemun-gu, Seoul 120-750, Korea

Received February 20, 2009; Revised April 13, 2009; Accepted April 29, 2009

Abstract

The aim of this study was to investigate if dietary *Chlorella vulgaris* (chlorella) intake would be effective on cadmium (Cd) detoxification in rats fed dietary Cd. Fourteen-week old male Sprague-Dawley (SD) rats weighing 415.0 ± 1.6 g were randomly divided into two groups and fed slightly modified American Institute of Nutrition-93 Growing (AIN-93G) diet without (n=10) or with (n=40) dietary Cd (200 ppm) for 8 weeks. To confirm alteration by dietary Cd intake, twenty rats fed AIN-93G diet without (n=10) and with (n=10) dietary Cd were sacrificed and compared. Other thirty rats were randomly blocked into three groups and fed slightly modified AIN-93G diets replacing 0 (n=10), 5 (n=10) or 10% (n=10) chlorella of total kg diet for 4 weeks. Daily food intake, body weight change, body weight gain/calorie intake, organ weight (liver, spleen, and kidney), perirenal fat pad and epididymal fat pad weights were measured. To examine Cd detoxification, urinary Cd excretion and metallothionein (MT) concentrations in kidney and intestine were measured. Food intake, calorie intake, body weight change, body weight gain/calorie intake, organ weight and fat pad weights were decreased by dietary Cd intake. Urinary Cd excretion and MT concentrations in kidney and small intestine were increased by dietary Cd. After given Cd containing diet, food intake, calorie intake, body weight change, body weight gain/calorie intake, organ weights and fat pad weights were not influenced by dietary chlorella intake. Renal MT synthesis tended to be higher in a dose-dependent manner, but not significantly. And chlorella intake did not significantly facilitate renal and intestinal MT synthesis and urinary Cd excretion. These findings suggest that, after stopping cadmium supply, chlorella supplementation, regardless of its percentage, might not improve cadmium detoxification from the body in growing rats.

Key Words: *Chlorella vulgaris*, cadmium excretion, metallothionein, Sprague-Dawley rats

Introduction

Cadmium (Cd) is a highly toxic environmental heavy metal and classified by International Agency for Research on Cancer of USA (IARC) as a human carcinogen (IRAC, 1993). Occurrence of toxic effects by Cd depends on the route, the concentration and the exposure time (Lopez *et al.*, 2006). Occupational exposure to cadmium, such as working with cadmium containing pigments, plastics, glass, metal alloys and electrode material in nickel-cadmium batteries, and non-occupational exposures such as food, water and cigarette smoke induce uptake of Cd from the environment into the body through pulmonary and enteral pathways (Waisberg *et al.*, 2003). Several population studies showed that not only occupationally exposed workers but also environmentally exposed populations can experience moderate to severe health problems due to Cd toxicity (Akesson *et al.*, 2006; Jarup, 2002; Nordberg, 2004; Sandy *et al.*, 2007; WHO, 2000). The detailed mechanisms for the toxic effects of cadmium are still largely unknown. However, it is thought to involve interferences with antioxidant enzymes (Hussain *et al.*,

1987), alteration in thiol proteins (Belyaeva & Korotkov, 2003; Li *et al.*, 1993), inhibition of energy metabolism (Muller, 1986), alteration in DNA structure (Coogan *et al.*, 1992) and affectation of some enzyme activities (Casalino *et al.*, 2002; Jay *et al.*, 1991; Watjen *et al.*, 2001).

Chlorella vulgaris (CV) is unicellular marine algae. Chlorella contains about 55~67% protein, 1~4% chlorophyll, 9~18% dietary fiber, and an amount of minerals and vitamins (Kunimasa *et al.*, 1999). These algae are considered to be highly resistant to heavy metals including Cd. It can also chelate heavy metal ions such as Cd (Guzman *et al.*, 2001). These effects are thought to be mediated through metal ion-protein interactions of a variety of cellular targets. And in some cases, the metal ions-protein complex form is considered to participate in detoxification mechanisms (Goyer, 1983; Goyer, 1984; Yoshida *et al.*, 2006). Chlorella is widely used as a food supplement in the forms of tablets, capsules, extract liquid or a food additive in Japan (Merchant & Andre, 2001). Numerous experimental studies have demonstrated various pharmacological effects of CV and CV extract, which include improvement in promotion of dioxin

* This work was supported by the grants from the Ministry of Education and Human Resources Development for the second stage of Brain Korea 21 Project in 2007.

[§] Corresponding Author: Mi Kyung Kim, Tel. 82-2-3277-3087, Fax. 82-2-3277-2862, Email. mkk@ewha.ac.kr

excretion (Yoshida *et al.*, 2006), growth of cells, animals and plants (Han *et al.*, 2002), and growth and development of children (Lee & Lee, 2002).

In several studies (Kim *et al.*, 2003b; Kunimasa *et al.*, 1999; Morita *et al.*, 1999; Shim, 2008), when rats fed dietary chlorella simultaneously with cadmium or dioxin, chlorella intake might stimulate cadmium excretion in urine and feces, and decrease synthesis of metallothionein (MT) in liver, kidney and small intestine. However, these events mostly resulted from preventing the absorption of Cd by chlorella intake. When *Chlorella vulgaris* is exposed to heavy metals, metal binding MT-like proteins in the cell were induced and these proteins might be involved in the detoxification of heavy metals. (Huang *et al.*, 2009) In this study, therefore, it was wondered that after cadmium is absorbed, to the exclusion of excretion in feces, if dietary chlorella intake could have some effects on betterment of growth and Cd detoxification. The overall goal of this study was conducted to investigate whether dietary *Chlorella vulgaris* (chlorella) intake would be effective on growth and Cd detoxification in rats already stressed with 8-week, high-level Cd exposure.

Materials and Methods

Experimental animals and diets

Fifty 14-week-old male Sprague Dawley rats (CD (SD) IGS, Outbred, Charles River Origin; Jung-Ang Lab. Animal, Inc., Korea) were placed in individual stainless steel wire-mesh cages in a climate-controlled room. The room had a 12:12 hr light-dark cycle, a temperature of 22-24°C and a relative humidity of 45 ± 5%. The rats were fed a pellet diet (Samyang Co., Korea) for the first seven days (adaptation period). The rats weighed 415.03 ± 1.60 g (mean ± standard error) at the end of the adaptation period.

The rats were randomly divided into Cd-free group (n=10) and Cd group (n=40). They were fed the modified American Institute of Nutrition (AIN)-93G diet without or with 200 ppm cadmium (CdCl₂ 406 ppm, 0.04% of kg diet) for 8 weeks. For confirming cadmium-induced stress applied, all rats of the Cd free group and 10 rats of the Cd group were sacrificed. The remaining thirty rats in Cd group were then stratified according to body weight and randomly blocked into three treatment groups according to different levels of chlorella (0, 5, or 10% of kg diet, w/w) in Table 1 for the experimental period, which lasted for 4 weeks. The chlorella sample used in this study (*Chlorella vulgaris*) was obtained from Daesang Co. (Korea). The diets were formulated according to the nutrient content of the AIN-93G diet.

This study was conducted at the nutrition laboratory of Ewha Womans University, in compliance with the Guide for the Care and Use of Laboratory Animals. During the experimental period, the rats were allowed *ad libitum* access to the experimental diets and distilled water. For preventing contamination of other minerals,

Table 1. Composition of experimental diets (units:g/kg diet)

Ingredients	Groups	Cd +		Cd -	
		chlorella 0%	chlorella 0%	chlorella 5%	chlorella 10%
Com Starch		396.78	400.2	380.19	360.18
Dextrinized cornstarch		132	132	125.4	118.8
Sucrose		100	100	95	90
Casein (>85% protein)		200	200	190	180
Soybean oil		70	70	66.5	63
Fiber		50	50	47.5	45
Mineral mix ¹⁾		35	35	33.25	31.5
Vitamin mix ²⁾		10	10	9.5	9
Choline bitartrate		2.5	2.5	2.375	2.25
tert-Butyl hydroquinone		0.014	0.014	0.0133	0.0126
Chlorella powder		0	0	50	100
Cadmium Chloride		0.406	0	0	0
Total amount		1000	1000	1000	1000
Total calories (kcal)		3721.668	3733.98	3733.48	3732.98
Energy ratio (%)	Carbohydrate	62.0	62.1	59.2	56.3
	Protein	19.2	19.2	21.5	23.8
	Fat	16.9	16.9	17.6	18.3

¹⁾ Mineral mix (AIN-93G-MIX) (g/kg mixture): anhydrous calcium carbonate, 357; monobasic potassium phosphate, 196; sodium chloride, 74; potassium sulfate, 46.6; tripotassium citrate monohydrate, 70.78; magnesium oxide, 24; ferric citrate, 6.06; zinc carbonate, 1.65; manganous carbonate, 0.63; cupric carbonate, 0.3; potassium iodate, 0.01; anhydrous sodium selenate, 0.01025; ammoniumparamolybdate 4-hydrate, 0.00795; sodium metasilicate 9-hydrate, 1.45; chromium potassium sulfate 12-hydrate, 0.275; boric acid, 0.0815; sodium fluoride, 0.0635; nickel carbonate, 0.0318; lithium chloride, 0.0174; ammonium vanadate, 0.0066; powered sucrose 221.026

²⁾ Vitamin mix (AIN-93-VX) (g/kg mixture) : Niacin 3, Calcium Pantothenate 1.60, Pyridoxine HCl 0.70, Thiamine HCl 0.60, Riboflavin 0.60, Folic Acid 0.20, Biotin 0.02, Vitamin E Acetate (500 IU/g) 15, Vitamin B₁₂ (0.1%) 2.50, Vitamin A Palmitate (500,000 IU/g) 0.80, Vitamin D₃ (400,000 IU/g) 0.25, Vitamin K₁/Dextrose Mix (10 mg/g) 7.50, Sucrose 967.23

all instruments were treated with 0.4% ethylenediaminetetraacetic acid (EDTA) solution and washed with distilled deionized water.

Specimen collection and analytical method

Urine was collected in an acid-washed bottle treated with a few drops of toluene as preservatives using the metabolic cages during the final three days of experimental period. After collection, the bottle was filled up to 100 ml with deionized water and centrifuged at 7,000 rpm for 10 min (Supra 22K High speed centrifuge, Hanil, Korea). Aliquots of the samples were stored at -20°C until determination of Cd concentration. Before sacrificing, animals were deprived of food for 12 hrs and sacrificed after anesthetization with diethyl ether. The liver, kidney, spleen, intestine, epididymal fat pad and perirenal fat pad were removed and weighed. The kidney and the intestine were perfused with saline and stored at -80°C until analysis for MT determination.

Cd concentration in urine was measured by the method of Zinterhofer (Zinterhofer *et al.*, 1971). Urine was adjusted to pH 5.5. 10% Triton X-100, 2% ADPC (Ammonium pyrrolidinedithiocarbamate) and MIBK (4-Methyl-2-pentanone, methyl isobutyl ketone) were added to urine one after another. The

solutions were mixed well, and centrifuged at 3,500 rpm for 20 min at RT (Supra 22K high speed centrifuge, Hanil, Korea). MIBK layers were treated with nitric acid and determined using AAS (Atomic Absorption Spectrophotometer, Model 6701F, SHIMADU CO., Japan). The analyses were performed at the 228.8 nm resonance line.

Determination of MT was measured by the methods of cadmium/hemoglobin affinity assay (Eaton & Toal, 1982; Eaton & Cherian, 1991). One gram of small intestine or 0.5 g of kidney was homogenized in 1 vol. of 10 mM Tris-HCl buffer (pH 7.4, 4°C) and centrifuged at 10,000 rpm for 10 min at 4°C (Micro 17R, Micro Refrigerated Centrifuge, Hanil, Korea). The supernatant fraction was heated for 2 min in a boiling water bath. The heated samples were then centrifuged at 10,000 rpm for 2 min to remove precipitated proteins. The samples were mixed with 0.26 mM CdCl₂ solution and incubated at RT for 15 min. The metal binding sites of MT were saturated with Cd during this incubation step. For removal of excessive Cd, it was precipitated by addition of rat RBC hemolysate, which was prepared following the method of Onosaka and Cherian (Onosaka & Cherian, 1982), and heat treatment in a boiling water bath for 2min. The heat treated Cd-bound hemoglobin was removed by centrifugation at 10,000 rpm for 2 min at 4°C. The amount of Cd in the heated supernatant fraction is converted from the amount of MT-bound Cd and was determined in using AAS. The analyses were performed at the 228.8 nm resonance line. The concentration of MT in each tissue was calculated by assuming that 7 g-atom of Cd is bound to each mole of thionein which has a molecular weight of 6050 by amino acid analysis.

Statistical analysis

All statistical analyses were performed by the SAS program package version 9.1. All results were expressed as mean ± standard error (SE). The data between Cd fed and Cd non-fed group were analyzed by unpaired t-test. The data between chlorella fed and chlorella non-fed group was analyzed by one-way analysis of variance (ANOVA) and differences among experimental groups were evaluated using Duncan's multiple range tests at the P<0.05 significant level. All analyses were performed in duplicate or triplicate.

Results

Food intake, body weight change and energy efficiency ratio

Daily food intake, calorie intake, body weight change and body weight gain/calorie intake (energy efficiency ratio, EER) were shown in Table 2 and Table 5. Daily food intake, calorie intake, body weight change and EER in rats fed Cd-containing diet (Cd+ group) were significantly lower than those in rats fed Cd free diet (Cd- group). Daily food intake, calorie intake, body weight

Table 2. Food intake, calorie intake, body weight change and energy efficiency ratio in rats fed diets with different level of cadmium

Group ¹⁾	Food Intake (g/day)	Calorie Intake (cal/day)	Body weight change (g/8 week)	EER ⁴⁾
Cd-	24.09 ± 0.66 ²⁾³⁾	89.95 ± 2.45 ^a	169.81 ± 8.94 ^a	1.88 ± 0.05 ^a
Cd+	20.70 ± 1.15 ^b	77.01 ± 4.27 ^b	46.51 ± 6.35 ^b	0.60 ± 0.09 ^b

¹⁾ Cd- : Cadmium 0 ppm with 0% (w/w) chlorella powder

Cd+ : Cadmium 200 ppm with 0% (w/w) chlorella powder

²⁾ Mean ± standard error (S.E)

³⁾ Values within a column with different letters are significantly different at $\alpha=0,05$ level by unpaired t-test.

⁴⁾ Body weight gain (g/d) /calorie intake (kcal/d) means energy efficiency ratio (EER).

Table 3. Organ and tissue weights in rats fed diets with different level of cadmium (Unit : g/kg BW)

Group ¹⁾	Liver	Kidney	Spleen	Perirenal fat pad	Epididymal fat pad
Cd-	26.60 ± 0.49 ²⁾³⁾	5.25 ± 0.13 ^a	1.24 ± 0.06 ^a	20.47 ± 1.61 ^a	32.63 ± 1.83 ^a
Cd+	24.24 ± 0.57 ^b	5.85 ± 0.13 ^b	1.48 ± 0.04 ^b	8.64 ± 0.87 ^b	11.23 ± 1.33 ^b

¹⁾ See Table 2.

²⁾ Mean ± standard error (S.E)

³⁾ Values within a column with different letters are significantly different at $\alpha=0,05$ level by unpaired t-test.

change and EER in CV-5 group were tended to be higher than those in CV-0 and CV-10 groups, though not significantly.

Organ and tissue weights

Organ (liver, kidney and spleen) and tissue (epididymal and perirenal fat pad) weights per kg body weight are shown in Table 3 and Table 6. Weight of liver, kidney and spleen in rats fed Cd-containing diet was significantly lower than those in rats fed Cd free diet. Both perirenal and epididymal fat pad weights in Cd fed group were significantly lower than those of Cd- group. Organ (liver, kidney, and spleen) and epididymal fat pad weights in CV-5 group were tended to be lower than those in CV-0 and CV-10 groups. There was no observable difference in perirenal fat pad weight between chlorella untreated group(CV-0) and chlorella fed groups (CV-5 and CV-10).

Based on the above results, the rats fed Cd containing diet were exposed to some stress. However, in rats fed chlorella containing diet, food intake, recovery of suppressed body weight gain, and organ and tissue weights were not affected compared with those fed chlorella free diet.

Cadmium concentration in urine and metallothionein (MT) concentrations in kidney and small intestine

Urinary Cd concentration and MT concentrations in kidney and small intestine were shown in Table 4 and Table 7. Urinary Cd concentration in Cd fed group was significantly higher than that in Cd non-fed group. Renal and intestinal MT concentrations in the rats fed Cd containing diet were significantly higher than those in the rats fed Cd free diet, as well. Though not significantly, urinary Cd concentration in CV-5 group was tended to be lower than CV-0 and CV-10 groups. Renal MT

Table 4. Urinary cadmium concentration and metallothionein concentrations in kidney and small intestine in rats fed diets with different level of cadmium

Group ¹⁾	Urinary Cd ($\mu\text{g/day}$)	Renal MT ($\mu\text{g/g wet wt}$)	Intestinal MT ($\mu\text{g/g wet wt}$)
Cd-	0.12 \pm 0.07 ²⁾³⁾	2.56 \pm 0.52 ^b	0.56 \pm 0.11 ^b
Cd+	4.92 \pm 0.80 ^a	14.80 \pm 1.64 ^a	1.87 \pm 0.19 ^a

¹⁾ See Table 2.²⁾ Mean \pm standard error (S,E)³⁾ Values within a column with different letters are significantly different at $\alpha=0,05$ level by by unpaired t-test.**Table 5.** Food intake, calorie intake, body weight change and energy efficiency ratio in rats fed diets with different level of chlorella powder

Group ¹⁾	Food Intake (g/day)	Calorie Intake (cal/day)	Body weight change (g/4 week)	EER ⁴⁾
CV-0	25.47 \pm 0.83 ^{2)NS3)}	98.86 \pm 3.11 ^{NS}	80.46 \pm 5.33 ^{NS}	0.81 \pm 0.03 ^{NS}
CV-5	27.41 \pm 0.92	101.46 \pm 3.40	92.99 \pm 8.64	0.90 \pm 0.06
CV-10	26.71 \pm 0.84	98.01 \pm 3.08	82.37 \pm 7.46	0.82 \pm 0.06

¹⁾ CV-0: No cadmium with 0% (w/w) chlorella powder

CV-5: No cadmium with 5% (w/w) chlorella powder

CV-10: No cadmium with 10% (w/w) chlorella powder

²⁾ Mean \pm standard error (S,E)³⁾ Values within a column are not significant at $\alpha=0,05$ level by Duncan's multiple range test.⁴⁾ Body weight gain (g/d) /calorie intake (kcal/d) means energy efficiency ratio (EER).

concentration was tended to be higher in a dose-dependent manner in chlorella fed rats compared to chlorella untreated rats. Intestinal MT concentration in CV-5 group was higher than those in CV-0 and CV-10 groups.

There were no significant differences in urinary Cd and renal and intestinal MT concentrations among the groups (CV-0, CV-5 and CV-10 groups).

Discussion

Growth inhibition is the general symptom of Cd exposure (Choi & Rhee, 2001; Kim *et al.*, 1999; Shim, 2008; Yoon & Rhee, 1994). In the present study, the rats fed Cd-containing diet had significantly lower daily food intake, body weight gain and organ and tissue weights. These results were regarded that Cd decreases food intake by reducing appetite, calorie intake along with suppressed food intake, and body weight change and EER by affecting nutritional absorption and metabolism (Itokawa *et al.*, 1974).

Urinary Cd excretion and MT concentrations of kidney and small intestine were significantly increased with dietary Cd intake (Table 4). This result is consistent with other studies. In the previous study, compared with Cd non-fed group, urinary Cd excretion and MT concentrations in those organs were significantly increased in dietary 160 ppm Cd fed group (Shim, 2008). The other study showed that urinary Cd excretion and MT concentrations in kidney were significantly increased in rats exposed to 500 ppm Cd in drinking water for 20 weeks (Kim *et al.*, 2003a). MT content was increased in several tissues (liver,

Table 6. Organ and tissue weights in rats fed diets with different level of chlorella powder (Unit : g/kg BW)

Group ¹⁾	Liver	Kidney	Spleen	Perirenal fat pad	Epididymal fat pad
CV-0	28.35 \pm 0.82 ²⁾³⁾	5.87 \pm 0.15 ^{NS4)}	1.47 \pm 0.08 ^{NS}	22.01 \pm 0.93 ^{NS}	27.41 \pm 1.11 ^{NS}
CV-5	25.53 \pm 0.76 ^b	5.66 \pm 0.09	1.37 \pm 0.04	22.39 \pm 1.57	25.75 \pm 1.53
CV-10	27.21 \pm 0.43 ^{ab}	6.11 \pm 0.14	1.53 \pm 0.05	24.75 \pm 1.32	27.46 \pm 1.39

¹⁾ See Table 5.²⁾ Mean \pm standard error (S,E)³⁾ Values within a column with different letters are significantly different at $\alpha=0,05$ level by Duncan's multiple range test.⁴⁾ Values within a column are not significant at $\alpha=0,05$ level by Duncan's multiple range test.**Table 7.** Urinary cadmium concentration and metallothionein concentrations in kidney and small intestine in rats fed diets with different level of chlorella powder

Group ¹⁾	Urinary Cd ($\mu\text{g/day}$)	Renal MT ($\mu\text{g/g wet wt}$)	Intestinal MT ($\mu\text{g/g wet wt}$)
CV-0	0.32 \pm 0.09 ^{2)NS3)}	18.33 \pm 2.45 ^{NS}	2.18 \pm 0.38 ^{NS}
CV-5	0.16 \pm 0.05	18.94 \pm 2.61	3.41 \pm 0.67
CV-10	0.35 \pm 0.01	19.11 \pm 2.93	1.82 \pm 0.23

¹⁾ See Table 5.²⁾ Mean \pm standard error (S,E)³⁾ Values within a column are not significant at $\alpha=0,05$ level by Duncan's multiple range test.

kidney, heart and lung) in different species upon time-dependent Cd exposure (Bobillier-Chaumont *et al.*, 2006; Bonda *et al.*, 2004; Lange *et al.*, 2002).

Accumulation of Cd makes such a dangerous and minimizing effect on body functions leading to kidney toxicity (Shaikh *et al.*, 1999). Many works have been demonstrated that natural compounds or substances along with cadmium intake could facilitate cadmium excretion (Choi & Rhee, 2001; Kim *et al.*, 1999; Kim *et al.*, 2003a; Kim *et al.*, 2003b; Shim, 2008; Yoon & Rhee, 1994). These studies could not be convinced that chlorella supplementation after accumulation of absorbed Cd might also be effective in Cd detoxification.

Absorbed Cd is eliminated through MT from the organism mainly *via* urine. The amount of Cd excreted daily in urine is, however, very small: It represents only about 0.005-0.01% of the total body burden which corresponds to a biological half-life for Cd of about 20-40 years (WHO, 2000). MT has attracted attention as a protein possibly related to metabolism and detoxification of heavy metals because it has lower affinity to the essential metal zinc but high affinity to toxic heavy metals such as Cd and Hg (Kagi & Vellee, 1960). MT was noted first for its function as a heavy metal detoxicating protein (Brouwer *et al.*, 1993). Metal binding MT-like proteins in the *Chlorella vulgaris* cell were induced by exposing heavy metals and these proteins might be involved in the detoxification of those metals. (Huang *et al.*, 2009) This showed that chlorella intake have the possibility for enhancing synthesis of metal binding MT-like proteins by capturing absorbed heavy metals in the body. Accordingly, if chlorella intake facilitates urinary Cd excretion by the above mentioned mechanism, it would be helpful in the detoxification of Cd. Thus, the aim of this study was to evaluate

the effect of chlorella on cadmium detoxification after dietary Cd-induced stress in growing rats.

After stopping the feeding dietary Cd, daily food intake, calorie intake, body weight change, FER and organ weight were not affected by dietary chlorella intake. This outcome was similar to the results in several studies (Hwang, 2006; Shim, 2008) that there were no significant differences in food intake, body weight change, and organ and tissue weights among three experimental groups (chlorella non-fed group, and 5% or 10% chlorella fed group). This finding was thought that, although chlorella reported the growth stimulation in itself, cells, animals, plants, infant and child (Han *et al.*, 2002; Kim *et al.*, 2003b), After exposing dietary Cd, chlorella intake, regardless of its percentage, would not promote improvement of suppressed growth in rats.

Though not significant, renal MT concentration tended to rise dose-dependently. *In vivo* studies showed that Cd and high protein containing diet increased MT synthesis in rat organs (Kwon & Kim, 1992; Park & Kim, 1996; Revis, 1981). Since chlorella contains 55~67% protein and an amount of minerals (Morita *et al.*, 1999), even though Cd acted as an activator of MT synthesis during the cadmium feeding period, MT *de novo* synthesis would be promoted during chlorella intake after given Cd stimuli, although not significant.

The effect of dietary chlorella intake on intestinal MT concentration was not a dose-dependent manner. Gropper *et al.* (2004) reported that divalent metal ions share their transporters and excessively ingested divalent cations such as Cu, Fe and Zn impede their absorption with each other. For high concentrations of nutrients such as minerals interfere with the absorption of each other, excessive chlorella intake, in the case of the diet containing 10% chlorella, seemed to inhibit MT synthesis.

Urinary cadmium concentration was not affected by dietary chlorella intake. Several studies presented that simultaneous intake of Cd and chlorella facilitated urinary Cd excretion and this finding resulted from inhibition of absorption but not from facilitation of excretion (Kim *et al.*, 2003b; Shim, 2008). And a few studies presented that, when renal Cd-MT form was increased, urinary Cd excretion was facilitated (Friberg, 1984; Park & Kim, 1996). Because MT concentration in kidney was somewhat increased by chlorella intake, but not observably (Table 7), its concentration did not lead to increase urinary Cd excretion. Based on above explanation, if simultaneous intake of chlorella and Cd did not happen, Cd was already accumulated in the body and chlorella supplementation could not promote Cd excretion *via* urine.

On the other hand, the exact mechanism of detoxification by chlorella intake was not able to be known. Since concentrations of Cd in tissues of liver, kidney or small intestine were not measured, contribution of chlorella in MT synthesis could not be figured out. Hence, in a succeeding study, it has to additionally evaluate measurements of free Cd concentrations in blood and organs, MT-bound Cd concentrations in organs and fecal Cd excretion.

Taken together, it is thought over that, after given Cd exposure, non-simultaneous dietary chlorella intake was not likely to promote growth in animal and facilitate Cd excretion *via* urine through detoxification of Cd-MT like proteins.

Literature cited

- Akesson A, Bjellerup P, Lundh T, Lidfeldt J, Nerbrand C, Samsioe G, Skerfving S & Vahter M (2006). Cadmium-induced effects on bone in a population-based study of women. *Environ Health Perspect* 114:830-834.
- Belyaeva EA & Korotkov SM (2003). Mechanism of primary Cd²⁺-induced rat liver mitochondria dysfunction: discrete modes of Cd²⁺ action on calcium and thiol-dependent domains. *Toxicol Appl Pharmacol* 192:56-68.
- Bobillier-Chaumont S, Maupoil V & Berthelot A (2006). Metallothionein induction in the liver, kidney, heart and aorta of cadmium and isoproterenol treated rats. *J Appl Toxicol* 26:47-55.
- Bonda E, Wlostowski T & Krasowska A (2004). Testicular toxicity induced by dietary cadmium is associated with decreased testicular zinc and increased hepatic and renal metallothionein and zinc in the bank vole (*Clethrionomys glareolus*). *Biometals* 17:615-624.
- Brouwer M, Hoexum-Brouwer T & Cashon RE (1993). A putative glutathione-binding site in CdZn-metallothionein identified by equilibrium binding and molecular-modelling studies. *Biochem J* 294:219-225.
- Casalino E, Calzaretto G, Sblano C & Landriscina C (2002). Molecular inhibitory mechanism of antioxidant enzymes in rat liver and kidney by cadmium. *Toxicology* 179:37-50.
- Choi JH & Rhee SJ (2001). Effects of green tea catechin on cadmium accumulation in chronic cadmium poisoned rats. *The Korean Journal of Nutrition* 34:384-392.
- Coogan TP, Bare RM & Waalkes MP (1992). Cadmium-induced DNA strand damage in cultured liver cells: reduction in cadmium genotoxicity following zinc pretreatment. *Toxicol Appl Pharmacol* 113:227-233.
- Eaton DL & Cherian MG (1991). Determination of metallothionein in tissue by cadmium-hemoglobin affinity assay. *Methods Enzymol* 205:83-88.
- Eaton DL & Toal BF (1982). Evaluation of the Cd/Hb affinity assay for the rapid determination of metallothionein in biological tissue. *Toxicol Appl Pharmacol* 66:134-142.
- Friberg L (1984). Cadmium and the kidney. *Environ Health Perspect* 54:1-11.
- Goyer RA (1983). Intracellular sites of toxic metals. *Neurotoxicology* 4:147-156.
- Goyer RA (1984). Metal-protein complexes in detoxification process. In: Brown SS (Eds.), *Clinical Chemistry and Clinical Toxicology*, 2:199-209. Academic Press, London. UK
- Gropper SS, Smith JL & Groff JL (2004) The antioxidant nutrients, reactive species, and disease. In: *Advanced nutrition and human metabolism 4th edition*, p.368-377. Thomson Wadsworth, California. USA
- Guzman S, Gato A & Calleja JM (2001). Antiinflammatory, analgesic and free radical scavenging activities of the marine microalgae *Chlorella stigmatophora* and *Phaeodactylum tricornutum*. *Phytother Res* 15:224-230.
- Han JG, Kang GG, Kim JK & Kim SH (2002). The present status

- and future of Chlorella. *Food Science and Industry* 35:64-69.
- Huang Z, Li L, Huang G, Yan Q, Shi B & Xu X (2009). Growth-inhibitory and metal-binding proteins in *Chlorella vulgaris* exposed to cadmium or zinc. *Aquat Toxicol* 91:54-61.
- Hussain T, Shukla GS & Chandra SV (1987). Effects of cadmium on superoxide dismutase and lipid peroxidation in liver and kidney of growing rats: in vivo and in vitro studies. *Pharmacol Toxicol* 60:355-359.
- Hwang YK, Choi HJ, Nan M, Yoo JD & Kim YH (2006). Effect of Chlorella on metallothionein synthesis and binding capacity of cadmium in cadmium poisoned rat liver and kidney. *Journal of Experimental and Biomedical Sciences* 12:23-27.
- International Agency for Research on Cancer (IARC) (1993). International agency for research on cancer monographs on the evaluation of the carcinogenic risks to humans. Beryllium, cadmium, mercury, and exposures in the glass manufacturing industry. *Intl Agency Res Cancer* 58:119-237. ARC Scientific Publications, Lyon. France
- Itokawa Y, Abe T, Tabei R & Tanaka S (1974). Renal and skeletal lesions in experimental cadmium poisoning: Histological and biological approaches. *Arch Environ Health* 28:149-154.
- Jarup L (2002). Cadmium overload and toxicity. *Nephrol Dial Transplant* 17:35-39.
- Jay D, Zamorano R, Munoz E, Gleason R & Boldu JL (1991). Study of the interaction of cadmium with membrane-bound succinate dehydrogenase. *J Bioenerg Biomembr* 23:381-389.
- Kagi JHR & Vellee BL (1960). Metallothionein: a cadmium- and zinc-containing protein from equine renal cortex. *J Biol Chem* 235:3460-3465.
- Kim HJ, Bae KH, Lee HJ, Eun JB & Kim MK (1999). Effect of hesperidin extracted from tangerine peel on Cd and lipid metabolism, and antioxidative capacity in rats. *The Korean Journal of Nutrition* 32:137-149.
- Kim MJ, Hong JH & Rhee SJ (2003a). Effect of vitamin E on cadmium accumulation and excretion in chronic cadmium poisoned rats. *The Korean Journal of Nutrition* 36:691-698.
- Kim YH, Hwang YK, Lee YW, Yun JY, Hwang JM & Yoo JD (2003b). Effect of chlorella diet supplementation on blood and urine cadmium levels in cadmium poisoned rats. *J Biomed Lab Sci* 9:133-137.
- Kunimasa M, Takahiko M, Takao I & Takashi Hasegawa (1999). Chlorella accelerates dioxin excretion in rats. *J Nutr* 129:1731-1736.
- Kwon OR & Kim MK (1992). Effect of dietary protein and calcium levels on metallothionein and histopathological changes during cadmium intoxication in rats. *The Korean Journal of Nutrition* 25:360-378.
- Lange A, Aussenil O & Segner H (2002). Alterations of tissue glutathione levels and metallothionein mRNA in rainbow trout during single and combined exposure to cadmium and zinc. *Comp Biochem Physiol C Toxicol Pharmacol* 131:231-243.
- Lee YK & Lee HG (2002). Industrial uses of algae. *Korean Journal of Applied Microbiology and Bioengineering* 15:19-24.
- Li W, Zhao Y & Chou IN (1993). Alterations in cytoskeletal protein sulfhydryls and cellular glutathione in cultured cells exposed to cadmium and nickel ions. *Toxicology* 77:65-79.
- Lopez E, Arce C, Oset-Gasque MJ, Canadas S & Gonzalez MP (2006). Cadmium induces reactive oxygen species generation and lipid peroxidation in cortical neurons in culture. *Free Radic Biol Med* 40:940-951.
- Merchant RE & Andre CA (2001). A review of recent clinical trials of the nutritional supplement *Chlorella pyrenoidosa* in the treatment of fibromyalgia, hypertension, and ulcerative colitis. *Altern Ther Health Med* 7:79-91.
- Morita K, Matsueda T, Iida T & Hasegawa T (1999). Chlorella accelerates dioxin excretion in rats. *J Nutr* 129:1731-1736.
- Muller L (1986). Consequences of cadmium toxicity in rat hepatocytes: mitochondrial dysfunction and lipid peroxidation. *Toxicology* 40:285-295.
- Nordberg GF (2004). Cadmium and health in the 21st century-historical remarks and trends for the future. *Biometals* 17:485-489.
- Onosaka S & Cherian MG (1982). Comparison of metallothionein determination by polarographic and cadmium-saturation methods. *Toxicol Appl Pharmacol* 63:270-274.
- Park JY & Kim MK (1996). Effect of dietary protein and cysteine levels on cadmium toxicity in rats. *The Korean Journal of Nutrition* 29:261-471.
- Revis NW (1981). The relationship of dietary protein to metallothionein and cadmium-induced renal damage. *Toxicology* 20:323-333.
- Sandy T, Ann G, John M, Karen S, Nele H, Ivo L & Emmy VK (2007). Low cadmium exposure triggers a biphasic oxidative stress response in mice kidneys. *Toxicology* 236:29-41.
- Shaikh ZA, Vu TT & Zaman K (1999). Oxidative stress as a mechanism of chronic cadmium-induced hepatotoxicity and renal toxicity and protection by antioxidants. *Toxicol Appl Pharmacol* 154:256-263.
- Shim JA (2008). Effect of Chlorella intake on Cadmium metabolism in rats. Master thesis. Ewha Womans University, Seoul. Republic of Korea
- Waisberg M, Joseph P, Hale B & Beyersmann D (2003). Molecular and cellular mechanisms of cadmium carcinogenesis. *Toxicology* 192:95-117.
- Watjen W, Benters J, Haase H, Schwede F, Jastorff B & Beyersmann D (2001). Zn²⁺ and Cd²⁺ increase the cyclic GMP level in PC12 cells by inhibition of the cyclic nucleotide phosphodiesterase. *Toxicology* 157:167-175.
- World Health Organisation (WHO) (2000). Chapter 6.3 Cadmium. In: *Air Quality Guidelines - second edition*, p.1-11. World Health Organization Regional Office for Europe, Copenhagen. Denmark
- Yoon YH & Rhee SJ (1994). Effect of Korean green tea, oolong tea and black tea beverage on the antioxidative detoxification in rats poisoned with cadmium. *The Korean Journal of Nutrition* 27:1007-1017.
- Yoshida N, Ikeda R & Okuno T (2006). Identification and characterization of heavy metal-resistant unicellular algae isolated from soil and its potential for phytoremediation. *Bioresour Technol* 97:1843-1849.
- Zinterhofer LTM, Jotolow PT & Fappiano A (1971). Atomic absorption determination of lead in blood and urine in the presence of EDTA. *J Lab Clin Med* 78:664-674.