

RESEARCH

Open Access



# Prepregnancy body mass index, gestational weight gain, and maternal prepartum inflammation in normal pregnancies: findings from a Chinese cohort

Kuanrong Li<sup>1\*</sup>, Chuanzi Yang<sup>1</sup>, Jiaying Fan<sup>2</sup>, Xiaojun Li<sup>1</sup>, Chongjuan Gu<sup>2</sup> and Huishu Liu<sup>2\*</sup>

## Abstract

**Background:** Obesity has been linked to systemic inflammation in population studies.

**Objective:** To examine the associations of prepregnancy body mass index (pBMI) and total gestational weight gain (tGWG) with maternal prepartum low-grade inflammation (LGI) and clinically significant inflammation (CSI) defined by serum C-reactive protein (CRP) concentration.

**Methods:** Five thousand four hundred seventy-six Chinese women with uncomplicated pregnancies and recorded data on pBMI and prepartum body weight were included in this study. Blood samples were drawn before delivery for high-sensitivity CRP assay. Inadequate, optimal, and excessive tGWG were defined using the Institute of Medicine's recommendation. Multivariable Poisson regressions were used to estimate relative risks (RRs) for having prepartum LGI and CSI (defined as CRP concentration 3–10 and > 10 mg/L, respectively) across pBMI and tGWG categories.

**Results:** The mean pBMI, mean tGWG, and median maternal prepartum CRP concentration were 20.4 kg/m<sup>2</sup>, 13.9 kg, and 3.3 mg/L, respectively. The prevalence of prepartum CSI and LGI was 7.2% and 47.8%. The adjusted RRs (95% confidence interval) of CSI for normal (18.5–24.9 kg/m<sup>2</sup>) and high ( $\geq 25$  kg/m<sup>2</sup>) vs. low pBMI (< 18.5 kg/m<sup>2</sup>) were 1.35 (1.05–1.74) and 2.28 (1.53–3.39), respectively. The respective adjusted RRs of LGI were 1.19 (1.11–1.28) and 1.59 (1.42–1.77). The adjusted RRs for excessive vs. optimal tGWG was 1.18 (0.94–1.48) for CSI and 1.14 (1.07–1.21) for LGI.

**Conclusions:** Prepregnancy overweight/obesity and excessive tGWG increase the risk of maternal prepartum systemic inflammation, which further highlights the importance of weight management before and during pregnancy.

**Keywords:** Prepregnancy body mass index, Gestational weight gain, C-reactive protein, Systemic inflammation, Cohort study

## Background

C-reactive protein (CRP) is an acute-phase reactant synthesized by the hepatocytes in response to both infectious and non-infectious factors, and elevated CRP level is considered to reflect systemic inflammation [1, 2]. In general populations, positive associations have been found between increased adiposity and serum CRP concentration [3–6]. This supports the concept that obesity contributes to a state of low-grade systemic inflammation

\*Correspondence: lik@gwcmc.org; liuhuishu@gwcmc.org

<sup>1</sup> Clinical Epidemiology Group, Department of Clinical Research, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, 9 Jinsui Road, Guangzhou 510623, China

<sup>2</sup> Department of Gynecology and Obstetrics, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, 9 Jinsui Road, Guangzhou 510623, China



by promoting the production of a wide variety of pro-inflammatory cytokines [7].

The positive associations between prepregnancy body mass index (pBMI) and the risks of gestational diabetes and preeclampsia are well known [8–10], and both morbidities have been associated with elevated maternal serum CRP levels [11–13]. Several studies have also reported direct associations between prepregnancy obesity and elevated gestational CRP levels, and have suggested that inflammation is likely to be a mediator between prepregnancy obesity and adverse pregnancy outcomes [14–17].

It seems clear that women with prepregnancy overweight/obesity are prone to excessive gestational weight gain (GWG) [18–22]; it is thus of interest to examine whether excessive GWG, after taking into account prepregnancy adiposity, is an independent risk factor driving maternal systemic inflammation. Previous studies addressing this question have produced conflicting findings [23–25].

The goal of this study was to examine the associations of pBMI and total GWG (tGWG) with maternal prepartum serum CRP concentration in a large cohort of Chinese women who had undergone apparently healthy pregnancies.

## Methods

### Study population and data collection

This study was conducted at Guangzhou Women and Children's Medical Center, and was based on a cohort initially including 22,455 pregnant women who delivered singletons following spontaneous pregnancy between 2016 and 2020. Data on pBMI, tGWG, and maternal prepartum serum high-sensitivity CRP concentration (measured on the same day of delivery or on the day preceding delivery) were available in these women. Exclusion criteria included women with the following characteristics: 1) pre-existing diabetes mellitus, hypertension, cardiovascular disease, renal diseases, liver diseases, thyroid diseases, or chronic infectious diseases (e.g. hepatitis B) ( $n=5226$ ); 2) preeclampsia, gestational diabetes mellitus, or any acute or chronic infectious diseases ascertained during the course of the index pregnancy ( $n=9386$ ); 3) premature rupture of membranes and placental problems such as placenta previa ( $n=8925$ ) in the index pregnancy; 4) fetal/neonatal structural abnormalities ( $n=1085$ ); 5) non-term ( $<37$  or  $\geq 42$  gestational weeks) deliveries ( $n=903$ ); 6) neonate 5-min Apgar score  $<7$  ( $n=22$ ); and 7) abnormal neonate birth weight ( $<2500$  or  $>4000$  g,  $n=1329$ ). We further excluded women with serum CRP concentration measured after delivery ( $n=426$ ) and women with seemingly erroneous values on body weight before delivery ( $<30$  or  $>150$  kg,  $n=68$ ) and

on total gestational weight gain (determined by the lower and upper 0.5 percentiles, i.e.,  $\leq 0$  or  $\geq 29$  kg,  $n=181$ ). After these exclusions, a total of 5476 women remained for final analysis. The study participant flowchart was provided in the [Supplementary file](#).

Maternal demographics and reproductive history information as well as self-reported prepregnancy weight were collected at the antenatal booking visit. All women underwent ultrasound examinations performed at 11–13 gestational weeks. The crown-rump length from this ultrasound was used to determine gestational age. Maternal weight before delivery was obtained from the admission assessment form on labor and delivery. Maternal non-fasting venous blood samples, which were usually drawn shortly after admission, were collected in serum separator tubes and then analyzed in our laboratory. Serum CRP concentration was quantified using turbidimetric inhibition immunoassay kit (Strong Biotechnologies, Beijing, China).

This study was approved by the Guangzhou Women and Children's Medical Center ethics committee (2021-030A01). Because this was a retrospective, chart review study, the ethics committee determined that patient informed consent was not required.

### Statistical analysis

Study participants' characteristics were described using mean and standard deviation for continuous variables and using frequency and percentage for categorical ones. The relationships of pBMI and tGWG with log-transformed maternal prepartum serum CRP concentration were analyzed using linear regression, with and without adjustment for other maternal and fetal characteristics. The linear models were extended with restricted cubic spline terms to identify possible non-linearity.

Maternal prepartum serum CRP concentration was then categorized into  $<3$  mg/L,  $3-10$  mg/L, and  $>10$  mg/L. This study defined the last two categories as low-grade inflammation (LGI) and clinically significant inflammation (CSI), consistent with other studies [6, 26–29]. Multivariable Poisson regression models were built to estimate the relative risks (RRs) of having LGI and CSI across pBMI and tGWG categories, which were defined using the WHO criteria (pBMI  $<18.5$ ,  $18.5-24.9$ , and  $\geq 25.0$  kg/m<sup>2</sup>) and the approximate tertile cut-offs (tGWG  $<12.0$ ,  $12.0-14.9$ , and  $\geq 15.0$  kg), respectively. The multivariable Poisson models adjusted for maternal age, parity, and gestational length. When applicable, pBMI and tGWG were mutually adjusted for. Test for trend was performed by substituting the actual CRP values with their median for each pBMI or tGWG category and then introducing the medians into the model as a continuous variable. Please note that women with

prepartum CSI were excluded when modeling the risk of prepartum LGI.

For tGWG, we also adopted the GWG recommendation proposed by the Institute of Medicine (IOM) to categorize women into three groups, i.e., inadequate, optimal, and excessive GWG. According to the IOM recommendation, the normal tGWG is 12.5 – 18, 11.5 – 16, 7 – 11.5, and 5 – 9 kg for women with pBMI < 18.5, 18.5 – 24.9, 25 – 29.9, and  $\geq 30$  kg/m<sup>2</sup>, respectively [30]. Two additional Poisson regression models were built to examine whether nonoptimal tGWG is associated with different risks of LGI and CSI compared with optimal tGWG.

Two-sided  $P < 0.05$  was considered statistically significant. The R software (version 4.0.2, R foundation, Vienna, Austria) was used to conduct the statistical analyses.

## Results

Among 5476 women, 394 (7.2%) and 2,617 (47.8%) met the criteria for CSI and LGI. The median prepartum CRP concentration for the entire cohort was 3.3 mg/L. Table 1 shows the baseline characteristics of the cohort. Women with prepartum CSI were statistically significantly younger, more likely to be nulliparous and to have higher pBMI than women without. The highest tertile of tGWG ( $\geq 15$  kg) was more frequent among women with than without CSI (43.9% vs. 39.7%).

In both the unadjusted and adjusted linear regression analyses, pBMI and tGWG demonstrated statistically significant positive associations with maternal prepartum CRP concentration (Table 2). Restricted cubic spline analyses revealed no statistically significant non-linearity.

The associations of pBMI and tGWG with maternal prepartum CSI and LGI risks are presented in Table 3. Compared with low pBMI (< 18.5 kg/m<sup>2</sup>), normal pBMI (18.5 – 24.9 kg/m<sup>2</sup>) and overweight/obesity ( $\geq 25.0$  kg/m<sup>2</sup>) were statistically significantly associated with an increased risk of maternal prepartum CSI, with an RR (95% confidence interval [CI]) of 1.35 (1.05–1.74) and 2.28 (1.53–3.39), respectively ( $P_{\text{trend}} < 0.01$ ), whereas tGWG tertiles showed no statistically significant associations. Overweight/obesity and normal BMI vs. low pBMI were also statistically significantly associated with an increased risk of maternal prepartum LGI [RR (95% CI): 1.19 (1.11–1.28), 1.59 (1.42–1.77);  $P_{\text{trend}} < 0.01$ ]. For tGWG tertiles, a statistically significant positive association was seen for the highest ( $\geq 15.0$  kg) vs. the lowest tertile (< 12.0 kg) [RR (95% CI): 1.17 (1.09–1.25)]. No statistically significant interaction was detected between pBMI and tGWG (data not shown).

As shown in Table 4, both excessive and inadequate tGWG vs. normal tGWG demonstrated positive but statistically non-significant associations with maternal prepartum CSI [RR (95% CI): 1.10 (0.88–1.39); 1.18

**Table 1** Baseline characteristics of study participants, overall and by maternal prepartum CSI status

	Total (n = 5476)	Maternal prepartum CSI <sup>a</sup>		P <sup>b</sup>
		No (n = 5082)	Yes (n = 394)	
Maternal age, mean (SD)	30.4 (4.0)	30.5 (4.0)	29.9 (3.9)	0.006
Parity category, n (%)				
Nulliparous	2512 (45.9)	2303 (45.3)	209 (53.0)	
Primiparous	2819 (51.5)	2645 (52.0)	174 (44.2)	
Multiparous	145 (2.6)	134 (2.7)	11 (2.8)	0.010
pBMI (kg/m <sup>2</sup> ), mean (SD)	20.4 (2.5)	20.4 (2.5)	21.0 (3.0)	< 0.001
pBMI category, n (%)				
< 18.5 kg/m <sup>2</sup>	1193 (21.8)	1123 (22.1)	70 (17.8)	
18.5 – 24.9 kg/m <sup>2</sup>	3986 (72.8)	3696 (72.7)	290 (73.6)	
$\geq 25$ kg/m <sup>2</sup>	297 (5.4)	263 (5.2)	34 (8.6)	< 0.001
tGWG (kg), mean (SD)	13.9 (4.0)	13.8 (3.9)	14.0 (4.2)	0.664
tGWG tertiles, n (%)				
< 12 kg	1579 (28.8)	1461 (28.8)	118 (30.0)	
12 – 14.9 kg	1705 (31.2)	1602 (31.5)	103 (26.1)	
$\geq 15$ kg	2192 (40.0)	2019 (39.7)	173 (43.9)	0.075
Gestational length (weeks), mean (SD)	39.4 (0.9)	39.4 (0.9)	39.5 (0.9)	0.528

<sup>a</sup> Maternal prepartum CSI was defined as maternal prepartum serum CRP concentration > 10 mg/L

<sup>b</sup> P values were calculated by t-test for continuous variables and  $\chi^2$  test for categorical variables

CRP C-reactive protein; CSI clinically significant inflammation; pBMI prepregnancy body mass index; SD standard deviation; tGWG total gestational weight gain

(0.94–1.48)]. Compared with normal tGWG, inadequate tGWG was associated with a statistically significantly decreased risk of maternal prepartum LGI risk [OR (95% CI): 0.91 (0.85–0.98)], while excessive tGWG showed a statistically significant but positive association [OR (95% CI): 1.14 (1.07–1.21)].

## Discussion

In this study of women who had undergone apparently healthy pregnancies, pBMI and tGWG were both positively associated with maternal prepartum serum CRP concentration. Furthermore, higher pBMI was statistically significantly associated with increased prepartum CSI and LGI risks. Excessive tGWG, as defined by the IOM recommendation, was independently and statistically significantly associated with an increased LGI risk.

The association between obesity and maternal CRP level has been observed in pregnant women [14–17]. Although normal pregnancy is known as a process involving inflammation [31], elevated inflammation is

**Table 2** Unadjusted and adjusted associations of pBMI and tGWG with log-transformed maternal prepartum serum CRP concentration (n = 5476)

	Unadjusted model		Adjusted model <sup>a</sup>		Adjusted RCS model <sup>a</sup>	
	β	P	β	P	β	P
pBMI	4.54 × 10 <sup>-2</sup>	<0.01	5.69 × 10 <sup>-2</sup>	<0.01	5.68 × 10 <sup>-2</sup>	<0.01
tGWG	1.40 × 10 <sup>-2</sup>	<0.01	1.69 × 10 <sup>-2</sup>	<0.01	1.28 × 10 <sup>-2</sup>	0.297
pBMI RCS term					-1.85 × 10 <sup>-6</sup>	0.985
tGWG RCS term					3.28 × 10 <sup>-5</sup>	0.735

<sup>a</sup> pBMI and tGWG were mutually adjusted for in the multivariable models. The multivariable models were also adjusted for maternal age, parity, and gestational length  
CRP C-reactive protein; pBMI prepregnancy body mass index; RCS restricted cubic spline; tGWG total gestational weight gain

**Table 3** Adjusted associations of pBMI and tGWG with maternal prepartum CSI and LGI (n = 5476)

	No. prepartum inflammation/total no	RR (95% CI) <sup>a</sup>	P	P <sub>trend</sub>
<i>Maternal prepartum CSI (n = 394)<sup>b</sup></i>				
pBMI				
< 18.5 kg/m <sup>2</sup>	70/1193	1.00		
18.5 – 24.9 kg/m <sup>2</sup>	290/3986	1.35 (1.05–1.74)	0.020	
≥ 25.0 kg/m <sup>2</sup>	34/297	2.28 (1.53–3.39)	<0.001	<0.001
tGWG tertiles				
< 12.0 kg	118/1579	1.00		
12.0 – 14.9 kg	103/1705	0.83 (0.64–1.07)	0.150	
≥ 15.0 kg	173/2192	1.07 (0.86–1.35)	0.534	0.339
<i>Maternal prepartum LGI (n = 2617)<sup>c</sup></i>				
pBMI				
< 18.5 kg/m <sup>2</sup>	507/1123	1.00		
18.5 – 24.9 kg/m <sup>2</sup>	1935/3696	1.19 (1.11–1.28)	<0.001	
≥ 25.0 kg/m <sup>2</sup>	175/263	1.59 (1.42–1.77)	<0.001	<0.001
tGWG tertiles				
< 12.0 kg	706/1461	1.00		
12.0 – 14.9 kg	790/1602	1.04 (0.96–1.12)	0.312	
≥ 15.0 kg	1121/2019	1.17 (1.09–1.25)	<0.001	<0.001

<sup>a</sup> pBMI and tGWG were mutually adjusted for in the multivariable models. The multivariable models also adjusted for maternal age, parity, and gestational length

<sup>b</sup> Maternal prepartum CSI was defined as maternal prepartum serum CRP concentration > 10 mg/L

<sup>c</sup> Maternal prepartum LGI was defined as maternal prepartum serum CRP concentration ≥ 3.0 and ≤ 10 mg/L. Women with prepartum CSI were excluded  
CI Confidence interval, CRP C-reactive protein, CSI clinically significant inflammation, LGI low-grade inflammation, pBMI prepregnancy body mass index, RR relative risk, tGWG total gestational weight gain

considered pathogenic in the development of preeclampsia, retarded fetal growth, and many other maternal and fetal complications [31–33]. In order to identify elevated CRP levels, reference CRP intervals

**Table 4** Adjusted association of tGWG categorized by the IOM recommendation with maternal prepartum CSI and LGI (n = 5476)

	No. prepartum inflammation/total no	RR (95% CI) <sup>a</sup>	P
<i>Maternal prepartum CSI (n = 394)<sup>b</sup></i>			
Normal tGWG			
177/2707	1.00		
Inadequate tGWG			
104/1446	1.10 (0.88–1.39)	0.404	
Excessive tGWG			
113/1323	1.18 (0.94–1.48)	0.158	
<i>Maternal prepartum LGI (n = 2617)<sup>c</sup></i>			
Normal tGWG			
1277/2530	1.00		
Inadequate tGWG			
618/1342	0.91 (0.85–0.98)	0.011	
Excessive tGWG			
722/1210	1.14 (1.07–1.21)	<0.001	

<sup>a</sup> The multivariable models adjusted for maternal age, parity, pBMI, and gestational length

<sup>b</sup> Maternal prepartum CSI was defined as maternal prepartum serum CRP concentration > 10 mg/L

<sup>c</sup> Maternal prepartum LGI was defined as maternal prepartum serum CRP concentration ≥ 3.0 and ≤ 10 mg/L. Women with prepartum CSI were excluded  
CI Confidence interval, CRP C-Reactive protein, CSI Clinically significant inflammation, IOM Institute of Medicine, LGI Low-grade inflammation, pBMI Prepregnancy body mass index, RR Relative risk, tGWG Gestational weight gain;

throughout normal pregnancies need to be established. In an ad hoc, small-scale Swedish study, the reference interval (i.e. the distance from 2.5 to 97.5 percentile) for maternal prepartum CRP concentration was 0.38 – 24.75 mg/L [34]. In our cohort, the corresponding interval was 0.56 – 16.7 mg/L. In one previous study including only 81 American women with normal pregnancy, the median CRP levels at or beyond 36 gestational weeks was 9 mg/L for women not in labor and 13 mg/L for women in labor [35]. In contrast, the median prepartum CRP concentration in the present study was considerably lower (3.3 mg/L), possibly owing to the low percentage of women with prepregnancy overweight or obesity. Despite a mean BMI of 25 kg/m<sup>2</sup>, another study reported remarkably low CRP

concentrations at 36–38 gestational weeks across all BMI categories among normal pregnancies (e.g., the mean CRP concentration was lower than 1.69 mg/L even for women with BMI  $\geq 30$  kg/m<sup>2</sup>) [36], somewhat contradicting the current knowledge that pregnancy by nature is a low-grade inflammatory state.

Several studies investigating the longitudinal changes of maternal CRP concentration throughout pregnancy have reported either increasing or decreasing trends [34, 37]. It is interesting that in the study by Larsson et al. [34], a notable decrease in maternal CRP concentration was seen prepartum (0–14 days before delivery) after an overall increasing trend until the 34–38 gestational weeks. Although our study measured maternal CRP on one single occasion when the CRP concentration is supposed to be low, we ascertained 47.8% women with LGI, confirming the extensive involvement of inflammation throughout pregnancy [31]. This study also suggests that the proinflammatory effect of prepregnancy adiposity persists as pregnancy proceeds, providing a possible explanation to the high CRP concentration in the colostrum from obese mothers and the presence of CRP in breast milk as a result of elevated maternal postpartum CRP concentration [38, 39]. However, this study is inconsistent with the study by Friss CM et al., where they found that the evident association between adiposity and maternal CRP concentration diminished toward the end of pregnancy [36].

Evidence obtained from non-pregnant populations indicates a positive association between weight gain and systemic inflammation characterized by elevated CRP concentration [3, 4, 40, 41]. In this study, we observed a similar positive association between tGWG and maternal prepartum CRP concentration, consisting with one prior study, in which per 1-kg increment in GWG was associated with a 3% increase in maternal CRP concentration [25]. However, the association was null in the study by Friss CM et al. [36]. In another study, tGWG showed positive associations with adiponectin and leptin but not with CRP [23].

Although we reported a positive association of tGWG with maternal prepartum CRP concentration and an increased LGI risk for women with excessive GWG, we could not determine a clear causal relationship, given the findings from prospective investigations suggesting that higher baseline CRP levels might promote the future weight gain [3, 42–44]. In order to rule out this reverse causality, prepregnancy or early-pregnancy CRP concentration should also be taken into account.

Our study's strengths include its relatively large sample size and careful selection of only pregnancies without significant medical comorbidities. However, a few limitations need to be acknowledged. First, this study

lacked some important data. As we discussed above, baseline maternal CRP concentration was not available in this study; thus we were unable to confirm the causal relationship between tGWG and maternal prepartum CRP level. We were also unable to adjust for some life-style factors in our multivariable analyses due to lack of data. For example, information on cigarette smoking and alcohol consumption was not collected, although they are believed to be relatively uncommon among Chinese women. In addition, we did not collect any data on other confounding factors including socioeconomic status, diet, and physical activity. According to previous studies [45–47], these factors might affect serum CRP concentration in non-pregnant populations, but we could not rule out their potential confounding effects in this study. Secondly, we followed the protocols of several previous studies to define LGI and CSI [26, 48, 49], but there is no clear consensus. In addition, as those studies measured CRP concentration in non-pregnant populations, and therefore their protocols might not be suitable for this study due to the typically elevated serum CRP levels in pregnant women. Thirdly, pBMI based on self-reported prepregnancy body weight was subject to recall bias and how it biased our results was determined by the pattern of misreporting. Lastly, the present study was conducted in a considerably lean cohort, which would limit the generalization of our results to populations where overweight and obesity are much more prevalent.

## Conclusions

This cohort study among Chinese women with uncomplicated pregnancies suggests that higher pBMI and excessive tGWG may increase the risk of maternal prepartum systemic inflammation, further highlighting the importance of weight management before and during pregnancy.

## Supplementary information

The online version contains supplementary material available at <https://doi.org/10.1186/s12884-022-04849-y>.

Additional file 1.

## Acknowledgements

The authors would like to thank Mrs. Juan Li for her administrative support.

## Authors' contributions

KL and HL designed the study. KL performed the statistical analyses and drafted the manuscript. CY and XL contributed to data collection. HL, JF and CG helped with the result interpretation. All authors critically reviewed the manuscript and gave their final approval for this submission.

## Funding

This study was financially supported by a joint project between Guangzhou Metropolitan Science Bureau and Guangzhou Women and Children's Medical

Center (grant reference number: 202102010245). The funder of this study had no role in study design, data analysis, and result interpretation and reporting.

#### Availability of data and materials

The data that support the findings of this study are available from the corresponding authors upon reasonable request.

#### Declarations

##### Ethics approval and consent to participate

This study was a part of a research project investigating gestational diabetes diagnosis and risk factors based on retrospective analysis of clinical data from electronic health records. The Guangzhou Women and Children's Medical Center ethics committee approved the study protocol on February 25, 2021 (ref. no. 2021-030A01), in which we requested to waive the collection of patient informed consents. The study was conducted in accordance with the Declaration of Helsinki.

##### Consent for publication

Not applicable.

##### Competing interests

The authors declare that they have no conflicts of interest.

Received: 29 July 2021 Accepted: 6 June 2022

Published online: 29 June 2022

#### References

- Gabay C, Kushner I. Acute-phase proteins and other systemic responses to inflammation. *N Engl J Med*. 1999;340(6):448–54.
- Young B, Gleeson M, Cripps AW. C-reactive protein: a critical review. *Pathology*. 1991;23(2):118–24.
- Barinas-Mitchell E, Cushman M, Meilahn EN, Tracy RP, Kuller LH. Serum levels of C-reactive protein are associated with obesity, weight gain, and hormone replacement therapy in healthy postmenopausal women. *Am J Epidemiol*. 2001;153(11):1094–101.
- Fransson EI, Batty GD, Tabak AG, Brunner EJ, Kumari M, Shipley MJ, Singh-Manoux A, Kivimaki M. Association between change in body composition and change in inflammatory markers: an 11-year follow-up in the Whitehall II Study. *J Clin Endocrinol Metab*. 2010;95(12):5370–4.
- Nguyen XM, Lane J, Smith BR, Nguyen NT. Changes in inflammatory biomarkers across weight classes in a representative US population: a link between obesity and inflammation. *J gastrointest surg*. 2009;13(7):1205–12.
- Visser M, Bouter LM, McQuillan GM, Wener MH, Harris TB. Elevated C-reactive protein levels in overweight and obese adults. *JAMA*. 1999;282(22):2131–5.
- Berg AH, Scherer PE. Adipose tissue, inflammation, and cardiovascular disease. *Circ Res*. 2005;96(9):939–49.
- Shin D, Song WO. Prepregnancy body mass index is an independent risk factor for gestational hypertension, gestational diabetes, preterm labor, and small- and large-for-gestational-age infants. *J Matern fetal Neonatal Med*. 2015;28(14):1679–86.
- Rosenberg TJ, Garbers S, Chavkin W, Chiasson MA. Prepregnancy weight and adverse perinatal outcomes in an ethnically diverse population. *Obstet Gynecol*. 2003;102(5 Pt 1):1022–7.
- Schummers L, Hutcheon JA, Bodnar LM, Lieberman E, Himes KP. Risk of adverse pregnancy outcomes by prepregnancy body mass index: a population-based study to inform prepregnancy weight loss counseling. *Obstet Gynecol*. 2015;125(1):133–43.
- Qiu C, Sorensen TK, Luthy DA, Williams MA. A prospective study of maternal serum C-reactive protein (CRP) concentrations and risk of gestational diabetes mellitus. *Paediatr Perinat Epidemiol*. 2004;18(5):377–84.
- Wolf M, Sandler L, Hsu K, Vossen-Smirnakis K, Ecker JL, Thadhani R. First-trimester C-reactive protein and subsequent gestational diabetes. *Diabetes Care*. 2003;26(3):819–24.
- Rebelo F, Schluskel MM, Vaz JS, Franco-Sena AB, Pinto TJ, Bastos FI, Adegoye AR, Kac G. C-reactive protein and later preeclampsia: systematic review and meta-analysis taking into account the weight status. *J Hypertens*. 2013;31(1):16–26.
- Shin D, Hur J, Cho EH, Chung HK, Shivappa N, Wirth MD, Hebert JR, Lee KW. Pre-Pregnancy Body Mass Index Is Associated with Dietary Inflammatory Index and C-Reactive Protein Concentrations during Pregnancy. *Nutrients*. 2017;9(4):351.
- Retnakaran R, Hanley AJ, Raif N, Connelly PW, Sermer M, Zinman B. C-reactive protein and gestational diabetes: the central role of maternal obesity. *J Clin Endocrinol Metab*. 2003;88(8):3507–12.
- Bodnar LM, Ness RB, Harger GF, Roberts JM. Inflammation and triglycerides partially mediate the effect of prepregnancy body mass index on the risk of preeclampsia. *Am J Epidemiol*. 2005;162(12):1198–206.
- Witteveen AB, Henrichs J, Bellers M, van Oenen E, Verhoeven CJ, Vrijkotte TGM. Mediating role of C-reactive protein in associations between pre-pregnancy BMI and adverse maternal and neonatal outcomes: the ABCD-study cohort. *The journal of maternal-fetal & neonatal medicine : the official journal of the European Association of Perinatal Medicine, the Federation of Asia and Oceania Perinatal Societies, the International Society of Perinatal Obstet*. 2022;35(15):2867–75.
- Krukowski RA, Bursac Z, McGehee MA, West D. Exploring potential health disparities in excessive gestational weight gain. *J Womens Health*. 2013;22(6):494–500.
- Restall A, Taylor RS, Thompson JM, Flower D, Dekker GA, Kenny LC, Poston L, McCowan LM. Risk factors for excessive gestational weight gain in a healthy, nulliparous cohort. *J obes*. 2014;2014:148391.
- Garay SM, Sumption LA, Pearson RM, John RM. Risk factors for excessive gestational weight gain in a UK population: a biopsychosocial model approach. *BMC Pregnancy Childbirth*. 2021;21(1):43.
- Heery E, Kelleher CC, Wall PG, McAuliffe FM. Prediction of gestational weight gain - a biopsychosocial model. *Public Health Nutr*. 2015;18(8):1488–98.
- Kowal C, Kuk J, Tamim H. Characteristics of weight gain in pregnancy among Canadian women. *Matern Child Health J*. 2012;16(3):668–76.
- Logan CA, Bornemann R, Koenig W, Reister F, Walter V, Fantuzzi G, Weyermann M, Brenner H, Genuone J, Rothenbacher D. Gestational weight gain and fetal-maternal adiponection, leptin, and CRP: results of two birth cohorts studies. *Sci Rep*. 2017;7:41847.
- Perrig W, Rifas-Shiman SL, Rich-Edwards JW, Stuebe AM, Oken E. Inflammation and weight gain in reproductive-aged women. *Ann Hum Biol*. 2016;43(1):91–5.
- Hrolfsdottir L, Schalkwijk CG, Birgisdottir BE, Gunnarsdottir I, Maslova E, Granstrom C, Strom M, Olsen SF, Halldorsson TI. Maternal diet, gestational weight gain, and inflammatory markers during pregnancy. *Obesity*. 2016;24(10):2133–9.
- Dhingra R, Gona P, Nam BH, D'Agostino RB Sr, Wilson PW, Benjamin EJ, O'Donnell CJ. C-reactive protein, inflammatory conditions, and cardiovascular disease risk. *Am J Med*. 2007;120(12):1054–62.
- Dinh KM, Kaspersen KA, Mikkelsen S, Pedersen OB, Petersen MS, Thorner LW, Hjalgrim H, Rostgaard K, Ullum H, Erikstrup C. Low-grade inflammation is negatively associated with physical health-related quality of Life in healthy individuals: results from the danish blood donor study (DBDS). *PLoS ONE*. 2019;14(3):e0214468.
- Rifai N, Ridker PM. Population distributions of C-reactive protein in apparently healthy men and women in the United States: implication for clinical interpretation. *Clin Chem*. 2003;49(4):666–9.
- Imhof A, Frohlich M, Loewel H, Helbecque N, Woodward M, Amouyel P, Lowe GD, Koenig W. Distributions of C-reactive protein measured by high-sensitivity assays in apparently healthy men and women from different populations in Europe. *Clin Chem*. 2003;49(4):669–72.
- Weight Gain During Pregnancy: Reexamining the Guidelines. Institute of Medicine (US) and National Research Council (US) Committee to Reexamine IOM Pregnancy Weight Guidelines. Kathleen M Rasmussen, Ann LYaktine<br>, editors. Washington (DC): National Academies Press (US); 2009. <https://pubmed.ncbi.nlm.nih.gov/20669500/>.
- Romero R, Gotsch F, Pineles B, Kusanovic JP. Inflammation in pregnancy: its roles in reproductive physiology, obstetrical complications, and fetal injury. *Nutr Rev*. 2007;65(12 Pt 2):S194–202.
- Palm M, Axelsson O, Wernroth L, Larsson A, Basu S. Involvement of inflammation in normal pregnancy. *Acta Obstet Gynecol Scand*. 2013;92(5):601–5.

33. Cotechini T, Graham CH. Aberrant maternal inflammation as a cause of pregnancy complications: A potential therapeutic target? *Placenta*. 2015;36(8):960–6.
34. Larsson A, Palm M, Hansson LO, Basu S, Axelsson O. Reference values for alpha1-acid glycoprotein, alpha1-antitrypsin, albumin, haptoglobin, C-reactive protein, IgA, IgG and IgM during pregnancy. *Acta Obstet Gynecol Scand*. 2008;87(10):1084–8.
35. Watts DH, Krohn MA, Wener MH, Eschenbach DA. C-reactive protein in normal pregnancy. *Obstet Gynecol*. 1991;77(2):176–80.
36. Friis CM, Paasche Roland MC, Godang K, Ueland T, Tanbo T, Bollerslev J, Henriksen T. Adiposity-related inflammation: effects of pregnancy. *Obesity*. 2013;21(1):E124–130.
37. Christian LM, Porter K. Longitudinal changes in serum proinflammatory markers across pregnancy and postpartum: effects of maternal body mass index. *Cytokine*. 2014;70(2):134–40.
38. Fujimori M, Franca EL, Morais TC, Fiorin V, de Abreu LC, Honorio-Franca AC. Cytokine and adipokine are biofactors can act in blood and colostrum of obese mothers. *BioFactors*. 2017;43(2):243–50.
39. Whitaker KM, Marino RC, Haapala JL, Foster L, Smith KD, Teague AM, Jacobs DR, Fontaine PL, McGovern PM, Schoenfeld TC, et al. Associations of Maternal Weight Status Before, During, and After Pregnancy with Inflammatory Markers in Breast Milk. *Obesity*. 2017;25(12):2092–9.
40. Saito I, Yonemasu K, Inami F. Association of body mass index, body fat, and weight gain with inflammation markers among rural residents in Japan. *Circ J*. 2003;67(4):323–9.
41. Thompson AL, Koehler E, Herring AH, Paynter L, Du S, Zhang B, Popkin B, Gordon-Larsen P. Weight Gain Trajectories Associated With Elevated C-Reactive Protein Levels in Chinese Adults. *J Am Heart Assoc*. 2016;5(9):e003262.
42. Barzilay JI, Forsberg C, Heckbert SR, Cushman M, Newman AB. The association of markers of inflammation with weight change in older adults: the Cardiovascular Health Study. *Int J Obes*. 2006;30(9):1362–7.
43. Engstrom G, Hedblad B, Stavenow L, Lind P, Janzon L, Lindgarde F. Inflammation-sensitive plasma proteins are associated with future weight gain. *Diabetes*. 2003;52(8):2097–101.
44. Holz T, Thorand B, Doring A, Schneider A, Meisinger C, Koenig W. Markers of inflammation and weight change in middle-aged adults: results from the prospective MONICA/KORA S3/F3 study. *Obesity*. 2010;18(12):2347–53.
45. Nazmi A, Victora CG. Socioeconomic and racial/ethnic differentials of C-reactive protein levels: a systematic review of population-based studies. *BMC Public Health*. 2007;7:212.
46. Hart MJ, Torres SJ, McNaughton SA, Milte CM. A Dietary Inflammatory Index and associations with C-reactive protein in a general adult population. *Eur J Nutr*. 2021;60(7):4093–106.
47. Kasapis C, Thompson PD. The effects of physical activity on serum C-reactive protein and inflammatory markers: a systematic review. *J Am Coll Cardiol*. 2005;45(10):1563–9.
48. Woloshin S, Schwartz LM. Distribution of C-reactive protein values in the United States. *N Engl J Med*. 2005;352(15):1611–3.
49. Kushner I, Samols D, Magrey M. A unifying biologic explanation for “high-sensitivity” C-reactive protein and “low-grade” inflammation. *Arthritis Care Res*. 2010;62(4):442–6.

## Publisher's note

Springer nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

