


CONTEMPORARY REVIEW

# In-Depth Extracorporeal Cardiopulmonary Resuscitation in Adult Out-of-Hospital Cardiac Arrest

Mark Dennis , MBBS(Hons), PhD; Sean Lal, MBBS, PhD; Paul Forrest, MBChB; Alistair Nichol, MB BCh, PhD; Lionel Lamhaut, MD, PhD; Richard J. Totaro, MBBS; Brian Burns, MB BCh, MSc; Claudio Sandroni, MD

**ABSTRACT:** The use of extracorporeal cardiopulmonary resuscitation (E-CPR) for the treatment of patients with out-of-hospital cardiac arrest who do not respond to conventional cardiopulmonary resuscitation (CPR) has increased significantly in the past 10 years, in response to case reports and observational studies reporting encouraging results. However, no randomized controlled trials comparing E-CPR with conventional CPR have been published to date. The evidence from systematic reviews of the available observational studies is conflicting. The inclusion criteria for published E-CPR studies are variable, but most commonly include witnessed arrest, immediate bystander CPR, an initial shockable rhythm, and an estimated time from CPR start to establishment of E-CPR (low-flow time) of <60 minutes. A shorter low-flow time has been consistently associated with improved survival. In an effort to reduce low-flow times, commencement of E-CPR in the prehospital setting has been reported and is currently under investigation. The provision of an E-CPR service, whether hospital based or prehospital, carries considerable cost and technical challenges. Despite increased adoption, many questions remain as to which patients will derive the most benefit from E-CPR, when and where to implement E-CPR, optimal post-arrest E-CPR care, and whether this complex invasive intervention is cost-effective. Results of ongoing trials are awaited to determine whether E-CPR improves survival when compared with conventional CPR.

**Key Words:** cardiac arrest ■ cardiopulmonary resuscitation ■ ECPR ■ extracorporeal circulation

Every year, ~350 000 people in the United States<sup>1</sup> and 275 000 in Europe<sup>2</sup> experience an out-of-hospital cardiac arrest (OHCA). Overall survival to discharge from OHCA resuscitated with conventional cardiopulmonary resuscitation (C-CPR) and advanced life support protocols including defibrillation, is reported to be between 2% and 15%.<sup>1,3</sup> These poor survival rates are consistent across geographical locations and have had only modest improvement over time.<sup>4</sup>

In patients with OHCA that is refractory to C-CPR and in whom the cause of the OHCA is potentially reversible, extracorporeal cardiopulmonary resuscitation (E-CPR) can provide a bridge to definitive treatment and recovery. The use of E-CPR has increased 10-fold in the past 10 years<sup>5</sup> and many observational

studies have shown encouragingly high survival rates.<sup>6,7</sup> However, no randomized controlled trials of E-CPR versus C-CPR have been published to date, thus the benefits of E-CPR over C-CPR remain unproven. This review examines the current literature on E-CPR for OHCA, including patient selection, implementation models, post-arrest care, cost-effectiveness, and efficacy. We also highlight areas of ongoing research on E-CPR and potential future developments.

## EXTRACORPOREAL MEMBRANE OXYGENATION AND E-CPR

Extracorporeal membrane oxygenation (ECMO) is a combination of a blood pump and an oxygenator that

Correspondence to: Mark Dennis, MBBS(Hons), PhD, Cardiology Department, Royal Prince Alfred Hospital, Missenden Road, Camperdown, New South Wales 2050 Australia. Email: mark.dennis@sydney.edu.au

Supplementary Materials for this article is available at <https://www.ahajournals.org/doi/suppl/10.1161/JAHA.120.016521>

For Disclosures, see page 8.

© 2020 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

JAHA is available at: [www.ahajournals.org/journal/jaha](http://www.ahajournals.org/journal/jaha)

### Nonstandard Abbreviations and Acronyms

<b>APACAR2A</b>	Comparative Study Between a Pre-hospital and an In-hospital Circulatory Support Strategy (ECMO) in Refractory Cardiac Arrest
<b>BLENDER</b>	Blend to Limit Oxygen in ECMO: A Randomised Controlled Registry Trial
<b>C-CPR</b>	conventional cardiopulmonary resuscitation
<b>CPR</b>	cardiopulmonary resuscitation
<b>ECMO</b>	extracorporeal membrane oxygenation
<b>E-CPR</b>	extracorporeal cardiopulmonary resuscitation
<b>EMS</b>	emergency medical service
<b>EXACT</b>	Reduction of Oxygen After Cardiac Arrest trial
<b>OHCA</b>	out-of-hospital cardiac arrest
<b>OR</b>	odds ratio
<b>RCA</b>	refractory cardiac arrest
<b>ROSC</b>	return of spontaneous circulation
<b>Sub 30</b>	Feasibility of pre-hospital ECPR implementation in under 30 minutes trial
<b>TAME</b>	Targeted Therapeutic Mild Hypercapnia After Resuscitated Cardiac Arrest

can be used to support either pulmonary or both pulmonary and cardiac function (venovenous or venoarterial configurations, respectively). E-CPR is used to describe the use of venoarterial ECMO in refractory cardiac arrest (RCA). The increased adoption of E-CPR has been facilitated by the advent of small, portable ECMO devices and circuit improvements. Attributable to advances in ECMO technology, it is possible to deliver either partial or full cardiorespiratory support for weeks or months, if required.

An ECMO circuit consists of a centrifugal pump and a membrane oxygenator for oxygen delivery, CO<sub>2</sub> removal, and temperature management. In E-CPR, venoarterial ECMO is established while cardiopulmonary resuscitation (CPR) is ongoing. The drainage (access) cannula is placed into the inferior vena cava via the femoral vein, and the “return” cannula is inserted into the femoral artery to the level of the common iliac artery (Figure 1). E-CPR for OHCA is usually provided as part of a “bundle of care,”<sup>8,9</sup> which includes early patient transfer to hospital, mechanical CPR to provide effective CPR during transport, early initiation of

E-CPR, and early definitive treatment (eg, coronary angioplasty [Figure 2]).

### RATIONALE FOR E-CPR


C-CPR either by manual or mechanical chest compression is the mainstay of circulatory support during cardiac arrest. However, even with optimal technique, C-CPR delivers only 15% to 25% of normal cardiac output<sup>10</sup> (or a cardiac index of  $\approx 0.6$  L/min per m<sup>2</sup>)<sup>11</sup> with consequent rapid development of ischemic damage to vital organs (this is commonly referred to as a *low-flow* state). Conversely, E-CPR can provide near-normal levels of cerebral and end organ perfusion.<sup>12</sup> The ability to provide full cerebral and end organ blood supply, even for days or weeks, with E-CPR has enabled a paradigm shift in cardiac arrest—preservation of the brain while awaiting return of spontaneous circulation (ROSC), definitive care, and cardiac recovery.

### INDICATIONS AND PATIENT SELECTION FOR E-CPR

To date, there are no unanimously accepted indications for E-CPR. The consensus-based French Ministry of Health guidelines<sup>13</sup> recommend considering E-CPR for RCA when  $\geq 1$  of the following conditions are met: (1) known reversible cause (eg, drug intoxication or hypothermia), likely to require prolonged life support; (2) signs of life during CPR; (3) no-flow (the period of time from cardiac arrest to the commencement of CPR) and low-flow durations  $< 5$  and 100 minutes, respectively; (3) an initial shockable rhythm; and (4) an end-tidal CO<sub>2</sub>  $> 10$  mm Hg. However, these guidelines have not been prospectively validated. Moreover, few subsequent studies on E-CPR for OHCA have adopted all of these indications and some have adopted none of them.<sup>14</sup>

The inclusion criteria used in many E-CPR studies include age 18 to 65 years or 18 to 70 years, witnessed RCA, immediate bystander CPR, initial shockable rhythm, access to immediate coronary angiography, and an anticipated low-flow period (ie, the interval from C-CPR commencement to E-CPR) of  $< 60$  minutes.<sup>8</sup>

A systematic review and meta-analysis of 841 E-CPR-treated patients with OHCA (Debaty et al<sup>15</sup>) for prognostic factors or neurological outcome and survival demonstrated that an initial shockable rhythm was associated with twice the odds (odds ratio [OR], 2.20; 95% CI, 1.30–3.72) of favorable outcomes (survival with good neurological function) than nonshockable rhythms. Other factors associated with favorable outcomes were low-flow duration (54 versus 64 minutes) (OR, 0.90; 95% CI, 0.81–0.99 [ $P=0.04$ ]), higher arterial




**Figure 1. Venoarterial extracorporeal membrane oxygenation (ECMO) using a peripheral configuration.**

Venous blood (blue) is drained via a cannula positioned at the inferior vena cava to the right atrial junction and passes through the extracorporeal membrane where oxygenation and CO<sub>2</sub> removal occurs. The now oxygenated blood (red) is returned via a “return” cannula position in the common iliac artery or descending aorta. The distal perfusion catheter, applied after ECMO support is established, is inserted into the superficial femoral artery distal to the insertion point of the femoral return cannula, and it supplies oxygenated blood to the distal limb to prevent distal limb ischemia.

pH (difference, 0.12; 95% CI, 0.03–0.22 [ $P=0.01$ ]), and lower serum lactate (difference,  $-3.52$  mmol/L; 95% CI,  $-5.05$  to  $-1.99$  [ $P<0.001$ ]). Bystander CPR was also associated with higher summary odds for favorable outcome (OR, 2.81; 95% CI 0.95–8.32), although this was not statistically significant.

Age has been variably reported as a negative prognostic marker for E-CPR. Goto et al<sup>16</sup> found that age

older than 70 years was associated with poor outcomes. However, Yu et al<sup>17</sup> found that age older than 75 years was not predictive of poor outcome if the low-flow duration was  $<60$  minutes. Debaty et al<sup>15</sup> also did not find an association between adverse prognosis and advancing age.<sup>15</sup> Although age is not currently considered a contraindication to E-CPR, most studies report a median age of patients undergoing E-CPR younger than



**Figure 2. Key steps in extracorporeal cardiopulmonary resuscitation (E-CPR) for out-of-hospital cardiac arrest (OHCA).**<sup>8,9</sup> Key initial cardiac arrest variables of witnessed arrest, immediate bystander cardiopulmonary resuscitation (CPR) and shockable rhythm are currently required to qualify for a potential E-CPR-eligible OHCA in most trials and services. If the arrest is refractory<sup>8,9</sup> to advanced life support measures and E-CPR is available at a nearby hospital, the patient is transferred to the nearest E-CPR-capable hospital. Many services utilize mechanical CPR to enable ongoing chest compression during transfer. Prearrival notification to the accepting hospital by emergency medical services (EMS) is often made. On arrival to the hospital, cannulation, establishment of extracorporeal membrane oxygenation, and subsequent definitive treatment of the underlying cause of the arrest is made. Standard postcardiac arrest care is implemented in the intensive care unit (ICU). \*End-tidal CO<sub>2</sub> (ETCO<sub>2</sub>) and lactate (on arrival to hospital) are variably used as inclusion criteria. <sup>§</sup>Varying definitions of when an arrest becomes "refractory" exist including no return of spontaneous circulation (ROSC) after 30 minutes of conventional CPR.<sup>9</sup>

60 years<sup>14</sup>—lower than the median age of 64 years reported in the largest OHCA registries on adult OHCA.<sup>18</sup>

Other common inclusion criteria for E-CPR include signs of life and end-tidal CO<sub>2</sub> level >10 mm Hg on arrival to the emergency department.<sup>7,19</sup> Their value as inclusion criteria in E-CPR is yet to be systematically assessed.

### WHEN SHOULD THE TRANSITION TO E-CPR OCCUR?

The optimal time point to transition from C-CPR to E-CPR strategy is yet to be determined. Implementation of E-CPR too early may unnecessarily expose patients with C-CPR in whom ROSC may potentially occur to an expensive procedure with significant additional risks, whereas delaying E-CPR implementation may jeopardize the potential benefit from the intervention by increasing the risk of critical brain and end organ injury from prolonged hypoperfusion.

After more than 35 minutes of C-CPR without subsequent E-CPR, <1% of patients with OHCA overall will have spontaneous ROSC and survive with a favorable neurological outcome.<sup>20</sup> Therefore the decision to commence E-CPR needs to occur before this point. In an analysis of North American emergency medical services (EMS) data,<sup>21</sup> transport to an E-CPR center between 8 to 24 minutes of EMS resuscitation had the highest sensitivity and specificity for favorable outcome, with 16 minutes being the optimal time.<sup>21</sup>

In a retrospective Korean propensity-matched analysis (444 patients with C-CPR and 55 patients with E-CPR), Kim et al<sup>22</sup> reported that E-CPR outcomes became superior to C-CPR after 21 minutes of resuscitation. Additionally, E-CPR was associated with improved neurologically intact survival from RCA (21% versus 0% for 41–60 minutes of C-CPR). These data suggest that E-CPR may be considered after 10 to 20 minutes of unsuccessful C-CPR<sup>8,23</sup> in selected patients.<sup>8,24</sup> This, however, presents significant logistical challenges. In order to meet these targets, EMS teams may need to alter their protocols to facilitate shorter on-scene times and adopt mechanical C-CPR to facilitate effective CPR during transportation, ie, moving from a “stay and treat” to a “load and go” strategy. An alternative is implementation of E-CPR at the site of the cardiac arrest—*prehospital E-CPR*—which is discussed below. Any of these options would represent a major change for many EMS systems.

### E-CPR VERSUS C-CPR

While no published randomized control trials of E-CPR versus C-CPR have been published to date, a small number of propensity-matched observational studies have been completed. In a study of 162 adult patients (E-CPR=53, C-CPR=109)<sup>25</sup> with witnessed cardiac arrest of cardiac origin who had undergone CPR for longer than 20 minutes, the intact



survival rate was higher in the E-CPR group than in the propensity-matched C-CPR group (29.2% [7/24] versus 8.3% [2/24], log-rank  $P=0.018$ ). The Save-J (Study of Advanced Cardiac Life Support for Ventricular Fibrillation With Extracorporeal Circulation in Japan)<sup>26</sup> prospectively compared E-CPR and C-CPR outcomes in 454 patients with OHCA who had an initial shockable rhythm and no ROSC until at least 15 minutes after arrival to the hospital. Favorable neurological outcome at 6 months was found in 29 of 260 patients with E-CPR (11.2%) versus 5 of 194 patients (2.6%) in the C-CPR group ( $P=0.001$ ).

Most recently, a large Parisian registry<sup>27</sup> compared C-CPR- and E-CPR-treated patients with OHCA. Overall survival was 8.4% in 525 patients treated with E-CPR and 8.6% in 12 666 patients treated with C-CPR ( $P=0.91$ ). After multivariable regression analysis, the provision of E-CPR was not associated with increased survival (OR, 1.3; 95% CI, 0.8–2.1 [ $P=0.24$ ]). A propensity score-matched analysis of 429 pairs for survival adjusted for major confounders did not find a significant difference in survival between the C-CPR and E-CPR groups (OR, 0.8; 95% CI, 0.5–1.3 [ $P=0.41$ ]). However, despite the large size of that study, several limitations exist. First, there was no specific time to transition to E-CPR strategy nor strict inclusion criteria for E-CPR. This resulted in only 27% of patients having a shockable rhythm and 39% of E-CPR-treated patients receiving CPR for over 90 minutes. Further, E-CPR was initiated at the discretion of treating clinicians, thereby introducing selection bias by known or unknown confounders, which may not have been corrected for in propensity and regression analysis.

Bartos et al<sup>28</sup> retrospectively compared 160 consecutive patients with OHCA with refractory ventricular tachycardia/ventricular fibrillation treated with E-CPR as per predefined University of Minnesota protocol<sup>29</sup> with 654 patients who received standard CPR as part of the Amiodarone, Lidocaine or Placebo in Out-of-Hospital Cardiac Arrest Study (ALPS)<sup>30</sup> using multivariable analysis. Despite E-CPR-treated patients having longer mean duration of CPR (60 versus 35 minutes,  $P<0.001$ ), the E-CPR cohort was associated with better neurological outcomes at all CPR durations <60 minutes (33% E-CPR group versus 23%,  $P=0.01$ ). No patients treated with C-CPR survived if their period of resuscitation was >40 minutes compared with 25% (9/26) in the E-CPR group.

While some systematic reviews and meta-analyses suggested that E-CPR may improve survival or neurological outcome,<sup>31–36</sup> the most recent and comprehensive systematic review from Holmberg et al<sup>14</sup> yielded mixed results. This review included 12 studies on adult OHCA reporting survival to discharge. Of these, 7 showed higher odds of survival with E-CPR versus

C-CPR, while 5 showed lower odds. Among 8 studies reporting survival with good neurological outcome, 7 favored E-CPR. A meta-analysis was not performed because of high study heterogeneity.

All systematic reviews of E-CPR have rated the quality of evidence as very low. The inability of the available reviews to draw consistent strong conclusions on E-CPR for OHCA is a direct result of significant limitations of the individual studies reviewed. Most reviewed studies were small, single-center, and from diverse locations with different EMS systems, all of which limit the quality of the comparative data. Furthermore, inclusion criteria for E-CPR differ between studies, and other unmeasured confounders may limit the internal and external validity of results. Finally, several E-CPR studies have assessed a bundle of care that includes expedited transport to hospital, mechanical CPR, and immediate coronary angiography. While these have not been shown to be convincingly beneficial in their respective own right,<sup>37–39</sup> it is possible that a whole E-CPR “bundle” contributes more to outcomes than E-CPR in isolation.

Based on the low quality of the current E-CPR literature, in its recent Focused Update on Advanced Cardiovascular Life Support,<sup>40</sup> the American Heart Association concluded that there was insufficient evidence to recommend the routine use of E-CPR but that it should be considered as rescue therapy for selected patients when it can be rapidly implemented and supported by skilled providers (class 2b, level of evidence C).

## Prehospital E-CPR

Implementation of E-CPR at the cardiac arrest scene could potentially decrease the time to E-CPR initiation. In a cohort study from Paris,<sup>19</sup> two periods of different E-CPR management strategies were compared. The first period ( $n=114$  patients) included a mandatory 30-minute interval of C-CPR before either transport to hospital (if within 20 minutes range) or initiation of prehospital E-CPR. With this strategy, low-flow duration was  $93\pm 27$  minutes, with 8% neurologically intact survival. In the second period ( $n=42$  patients), a more aggressive prehospital E-CPR strategy and more defined inclusion criteria were implemented, with a dedicated on-call E-CPR team that was dispatched to all patients with OHCA younger than 70 years. Attempts to perform E-CPR commenced after 20 minutes of unsuccessful C-CPR, with the aim of initiating pump flow within 60 minutes from the onset of cardiac arrest. After the implementation of this strategy, the mean low-flow interval was reduced by 20 minutes and neurologically intact survival improved to 29% (21% absolute increase,  $P<0.001$ ). In the propensity-matched analysis of the 2

cohorts, prehospital E-CPR was associated with significantly reduced low-flow duration and higher rates of ROSC compared with hospital-based E-CPR, even if it was not an independent predictor of survival to discharge.

There are several prospective studies (either in progress or planned) to assess the feasibility and efficacy of prehospital E-CPR. For example, the London Sub 30 (Feasibility Study of a Pre-Hospital Extra-Corporeal Membrane Oxygenation (ECMO) Capable Advanced Resuscitation Team at Achieving Blood Flow Within 30 Minutes in Patients With Refractory Cardiac Arrest) study (NCT03700125)<sup>19</sup> will attempt to establish E-CPR within 30 minutes of OHCA using mobile prehospital E-CPR teams in London. The ongoing Paris APACAR2 (A Comparative Study Between a Pre-hospital and an In-hospital Circulatory Support Strategy (ECMO) in Refractory Cardiac Arrest) (NCT02527031)<sup>41</sup> is randomizing patients with OHCA to receive either prehospital or hospital E-CPR, depending on their location and predicted transport time to hospital. Patients with prehospital E-CPR will receive E-CPR between 20 and 30 minutes of C-CPR. Participants randomized to hospital-based E-CPR will be transferred to the hospital with mechanical CPR. Selection criteria include no-flow duration <5 minutes, age 18 to 65 years, refractory arrest (defined as 20 minutes of C-CPR), and either shockable rhythm or signs of life during resuscitation.

## POST-E-CPR RESUSCITATION CARE

Intensive care management of patients with E-CPR is complex and involves a multidisciplinary approach. There are few data on how patients with E-CPR should be managed, and conventional postarrest protocols are generally utilized. The optimal management of arterial oxygen (partial pressure of O<sub>2</sub>) and CO<sub>2</sub> (partial pressure of CO<sub>2</sub>) post-OHCA has not been identified in patients with C-CPR or patients with E-CPR. Arterial hyperoxia is common postcardiac arrest (with or without E-CPR) and has been associated with unfavorable outcomes in some observational C-CPR studies.<sup>42–44</sup>

Following the initiation of other forms of ECMO, a rapid reduction in partial pressure of CO<sub>2</sub> has been associated with adverse neurological outcomes.<sup>45</sup> It has been postulated that this could be caused by an acute reduction in cerebral blood flow.<sup>45</sup> However, in a randomized study<sup>46</sup> of patients with C-CPR and ROSC comparing low-normal versus high-normal partial pressure of CO<sub>2</sub> and normoxia versus moderate hyperoxia, no significant difference in levels of neurological biomarkers was found.<sup>46</sup> This study

was underpowered to identify differences in clinical outcomes.

The randomized controlled trials of reduction of oxygen administration to target an oxygen saturation of 90% to 94%, compared with 98% to 100%, as soon as feasible following successful resuscitation from OHCA (EXACT [Reduction of Oxygen After Cardiac Arrest trial]—NCT03138005) and of oxygen management in patients with ECMO (BLENDER [Blend to Limit Oxygen in ECMO: A Randomised Controlled Registry Trial]—NCT03841084), comparing clinical outcomes in patients with ECMO randomized to receive either 100% oxygen to the oxygenator versus a restrictive strategy, as well as the upcoming TAME [Targeted Therapeutic Mild Hypercapnia After Resuscitated Cardiac Arrest] study (NCT03114033) comparing mild hypercapnia versus normocapnia in C-CPR OHCA, may help guide future CO<sub>2</sub> and oxygen management.

In patients undergoing E-CPR, systemic anticoagulation is required to prevent thrombotic complications from the patient–circuit interface. However, this increases the risk of bleeding, and in patients with E-CPR, significant bleeding has been reported in up to 70% of cases and is associated with worse outcomes.<sup>47</sup> Unfractionated heparin is the most commonly used anticoagulant during ECMO. This is routinely monitored using either activated clotting time, partial thromboplastin time, or anti-Xa activity, the former 2 of which can be measured as point-of-care tests.<sup>48,49</sup> More recently, thromboelastography and thromboelastometry (rotational thromboelastometry) have been investigated as alternative tests for monitoring anticoagulation during ECMO. Unlike activated partial thromboplastin time, which is plasma-based, rotational thromboelastometry and thromboelastography are point-of-care viscoelastic tests that assess multiple coagulation functions in whole blood, including platelet function, fibrinogen function, and fibrinolysis. Preliminary evidence indicates that viscoelastic tests are feasible for anticoagulation management during extracorporeal circulation and are associated with reduced heparin consumption compared with activated partial thromboplastin time–based protocols.<sup>50</sup> Thrombin inhibitors (such as bivalirudin) are mainly used when heparin is contraindicated or, as low antithrombin 3 levels are common on ECMO,<sup>51</sup> in heparin resistance and may enable more tightly controlled anticoagulation<sup>52</sup> in patients undergoing extracorporeal circulation.

## COST-EFFECTIVENESS OF E-CPR

Significant training, skill, equipment, and expense is required to provide an effective E-CPR service. Three recent studies from North America, Australia, and

**Table. Upcoming Randomized Controlled Trials on E-CPR**

Trial Name and Registration Number	Sample Size	Location	Inclusion Criteria	Estimated Completion
INCEPTION (NCT03101787) <sup>62</sup> Standard arm: standard treatment as per ERC guidelines. Intervention: preclinical resuscitation by EMS and transport to the ED with ongoing mechanical cardiopulmonary resuscitation. Initiation of E-CPR in the ED	110	The Netherlands	<ul style="list-style-type: none"> <li>Age 18–70 y</li> <li>Witnessed OHCA</li> <li>Initial rhythm of ventricular fibrillation or ventricular tachycardia when external defibrillation administered</li> <li>Bystander CPR</li> <li>No ROSC within 15 min of C-CPR</li> </ul>	July 2021
EROCA (NCT03065647) <sup>63</sup> Standard arm: BLS and ACLS by EMS per existing protocols at the scene of the cardiac arrest. Intervention arm: patients with OHCA refractory to initial BLS and ACLS will be transported by EMS with ongoing mechanical CPR and ACLS to an ED capable of initiating E-CPR. E-CPR instituted within ED	30	Michigan, United States	<ul style="list-style-type: none"> <li>Age 18–70 y</li> <li>OHCA of presumed nontraumatic cause</li> <li>Predicted arrival time at E-CPR-capable hospital within 1 h</li> <li>Witnessed arrest or initial shockable rhythm</li> <li>Persistent cardiac arrest after initial cardiac rhythm analysis and shock</li> </ul>	December 2020
Prague OHCA Study (NCT01511666) <sup>64</sup> Standard arm: patients in the standard arm will be further managed as per ERC guidelines. Coronary angiography will be performed only if indicated according to routine practice, and mild therapeutic hypothermia will be instituted as soon as possible as per guidelines. Intervention arm: (hyperinvasive) immediate institution of a mechanical chest compression device and prehospital intra-arrest cooling by RhinoChill device (BeneChill). Directly transferred to cardiac center catheterization laboratory under continuous CPR. After admission to cardiac catheterization laboratory for coronary angiography +/- E-CPR	170	Prague, Czech Republic	<ul style="list-style-type: none"> <li>Age 18–65 y</li> <li>Witnessed OHCA of presumed cardiac cause</li> <li>Minimum of 5 min of advanced life support performed by emergency medical team without sustained ROSC</li> <li>ECMO team and bed-capacity in cardiac center available</li> </ul>	December 2020
APACAR2 (NCT02527031) <sup>65</sup> Patients with refractory OHCA—defined by the failure of EMS to resuscitate at the 20th min of cardiac arrest with a minimum of 3 automatic external defibrillations or equivalent analysis will be randomized to prehospital arm: E-CPR in prehospital setting—implementation of E-CPR support at site of cardiac arrest and then transfer to hospital; or in-hospital arm: transfer to hospital for E-CPR implementation in hospital setting	210	Paris, France	<ul style="list-style-type: none"> <li>Adults older than 18 y and those younger than 65 y</li> <li>Refractory cardiac arrest (defined by the failure of professionals to resuscitate at the 20th min of cardiac arrest with a minimum of 3 defibrillator shocks)</li> <li>Beginning of C-CPR within the first 5 min after cardiac arrest (no-flow duration &lt;5 min) with shockable rhythm or the presence of signs of life during resuscitation (any rhythm)</li> <li>Medical cause of the cardiac arrest</li> <li>End-tidal CO<sub>2</sub> &gt;10 mm Hg at the time of inclusion</li> <li>Absence of major comorbidities and E-CPR team available on-site within 40 min of cardiac arrest</li> </ul>	March 2020

ACLS indicates advanced cardiac life support; APACAR2, A Comparative Study Between a Pre-hospital and an In-hospital Circulatory Support Strategy (ECMO) in Refractory Cardiac Arrest; BLS, basic life support; CPR, cardiopulmonary resuscitation; C-CPR, conventional cardiopulmonary resuscitation; E-CPR, extracorporeal cardiopulmonary resuscitation; ED, emergency department; EMS, emergency medical services; ERC, European Resuscitation Council; EROCA, E-CPR for Refractory Out-Of-Hospital Cardiac Arrest; INCEPTION, Early Initiation of Extracorporeal Life Support in Refractory OHCA; OHCA, out-of-hospital cardiac arrest; and ROSC, return of spontaneous circulation.

Japan (Table S1) reported that E-CPR was within accepted cost thresholds for advanced medical interventions.<sup>53–55</sup> These costs ranged between \$16 000 to \$52 000/quality-adjusted life-years, depending on geographical location. E-CPR may be cost-effective for several reasons; patients with E-CPR are generally younger than patients with C-CPR and have good neurological outcomes with minimal residual disabilities and long-life expectancy postevent. In these studies, however, the maximum follow-up period was only 1 year,<sup>55</sup> and long-term outcome data

are needed in future cost-effectiveness analyses. The cost-effectiveness of a prehospital E-CPR strategy is unknown.

The rates of death by neurological criteria in patients resuscitated with E-CPR are ≈3 times higher than in those resuscitated with C-CPR, resulting in an increased potential for organ donation.<sup>56,57</sup> As of yet, the potential for increased organ donation has not been considered in cost-effectiveness modeling of E-CPR programs and presents some ethical questions.

## FUTURE DIRECTIONS OF E-CPR

Using most commonly used inclusion criteria (ie, aged 18 to 65 years, witnessed cardiac arrest, CPR started within 10 minutes, absence of asystole as the first cardiac rhythm on EMS arrival), it is estimated that only 4% to 11%<sup>58–60</sup> of all patients with OHCA would be potentially eligible for E-CPR.<sup>60</sup> The relative infrequency of RCAs meeting current E-CPR inclusion criteria (even in tertiary referral centers)<sup>27,59,60</sup> inherently limits large sample cohorts and therefore the ability to accurately plan recruitment timelines. Moreover, given the known poor outcomes with extended C-CPR, with neurologically intact outcomes of <1% at ≈35 to 40 minutes,<sup>20,61</sup> some clinicians have raised ethical concerns regarding randomization of patients to continued C-CPR (when it may already be deemed to have failed) and E-CPR is available. However, a small number of randomized controlled trials are currently comparing E-CPR with C-CPR (Table).<sup>62–65</sup>

Although difficult to conduct, large-scale multi-center studies of E-CPR are needed to address the significant uncertainties regarding efficacy, case selection, timing, and prognostic markers versus C-CPR. In future studies, standardized definitions of key variables including what constitutes an RCA and CPR duration or arrest to ECMO flow times is required to enable more robust comparisons of outcomes. Moreover, documentation and adjustment for other known prognostic variables including pre-cardiac arrest performance status and intracardiac arrest variables (eg, end-tidal CO<sub>2</sub>, lactate, pH, and potassium) is required.<sup>14</sup>

An initial shockable rhythm is a common entry criterion for most E-CPR programs. This is present in only about one third of OHCA cases<sup>66</sup> and has a significantly better outcome than nonshockable rhythms. However, it is possible that patients with initial nonshockable rhythms may also benefit from E-CPR if other favorable prognostic markers are present.<sup>67</sup>

Should E-CPR be proven to improve survival in selected patients with OHCA, there would be significant organizational and economic implications for health systems to ensure optimal equity of access.

High-volume venovenous ECMO centers have better outcomes than low-volume centers.<sup>68</sup> Given the high complexity of patients with E-CPR, managing them in major centers with significant experience is reasonable. A composite model of having hospital centers that establish E-CPR support, then transfer the patient to a larger center for continued care, has been proposed in a “hub and spoke”-type format. However, this model is yet to be tested in any large-scale settings and may require additional resources to facilitate

the transfer of critically unwell patients with E-CPR between locations.

## CONCLUSIONS

E-CPR for refractory OHCA may improve outcomes in carefully selected patients. However, the evidence base for E-CPR efficacy, patient selection, and post-cardiac arrest management is poor. Much work is needed in this area and large-scale high-quality trials, although challenging, should be made a priority.

## ARTICLE INFORMATION

### Affiliations

From the Sydney Medical School, University of Sydney, Australia (M.D., S.L., P.F.); Departments of Cardiology (M.D., S.L.), Anaesthesia (P.F.), and Intensive Care (R.J.T.), Royal Prince Alfred Hospital, Sydney, Australia; School of Public Health and Preventive Medicine, Monash University, Melbourne, Australia (A.N.); Department of Intensive Care, The Alfred Hospital, Melbourne, Australia (A.N.); INSERM U970 Team 4 “Sudden Death Expertise Center”, Paris, France (L.L.); Paris Descartes University, Paris, France (L.L.); SAMU de Paris-DAR Necker University Hospital-Assistance Public Hopitaux de Paris, Paris, France (L.L.); Greater Sydney Area Helicopter Emergency Medical Service, New South Wales, Ambulance Service, Sydney, Australia (B.B.); Istituto Anestesiologia e Rianimazione Università Cattolica del Sacro Cuore – Policlinico Universitario Agostino Gemelli – IRCCS, Rome, Italy (C.S.); University College Dublin-Clinical Research Centre, St Vincent’s University Hospital, Dublin, Ireland (A.N.).

### Disclosures

None.

### Supplementary Materials

Table S1

References 54–56

## REFERENCES

1. Benjamin EJ, Muntner P, Alonso A, Bittencourt MS, Callaway CW, Carson AP, Chamberlain AM, Chang AR, Cheng S, Das SR, et al. Heart disease and stroke statistics—2019 update: a report from the American Heart Association. *Circulation*. 2019;139:e56–e528.
2. Grasner JT, Lefering R, Koster RW, Masterson S, Bottiger BW, Herlitz J, Wnent J, Tjelmeland IB, Ortiz FR, Maurer H, et al. EuReCa ONE-27 Nations, ONE Europe, ONE Registry: a prospective one month analysis of out-of-hospital cardiac arrest outcomes in 27 countries in Europe. *Resuscitation*. 2016;105:188–195.
3. Berdowski J, Berg RA, Tijssen JG, Koster RW. Global incidences of out-of-hospital cardiac arrest and survival rates: systematic review of 67 prospective studies. *Resuscitation*. 2010;81:1479–1487.
4. Rea TD, Eisenberg MS, Becker LJ, Murray JA, Hearne T. Temporal trends in sudden cardiac arrest: a 25-year emergency medical services perspective. *Circulation*. 2003;107:2780–2785.
5. Richardson AS, Schmidt M, Bailey M, Pellegrino VA, Rycus PT, Pilcher DV. ECMO cardio-pulmonary resuscitation (ECPR), trends in survival from an international multicentre cohort study over 12-years. *Resuscitation*. 2017;112:34–40.
6. Dennis M, Buscher H, Gattas D, Burns B, Habig K, Bannon P, Patel S, Buhr H, Reynolds C, Scott S, et al. Prospective observational study of mechanical cardiopulmonary resuscitation, extracorporeal membrane oxygenation and early reperfusion for refractory cardiac arrest in Sydney: the 2CHEER study. *Crit Care Resusc*. 2020;22:26–34.
7. Bartos JA, Carlson K, Carlson C, Raveendran G, John R, Aufderheide TP, Yannopoulos D. Surviving refractory out-of-hospital ventricular fibrillation cardiac arrest: critical care and extracorporeal membrane oxygenation management. *Resuscitation*. 2018;132:47–55.



8. Hutin A, Abu-Habsa M, Burns B, Bernard S, Bellezzo J, Shinar Z, Torres EC, Gueugniaud PY, Carli P, Lamhaut L. Early ECPR for out-of-hospital cardiac arrest: best practice in 2018. *Resuscitation*. 2018;130:44–48.
9. Guglin M, Zucker MJ, Bazan VM, Bozkurt B, El Banayosy A, Estep JD, Gurley J, Nelson K, Malyala R, Panjra GS, et al. Venoarterial ECMO for adults: JACC scientific expert panel. *J Am Coll Cardiol*. 2019;73:698–716.
10. Lurie KG, Nemergut EC, Yannopoulos D, Sweeney M. The physiology of cardiopulmonary resuscitation. *Anesth Analg*. 2016;122:767–783.
11. Delguercio LR, Feins NR, Cohn JD, Coomaraswamy RP, Wollman SB, State D. Comparison of blood flow during external and internal cardiac massage in man. *Circulation*. 1965;31(suppl 1):171–180.
12. Moller-Helgestad OK, Hyldebrandt JA, Banke A, Rud CS, Udesen NLJ, Linde L, Okkels-Jensen L, Schmidt H, Ravn HB, Moller JE. Impella CP or VA-ECMO in profound cardiogenic shock: left ventricular unloading and organ perfusion in a large animal model. *EuroIntervention*. 2019;14:e1585–e1592.
13. Conseil francais de reanimation c, Societe francaise d'anesthesie et de r, Societe francaise de c, Societe francaise de chirurgie thoracique et c, Societe francaise de medecine du, Societe francaise de p, Groupe francophone de reanimation et d'urgence p, Societe francaise de p, Societe de reanimation de langue f. Guidelines for indications for the use of extracorporeal life support in refractory cardiac arrest. French Ministry of Health. *Ann Fr Anesth Reanim*. 2009;28:182–190.
14. Holmberg MJ, Geri G, Wiberg S, Guerguerian AM, Donnino MW, Nolan JP, Deakin CD, Andersen LW; International Liaison Committee on Resuscitation's Advanced Life S, Pediatric Task F. Extracorporeal cardiopulmonary resuscitation for cardiac arrest: a systematic review. *Resuscitation*. 2018;131:91–100.
15. Debaty G, Babaz V, Durand M, Gaide-Chevronnay L, Fournel E, Blancher M, Bouvaist H, Chavanon O, Maignan M, Bouzat P, et al. Prognostic factors for extracorporeal cardiopulmonary resuscitation recipients following out-of-hospital refractory cardiac arrest. A systematic review and meta-analysis. *Resuscitation*. 2017;112:1–10.
16. Goto T, Morita S, Kitamura T, Natsukawa T, Sawano H, Hayashi Y, Kai T. Impact of extracorporeal cardiopulmonary resuscitation on outcomes of elderly patients who had out-of-hospital cardiac arrests: a single-centre retrospective analysis. *BMJ Open*. 2018;8:e019811.
17. Yu HY, Wang CH, Chi NH, Huang SC, Chou HW, Chou NK, Chen YS. Effect of interplay between age and low-flow duration on neurologic outcomes of extracorporeal cardiopulmonary resuscitation. *Intensive Care Med*. 2019;45:44–54.
18. Chan PS, McNally B, Tang F, Kellermann A; Group CS. Recent trends in survival from out-of-hospital cardiac arrest in the United States. *Circulation*. 2014;130:1876–1882.
19. Lamhaut L, Hutin A, Puymirat E, Jouan J, Raphalen JH, Jouffroy R, Jaffry M, Dagron C, An K, Dumas F, et al. A pre-hospital extracorporeal cardio pulmonary resuscitation (ECPR) strategy for treatment of refractory out hospital cardiac arrest: an observational study and propensity analysis. *Resuscitation*. 2017;117:109–117.
20. Goto Y, Funada A, Goto Y. Relationship between the duration of cardiopulmonary resuscitation and favorable neurological outcomes after out-of-hospital cardiac arrest: a prospective, nationwide, population-based cohort study. *J Am Heart Assoc*. 2016;5:e002819. DOI: 10.1161/JAHA.115.002819
21. Grunau B, Reynolds J, Scheuermeyer F, Stenstrom R, Stub D, Pennington S, Cheskes S, Ramanathan K, Christenson J. Relationship between time-to-ROSC and survival in out-of-hospital cardiac arrest ECPR candidates: when is the best time to consider transport to hospital? *Prehosp Emerg Care*. 2016;20:615–622.
22. Kim SJ, Jung JS, Park JH, Park JS, Hong YS, Lee SW. An optimal transition time to extracorporeal cardiopulmonary resuscitation for predicting good neurological outcome in patients with out-of-hospital cardiac arrest: a propensity-matched study. *Crit Care*. 2014;18:535.
23. Abrams D, Garan AR, Abdelbary A, Bacchetta M, Bartlett RH, Beck J, Belohlavek J, Chen YS, Fan E, Ferguson ND, et al. Position paper for the organization of ECMO programs for cardiac failure in adults. *Intensive Care Med*. 2018;44:717–729.
24. Wengenmayer T, Rombach S, Ramshorn F, Biever P, Bode C, Duerschmied D, Staudacher DL. Influence of low-flow time on survival after extracorporeal cardiopulmonary resuscitation (ECPR). *Crit Care*. 2017;21:157.
25. Maekawa K, Tanno K, Hase M, Mori K, Asai Y. Extracorporeal cardiopulmonary resuscitation for patients with out-of-hospital cardiac arrest of cardiac origin: a propensity-matched study and predictor analysis. *Crit Care Med*. 2013;41:1186–1196.
26. Sakamoto T, Morimura N, Nagao K, Asai Y, Yokota H, Nara S, Hase M, Tahara Y, Atsumi T; Group S-JS. Extracorporeal cardiopulmonary resuscitation versus conventional cardiopulmonary resuscitation in adults with out-of-hospital cardiac arrest: a prospective observational study. *Resuscitation*. 2014;85:762–768.
27. Bougouin W, Dumas F, Lamhaut L, Marjion E, Carli P, Combes A, Pirracchio R, Aissaoui N, Karam N, Deye N, et al. Extracorporeal cardiopulmonary resuscitation in out-of-hospital cardiac arrest: a registry study. *Eur Heart J*. 2020;24:ehz75. [Epub ahead of print].
28. Bartos JA, Grunau B, Carlson C, Duval S, Ripeckyj A, Kalra R, Raveendran G, John R, Conterato M, Frascione RJ, et al. Improved survival with extracorporeal cardiopulmonary resuscitation despite progressive metabolic derangement associated with prolonged resuscitation. *Circulation*. 2020;141:877–886.
29. Yannopoulos D, Bartos JA, Martin C, Raveendran G, Missov E, Conterato M, Frascione RJ, Trembley A, Sipprell K, John R, et al. Minnesota resuscitation consortium's advanced perfusion and reperfusion cardiac life support strategy for out-of-hospital refractory ventricular fibrillation. *J Am Heart Assoc*. 2016;5:e003732. DOI: 10.1161/JAHA.116.003732
30. Kudenchuk PJ, Daya M, Dorian P; Resuscitation Outcomes Consortium I. Amiodarone, lidocaine, or placebo in out-of-hospital cardiac arrest. *N Engl J Med*. 2016;375:802–803.
31. Grunau B, Hornby L, Singal RK, Christenson J, Ortega-Deballon I, Shemie SD, Bashir J, Brooks SC, Callaway CW, Guadagno E, et al. Extracorporeal cardiopulmonary resuscitation for refractory out-of-hospital cardiac arrest: the state of the evidence and framework for application. *Can J Cardiol*. 2018;34:146–155.
32. Ahn C, Kim W, Cho Y, Choi KS, Jang BH, Lim TH. Efficacy of extracorporeal cardiopulmonary resuscitation compared to conventional cardiopulmonary resuscitation for adult cardiac arrest patients: a systematic review and meta-analysis. *Sci Rep*. 2016;6:34208.
33. Ortega-Deballon I, Hornby L, Shemie SD, Bhanji F, Guadagno E. Extracorporeal resuscitation for refractory out-of-hospital cardiac arrest in adults: a systematic review of international practices and outcomes. *Resuscitation*. 2016;101:12–20.
34. Kim SJ, Kim HJ, Lee HY, Ahn HS, Lee SW. Comparing extracorporeal cardiopulmonary resuscitation with conventional cardiopulmonary resuscitation: a meta-analysis. *Resuscitation*. 2016;103:106–116.
35. Wang GN, Chen XF, Qiao L, Mei Y, Lv JR, Huang XH, Shen B, Zhang JS. Comparison of extracorporeal and conventional cardiopulmonary resuscitation: a meta-analysis of 2 260 patients with cardiac arrest. *World J Emerg Med*. 2017;8:5–11.
36. Twohig CJ, Singer B, Grier G, Finney SJ. A systematic literature review and meta-analysis of the effectiveness of extracorporeal-CPR versus conventional-CPR for adult patients in cardiac arrest. *J Intensive Care Soc*. 2019;20:347–357.
37. Brooks SC, Hassan N, Bigham BL, Morrison LJ. Mechanical versus manual chest compressions for cardiac arrest. *Cochrane Database Syst Rev*. 2014;2:CD007260.
38. Patterson T, Perkins GD, Joseph J, Wilson K, Van Dyck L, Robertson S, Nguyen H, McConkey H, Whitbread M, Fothergill R, et al. A randomised trial of expedited transfer to a cardiac arrest centre for non-ST elevation ventricular fibrillation out-of-hospital cardiac arrest: the ARREST pilot randomised trial. *Resuscitation*. 2017;115:185–191.
39. Lemkes JS, Janssens GN, van der Hoeven NW, Jewbali LSD, Dubois EA, Meuwissen M, Rijpstra TA, Bosker HA, Blans MJ, Bleeker GB, et al. Coronary angiography after cardiac arrest without ST-segment elevation. *N Engl J Med*. 2019;380:1397–1407.
40. Panchal AR, Berg KM, Hirsch KG, Kudenchuk PJ, Del Rios M, Cabanas JG, Link MS, Kurz MC, Chan PS, Morley PT, et al. 2019 American Heart Association focused update on advanced cardiovascular life support: use of advanced airways, vasopressors, and extracorporeal cardiopulmonary resuscitation during cardiac arrest: an update to the American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation*. 2019;140:e881–e894.
41. Lamhaut LL. A comparative study between a pre-hospital and an in-hospital circulatory support strategy (ECMO) in pdf. *Clinicaltrials.gov*. 2015. Accessed January 21, 2020.
42. Kilgannon JH, Jones AE, Shapiro NI, Angelos MG, Milcarek B, Hunter K, Parrillo JE, Trzeciak S; Emergency Medicine Shock Research Network

- I. Association between arterial hyperoxia following resuscitation from cardiac arrest and in-hospital mortality. *JAMA*. 2010;303:2165–2171.
43. Kilgannon JH, Jones AE, Parrillo JE, Dellinger RP, Milcarek B, Hunter K, Shapiro NI, Trzeciak S; Emergency Medicine Shock Research Network I. Relationship between supranormal oxygen tension and outcome after resuscitation from cardiac arrest. *Circulation*. 2011;123:2717–2722.
  44. Bellomo R, Bailey M, Eastwood GM, Nichol A, Pilcher D, Hart GK, Reade MC, Egi M, Cooper DJ; Study of Oxygen in Critical Care G. Arterial hyperoxia and in-hospital mortality after resuscitation from cardiac arrest. *J Extra Corpor Technol*. 2013;45:26–32.
  45. Bembea MM, Lee R, Masten D, Kibler KK, Lehmann CU, Brady KM, Easley B. Magnitude of arterial carbon dioxide change at initiation of extracorporeal membrane oxygenation support is associated with survival. *J Extra Corpor Technol*. 2013;45:26–32.
  46. Jakkula P, Reinikainen M, Hastbacka J, Loisa P, Tiainen M, Pettila V, Toppila J, Lahde M, Backlund M, Okkonen M, et al. Targeting two different levels of both arterial carbon dioxide and arterial oxygen after cardiac arrest and resuscitation: a randomised pilot trial. *Intensive Care Med*. 2018;44:2112–2121.
  47. Otani T, Sawano H, Natsukawa T, Matsuoka R, Nakashima T, Takahagi M, Hayashi Y. D-dimer predicts bleeding complication in out-of-hospital cardiac arrest resuscitated with ECMO. *Am J Emerg Med*. 2018;36:1003–1008.
  48. Aubron C, Cheng AC, Pilcher D, Leong T, Magrin G, Cooper DJ, Scheinkestel C, Pellegrino V. Factors associated with outcomes of patients on extracorporeal membrane oxygenation support: a 5-year cohort study. *Crit Care*. 2013;17:R73.
  49. ELSO anticoagulation guideline. Available at: <https://www.elseo.org/portals/0/files/elseoanticoagulationguideline8-2014-table-contents.pdf>. Accessed March 22, 2020.
  50. Panigada M, E Iapichino G, Brioni M, Panarello G, Protti A, Grasselli G, Occhipinti G, Novembrino C, Consonni D, Arcadipane A, et al. Thromboelastography-based anticoagulation management during extracorporeal membrane oxygenation: a safety and feasibility pilot study. *Ann Intensive Care*. 2018;8:7.
  51. Byrnes JW, Swearingen CJ, Prodhan P, Fiser R, Dyamenahalli U. Antithrombin III supplementation on extracorporeal membrane oxygenation: impact on heparin dose and circuit life. *ASAIO J*. 2014;60:57–62.
  52. Ranucci M, Ballotta A, Kandil H, Isgro G, Carlucci C, Baryshnikova E, Pistuddi V; Surgical, Clinical Outcome Research G. Bivalirudin-based versus conventional heparin anticoagulation for postcardiotomy extracorporeal membrane oxygenation. *Crit Care*. 2011;15:R275.
  53. Dennis M, Zmudzki F, Burns B, Scott S, Gattas D, Reynolds C, Buscher H, Forrest P; Sydney ERIG. Cost effectiveness and quality of life analysis of extracorporeal cardiopulmonary resuscitation (ECPR) for refractory cardiac arrest. *Resuscitation*. 2019;139:49–56.
  54. Kawashima T, Uehara H, Miyagi N, Shimajiri M, Nakamura K, Chinen T, Hatano S, Nago C, Chiba S, Nakane H, et al. Impact of first documented rhythm on cost-effectiveness of extracorporeal cardiopulmonary resuscitation. *Resuscitation*. 2019;140:74–80.
  55. Bharmal MI, Venturini JM, Chua RFM, Sharp WW, Beiser DG, Tabit CE, Hirai T, Rosenberg JR, Friant J, Blair JEA, et al. Cost-utility of extracorporeal cardiopulmonary resuscitation in patients with cardiac arrest. *Resuscitation*. 2019;136:126–130.
  56. Casadio MC, Coppo A, Vargiolu A, Villa J, Rota M, Avalli L, Citerio G. Organ donation in cardiac arrest patients treated with extracorporeal CPR: a single centre observational study. *Resuscitation*. 2017;118:133–139.
  57. Sandroni C, D'Arrigo S, Callaway CW, Cariou A, Dragancea I, Taccone FS, Antonelli M. The rate of brain death and organ donation in patients resuscitated from cardiac arrest: a systematic review and meta-analysis. *Intensive Care Med*. 2016;42:1661–1671.
  58. Poppe M, Weiser C, Holzer M, Sulzgruber P, Datler P, Keferbock M, Zeiner S, Lobmeyr E, van Tulder R, Ziegler A, et al. The incidence of "load&go" out-of-hospital cardiac arrest candidates for emergency department utilization of emergency extracorporeal life support: a one-year review. *Resuscitation*. 2015;91:131–136.
  59. Grunau B, Scheuermeyer FX, Stub D, Boone RH, Finkler J, Pennington S, Carriere SA, Cheung A, MacRedmond R, Bashir J, et al. Potential candidates for a structured Canadian ECPR program for out-of-hospital cardiac arrest. *CJEM*. 2016;18:453–460.
  60. Reynolds JC, Grunau BE, Elmer J, Rittenberger JC, Sawyer KN, Kurz MC, Singer B, Proudfoot A, Callaway CW. Prevalence, natural history, and time-dependent outcomes of a multi-center North American cohort of out-of-hospital cardiac arrest extracorporeal CPR candidates. *Resuscitation*. 2017;117:24–31.
  61. Reynolds JC, Frisch A, Rittenberger JC, Callaway CW. Duration of resuscitation efforts and functional outcome after out-of-hospital cardiac arrest: when should we change to novel therapies? *Circulation*. 2013;128:2488–2494.
  62. Bol ME, Suverein MM, Lorusso R, Delnoij TSR, Brandon Bravo Bruinsma GJ, Otterspoor L, Kuijpers M, Lam KY, Vlaar APJ, Elzo Kraemer CV, et al. Early initiation of extracorporeal life support in refractory out-of-hospital cardiac arrest: design and rationale of the inception trial. *Am Heart J*. 2019;210:58–68.
  63. E-CPR for refractory out-of-hospital cardiac arrest (EROCA). 2020. Available at: <https://clinicaltrials.gov/ct2/show/NCT03065647>. Accessed January 21, 2020.
  64. Hyperinvasive approach in cardiac arrest. 2020. Available at: <https://clinicaltrials.gov/ct2/show/NCT01511666>.
  65. A comparative study between a pre-hospital and an in-hospital circulatory support strategy (ECMO) in refractory cardiac arrest (APACAR2). 2020. Available at: <https://clinicaltrials.gov/ct2/show/NCT02527031>.
  66. Rajan S, Folke F, Hansen SM, Hansen CM, Kragholm K, Gerds TA, Lippert FK, Karlsson L, Moller S, Kober L, et al. Incidence and survival outcome according to heart rhythm during resuscitation attempt in out-of-hospital cardiac arrest patients with presumed cardiac etiology. *Resuscitation*. 2017;114:157–163.
  67. Debaty G, Labarere J, Frascione RJ, Wayne MA, Swor RA, Mahoney BD, Domeier RM, Olinger ML, O'Neil BJ, Yannopoulos D, et al. Long-term prognostic value of gasping during out-of-hospital cardiac arrest. *J Am Coll Cardiol*. 2017;70:1467–1476.
  68. Barbaro RP, Odetola FO, Kidwell KM, Paden ML, Bartlett RH, Davis MM, Annich GM. Association of hospital-level volume of extracorporeal membrane oxygenation cases and mortality. Analysis of the extracorporeal life support organization registry. *Am J Respir Crit Care Med*. 2015;191:894–901.

# **SUPPLEMENTAL MATERIAL**

**Table S1. Summary of E-CPR cost effectiveness studies.**

	Location	Pt No.	Cost per ECPR patient (USD) to hospital discharge or death	Cost per survivor*	Cost per QALY	ICER per QALY
Bharmal et al <sup>56</sup>	North America	32	\$125,683 (IQR \$49,751-\$206,341)	NA	\$56,156	NA
Dennis et al <sup>54</sup>	Australia	62	\$52,615 (SD \$53, 016)	\$88,167	NA	\$17,648
Kawashima et al <sup>55</sup>	Japan	120	\$39,633 VT/VF, \$35,609 ASYS/PEA	\$213,656 PEA/ASYS, \$101,669 VT/VF	\$11,081 VT/VF, \$29,447 ASYS/PEA	\$16,246

\* Cost per survivor calculated as total hospital cost of all patients divided by number of survivors